

B–N Coupling

Platinum-Templated Coupling of B=N Units: Synthesis of BNB Analogues of 1,3-Dienes and a Butatriene

Carina Brunecker, Merle Arrowsmith, Felipe Fantuzzi, and Holger Braunschweig*

Abstract: The 1:2 reaction of $[\mu\text{-(dmpm)Pt(nbe)}]_2$ (dmpm = bis(dimethylphosphino)methane, nbe = norbornene) with $\text{Cl}_2\text{BNR}(\text{SiMe}_3)$ ($\text{R} = \text{tBu}$, SiMe_3) yields unsymmetrical (*N*-aminoboryl)aminoboryl Pt^{II} complexes by B–N coupling via ClSiMe_3 elimination. A subsequent intramolecular ClSiMe_3 elimination from the *tBu*-derivative leads to cyclization of the BNB unit, forming a unique 1,3,2,4-diazadiboretidin-2-yl ligand. In contrast, the analogous reaction with $\text{Br}_2\text{BN}(\text{SiMe}_3)_2$ leads, via a twofold BrSiMe_3 elimination, to a Pt^{II} A-frame complex bridged by a linear BNB isostere of butatriene. Structural and computational data confirm π electron delocalization over the entire BNB unit.

The replacement of C=C double bonds in organic molecules by isosteric covalent B=N units is not only interesting from a fundamental point of view, but also opens up the exploration of a vast hybrid organic–inorganic chemical space. While the typical B=N double bond (1.39 Å)^[1] is only marginally longer than a C=C double bond (1.34 Å, Figure 1), the intrinsic strong polarization of B–N bonds imparts very different electronic properties and stability to the resulting molecules and materials, which can be exploited for new applications in materials science, catalysis, and medicinal chemistry.

Since the landmark synthesis of borazine by Stock and Pohland in 1926 (Figure 1, I),^[2] new synthetic methodologies have enabled access to an ever-increasing variety of B=N/C=C-isosteric compounds and materials, including boron nitride^[3] and borocarbonitride ($\text{B}_x\text{C}_y\text{N}_z$) nanomaterials,^[4] hybrid organic–inorganic BN-doped conjugated polymers,^[5] (poly)aromatic compounds,^[6] and aromatic small molecules.^[7] However, well-defined acyclic conjugated BN chains, such as

How to cite: *Angew. Chem. Int. Ed.* **2021**, *60*, 16864–16868
International Edition: doi.org/10.1002/anie.202106161
German Edition: doi.org/10.1002/ange.202106161





Figure 1. Conjugated organic systems and their all-BN isosteres.

poly(iminoboranes) (III) or BN-based cumulenes (IV), remain difficult to access. The intuitive synthetic routes to III via the polymerization of iminoborane ($\text{RB}=\text{NR}'$) precursors^[8] or the dehydrocoupling of amine borane ($\text{H}_2\text{RB}\cdot\text{NH}_2\text{R}'$) precursors^[9] are in practice marred by the formation of cyclic oligomers such as I and II. The most efficient access to higher oligo(iminoboranes) is by B–N coupling of chloroborane and silylamine precursors via ClSiMe_3 elimination.^[10] The group of Helten has used this methodology to synthesize the first well-defined oligo(iminoboranes) (V) by polycondensation of 1,3-bis(trimethylsilyl)-1,3,2-diazaborolidine precursors with dichloro-(organo)boranes (Scheme 1a).^[11] Our group has also reported the coupling of two $\text{Cl}_2\text{BN}(\text{SiMe}_3)_2$ molecules at $[(\text{C}_5\text{H}_5)_2\text{Ru}(\text{CO})_2]\text{Na}$ with elimination of NaCl and ClSiMe_3 , yielding the (*N*-aminoboryl)aminoboryl complex VI (Scheme 1b).^[12]

We have recently reported the synthesis of the boranedial A-frame complexes **2-X^Y** from the twofold oxidative addition of dihaloborane precursors (X_2BY , $\text{X} = \text{Cl}, \text{Br}, \text{I}$; $\text{Y} = \text{X}$, alkyl, aryl, amino) to the bis(dimethylphosphino)methane (dmpm)-bridged Pt^0 complex **1** (Scheme 2a).^[13] Inspired also by the metal-templated coupling of two BN units at ruthenium in complex VI (Scheme 1b),^[12] we now report the use of the $\text{Pt}_2(\text{dmpm})_2$ scaffold as a template for the coupling of B=N units derived from the coupling of dihalo(silylamino)boranes




Scheme 1. Examples of syntheses of oligo(iminoboranes) by B–N coupling via ClSiMe_3 elimination.

[*] C. Brunecker, Dr. M. Arrowsmith, Dr. F. Fantuzzi, Prof. Dr. H. Braunschweig
Institute for Inorganic Chemistry
Julius-Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg (Germany)
and
Institute for Sustainable Chemistry & Catalysis with Boron
Julius-Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg (Germany)
E-mail: h.braunschweig@uni-wuerzburg.de

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202106161>.

© 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.




Scheme 2. Synthesis of boranediyl-bridged diplatinum A-frame complexes.

($X_2BNR(SiMe_3)$, $X = Cl, Br, I$; $R = tBu, SiMe_3$) by elimination of $XSiMe_3$, ultimately leading to the isolation of the first BNBN-cumulene, isosteric with butatriene.

Whereas the reaction of complex **1** with Cl_2BNMe_2 yields the aminoboranediyl-bridged A-frame complex **2-Cl^{NMe2}** (Scheme 2a), the reactions of **1** with $Cl_2BNR(SiMe_3)$ ($R = tBu, SiMe_3$) always proceeded in a 1:2 ratio. The resulting products **3^{tBu}** and **3^{SiMe3}**, which precipitated as pale yellow solids, both display two broad ^{11}B NMR resonances, at 53 (fwhm ≈ 1280 Hz, PtB) and 32 ppm (fwhm ≈ 880 Hz, N_2BCl) for **3^{tBu}**, and 57 (fwhm ≈ 1990 Hz, PtB) and 33 ppm (fwhm ≈ 750 Hz, N_2BCl) for **3^{SiMe3}** (Scheme 3a). Complexes **3^R** are reminiscent of complex **VI** (Scheme 1b), which shows similar ^{11}B NMR resonances at 60.3 and 35.0 ppm.^[12] The $^{31}P\{^1H\}$ spectra of **3^R** show two multiplets with higher-order satellites in a 1:1 ratio, at -14.3 ($^1J_{P-Pt} = 3195$ Hz, P_2PtCl) and -29.9 ppm ($^1J_{P-Pt} = 2733$ Hz, P_2PtB) for **3^{tBu}**, and -14.3

($^1J_{P-Pt} = 3150$ Hz, P_2PtCl) and -29.6 ppm ($^1J_{P-Pt} = 2708$ Hz, P_2PtB) for **3^{SiMe3}**. X-ray crystallographic analyses of single crystals of **3^{tBu}** confirmed the coupling of the two BN units at one platinum center (Figure 2). Due to systematic rotational disorder of the terminal $B(Cl)NtBu(SiMe_3)$ moiety, structural parameters cannot be fully discussed. The Pt–Pt distance of 2.7067(6) Å, however, is clearly indicative of Pt–Pt bonding. The Pt2–B1 bond length of 2.039(6) Å is within the typical range for square planar platinum amino(chloro)boryl complexes (2.00–2.85 Å), while the B1–N1 bond of 1.421(8) Å is slightly longer than in these complexes (ca. 1.39 Å)^[14] due to the additional π electron delocalization over the entire BNBN unit in **3^{tBu}**.

Complex **3^{SiMe3}** could not be fully characterized as it decomposed rapidly in solution into $ClSiMe_3$ and a number of $dmpm$ -containing platinum complexes, the known complex $[\mu-(dmpm)PtCl]_2$ (**5-Cl**: $\delta(^{31}P) = -19.3$ ppm, $^1J_{P-Pt} = 2650$ Hz)^[13a] being the major decomposition product (Scheme 3b, see Figure S18 in the SI). The fate of the remaining $[BNSiMe_3]_2$ fragment could not be determined as the ^{11}B NMR spectrum of the final product mixture was silent, and a colorless by-product, insoluble in all common organic solvents, was formed.^[15] In contrast, **3^{tBu}** was stable in solution at room temperature but selectively converted to **4^{tBu}** at 80 °C by intramolecular cyclization of the BNBN moiety under



Scheme 3. Reactions of complex **1** with $Cl_2BNR(SiMe_3)$ ($R = tBu, SiMe_3$). Isolated yields in parentheses.





Figure 2. Crystallographically derived molecular structures of (from left to right) **3^{tBu}** (least disordered one of the two molecules of **3^{tBu}** in the asymmetric unit), **4^{tBu}**, and **6**.^[26] Thermal ellipsoids at 50% probability. Thermal ellipsoids of ligand periphery and hydrogen atoms omitted for clarity. Only the major part of the disorders in **3^{tBu}** (terminal $B(Cl)NtBu(SiMe_3)$ moiety) and **4^{tBu}** (entire $(BNtBu)_2Cl$ moiety and one $dmpm$ ligand) is shown. Due to the restraints applied to these disorders during refinement, the structural parameters of **3^{tBu}** and **4^{tBu}** may not be fully discussed. Selected bond lengths (Å) and angles ($^\circ$) for **3^{tBu}**: Cl1–Pt1 2.4939(13), Pt1–Pt2 2.7067(6), Pt–P 2.2446(14)–2.2651(14), Pt2–B1 2.039(6), B1–N1 1.421(8), Cl1–Pt1–Pt2 172.32(3), P1–Pt2–B1 174.14(16), $\Sigma(\angle B1)$ 360.0(4), torsion angles P1–Pt1–Pt2–P2 $-47.8(4)$, P3–Pt1–Pt2–P4 $-54.32(5)$; for **4^{tBu}**: Cl1–Pt1 2.535(3), Pt1–Pt2 2.7214(7); for **6**: Pt1...Pt2 3.2397(3), Pt1–B1 2.028(6), Pt2–B1 2.021(6), Pt1–Br1 2.6098(6), Pt2–Br2 2.6363(6), Pt–P 2.2679(15)–2.2913(14), B1–N1 1.396(7), N1–B2 1.237(8), B2–N2 1.388(8), Pt1–B1–Pt2 106.3(3), B1–N1–B2 173.8(6), N1–B2–N2 171.3(7), torsion angles P1–Pt1–Pt2–P2 $-12.29(5)$, P3–Pt1–Pt2–P4 $-23.83(5)$.


ClSiMe₃ elimination (Scheme 3c). This reaction is analogous to the cyclization of RCIB–N(*t*Bu)–B(Cl)–N*t*Bu(SiMe₃) (R = NMe₂, NEt₂, Et, *i*Bu) to 1,3,2,4-diazadiboretidines by ClSiMe₃ elimination, reported by Paetzold in 1988.^[16] The ¹¹B NMR spectrum of **4**^{*t*Bu} is nearly identical to that of **3**^{*t*Bu}, displaying two broad resonances at 54 (fwhm ≈ 1480 Hz, PtB) and 32 ppm (fwhm ≈ 470 Hz, N₂BCl). The conversion of **3**^{*t*Bu} to **4**^{*t*Bu} is evidenced more clearly by changes in the ³¹P{¹H} spectrum, which shows two new 1:1 multiplets with higher-order satellites, both shifted ca. 2 ppm downfield from **3**^{*t*Bu}, at –12.8 (¹J_{P–Pt} = 3198 Hz, P₂PtCl) and –27.6 ppm (¹J_{P–Pt} = 2632 Hz, P₂PtB), the ¹J_{P–Pt} coupling constant of the latter being ca. 100 Hz smaller than in **3**^{*t*Bu}. Crystallization attempts of **4**^{*t*Bu} always yielded pseudo-merohedrally twinned crystals (see solid-state structure in Figure 2), in which the BNBn heterocycle presents a twofold disorder by rotation of about the Pt₂–B1 bond, thus precluding any discussion of bond lengths and angles in this unit. Despite the well-established chemistry of 1,3,2,4-diazadiboretidines as η⁴-ligands for transition metals,^[17] **3**^{*t*Bu} represents a hitherto unknown binding mode of this type of ligand as an anionic η¹-ligand via coordination at boron. In solution at room temperature, compound **4**^{*t*Bu} decomposed very slowly but selectively over a period of several weeks to complex **5-Cl** and an unidentified intractable colorless solid, by formal loss of “[BN(*t*Bu)]₂” (Scheme 3d).^[15]

To our surprise the reaction of **1** with Br₂BN(SiMe₃)₂ resulted instead in the formation of the A-frame complex **6**, isolated as a yellow solid in 46% yield (Scheme 4).^[18] The ¹¹B NMR spectrum of **6** displays two broad resonances at ca. 57 (fwhm ≈ 1510 Hz) and 26 ppm (fwhm ≈ 690 Hz), the former being attributed to the platinum-bound boron nucleus by analogy with the ¹¹B NMR shift of the related dimethylaminoborane-bridged A-frame complex **2-Br**^{NMe₂} (δ(¹¹B) = 52 ppm),^[13] the latter to the dicoordinate NBN boron nucleus. The ³¹P{¹H} NMR spectrum showed a singlet at –7.1 ppm, close to that of **2-Br**^{NMe₂} (δ(³¹P) = –5.6 ppm), with a higher-order satellite splitting pattern typical for A-frame complexes (¹J_{P–Pt} = 3568 Hz, ³J_{P–Pt} = 272 Hz, ¹J_{Pt–Pt} = 1826 Hz). ¹¹B and ³¹P{¹H} NMR-spectroscopic monitoring of the reaction showed no sign of formation of the bromide analogue of **3**^{SiMe₃}.

We propose that the formation of complexes **3**^R and **6** proceeds via a same intermediate η¹-(silylamino)haloboryl complex **Int-X**^R formed by the oxidative addition of X₂BNR(SiMe₃) to **1** (Scheme 5).^[19] This step can be followed either by B–N coupling with a second equivalent X₂BNR(SiMe₃) via



Scheme 4. Reaction of complex **1** with BBr₂N(SiMe₃)₂. Isolated yield in parentheses.



Scheme 5. Proposed mechanism of formation of **3**^R and **6** via the common intermediate **Int-X**^R.

XSiMe₃ elimination (reaction rate constant *k*_a) to form an η¹-(*N*-aminoboryl)aminoboryl complex analogous to **3**^R, or by the oxidative addition of the second B–X bond of the silylamino(halo)boryl ligand to platinum to form the (silylamino)boranediyl A-frame complex **2-X**^{NR(SiMe₃)} (reaction rate constant *k*_b). For R = SiMe₃, the latter then undergoes twofold XSiMe₃ elimination with a second equivalent of X₂BN(SiMe₃)₂ to form complex **6**. The selectivity of the reaction is therefore determined by the relative values of the reaction rate constants *k*_a and *k*_b; for X = Cl the rate of B–N coupling outperforms that of oxidative addition of B–Cl to Pt, leading to the exclusive formation of **3**^R, the opposite being the case for X = Br, leading to the exclusive formation of **6**.

The solid-state structure of **6** (Figure 2) confirmed the formation of the near-linear BNBn unit bridging the two platinum centers (B1–N1–B2 173.8(6), N1–B2–N2 171.3(7)°). While the Pt–B bond lengths of 2.028(6) and 2.021(6) Å are similar to those in complex **2-Br**^{NMe₂} (2.028(10), 2.042(9) Å), the A-frame structure itself is more strongly distorted from the ideal A-frame than in **2-Br**^{NMe₂}, as evident in the much shorter Pt...Pt distance (**6** 3.2397(3); **2-Br**^{NMe₂} 3.3003(4) Å) and larger P1/3–Pt1–Pt2–P2/4 torsion angles (**6** –12.29(5), –23.83(5)°; **2-Br**^{NMe₂} 4.96(7), 15.62(8)°).^[13] Furthermore, the B1–N1 and B2–N2 bond lengths of 1.396(7) and 1.388(8) Å are within the range of partial double bonds, whereas the central N1–B2 bond is significantly shorter (1.237(8) Å), corresponding to a partial triple bond.^[1] While the linear BNBn motif can be viewed formally as a 1-boryl-2-(amino)iminoborane, the delocalization of the π electron density apparent in the B–N bond lengths makes it structurally more akin to an all-BN isostere of a butatriene. Unlike butatriene, however, which is fully planar, the B1 and N2 planes form an angle of ca. 24°, which could result from the steric repulsion between the SiMe₃ groups and the dmpm ligands.

The electronic structure of **6** was further investigated using DFT and intrinsic bond orbital (IBO)^[20] calculations. The BNBn motif in the optimized structure of **6**, obtained at the M06^[21]-D3^[22]/cc-pVDZ^[23], aug-cc-pVDZ-PP[Pt]^[24] level of theory, shows a larger deviation from linearity (B1–N1–B2 161.3°, N1–B2–N2 176.2°) than that of the solid-state structure. Similar results were obtained with other density functionals (see details in the SI). In order to investigate the origin of this deviation, we performed computations on four truncated model systems, in which the PMe₂ and SiMe₃ groups were successively replaced with PH₂ and SiH₃ or H, respectively (see Figure S19 in the SI). In all of these cases, the BNBn

moiety was found to be linear (B1-N1-B2 and N1-B2-N2 178.8–180.0°). The distortion from linearity therefore seems to arise from the steric repulsion between the PMe_2 and SiMe_3 substituents, although the additional influence of crystal packing forces in the solid-state structure cannot be discounted. Furthermore, the calculated Mayer bond orders (MBOs)^[25] of the BNB motif in **4** (B1–N1: 1.38, N1–B2: 2.11, B2–N2: 1.32) are very similar to those obtained for the parent H_2BNBNH_2 system (B1–N1: 1.51, N1–B2: 2.13, B2–N2: 1.43), these values suggesting strong cumulenic character in both cases. Indeed, inspection of the IBOs of **6** (Figure 3 a) reveals that IBO-1 and IBO-3, which are orthogonal to the (Pt1-B1-Pt2) plane, are partially delocalized to the neighboring B2 and B1 atoms, evidencing deviation from the 1-boryl-2-(amino)iminoborane picture. This view is also supported by inspection of the canonical Kohn–Sham molecular orbitals (MOs) of **6** and H_2BNBNH_2 (Figure 3 b and S20 in the SI), where π electron delocalization over the entire BNB unit is observed. The description of **6** as a BNB analogue of butatriene is, therefore, fully supported by quantum chemical investigations.




Figure 3. a) Selected IBOs of **6**. b) The fully π -delocalized MOs of **6** (left, HOMO–30) and H_2BNBNH_2 (right, HOMO–3), highlighting the cumulenic character of their BNB motifs.

To conclude, we have shown that the $[\mu\text{-(dmpm)Pt}]_2$ framework acts as an effective template for the coupling of B=N units obtained by the intermolecular B–N coupling of dihalo(silylamino)boranes via halosilane elimination. For $\text{Cl}_2\text{BNR}(\text{SiMe}_3)$ precursors BN chain growth occurs at a side-on Pt^{I} complex, whereas for $\text{Br}_2\text{BN}(\text{SiMe}_3)_2$ an A-frame Pt^{II} complex bridged by a linear BNB unit is formed. Structural and computational analyses confirm a cumulenic motif isosteric with butatriene.

Acknowledgements

We thank the Deutsche Forschungsgemeinschaft for financial support of this project. F.F. thanks the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and the Alexander von Humboldt (AvH) Foundation for a CAPES-Humboldt postdoctoral fellowship. Open access funding enabled and organized by Projekt DEAL.

Conflict of Interest

The authors declare no conflict of interest.

Keywords: 1,3,2,4-diazadiboretidin-2-yl ligand · A-frame complex · B–N coupling · butatriene analogue · isosterism

- [1] S. Berski, Z. Latajkaa, A. J. Gordon, *New J. Chem.* **2011**, 35, 89–96.
- [2] A. Stock, E. Pohland, *Ber. Dtsch. Chem. Ges.* **1926**, 59, 2210–2223.
- [3] Recent reviews and book chapters: a) J. Wang, L. Zhang, L. Wang, W. Lei, Z.-S. Wu, *Energy Environ. Mater.* **2021**, <https://doi.org/10.1002/eem2.12159>; b) J. Yin, J. Li, Y. Hang, J. Yu, G. Tai, X. Li, Z. Zhang, W. Guo, *Small* **2016**, 12, 2942–2968; c) G. R. Bhimanapati, N. R. Glavin, J. A. Robinson, *2D Boron Nitride: Synthesis and Applications in 2D Materials* (Eds. F. Iacopi, J. J. Boeckl, C. Jagadish), Elsevier Science & Technology, Amsterdam, **2016**, pp. 101–148.
- [4] C. N. R. Rao, K. Pramoda, *Bull. Chem. Soc. Jpn.* **2019**, 92, 441–468.
- [5] Recent examples: a) S. Pang, Z. Wang, X. Yuan, L. Pan, W. Deng, H. Tang, H. Wu, S. Chen, C. Duan, F. Huang, Y. Cao, *Angew. Chem. Int. Ed.* **2021**, 60, 8813–8817; *Angew. Chem.* **2021**, 133, 8895–8899; b) H. Oubaha, N. Demitri, J. Rault-Berthelot, P. Dubois, O. Coulembier, D. Bonifazi, *J. Org. Chem.* **2019**, 84, 9101–9116; c) B. Thiedemann, P. J. Gliese, J. Hoffmann, P. G. Lawrence, F. D. Sonnichsen, A. Staubitz, *Chem. Commun.* **2017**, 53, 7258–7261; Most recent review: d) H. Helten, *Chem. Eur. J.* **2016**, 22, 12972–12982.
- [6] J. Huang, Y. Li, *Front. Chem.* **2018**, 6, 341; X.-Y. Wang, F.-D. Zhuang, R.-B. Wang, X.-C. Wang, X.-Y. Cao, J.-Y. Wang, J. Pei, *J. Am. Chem. Soc.* **2014**, 136, 3764–3767.
- [7] Recent reviews: a) C. R. McConnell, S.-Y. Liu, *Chem. Soc. Rev.* **2019**, 48, 3436–3453; b) Z. X. Giustra, S.-Y. Liu, *J. Am. Chem. Soc.* **2018**, 140, 1184–1194; c) G. Bélanger-Chabot, H. Braunschweig, D. K. Roy, *Eur. J. Inorg. Chem.* **2017**, 4353–4368; d) E. R. Abbeya, S.-Y. Liu, *Org. Biomol. Chem.* **2013**, 11, 2060–2069.
- [8] a) J. Kiesgen, J. Munster, P. Paetzold, *Chem. Ber.* **1993**, 126, 1559–1563; b) P. Paetzold, *Adv. Inorg. Chem.* **1987**, 31, 123–170.
- [9] Recent reviews: a) D. Han, F. Anke, M. Trose, T. Beweries, *Coord. Chem. Rev.* **2019**, 380, 260–286; b) A. L. Colebatch, A. S. Weller, *Chem. Eur. J.* **2019**, 25, 1379–1390; c) N. T. Coles, R. L. Webster, *Isr. J. Chem.* **2017**, 57, 1070–1081.
- [10] a) K. Ma, H.-W. Lerner, S. Scholz, J. W. Bats, M. Bolte, M. Wagner, *J. Organomet. Chem.* **2002**, 664, 94–105; b) H. Nöth, N. Storch, *Chem. Ber.* **1974**, 107, 1028–1037; c) H. Nöth, M. J. Sprague, *J. Organomet. Chem.* **1970**, 23, 323–327; d) H. Jenne, K. Niedenzu, *Inorg. Chem.* **1964**, 3, 68–70.
- [11] a) O. Ayhan, N. A. Riensch, C. Glasmacher, H. Helten, *Chem. Eur. J.* **2018**, 24, 5883–5894; b) O. Ayhan, T. Eckert, F. A.

- Plamper, H. Helten, *Angew. Chem. Int. Ed.* **2016**, *55*, 13321–13325; *Angew. Chem.* **2016**, *128*, 13515–13519.
- [12] H. Braunschweig, C. Kollann, K. W. Klinkhammer, *Eur. J. Inorg. Chem.* **1999**, 1523–1529.
- [13] a) C. Brunecker, M. Arrowsmith, J. H. Müssig, J. Böhnke, A. Stoy, M. Heß, A. Hofmann, C. Lenczyk, C. Lichtenberg, J. Ramler, A. Rempel, H. Braunschweig, *Dalton Trans.* **2021**, *50*, 3506–3515; b) C. Brunecker, J. H. Müssig, M. Arrowsmith, F. Fantuzzi, A. Stoy, J. Böhnke, A. Hofmann, R. Bertermann, B. Engels, H. Braunschweig, *Chem. Eur. J.* **2020**, *26*, 8518–8523.
- [14] a) J. P. H. Charmant, C. Fan, N. C. Norman, P. G. Pringle, *Dalton Trans.* **2007**, 114–123; b) D. Curtis, M. J. G. Lesley, N. C. Norman, A. G. Orpen, J. Starbuck, *J. Chem. Soc. Dalton Trans.* **1999**, 1687–1694.
- [15] This intractable by-product is likely to result from the polymerization of “[BNR]₂”.
- [16] K. H. Van Bonn, P. Schreyer, P. Paetzold, R. Boese, *Chem. Ber.* **1988**, *121*, 1045–1057.
- [17] a) P. Paetzold, K. Delpy, R. Boese, *Z. Naturforsch. B* **1988**, *43*, 839–845; b) G. Schmid, D. Kampmann, W. Meyer, R. Boese, P. Paetzold, K. Delpy, *Chem. Ber.* **1985**, *118*, 2418–2428; c) K. Delpy, H.-U. Meier, P. Paetzold, C. von Plotho, *Z. Naturforsch. B* **1984**, *39*, 1696–1701; d) K. Delpy, D. Schmitz, P. Paetzold, *Chem. Ber.* **1983**, *116*, 2994–2999.
- [18] Like all **2-Br^Y** complexes (see Scheme 1 a), complex **6** decomposes slowly in solution to [μ-(dmpm)PtBr]₂ (**5-Br**, see ref. [13]) and intractable polymeric “[BN(SiMe₃)]_n”.
- [19] Since the homocoupling of Cl₂BNR(SiMe₃) by ClSiMe₃ elimination does not proceed at room temperature, the coupling step has to occur after the oxidative addition of X₂BNR(SiMe₃) to **VII**.
- [20] G. Knizia, *J. Chem. Theory Comput.* **2013**, *9*, 4834–4843.
- [21] Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* **2008**, *120*, 215–241.
- [22] S. Grimme, J. Antony, S. Ehrlich, H. Krieg, *J. Chem. Phys.* **2010**, *132*, 154104.
- [23] a) T. H. Dunning, *J. Chem. Phys.* **1989**, *90*, 1007–1023; b) D. E. Woon, T. H. Dunning, *J. Chem. Phys.* **1993**, *98*, 1358–1371; c) A. K. Wilson, D. E. Woon, K. A. Peterson, T. H. Dunning, *J. Chem. Phys.* **1999**, *110*, 7667–7676.
- [24] D. Figgen, K. A. Peterson, M. Dolg, H. Stoll, *J. Chem. Phys.* **2009**, *130*, 164108.
- [25] a) I. Mayer, *Chem. Phys. Lett.* **1983**, *97*, 270–274; b) I. Mayer, *Int. J. Quantum Chem.* **1984**, *26*, 151–154.
- [26] Deposition Numbers 2081529 (**4^{Bu}**), 2081530 (**6**), 2081531 (**3^{Bu}**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service. .

Manuscript received: May 7, 2021

Accepted manuscript online: May 25, 2021

Version of record online: June 24, 2021