



Kent Academic Repository

Syeda, Samar M., Tsegai, Zewdi J., Cazenave, Marine, Dunmore, Christopher J., Skinner, Matthew M. and Kivell, Tracy L. (2022) *Reconstructing hand use in Australopithecus sediba and Homo naledi: mapping variation in cortical thickness across the proximal and intermediate phalanges*. In: *PaleoAnthropology. 12th Annual Meeting of the European Society for the Study of Human Evolution. Abstracts. Tübingen, 22-24 September 2022. Podium Presentation Session 3. 2022 (2)*. p. 588. *Paleoanthropology Society*

Downloaded from

<https://kar.kent.ac.uk/97670/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.48738/2022.iss2.809>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Poster Presentation Number 82, Session 2, Friday 18:15

Reconstructing hand use in *Australopithecus sediba* and *Homo naledi*: mapping variation in cortical thickness across the proximal and intermediate phalanges

Samar M. Syeda¹, Zewdi J. Tsegai¹, Marine Cazenave^{1,2,3}, Christopher J. Dunmore¹, Matthew M. Skinner^{1,4}, Tracy L. Kivell^{1,4}

1 - Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK · 2 - Division of Anthropology, American Museum of Natural History, New York, New York, USA · 3 - Department of Anatomy, Faculty of Health Sciences, University of Pretoria, South Africa · 4 - Centre for the Exploration of the Deep Human Journey, University of the Witwatersrand, Johannesburg, South Africa

The inferred diversity of manual behaviors within the hominin lineage is exemplified by the mosaic morphology of the hands of *Australopithecus sediba* and *Homo naledi* [1-2]. Derived morphological features present in both fossil hominins point to human-like manipulation abilities, while curved fingers with well-developed flexor sheath ridges (FSRs) suggest the continued use of their hands for climbing and grasping [1-2]. The internal bone structure may reflect habitual locomotor and/or manipulative loading of the fingers due to bone's ability to (re)model throughout life in response to mechanical load [3], and thus variation in phalangeal cortical bone distribution holds the potential for reconstructing hand use among fossil hominins.

We investigated the distribution of phalangeal cortical thickness in extant great apes, *A. sediba* and *H. naledi* to inform our reconstruction of fossil manual behaviours. The sample included microCT scans of the proximal and intermediate phalanges of rays II-V of *Gorilla gorilla* (N=21), *Homo sapiens* (N=37), *Pongo pygmaeus* (N=9), *Pan* sp. (N=24), *A. sediba*, and *H. naledi* (Hand 1). We used the R package morphomap [4] to map diaphyseal cortical thickness and conducted a principal component analysis (PCA) on the cortical thickness values to explore patterns across species.

Results show clear separation among extant taxa across all phalanges. The extant taxa are separated along PC1, with some overlap between the African apes reflecting a pattern of thickness localised to the FSRs, with a different pattern of overall thickness of the shaft in *Pongo* and in humans. PC2 reflects differences in the location of greatest thickness; in African apes this occurs in the FSRs throughout the shaft, whereas in *Pongo* this is distally located with some dorsal thickening, and in humans it is primarily in the dorsodistal region of the shaft. The proximal phalanges of *A. sediba* consistently fall close to the African apes, reflecting maximum thickness localised to the FSRs, while *H. naledi* consistently falls close to humans, reflecting thickening of the dorsal region of the shaft and the FSRs distally. The intermediate phalanges of both fossil species fall between the African ape and human distribution, reflecting a thickening of the dorsal region of the bone and the FSRs, which are located in the proximal-to-midshaft region. The distribution of the *A. sediba* intermediate phalanges differs from those of *H. naledi* in that the thickening on the palmar surface extends further distally. This is consistent with external morphology of *A. sediba* intermediate phalanges that possess unusually long and prominent FSRs.

Overall, we show that variation in phalangeal cortical bone distribution clearly separates extant hominoid taxa that differ in their locomotor and manipulative behaviours. Within this comparative context, *A. sediba* proximal phalanges are most similar to the African ape cortical signal and intermediate phalanges are intermediate between all extant taxa. Along with a high degree of phalangeal curvature and prominent FSRs, this morphology better distributes stress and reduces strain on the bone during flexed finger loading (e.g., during climbing or suspension) [5]. In contrast, while *H. naledi* phalanges are most similar to humans in their cortical distribution, this appears to contradict their high degree of curvature, which may be a functional indication of arboreal locomotion [5]. The *H. naledi* phalanges indicate a hand uniquely adapted for arboreal behaviours in the degree of curvature but not cortical distribution, differing to that of extant great apes and *A. sediba*.

For access to specimens we thank the following individuals/institutions: Max Planck Institute for Evolutionary Anthropology (C. Boesch), J.-J. Hublin (College de France); Museum für Naturkunde - Leibniz Institute for Evolution and Biodiversity Science (F. Mayer, C. Funk); Powell-Cotton Museum (I. Litvne); Royal Museum for Central Africa (E. Gilissen); University of Florence (J. Moggi-Cecchi, S. Bortoluzzi); Johann-Friedrich-Blumenbach-Institute for Zoology and Anthropology, Georg-August University, Göttingen (B. Grosskopf); Frankfurt Senckenberg Museum (V. Volpato); University of the Witwatersrand (L. Berger, B. Zipfel); Science Academy of the Czech Republic (J. Svoboda). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 819960 and No. 101025719).

References: [1] Kivell, T.L., Kibii, J.M., Churchill, S.E., Schmid, P., Berger, L.R., 2011. *Australopithecus sediba* hand demonstrates mosaic evolution of locomotor and manipulative abilities. *Science* 333(6048), 1411-1417. [2] Kivell, T.L., Deane, A.S., Tocheri, M.W., Orr, C.M., Schmid, P., Hawks, J., Berger, L.R., Churchill, S.E., 2015. The hand of *Homo naledi*. *Nature Communications*, 6(1), pp.1-9. [3] Ruff, C., Holt, B., Trinkaus, E., 2006. "Who's afraid of the big bad Wolff? "Wolff's law" and bone functional adaptation. *American Journal of Physical Anthropology* 129(4), 484-498. [4] Profico, A., Bondioli, L., Raia, P., O'Higgins, P., Marchi, D., 2021. morphomap: An R package for long bone landmarking, cortical thickness, and cross-sectional geometry mapping. *American journal of Physical Anthropology* 174(1), 129-139. [5] Nguyen, N.H., Pahr, D.H., Gross, T., Skinner, M.M., Kivell, T.L., 2014. Micro-finite element (μFE) modeling of the siamang (*Symphalangus syndactylus*) third proximal phalanx: the functional role of curvature and the flexor sheath ridge. *Journal of Human Evolution* 67, 60-75.