



Diversity, fragmentation, and connectivity across the UK amphibian and reptile data management landscape

Rebecca K. Turner^{1,2} · Richard A. Griffiths² · John W. Wilkinson³ · Angela M. Julian⁴ · Mike P. Toms⁵ · Nick J. B. Isaac¹

Received: 9 April 2022 / Revised: 20 October 2022 / Accepted: 30 October 2022
© The Author(s) 2022

Abstract

Large-scale biodiversity monitoring remains a challenge in science and policy. ‘Biodiversity Observation Networks’ provide an integrated infrastructure for monitoring biodiversity through timely discovery, access, and re-use of data, but their establishment relies on an in-depth understanding of existing monitoring effort. We performed a scoping review and network analysis to assess the scope of available data on amphibians and reptiles in the UK and catalogue the mobilisation of information across the data landscape, thereby highlighting existing gaps. The monitoring portfolio has grown rapidly in recent decades, with over three times as many data sources than there are amphibian and reptile species in the UK now available. We identified 45 active sources of ‘FAIR’ (‘Findable’, ‘Accessible’, ‘Interoperable’ and ‘Reusable’) data. The taxonomic, geographic and temporal coverage of datasets appears largely uneven and no single source is currently suitable for producing robust multispecies assessments on large scales. A dynamic and patchy exchange of data occurs between different recording projects, recording communities and digital data platforms. The National Biodiversity Network Atlas is a highly connected source but the scope of its data (re-)use is potentially limited by insufficient accompanying metadata. The emerging complexity and fragmented nature of this dynamic data landscape is likely to grow without a concerted effort to integrate existing activities. The factors driving this complexity extend beyond the UK and to other facets of biodiversity. We recommend integration and greater stakeholder collaboration behind a coordinated infrastructure for data collection, storage and analysis, capable of delivering comprehensive assessments for large-scale biodiversity monitoring.

Keywords Biodiversity data · Biodiversity observation network (BON) · Biodiversity monitoring · FAIR data · Citizen science · Network analysis

Communicated by Pedro Eisenlohr.

✉ Rebecca K. Turner
rkt4@kent.ac.uk

Extended author information available on the last page of the article

Introduction

Human activity is influencing biodiversity turnover across the globe (Dornelas et al. 2014; Keil et al. 2015; Kaarlejärvi et al. 2021). Monitoring biodiversity at large spatial and temporal scales is central to understanding the magnitude of change, and is important in conservation planning and resource allocation by decision-makers (Parr et al. 2002; Petersen et al. 2021; Thornhill et al. 2021). To understand species status and monitor change requires high-quality data. Data must provide sufficient taxonomic, temporal and geographic coverage to reliably inform evidence-based conservation (Wetzel et al. 2018). Given the economical and logistical constraints involved with monitoring, our understanding of biodiversity turnover would likely benefit from combining data originating from different sources. Modern statistical advancements can facilitate the integration of data to produce accurate assessments on the state of biodiversity (Isaac et al. 2020). However, effective monitoring for the long-term requires streamlining disparate efforts in the collection, storage and analysis of biodiversity data.

Historically, biological recording has been coordinated by institutions and carried out by volunteer recorders (Roy et al. 2012; Pocock et al. 2015). Nowadays, the popular practice of engaging volunteers in a scientific project is commonly regarded as ‘citizen science’ (Cohn 2008) which can generate ‘community contributed data’. Citizen (or ‘community’) scientists can assist with the collection of biodiversity data on large spatial and temporal scales that would otherwise not be feasible using small-scale studies (Cohn 2008; Pocock et al. 2017; Dobson et al. 2020; Thornhill et al. 2021). Citizen science projects can vary in their objectives and methodological approaches, and, as with any dataset, can be subject to errors and biases imposed by observational processes (Oliveira et al. 2016; Dobson et al. 2020). Once the characteristics of datasets are understood, constituent data can be handled for bias mitigation (Roy et al. 2012; Isaac et al. 2014; Dobson et al. 2020). For instance, sophisticated analytical tools have emerged in recent years to assist data users to assess (e.g., Boyd et al. 2021) and address (e.g., Bird et al. 2014; Geldmann et al. 2016) sampling errors and biases within observation datasets. Datasets derived from citizen science projects can therefore complement small-scale systematic monitoring and are important sources of observation data, usable by scientists and resource-managers for monitoring biodiversity change (Roy et al. 2012; Bonney et al. 2014; Burgess et al. 2017; McKinley et al. 2017; Tredick et al. 2017; Thornhill et al. 2021).

The value of citizen science, combined with a surge in technological innovation, has increased the diversity of projects that generate data in recent years (Kosmala et al. 2016; McKinley et al. 2017; Pocock et al. 2017; Thornhill et al. 2021). Data management has also evolved with the increased wealth of biodiversity data. Data flows from individuals (e.g. recorders, project organisers, stakeholder groups, consultants) to digital data platforms (e.g. web-based apps, e-infrastructure data portals, multi-dataset repositories) via an array of quality assurance techniques (e.g. automated data verification tools, validated datasets) applied at various levels before reaching data users (James 2011; Roy et al. 2012; Pocock et al. 2015). Collectively, such diversification has resulted in a rich data-gathering landscape. Despite these advances, significant challenges remain for collating and analysing data from different sources. These include issues associated with data confidentiality and fears that open data could endanger sensitive species and their habitats (e.g. persecution, or accidental damage to sites by naturalists wanting to see species); as well as a reluctance to share data that could be used for commercial purposes (Griffiths et al. 2015; Fox et al. 2019). Whilst a rich array of biodiversity data now exists, there remain significant barriers

to ensuring that it informs decision-making efficiently. Nonetheless, at a time where there has never been a greater wealth of data available, recent advances in computing mean that there is now also an accompanying suite of statistical tools available to researchers to maximise the re-use of multiple datasets. Hence, the potential to widen the reach of existing biodiversity monitoring efforts and increase the re-use of data through integration is rapidly building momentum.

Effective integration requires a framework that unifies disparate monitoring efforts (König et al. 2019). By its very nature, data integration is dependent upon data sharing. The sheer magnitude of all biodiversity on Earth means that no single institution can know more than a tiny proportion of it at any given time (Walters and Scholes 2017). Stakeholder collaboration in the collection, sharing and analysis of biodiversity data therefore has clear benefits for biodiversity monitoring. Whilst innovative technology can enhance data collection, database design, and data sharing (Roy et al. 2012), large-scale monitoring is difficult without a coordinated infrastructure (Walters and Scholes 2017). ‘Biodiversity Observation Networks’ (e.g., see Wetzel et al. 2018) are collaborative organisational structures for monitoring biodiversity through sharing data, resource and expertise by stakeholders (Walters and Scholes 2017). A network can thus increase the mobilisation of data for research and resource-management. Accordingly, more precise tracking of biodiversity across space and time is possible, thereby enriching the contribution of disparate monitoring efforts (Constable et al. 2010; Jetz et al. 2019). However, the success of a network relies on uptake of common practices across the sector (König et al. 2019). Examples of relevant practices include data standards such as Darwin Core (Wieczorek et al. 2012) for data storage and guiding principles such as the FAIR data principles (Wilkinson et al. 2016) to ensure that data may be ‘Findable’, ‘Accessible’, ‘Interoperable’ and ‘Reusable’, thus enabling the mobilisation of usable biodiversity data.

While advances in technology will facilitate sharing of ‘best practice’ for handling and mobilising biodiversity data, there will still be challenges associated with maximising the quality of data and overcoming biases. Understanding the differences amongst data sources; their taxonomic, temporal and geographic scope; and the flow of data between sources can identify gaps in existing monitoring portfolios. In turn, this can illuminate ways in which monitoring and data may be integrated (Petersen et al. 2021). For instance, the monitoring of amphibians and reptiles in the United Kingdom (UK) is carried out in a number of ways by a diverse recording community. There are thirteen species of amphibians and reptiles native to the UK, many of which have experienced recent declines (Humphreys et al. 2011; Wilkinson and Arnell 2013; Beebee and Ratcliffe 2018; Gardner et al. 2019). Whilst climate change and habitat degradation threaten the UK populations (Dunford and Berry 2012; Turner and Maclean 2022) and may be implicated in declines, formal assessments of status and national trends have largely relied on anecdotal evidence (Hayhow et al. 2019). The lack of empirical evidence is surprising given that such a limited range of species should be relatively simple to identify, and that several are also subject to legal protection with mandatory reporting requirements. Adopting an integrated, network approach is therefore likely to enhance the monitoring and conservation effort for these species.

As an essential first step towards understanding the value of existing data and opportunities for integration, we surveyed the UK amphibian and reptile data management landscape. We used a scoping review and network analysis to characterise and track the mobilisation of information across the data landscape. To our knowledge, this study is the first to use this approach in the context of biodiversity data exploration. We identified an array of existing sources of amphibian and reptile data, characterised the scope of data sources

using the available meta-data associated with each source, and highlighted limitations in the existing monitoring portfolio for tracking species national status and population trends. The network analysis illuminated the dynamics at play within an unrealised Biodiversity Observation Network. To this end, the aims of this review were to:

- (1) Identify existing sources of UK amphibian and reptile observation data.
- (2) Characterise the taxonomic, geographic and temporal scope of UK amphibian and reptile data sources and their corresponding sampling and dataset quality assurance procedures.
- (3) Catalogue the mobilisation of data between sources of UK amphibian and reptile observation data.
- (4) Identify gaps in the existing UK amphibian and reptile data management landscape.
- (5) Provide recommendations for achieving an integrated Biodiversity Observation Network.

Methodology

Search strategy

We used a scoping review framework (see Arksey and O'Malley 2005; Levac et al. 2010) to survey sources of amphibian and reptile observation data. This enabled us to identify sources that were not strictly locked within academic literature, thus reflecting the type of data suitable for integration across a network. We performed searches between 7 November 2020 and 15 January 2021. First, data sources were identified through consultations with three stakeholder organisations: the Amphibian and Reptile Group (ARG UK) network, the Amphibian and Reptile Conservation (ARC) Trust, and the British Trust for Ornithology (BTO). The ARG UK network and the ARC Trust are leading amphibian and reptile conservation charities that support various projects monitoring and conserving native amphibian and reptile populations across the UK. The BTO is mainstream UK conservation charity leading on the conservation and research of birds and other British wildlife. The BTO Garden BirdWatch scheme is the one of the largest biodiversity monitoring citizen science projects in the UK and generates thousands of amphibian and reptile observations annually. Following initial consultations, we performed a series of desk-based searches of electronic databases, internet search engines, registries of biological records data, and the grey literature. We searched Google search engine (www.google.co.uk) using the key terms "UK reptile amphibian data" and "UK biodiversity recording database", respectively. The first 100 results of each search were reviewed as potential sources of reptile and/or amphibian observation data. Next, we searched the National Biodiversity Network (NBN) Atlas (www.nbnatlas.org) using the 'advanced search' function to filter for relevant data partners and datasets using the [Species/Taxon (any)] field and searching the key term "reptile OR amphibian". We then searched the UK Environmental Observation Framework Catalogue (www.ukeof.org.uk) for relevant datasets using the key term "reptile OR amphibian". Relevant platforms were also identified from manual interrogation of those listed on the National Forum for Biological Recording (www.nfbr.org.uk) and the Biological Records Centre (BRC) (www.brc.ac.uk). Lastly, it was important to manually search key organisations' websites to identify data sources that may have been missed in other searches (Arksey and O'Malley 2005). Accordingly, we searched the websites of six biological recording

organisations: the ARC Trust (www.arc-trust.org), ARG UK (www.arguk.org), the British Herpetological Society (BHS) (www.thebhs.org), BTO (www.bto.org), Froglife (www.froglife.org) and Natural England (NE) (www.naturalengland-defra.opendata.arcgis.com). All desk-based searches were repeated between 22 and 24 May 2021 to identify additional sources missed during the initial search. The results of our search strategy are provided in full in the electronic supplementary information (see S1).

Data source selection

Data source selection was an iterative process which involved searching for sources of data, refining the search strategy, and reviewing sources for inclusion (Levac et al. 2010) (see Fig. 1). Where appropriate, we grouped sources according to their overarching ‘umbrella’ organisations or collectives as these were analogous in their purpose and operations. The inclusion criteria used in this review focused on capturing FAIR datasets (see Wilkinson et al. 2016) which included records of UK-native live, wild amphibians (Common frog *Rana temporaria*, Common toad *Bufo bufo*, Great crested newt *Triturus cristatus*, Natterjack toad *Epidalea calamita*, Northern pool frog *Pelophylax lessonae*, Palmate newt *Lissotriton helveticus*, Smooth newt *Lissotriton vulgaris*) and reptiles (European adder *Vipera berus*, Grass snake *Natrix helvetica*, Sand lizard *Lacerta agilis*, Slow-worm *Anguis fragilis*, Smooth snake *Coronella austriaca*, Viviparous lizard *Zootoca vivipara*). Data sources were included if they were discoverable using the search strategy (‘Findable’), contained available datasets (‘Accessible’) with broadly applicable content and language (‘Interoperable’), and had descriptive information available about the sampling methods used for generating data (‘Reusable’).

Data charting

We collated information on the characteristics of datasets from data source websites and through consultations with data publishers, where this could be arranged (incl. ARC Trust, ARG UK, BTO, Royal Society for the Protection of Birds, Froglife, BRC, and The Woodland Trust). We abstracted the metadata and sampling event information for each data source using a data charting technique to synthesise and interpret the information. Where available, we captured the following information using a standardised form: source name; publisher/organiser; background information and purpose; recorder characteristics; type of available data; temporal coverage (i.e., year of establishment and/or year that source became involved in amphibian and/or reptile data collection, storage and/or management); geographic coverage; taxonomic coverage; data quality assurance procedures; data transfer activity.

Methodological appraisal

We summarised data sources according to their characteristics, data generation procedures and dataset attributes. This included an assessment of the taxonomic, geographic and temporal scope of the dataset, recorder characteristics and any data quality assurance techniques used, as derived from the available metadata associated with each data source. We categorised data sources based on the structural traits of their equivalent datasets using




Fig. 1 Flowchart illustrating the process of searching for data sources and the selection and grouping of sources meeting criteria for inclusion in this scoping review

five-point Likert scale ranging from highly structured (A) to unstructured (E) data (see electronic supplementary information, S2).

Data analysis

We performed a network analysis to visualise the mobilisation of data between sources and identify prominent sources in the network. Data sources were represented in a network as nodes and data transfers were mapped as links, plotted using the *GGally* (Schloerke et al. 2021), *ggplot2* (Wickham 2016), *network* (Butts 2015), and *igraph* (Csardi and Nepusz 2016) R packages. Network metrics were computed using the ‘*networkD3*’ (Allaire et al. 2017) R package. The ‘degree’ of nodes reflected the number of directional links with other nodes. The average number of links to pass through a node was calculated as ‘betweenness’. ‘Betweenness centrality’ was computed as the number of instances in which a node fell on the shortest path between two other nodes, thus facilitating data transfer between sources. To identify influential sources in the network, eigenvector values were computed which took account of the ‘degree’ of nodes and their connectedness to other well-connected nodes. Nodes with high eigenvector values were centralised in the network and were, therefore, largely influential in the mobilisation of data across the data landscape. All data analysis procedures were carried out using R studio v4.0.2 (R Core Team 2021).

Results

Data source attributes, contributors and quality assurance

We identified 45 sources of UK amphibian and reptile observation data from the scoping review (see Table 1). These sources clustered into three typologies: ‘recording projects’ (n=26), ‘recording communities’ (n=4) and ‘digital data platforms’ (n=15). Recording projects reflected a coordinated data collection activity that followed a defined methodology (e.g. systematic or semi-structured monitoring, see Table 1) with a discrete taxonomic, geographic or temporal focus. Recording communities were organised groups of individuals that carried out the collection of data and coordinated the storage and sharing of data. Recording communities typically organised and participated in sampling events, though were not defined by methodological constraints, and hence we identified recording communities that were associated with several datasets. Digital data platforms represented online tools for the direct capture, storage, or export of records. The structure of datasets varied across data sources. Heterogeneous datasets (Group C) were the most widely available across sources (42%), particularly for digital data platforms, with component records an aggregation of verified and unverified opportunistic sightings and systematic survey data collected by a variety of recorders. There was comparable abundance of highly structured (Group A, 24%) and semi-structured datasets (Group B, 22%). Generally, these datasets consisted of validated and verified records that had been collected using pre-defined (semi-)systematic methodologies.

The recorders contributing to the various data sources ranged from novice citizen scientists to experienced species surveyors and a combination thereof. Nine recording projects solely recruited citizen scientists (of any ability) to collect data. Seven recording projects only recruited experienced (often licenced) species surveyors to collect data, particularly when European-protected species were the taxonomic focus of monitoring. When these

Table 1 Sources of United Kingdom amphibian and reptile observation data

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Add an Adder	Amphibian and Reptile Conservation Trust	2008	Recording project	A citizen science project to collect information about historical adder distributions across the UK	E ^o
ARC reserves surveys	Amphibian and Reptile Conservation Trust	Pre-1950	Recording project	Surveys to monitor the amphibian and reptile populations on reserves managed by the Amphibian and Reptile Conservation Trust	B ^o
Amphibian & Reptile Groups (ARGs/RAGs)	Amphibian & Reptile Groups of the UK	2005	Recording community	Volunteers participate in amphibian and reptile survey and monitoring and other conservation activities. Records are submitted to their county-based group	B ^o , [¶]
ARGWEB	Amphibian & Reptile Groups of the UK	2016	Digital data platform	An online suite of apps developed for volunteer amphibian and reptile groups in the UK and Ireland to manage their membership, stakeholder base, and manage their collection, storage and analysis of amphibian and reptile observation data	C ^o , [¶]
Beautiful Burial Grounds	Caring for God's Acre	2018	Recording project	A citizen science project to promote the conservation of churchyards and burial sites across England and Wales. The project records all taxa including amphibians and reptiles	C ^o
Big Spawn Count	Northern Ireland Environment Agency, National Museums Northern Ireland	2019	Recording project	A citizen science project for the wider public to survey waterbodies in search of frogspawn	C ^o

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Connecting the Dragons	Amphibian and Reptile Conservation Trust	2018	Recording project	A citizen science project to restore amphibian and reptile habitats, promote community engagement and monitor population changes in South Wales	B°
Dorset Fixed Survey Units	Amphibian and Reptile Conservation Trust	1950s	Recording project	Surveys to monitor reptile populations on reserves in Dorset, England, managed by the Amphibian and Reptiles Conservation Trust	B°
Dragon Finder App	Froglife	2016	Digital data platform	A mobile application aimed at the wider public to identify and record amphibians and reptiles	D°
Environmental Information Data Centre (EIDC)	Natural Environment Research Council	1994	Digital data platform	A service to manage and promote the re-use of valuable datasets concerned with the terrestrial and freshwater sciences in the UK	A°
Garden BirdWatch	British Trust for Ornithology	2008	Recording project	A citizen science project to monitor large-scale patterns in garden use by birds and selected other taxa (including amphibians and reptiles)	B°
Garden Dragon Watch	Amphibian and Reptile Conservation Trust	2020	Recording project	A citizen science project aimed at the wider public to record amphibians and reptiles in gardens in the UK	C°

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Garden Wildlife Health	Zoological Society of London, British Trust for Ornithology, Froglife, Royal Society for the Protection of Birds	2013	Recording project	A citizen science project monitoring spatial patterns and long-term trends in the diseases of British wildlife, with particular emphasis on the surveillance of birds, amphibians, reptiles and hedgehogs in Great Britain	C ^{o,f}
Gems in the Dunes	Amphibian and Reptile Conservation Trust	2018	Recording project	Part of the 'Back from the Brink' programme; the project focuses on rare species on the Sefton Coast of England including the Merseyside sand lizard and natterjack toad. The project aims to record the species, undertake habitat management and rehabilitation, and to promote outreach and education	B ^o
GeoPortal	Natural England	2013	Digital data platform	The central database for all public biodiversity data held and collected by Natural England	C ^{o,f}
Global Biodiversity Information Facility	Global Biodiversity Information Facility	2001	Digital data platform	An international network and data infrastructure to provide open access to data about all types of life on Earth	C ^o

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Great Crested Newt Class Level 1 Licence Returns	Natural England (statutory agency)	1981	Recording project	As part of the conditions of licensing, licenced Great crested newt surveyors submit annual reports on the results of their surveys to provide information on the distribution of this European protected species. These results have also been used to inform activities related to the District Level Licensing Scheme	B ^o
Great Crested Newt District-Level Licensing Scheme	Natural England (statutory agency), NatureSpace Partnership	2017	Recording project	An eDNA surveillance scheme to replace the need for developers to conduct pre-development surveys for Great crested newt. The results of surveys are used to map areas where great crested newt presence may warrant mitigation action by developers	A ^y
HerpMapper	The HerpMapper Project	2013	Digital data platform	A citizen science project to collate and share information about amphibian and reptile observations worldwide	C ^o
iNaturalist	California Academy of Sciences, National Geographic Society	2010	Digital data platform	A citizen science project for members of the online community to share wildlife observations and create data for science and conservation	B ^o

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
iRecord	Biological Records Centre	2013	Digital data platform	A wildlife recording platform based on Indicia for citizen scientists and biological recording schemes in the UK	C ^o
iSpot	The Open University	2012	Digital data platform	A citizen science project for members of the general public to get help with identifying anything in nature in the UK	D ^o
Lancashire Amphibian & Reptile Atlas	Lancashire Amphibian & Reptile Group	2010	Recording project	A collaborative project which aims to record amphibian and reptile data to support an atlas for Lancashire, Greater Manchester and Merseyside, England	C ^o
Living ARCive	Amphibian and Reptile Conservation Trust	2008	Digital data platform	The central database for all data held and collected by the Amphibian and Reptile Conservation Trust	C ^{o,f}
Living Record	Living record	2010	Digital data platform	An online recording tool for capturing opportunistic and site-based survey data across all taxa, including amphibian and reptiles in the UK	B ^o

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Local Environmental Records Centres (LERCs)	ALERC, Local Environmental Records Centres	1902	Digital data platform	Independent not-for-profit organisations that collect, collate and manage information on the natural environment for a defined geographic area. Information products and services are made accessible to a range of audiences including decision-makers, the public, and researchers. The Association of Local Environmental Records Centres (ALERC) is a membership organisation representing LERCs across the UK	C ^o
Local Nature Partnerships	Various	Pre-1950s	Recording community	Local collaborations operating to protect, enhance and monitor local biodiversity and habitats	C ^o
Make the Adder Count	Amphibian & Reptile Groups UK	2005	Recording project	A citizen science project to collate long-term, standardised, quantitative data on adder occupancy across the UK	A ^o
National Amphibian and Reptile Recording Scheme	Amphibian and Reptile Conservation Trust	2007	Recording project	A citizen science project to generate robust, baseline data for reptiles and amphibians across the UK	A ^o

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
National BioBlitz Network	Natural History Consortium	2009	Recording community	A dispersed network of citizen scientists participating in 24-h events to find and identify species of plants, animals, fungi, and other organisms in one patch	D ^o
National Biodiversity Network Atlas	National Biodiversity Network	2001	Digital data platform	An online data platform created to exchange biodiversity information	C ^o
Natterjack Toad Monitoring Programme	Amphibian and Reptile Conservation Trust	1986	Recording project	A project monitoring all Natterjack toad sites across the UK to establish baseline counts for each colony, monitor changes in populations, and undertake appropriate habitat management	A ^o
Nature's Calendar	The Woodland Trust	2011	Recording project	A citizen science project to monitor selected UK wildlife to understand their phenological responses to changes in weather and climate	C ^o
PondNet	Freshwater Habitats Trust	2012	Recording project	A citizen science project to monitor trends in both pond quality and widespread, localised and restricted pond species, including amphibians, to inform conservation actions	A ^{o,f}

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Pool Frog Monitoring Programme	Amphibian and Reptile Conservation Trust	2005	Recording project	Post-release monitoring of Pool frogs to assess breeding success, pond-use and changes in habitat quality, and estimate the survival of spawn/larvae, juveniles and adults	A°
Record Pool	Amphibian & Reptile Groups UK, Amphibian and Reptile Conservation Trust	2005	Digital data platform	An online recording tool available to the wider public and professional surveyors to collect opportunistic and survey data on reptiles and amphibians in the UK. Data is made available locally and nationally, for conservation purposes	C°
Reptiles & Amphibians of the UK (RAUK)	www.herpetofauna.co.uk	Early 2000s	Digital data platform	An online forum providing information on the identification and recording of UK herpetofauna	D°
Royal Society for the Protection of Birds Reserves Biodiversity Monitoring Programme	Royal Society for the Protection of Birds	Pre-1950s	Recording project	Surveys to monitor the biodiversity on reserves managed by the Royal Society for the Protection of Birds	C°
Sand Lizard Monitoring Programme	Amphibian and Reptile Conservation Trust	1961	Recording project	Monitoring scheme for populations of sand lizards and their habitats through licenced species surveys and engagement with stakeholders. Surveys are conducted on a three-yearly cycle	A°

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
Scottish Grass Snakes	Caledonian Conservation, Amphibian and Reptile Groups of the UK	2012	Recording project	A collaborative project to collate observational data on the distribution of grass snake in Scotland	C°
Snakes in the Heather	Amphibian and Reptile Conservation Trust	2019	Recording project	A citizen science project to conserve and monitor populations of Smooth snake throughout its range in Southern England	A°
Toads on Roads	Froglife	1980s	Recording project	Local toad crossings across the UK are registered with Froglife. Citizen scientists monitor crossing sites to rescue toads and collect data on numbers of amphibians and mortalities during the spring breeding season	A°
UK Environmental Change Network Frog Data	UK Centre for Ecology & Hydrology	1994	Recording project	A project to record the phenology of spawning in the common frog in selected ponds and ditches and to identify any significant changes to population 'health' and the environment	A°
What's In Your Pond?	Northern Ireland Environment Agency, National Museums Northern Ireland	2018	Recording project	A citizen science project that collects records on common frog and smooth newt in Northern Ireland for conservation purposes	B°

Table 1 (continued)

Source	Organiser/publisher	Year of establishment ^a	Type of data source	Aims & operations	Data type ^b and structure ^c
The Wildlife Trusts	The Wildlife Trusts	Pre-1950s	Recording community	An umbrella organisation seeking to promote environmental and biodiversity conservation, research and education. Each county-based Wildlife Trust is an independent charity with its own membership scheme. Wildlife recording is performed by citizen scientists and professional staff	C ^c

^a Dates relevant to when the source became active in amphibian and reptile data collection, storage and/or management

^b Data type key: #Human observations; #Environmental DNA

^c Qualitative data structure key:

A = Source datasets appeared highly structured and included detailed information content for records. Records were collected using pre-defined sampling procedures, often with repeated surveys at defined intervals and using specialist survey equipment. Grid references were validated for accuracy and the recorders had expertise in recording species data

B = The data source generated semi-structured datasets and included some information content for records. Records had either been verified by an expert or the species identification were likely to be accurate based on the methods of identification or the expertise of the recorder. Records may not have been collected according to a defined, repeat survey methodology nor specialist survey equipment, but often at least semi-structured surveys had been used. The recorders may or may not have been trained in survey methods or identification

C = Datasets generated by the data source appeared heterogeneous in nature. Datasets were aggregations of records generated from opportunistic sightings and systematic surveys. Verified data was likely to be accurate, though unverified data may be less accurate. The expertise of the recorder, the sampling event procedures and species identification checks varied across records or were unknown

D = Records comprised of entirely opportunistic sightings, with little-to-no information pertaining to the sampling event, the recorder or metadata. Records may have had little or no verification and validation. Photographs of observations may be available for some records

E = Records comprised of entirely opportunistic sightings that were not collected according to a pre-defined sampling methodology. There were no defined record verification or validation procedures and there were no details on the expertise of data recorders, nor information pertaining to species identification

species were the taxonomic focus of citizen science-based recording projects, citizen scientists accompanied licenced species surveyors to collect data on systematic surveys and received training in species survey methodologies and identification. Across all citizen science-based recording projects, six of the project organisers provided only identification guides, whereas five of the projects did not issue identification guides or any formal training to citizen scientists. Most digital data platforms included verification and validation procedures for quality control purposes. Typically, this encompassed verification by species experts (e.g. ‘County Recorders’). Record verification for some platforms also relied on community knowledge, whereby online communities of wildlife recorders provided identification suggestions to each other’s observations, and automated computer checks to flag (likely) errors to recorders entering data or to verifiers after records had been entered. Further information on the characteristics of each data source is provided in the electronic supplementary information (S3).

Taxonomic coverage

Sources of data for all native species of UK amphibians and reptiles were identified. As illustrated in Fig. 2, multispecies datasets, particularly for widespread species, featured extensively across sources. Recording communities only generated multispecies data. Digital data platforms also typically captured multispecies data, whilst a targeted taxonomic focus was more common amongst recording projects. Amphibians were the taxonomic




Fig. 2 Focal species of United Kingdom (UK) amphibian and reptile observation data sources. Widespread amphibians include Common frog (*Rana temporaria*), Common toad (*Bufo bufo*), Great crested newt (*Triturus cristatus*), Palmate newt (*Lissotriton helveticus*), Smooth newt (*Lissotriton vulgaris*). Other UK-native amphibians (currently with legislative reporting requirements) include Natterjack toad (*Epidalea calamita*) and Northern pool frog (*Pelophylax lessonae*). Widespread reptiles include European adder (*Vipera berus*), Grass snake (*Matrix helvetica*), Slow-worm (*Anguis fragilis*), Viviparous lizard (*Zootoca vivipara*). Other UK-native reptiles (currently with legislative reporting requirements) comprise Sand lizard (*Lacerta agilis*) and Smooth snake (*Coronella austriaca*)

focus of data sources more frequently than reptiles. Common frog had the best coverage across all data sources, as approximately three-quarters of all sources captured observation data for this species. Sources of data for great crested newts were also widely available and seven sources included eDNA records. European adder, sand lizard and grass snake were the focal species most frequently targeted amongst the reptile recording projects. No source specifically targeted the sole collection of data for palmate newt, slow-worm, smooth newt or viviparous lizard, though data for these species were captured by multispecies sources and can be obtained from sources targeting species with legislative reporting requirements.

Geographic coverage

There was division in the geographic availability of data sources included in this review (see electronic supplementary information, S5). Data sources pertained mostly to England ($n=41$), followed by Wales ($n=34$) and Scotland ($n=30$). Northern Ireland had the fewest sources of data ($n=24$). Data available through digital data platforms generally had the largest (national) geographic coverage. We note, however, that while many data sources had the potential for national coverage, their actual spatial extents were often more restricted to a number of targeted sites (see electronic supplementary information, S3). This was particularly evident for recording projects and for recording communities that were bound by a local (e.g. Vice-county) perimeter of operation (see electronic supplementary information, S3).

Temporal coverage

The number of data sources has fluctuated widely over the last century (see Fig. 3). Records of human observations were available from 1900 whilst eDNA records emerged from 2013 onwards. One digital data platform was active since the start of the 1900s. Recording communities emerged in the 1910s and the first (still active) recording projects emerged in, or shortly prior to, the 1950s. The largest increase in the total number of active sources was observed between the 2000s and the 2010s; rising from 12 sources at the end of the 1990s to 44 active sources by 2019. Despite the majority of sources emerging later towards the 2000s, digital data platforms usually included historical records prior to the platform's establishment. Recording projects generally ranged between 3 (IQ1) and 31 (IQ3) years, with a mode of 3 years. Historical records were also available through the ARC Reserves Surveys, reflecting some of the earliest records available through a recording project. The Natterjack Toad Monitoring Programme and the Sand Lizard Monitoring Programme had the longest periods of continuous monitoring of any recording project, spanning 40 and 37 years respectively. Recording communities also typically had extensive periods of activity, on average 66 years.

Network analysis

The network analysis illustrated the flow of data between sources (see Fig. 4). The analysis indicated that the UK amphibian and reptile monitoring portfolio is a dynamic and fragmented data landscape. Two isolated nodes, with no links to any other source, were identified in the network. All other sources had at least one link, but some appeared to only receive data and did not export data to other sources. The degree (' g ')




Fig. 3 The number of active sources of United Kingdom amphibian and reptile observation data per decade from 1900 to 2021. The number of sources is depicted as counts of digital data platforms, recording communities and recording projects since the source became active in amphibian and reptile data collection, storage and/or management

of nodes averaged to 4.6 links per data source though 53% of sources had two or fewer links. Digital data platforms generally had the highest number of connections. Overall, the NBN Atlas had the highest number of connections ($g = 21$), followed by the LERCs ($g = 19$) and the Living ARCive ($g = 17$). The ARGs/RAGs ($g = 13$) and Great Crested Newt Level 1 Licence Returns ($g = 6$) sources were the most connected recording community and recording project in the network, respectively. On average, data mobilised across 2.4 links between sources within the network. Digital data platforms often fell on the shortest path between other nodes in the network (betweenness centrality, ' bc ') and were highly influential over the mobilisation of data across the network (eigenvector centrality, ' ec '). The NBN Atlas ($bc = 309$), Living ARCive ($bc = 281$), and LERCs ($bc = 172$) had the highest bc across all sources, indicating that these sources, particularly the NBN Atlas, most frequently bridged the transfer of data between two other sources in the network. The LERCs had the highest ec overall, indicating that these were the most centralised sources of data within the network, with high connectedness to other centralised sources. Other centralised sources ($ec > 0.60$) in the network included the NBN Atlas ($ec = 0.90$), iRecord ($ec = 0.67$), ARGs/RAGs ($ec = 0.65$), and Record Pool ($ec = 0.64$).

evidence base is only likely to increase. Many of the factors driving this situation are pertinent to biodiversity monitoring more widely, so the problems and solutions are likely to be general. The integration of data in a unifying network infrastructure that streamlines fragmented monitoring may offer more precise, up-to-date biodiversity assessments over multiple scales.

The UK amphibian and reptile monitoring portfolio is a diverse data landscape comprising recording projects, recording communities and digital data platforms that collect, curate, and share data for all native species. The large number of sources is testament to a growing conservation community and should be celebrated. However, this diversity presents challenges for synthesis in research and decision-making processes at national scales. Digital data platforms are key for the mobilisation of data, particularly the LERCs and the NBN Atlas, which are highly connected and centralised sources in the data landscape. Collectively, the LERCs interacted with other important sources more frequently than the NBN Atlas alone which led to their aggregated position as the most centralised sources in the data landscape. Though the mobilisation of data from some LERCs can sometimes be restricted by paywalls, formatting incompatibility or due to constraints on data sensitivity and confidentiality. At its inception, the NBN Atlas sought to become; “the best wildlife information management structure”, by capturing, enhancing and mobilising wildlife data, making information widely available and engaging people about wildlife (NBN Trust 2014). We found that the NBN Atlas is highly connected to other data sources and is a central distributor of information, frequently bridging the transfer of data between other sources. This suggests that the NBN Atlas has been reasonably successful towards achieving its aims in collating and making data widely available. However, the full vision of the NBN Atlas may not yet be realised as we found that it lacked detailed metadata on the sampling protocols used to generate datasets. This information is essential for reusing data in other contexts. Taken together, our findings suggest that whilst the NBN Atlas is the most publicly accessible source and has the potential to reach its objective of becoming “the best wildlife information management structure”, it currently falls short due to insufficient metadata and lower rates of data sharing with other important data distributors than could be achieved.

It is important to stress that the high centrality metric used in our analysis does not directly relate to “the ‘best’ data source. Instead, we used this metric to highlight which sources are influential in the mobilisation of biodiversity data (Zhao and Zhang 2020). There are many advantages to diversity in species recording and data management. Multiple organisations working together can address more facets of biodiversity monitoring beyond the capacity of any standalone organisation. A variety of stakeholders also fulfil different roles within a nature conservation network; from bottom-up primary data generators, with detailed regional or taxonomic expertise, to top-down statutory monitoring and governance. It is encouraging that we observed a high reciprocity of data transfer between sources as this suggests that many organisations are promoting a FAIR and open data landscape. However, high mobilisation of data may affect the quality of available data as there are multiple levels at which information may be lost through data manipulation and interpretation by data users. We identified isolated sources and one-way links which may pose significant weaknesses in the network. Catastrophic data loss could occur for some species and regions if an organisation collapses or ceases to collect data into the future. Poorly connected sources may also be less likely to contribute to wider biodiversity conservation efforts than well-connected sources. Hence, sources of this nature may limit the mobilisation of data across a network, hampering future integration efforts and restricting the information available for research and for informing national policy.

Currently, none of the existing sources of UK amphibian and reptile data appear to provide sufficient baseline information for national monitoring of all species, though some sources may have adequate foundations to build on for specific species, regions and time periods. Digital data platforms and recording communities generally have wide taxonomic scope, acting as “catch-all buckets” for any available data across large temporal and spatial extents. However, data made available through digital data platforms tend to lack sufficient quality to make reliable inferences of biodiversity dynamics (Bayraktarov et al. 2019) as they typically contain only presence-only records. National platforms with information-rich abundance data is lacking, but incorporation of standards such as the Darwin Core ‘Event’ category (Wieczorek et al. 2012), which formalise the capture and presentation of sampling information across heterogeneous datasets would make this possible. Nonetheless, the large datasets of presence-only records available through digital data platforms can complement systematic surveys, filling some of the spatial and temporal gaps often associated with small-scale studies (Isaac et al. 2020). In isolation, however, these datasets usually contain a variety of data biases (Petrovan et al. 2020), which can lead to misleading conclusions if not recognised and accounted for (Isaac and Pocock 2015).

Structured datasets arising through systematic monitoring of multiple species can enable standalone assessments of biodiversity. We observed that systematic monitoring is often restricted to a selection of sites within regions (i.e., via convenience sampling as sites are managed by project coordinators). Systematic monitoring currently favours amphibians over reptiles and most structured datasets are limited to species with legislative reporting requirements. For instance, most of the existing suite of structured amphibian and reptile datasets are single-species and arise from the systematic monitoring of European-protected species coordinated solely by conservation organisations. We did find that citizen science-based recording projects frequently generated multispecies datasets, usually for species that are widespread in their occurrence. Though such data may contain sampling biases of varying degrees, particularly as large heterogeneous collections of records (Isaac and Pocock 2015), and there has been limited empirical analyses of these datasets with regard to amphibians and reptiles (though see Humphreys et al. 2011; Wilkinson and Arnell 2013). Where recording projects, however, have focussed on single-species monitoring, there have been relatively more empirical outcomes. For instance, common frog (Scott et al. 2008, great crested newt (Beebee 1997; Denoël 2012), common toad (Petrovan and Schmidt 2016), and adder (Gardner et al. 2019) have all featured in empirical studies and we observed that these species are a popular focus in recording projects. Whereas, quantitative assessments for palmate newt, smooth newt, slow-worm and viviparous lizard are largely lacking and we found that these species had the lowest rates of monitoring of all widespread species. In the case of palmate newt, this could in part be due to difficulties with identification or lower rates of occupancy nationally. Likewise, we observed a clear geographic bias for England and lower rates for Northern Ireland. Though differences in human population densities and regional taxonomic prevalence could explain these findings.

Advances in computing are likely to have led to a variety of means for collecting, validating, and verifying data, and therefore have likely contributed to an increase in the uptake of citizen science approaches in biodiversity monitoring in recent years (August et al. 2015; Pocock et al. 2017). For instance, eDNA has emerged as a viable tool for amphibian monitoring (Biggs et al. 2015), and we found seven sources of eDNA in our search, all initiated since 2013. Amphibian and reptile surveillance is also a primary frontier in several emerging ecological remote-sensing techniques such as camera trapping (Welbourne et al. 2017). In line with other accounts (e.g. James 2011; Roy et al. 2012; Pocock et al. 2015),

we observed that an array of data quality assurance techniques can be imposed on datasets before being made available to data users. We caution, however, that excessive manipulation of data by publishers may reduce the quality of available metadata depending on the format in which it is published. As is typical for sources of biodiversity data (Roy et al. 2012; Dobson et al. 2020; Thornhill et al. 2021), many UK amphibian and reptile datasets reflect heterogeneous collections of records originating from opportunistic and systematic surveys. However, we found that extracting specific data collection procedures from these sources was either challenging or impossible. By restricting the availability of sampling event information associated with datasets, the potential for reuse of existing biodiversity data may be constrained for several large data sources.

Biodiversity and conservation science is in the midst of adopting more formal and systematic approaches to evidence synthesis. Historically, evidence reviews in biodiversity science have had lower standards of reproducibility (Grames and Elphick 2020). We adapted the traditional scoping review framework (Arksey and O'Malley 2005) to suit the needs of this review to provide a rigorous and transparent approach to mapping a baseline account of FAIR UK amphibian and reptile observation data; permitting gaps in the current monitoring portfolio to come to light. We hope that this study may serve as a template for summarising sources of biodiversity data, enabling comparable assessments and appraisals of existing data for other taxa and environments. We grouped and evaluated some sources as collective units as the evaluation of their separate entities was not feasible. These represented branched organisations that operated as independent groups. Therefore, it is important to note that not every independent branch of grouped sources (i.e. the 'LERCs', 'ARGs/RAGs', 'Local Nature Partnerships', and 'Wildlife Trusts') will necessarily have links to all of the data sources identified in the network analysis. Nonetheless, we expect that mapping them in this way provides a typical depiction of the characteristics and wider mobilisation of data and across a biodiversity data management landscape.

Effective large-scale biodiversity monitoring requires integration of localised and fragmented monitoring efforts, thereby extending the capacity of any stand-alone programme, to address pressing science and conservation issues (Kühl et al. 2020). We recommend integration of datasets and coordinated monitoring for more comprehensive status and trends assessments. A discussion on the complementarities amongst data sources in this review is provided in the electronic supplementary information (see S4). To achieve an integrated monitoring portfolio, stakeholder collaboration within a unified infrastructure, such as a network, is paramount. Aligning pathways in shared, interoperable formats, combined with core monitoring, allows for robust analyses on the patterns of large-scale biodiversity change (Kühl et al. 2020). We conclude this review by providing recommendations to improve on current practice and achieve an integrated biodiversity monitoring portfolio.

First, to improve transparency and allow data to be used more widely, data publishers should seek to improve the 'interoperability' and 'reusability' of datasets by providing data in clear, interoperable formats [e.g., 'Darwin Core' (Wieczorek et al. 2012)] to align with data standards and ensure that important sampling event metadata accompany records in datasets. Second, we urge data publishers to provide clarity on how information is disseminated and shared between recorders, scheme organisers, scientists and decision-makers. As a minimum, this would provide information about the level of data duplication when combining datasets in a single analytical framework. Data should be presented in a way that would enable it to be traced back to its origin and allow data-users to ascertain how the data was collected. Examining all facets of network communication was not within the scope of this review, but clear channels of communication will be essential to enable an integrated network to generate and share information more effectively. Future work should

explore current practice for sharing information and evidence between data publishers and government bodies so that clear channels for sharing data and information can permeate across the landscape. Third, the development of a validated tool to assess the ‘structure’ of datasets would likely enable more timely identification of fit-for-purpose datasets. Finally, we advocate for the establishment of an effective realised Biodiversity Observation Network, co-developed by stakeholders, and the enhancement of existing centralised data infrastructures that take account of these recommendations for collating, characterising, and sharing biodiversity data.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10531-022-02502-w>.

Acknowledgements This work was supported by the Natural Environment Research Council and ARIES Doctoral Training Partnership [Grant Number NE/S007334/1]. We are grateful to the Amphibian and Reptile Conservation (ARC) Trust, Amphibian and Reptile Groups (ARG) UK, the British Trust for Ornithology (BTO), Royal Society for the Protection of Birds (RSPB), Froglife, Biological Records Centre (BRC), and the Woodland Trust for sharing information on their data and their invaluable expertise on the dynamics of the UK amphibian and reptile data management landscape. In particular, we would like to thank Rob Ward (ARC Trust), Yvette Martin (ARC Trust), Emma Gardner (UK Centre for Ecology & Hydrology), Kate Risley (formerly BTO), Rebecca McHugh (RSPB), Silviu Petrovan (University of Cambridge), Jenny Tse-Leon (Froglife), Steve Langham (Surrey ARG), David Roy (BRC), Martin Harvey (BRC) and Lorianne Whittle (Woodland Trust). We thank our anonymous reviewers for their constructive feedback on previous manuscript drafts. We extend thanks to the thousands of dedicated volunteers who collect data on amphibians and reptiles in the UK and hope that this review showcases the value of their efforts for UK amphibian and reptile conservation.

Author contributions All authors contributed to the study conception and design. Material preparation, data searches, synthesis and analysis were performed by RKT. The first draft of the manuscript was written by RKT and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding This work was supported by the Natural Environment Research Council and the ARIES DTP [Grant Number NE/S007334/1].

Declarations

Competing interests The authors have no conflicts of interest to declare.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Allaire JJ, Gandrud C, Russell K, Yetman CJ (2017) networkD3: D3 JavaScript Network Graphs from R
 Arksey H, O’Malley L (2005) Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 8:19–32. <https://doi.org/10.1080/1364557032000119616>
 August T, Harvey M, Lightfoot P et al (2015) Emerging technologies for biological recording. *Biol J Linn Soc* 115:731–749. <https://doi.org/10.1111/bj.12534>
 Bayraktarov E, Ehmke G, O’Connor J et al (2019) Do big unstructured biodiversity data mean more knowledge? *Front Ecol Evol*. <https://doi.org/10.3389/fevo.2018.00239>

- Beebee TJC (1997) Changes in dewpond numbers and amphibian diversity over 20 years on chalk downland in Sussex, England. *Biol Conserv* 81:215–219. [https://doi.org/10.1016/S0006-3207\(97\)00002-5](https://doi.org/10.1016/S0006-3207(97)00002-5)
- Beebee TJC, Ratcliffe S (2018) Inferring status changes of three widespread British reptiles from NBN Atlas records. *Herpetol Bull* 143:18–22
- Biggs J, Ewald N, Valentini A et al (2015) Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biol Conserv* 183:19–28. <https://doi.org/10.1016/j.biocon.2014.11.029>
- Bird TJ, Bates AE, Lefcheck JS et al (2014) Statistical solutions for error and bias in global citizen science datasets. *Biol Conserv* 173:144–154. <https://doi.org/10.1016/j.biocon.2013.07.037>
- Bonney R, Shirk JL, Phillips TB et al (2014) Next Steps for citizen science. *Science* 343:1436–1437. <https://doi.org/10.1126/science.1251554>
- Boyd RJ, Powney GD, Carvell C, Pescott OL (2021) occAssess: an R package for assessing potential biases in species occurrence data. *Ecol Evol* 11:16177–16187. <https://doi.org/10.1002/ece3.8299>
- Burgess HK, DeBey LB, Froehlich HE et al (2017) The science of citizen science: exploring barriers to use as a primary research tool. *Biol Conserv* 208:113–120. <https://doi.org/10.1016/j.biocon.2016.05.014>
- Butts C (2015) network: Classes for Relational Data. The Statnet Project (<http://www.statnet.org>). R package version 1.13.0.1. <https://CRAN.R-project.org/package=network>
- Cohn JP (2008) Citizen science: can volunteers do real research? *Bioscience* 58:192–197. <https://doi.org/10.1641/B580303>
- Constable H, Guralnick R, Wiczorek J et al (2010) VertNet: a new model for biodiversity data sharing. *PLOS Biol* 8:e1000309. <https://doi.org/10.1371/journal.pbio.1000309>
- Csardi G, Nepusz T (2016) The igraph software package for complex network research
- Denoël M (2012) Newt decline in Western Europe: highlights from relative distribution changes within guilds. *Biodivers Conserv* 21:2887–2898. <https://doi.org/10.1007/s10531-012-0343-x>
- Dobson ADM, Milner-Gulland EJ, Aebischer NJ et al (2020) Making messy data work for conservation. *One Earth* 2:455–465. <https://doi.org/10.1016/j.oneear.2020.04.012>
- Dornelas M, Gotelli NJ, McGill B et al (2014) assemblage time series reveal biodiversity change but not systematic loss. *Science* 344:296–299. <https://doi.org/10.1126/science.1248484>
- Dunford R, Berry P (2012) Climate change modelling of English amphibians and reptiles: report to Amphibian and Reptile Conservation Trust (ARC-Trust)
- Fox R, Bourn NAD, Dennis EB et al (2019) Opinions of citizen scientists on open access to UK butterfly and moth occurrence data. *Biodivers Conserv* 28:3321–3341. <https://doi.org/10.1007/s10531-019-01824-6>
- Gardner E, Julian A, Monk C, Baker J (2019) Make the adder count: population trends from a citizen science survey of UK adders. *Herpetol J* 29:57–70. <https://doi.org/10.33256/hj29.1.5770>
- Geldmann J, Heilmann-Clausen J, Holm TE et al (2016) What determines spatial bias in citizen science? Exploring four recording schemes with different proficiency requirements. *Divers Distrib* 22:1139–1149. <https://doi.org/10.1111/ddi.12477>
- Grames EM, Elphick CS (2020) Use of study design principles would increase the reproducibility of reviews in conservation biology. *Biol Conserv* 241:108385. <https://doi.org/10.1016/j.biocon.2019.108385>
- Griffiths RA, Foster J, Wilkinson JW, Sewell D (2015) Science, statistics and surveys: a herpetological perspective. *J Appl Ecol* 52:1413–1417. <https://doi.org/10.1111/1365-2664.12463>
- Hayhow DB, Eaton MA, Stanbury AJ et al (2019) The State of Nature 2019. The State of Nature partnership
- Humphreys E, Toms M, Newson S et al (2011) An examination of reptile and amphibian populations in gardens, the factors influencing garden use and the role of a “Citizen Science” approach for monitoring their populations within this habitat, BTO Research Report No. 572
- Isaac NJB, Pocock MJO (2015) Bias and information in biological records. *Biol J Linn Soc* 115:522–531. <https://doi.org/10.1111/bij.12532>
- Isaac NJB, van Strien AJ, August TA et al (2014) Statistics for citizen science: extracting signals of change from noisy ecological data. *Methods Ecol Evol* 5:1052–1060. <https://doi.org/10.1111/2041-210X.12254>
- Isaac NJB, Jarzyna MA, Keil P et al (2020) Data integration for large-scale Mmodels of species distributions. *Trends Ecol Evol*. <https://doi.org/10.1016/j.tree.2019.08.006>
- James T (2011) Improving Wildlife Data Quality. NBN Trust, Nottingham
- Jetz W, McGeoch MA, Guralnick R et al (2019) Essential biodiversity variables for mapping and monitoring species populations. *Nat Ecol Evol* 3:539–551. <https://doi.org/10.1038/s41559-019-0826-1>
- Kaarlejärvi E, Salemaa M, Tonteri T et al (2021) Temporal biodiversity change following disturbance varies along an environmental gradient. *Glob Ecol Biogeogr* 30:476–489. <https://doi.org/10.1111/geb.13233>

- Keil P, Storch D, Jetz W (2015) On the decline of biodiversity due to area loss. *Nat Commun* 6:8837. <https://doi.org/10.1038/ncomms9837>
- König C, Weigelt P, Schrader J et al (2019) Biodiversity data integration—the significance of data resolution and domain. *PLoS Biol* 17:e3000183. <https://doi.org/10.1371/journal.pbio.3000183>
- Kosmala M, Wiggins A, Swanson A, Simmons B (2016) Assessing data quality in citizen science. *Front Ecol Environ* 14:551–560. <https://doi.org/10.1002/fee.1436>
- Kühl HS, Bowler DE, Bösch L et al (2020) Effective biodiversity monitoring needs a culture of integration. *One Earth*. <https://doi.org/10.1016/j.oneear.2020.09.010>
- Levac D, Colquhoun H, O'Brien KK (2010) Scoping studies: advancing the methodology. *Implement Sci* 5:69. <https://doi.org/10.1186/1748-5908-5-69>
- McKinley DC, Miller-Rushing AJ, Ballard HL et al (2017) Citizen science can improve conservation science, natural resource management, and environmental protection. *Biol Conserv* 208:15–28. <https://doi.org/10.1016/j.biocon.2016.05.015>
- NBN Trust (2014) NBN strategy 2015–2020 first draft for consultation 2015
- Oliveira U, Paglia AP, Brescovit AD et al (2016) The strong influence of collection bias on biodiversity knowledge shortfalls of Brazilian terrestrial biodiversity. *Divers Distrib* 22:1232–1244. <https://doi.org/10.1111/ddi.12489>
- Parr TW, Ferretti M, Simpson IC et al (2002) Towards a long-term integrated monitoring programme in Europe: network design in theory and practice. *Environ Monit Assess* 78:253–290. <https://doi.org/10.1023/A:1019934919140>
- Petersen TK, Speed JDM, Grøtan V, Austrheim G (2021) Species data for understanding biodiversity dynamics: the what, where and when of species occurrence data collection. *Ecol Solut Evid* 2:e12048. <https://doi.org/10.1002/2688-8319.12048>
- Petrovan SO, Schmidt BR (2016) Volunteer conservation action data reveals large-scale and long-term negative population trends of a widespread amphibian, the common toad (*Bufo bufo*). *PLoS ONE* 11:e0161943. <https://doi.org/10.1371/journal.pone.0161943>
- Petrovan SO, Vale CG, Sillero N (2020) Using citizen science in road surveys for large-scale amphibian monitoring: are biased data representative for species distribution? *Biodivers Conserv* 29:1767–1781. <https://doi.org/10.1007/s10531-020-01956-0>
- Pocock MJO, Roy HE, Preston CD, Roy DB (2015) The Biological Records Centre: a pioneer of citizen science. *Biol J Linn Soc* 115:475–493. <https://doi.org/10.1111/bij.12548>
- Pocock MJO, Tweddle JC, Savage J et al (2017) The diversity and evolution of ecological and environmental citizen science. *PLoS ONE* 12:e0172579. <https://doi.org/10.1371/journal.pone.0172579>
- R Core Team (2021) R: a language and environment for statistical computing
- Roy HE, Pocock MJO, Preston CD et al (2012) Understanding Citizen Science & Environmental Monitoring. NERC Centre for Ecology & Hydrology and Natural History Museum, Final Report on behalf of UK-EOF
- Schloerke B, Cooke D, Larmarange J et al (2021) GGally: extension to “ggplot2”
- Scott WA, Pithart D, Adamson JK (2008) Long-Term United Kingdom Trends in the Breeding Phenology of the Common Frog, *Rana temporaria*. *J Herpetol* 42:89–96. <http://www.jstor.org/stable/40060486>
- Thornhill I, Cornelissen JHC, McPherson JM et al (2021) Towards ecological science for all by all. *J Appl Ecol* 58:206–213. <https://doi.org/10.1111/1365-2664.13841>
- Tredick CA, Lewison RL, Deutschman DH et al (2017) A rubric to evaluate citizen-science programs for long-term ecological monitoring. *Bioscience* 67:834–844. <https://doi.org/10.1093/biosci/bix090>
- Turner RK, Maclean IMD (2022) Microclimate-driven trends in spring-emergence phenology in a temperate reptile (*Vipera berus*): evidence for a potential “climate trap”? *Ecol Evol* 12:e8623. <https://doi.org/10.1002/ece3.8623>
- Walters M, Scholes RJ (eds) (2017) The GEO Handbook on Biodiversity Observation Networks. Springer International Publishing, Cham
- Welbourne DJ, Paull DJ, Claridge AW, Ford F (2017) A frontier in the use of camera traps: surveying terrestrial squamate assemblages. *Remote Sens Ecol Conserv* 3:133–145. <https://doi.org/10.1002/rse2.57>
- Wetzel FT, Bingham HC, Groom Q et al (2018) Unlocking biodiversity data: prioritization and filling the gaps in biodiversity observation data in Europe. *Biol Conserv* 221:78–85. <https://doi.org/10.1016/j.biocon.2017.12.024>
- Wickham H (2016) ggplot2: elegant graphics for data analysis
- Wieczorek J, Bloom D, Guralnick R et al (2012) Darwin core: an evolving community-developed biodiversity data standard. *PLoS ONE* 7:e29715. <https://doi.org/10.1371/journal.pone.0029715>
- Wilkinson JW, Arnell AP (2013) NARRS Report 2012: Establishing the Baseline (HWM Edition). ARC Research Report 13/01

- Wilkinson MD, Dumontier M, IjJ A et al (2016) The FAIR Guiding Principles for scientific data management and stewardship. *Sci Data* 3:160018. <https://doi.org/10.1038/sdata.2016.18>
- Zhao Y, Zhang H (2020) Eigenvalues make the difference—a network analysis of the Chinese Super League. *Int J Sports Sci Coach* 15:184–194. <https://doi.org/10.1177/1747954120908822>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Rebecca K. Turner^{1,2}  · Richard A. Griffiths²  · John W. Wilkinson³  ·
Angela M. Julian⁴ · Mike P. Toms⁵  · Nick J. B. Isaac¹ 

¹ UK Centre for Ecology & Hydrology, Wallingford OX10 8BB, Oxfordshire, UK

² Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation, University of Kent, Canterbury, Kent, UK

³ Amphibian & Reptile Conservation Trust, 744 Christchurch Road, Bournemouth BH7 6BZ, Dorset, UK

⁴ Amphibian and Reptile Groups of the UK, 82 Gidley Way, Horspath, Oxford OX33 1TG, UK

⁵ British Trust for Ornithology, The Nunnery, Thetford IP24 2PU, Norfolk, UK