



Kent Academic Repository

Prey, Sven, Herok, Christoph, Fantuzzi, Felipe, Bolte, Michael, Lerner, HansWolfram, Engels, Bernd and Wagner, Matthias (2022) *Multifaceted behavior of a doubly reduced arylborane in B–H-bond activation and hydroboration catalysis*. Chemical Science . ISSN 2041-6520.

Downloaded from

<https://kar.kent.ac.uk/98799/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1039/D2SC05518J>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. E. Prey, C. Herok, F. Fantuzzi, M. Bolte, H. Lerner, B. Engels and M. Wagner, *Chem. Sci.*, 2022, DOI: 10.1039/D2SC05518J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Multifaceted behavior of a doubly reduced arylborane in B–H bond activation and hydroboration catalysis

Sven E. Prey,^{a,†} Christoph Herok,^{b,†} Felipe Fantuzzi,^{b,c} Michael Bolte,^a Hans-Wolfram Lerner,^a Bernd Engels^{b*} and Matthias Wagner^{a*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Alkali-metal salts of 9,10-dimethyl-9,10-dihydro-9,10-diboraanthracene ($M_2[DBA-Me_2]$; $M^+ = Li^+, Na^+, K^+$) activate the H–B bond of pinacolborane (HBpin) in THF already at room temperature. For $M^+ = Na^+, K^+$, the addition products $M_2[4]$ are formed, which contain one new H–B and one new B–Bpin bond; for $M^+ = Li^+$, the H^- ion is instantaneously transferred from the DBA-Me₂ unit to another equivalent of HBpin to afford Li[5]. Although Li[5] might commonly be considered a [Bpin]–adduct of neutral DBA-Me₂, it donates a [Bpin]⁺ cation to Li[SiPh₃], generating the silyl borane Ph₃Si–Bpin; Li₂[DBA-Me₂] with an aromatic central B₂C₄ ring acts as the leaving group. Furthermore, Li₂[DBA-Me₂] catalyzes the hydroboration of various unsaturated substrates with HBpin in THF. Quantum-chemical calculations complemented by *in situ* NMR spectroscopy revealed two different mechanistic scenarios that are governed by the steric demand of the substrate used: In the case of the bulky Ph(H)C=NtBu, the reaction requires elevated temperatures of 100 °C, starts with H–Bpin activation which subsequently generates Li[BH₄], so that the mechanism eventually turns into “hidden borohydride catalysis”. Ph(H)C=NPh, Ph₂C=O, Ph₂C=CH₂, and *i*PrN=C=N*i*Pr undergo hydroboration already at room temperature. Here, the active hydroboration catalyst is the [4+2] cycloadduct between the respective substrate and Li₂[DBA-Me₂]: In the key step, attack of HBpin on the bridging unit opens the bicyclo[2.2.2]octadiene scaffold and gives the activated HBpin adduct of the Lewis-basic moiety that was previously coordinated to the DBA-B atom.

Introduction

The activation of chemical bonds by main-group compounds is not only conceptually appealing, but also holds great application potential. A prominent class of p-block catalysts is that of Frustrated Lewis Pairs (FLPs), which contain suitable combinations of sterically encumbered Lewis acids (LA) and bases (LB). Together, these functional units contribute the vacant orbital and electron lone pair which, in classical catalysis, are provided by one single transition metal center with partially filled d orbitals.^{1,2} More recently, alternative systems based on doubly boron-doped (hetero)arenes have been reported.^{3,4} Common to all of them is a central six-membered ring featuring an aromatic π -electron system and two mutually cooperating B atoms integrated therein at opposite positions. As an example, Kinjo et al. disclosed that the 1,3,2,5-diazadiborinine **A** (Fig. 1) adds the single bond of H₃C–OSO₂CF₃, the double bonds of alkenes and carbonyls, as

well as the triple bonds of alkynes across its B sites. As mode of action, Kinjo postulated that **A** has a B(I)/B(III) mixed-valence character,^{5–7} which would render it formally analogous to an LB/LA FLP. A related 1,4,2,5-diazadiborinine activates H–H, B–H, Si–H, and P–H bonds. Here, it was proposed that the two chemically equivalent B atoms “act as both nucleophilic and electrophilic centers, demonstrating ambiphilic nature”.^{8,9} Catalytic cycles based on these diazadiborinines have not yet been described, with one notable exception: *in situ*-generated [4+2] cycloadducts of **A** and Ph(Me)C=O or H₂C=CH₂ act as “electrostatic catalysts” to promote hydroboration of carbonyl compounds with pinacolborane (HBpin; see Fig. 3 below).⁷ Wagner and coworkers have introduced 9,10-dihydro-9,10-diboraanthracene (DBA) dianions [DBA-R₂]²⁻ into catalysis (Fig. 1; R = H, Me). These species have indistinguishable, sterically accessible B atoms along with high-lying HOMO-energy levels and are therefore particularly reactive. Apart from reacting with H–H and C–H bonds, [DBA-R₂]²⁻ dianions undergo facile [4+2] cycloaddition with various unsaturated organic molecules.^{10–14}

The HOMO and LUMO of [DBA-R₂]²⁻ have the same local symmetries about the B atoms as, respectively, the LUMO and HOMO of H₂, and quantum-chemical calculations suggest a concerted, transition metal-like (TM-like) bond-cleavage pathway (Fig. 1).¹¹ H₂ activation by [DBA-R₂]²⁻ is at the core of two recently developed catalytic hydrogenation- and H–transfer cycles.¹³ In addition, it has been shown that [DBA-R₂]²⁻

^a Institut für Anorganische und Analytische Chemie, Goethe-Universität Frankfurt, D-60438 Frankfurt am Main, Germany

^b Institut für Physikalische und Theoretische Chemie, Julius-Maximilians-Universität Würzburg, D-97074 Würzburg, Germany

^c School of Chemistry and Forensic Science, University of Kent, Canterbury CT2 7NH, United Kingdom

† These authors contributed equally.

Electronic Supplementary Information (ESI) available: Synthetic procedures, NMR spectra, X-ray crystallographic data and computational details. CCDC 2207512–2207520. See DOI: 10.1039/x0xx00000x






Fig. 1 Top: 1,3,2,5-Diazadiborinine **A** can be regarded as B(I)/B(III) mixed-valence system, which activates substrate molecules in a similar manner to LB/LA-FLPs. Bottom: The frontier orbital symmetries of the 9,10-dihydro-9,10-diboraanthracene (DBA) dianion $[\text{DBA-R}_2]^{2-}$ allow concerted transition metal-like (TM-like) reactions.

can catalyze the disproportionation of CO_2 to CO and $[\text{CO}_3]^{2-}$.^{12,15}

In a joint experimental and theoretical effort, we are herein unveiling the capacity of the $[\text{DBA-R}_2]^{2-}$ platform to activate B–H bonds and catalyze hydroborations. Besides our system, there are also numerous FLPs that catalyze hydroborations. Various modes of action have been discussed.¹⁶ The hydroboration reaction is therefore an ideal tool to put into context the behavior of $[\text{DBA-R}_2]^{2-}$ dianions with those of other main group catalysts and to learn more about the subtleties of subvalent boron species.

Three key results are disclosed: (i) HBpin activation by $[\text{DBA-R}_2]^{2-}$ is a means of forming new B–B bonds. (ii) Depending on the steric bulk of the substrate, the hydroboration mechanism differs between “hidden borohydride catalysis”¹⁷ and actual DBA-driven catalysis. (iii) To gain accurate theoretical insights into the mechanisms, one must explicitly include the counter cations and their coordinating solvent molecules in the calculations, since simple continuum approaches are too imprecise.

Results and discussion


All reactions were performed in (deuterated) tetrahydrofuran (THF, THF- d_8).

Activation of HBpin by $\text{M}_2[\text{DBA-R}_2]$

Pinacolborane (HBpin) was selected as the hydroboration reagent for the following reasons: (i) in contrast to, e.g., B_2H_6 or $(9\text{-BBN})_2$ (BBN = 9-borabicyclo[3.3.1]nonane), HBpin is strictly monomeric¹⁸ and does not form solvent adducts in THF, which facilitates the theoretical assessment of its

reactivity. (ii) The alkyl boronic ester products ($\text{R}'\text{-Bpin}$) are comparatively stable to air and moisture, easy to purify, and have a wide range of applications.¹⁹ The aforementioned assets are the result of pronounced O=B π donation, decreasing the electrophilicity of the B atom. On the other hand, less Lewis-acidic boranes are less prone to spontaneous addition to unsaturated substrates, so that a catalyst is often required. Thus, hydroboration reactions with HBpin provide the ideal setting to further explore the scope of $\text{M}_2[\text{DBA-R}_2]$ -mediated reactions.

Our investigations into the activation of HBpin by $\text{M}_2[\text{DBA-R}_2]$ were carried out with $\text{M}^+ = \text{Li}^+ - \text{K}^+$ as counter cations and $\text{R} = \text{H}$, Me as B-bonded substituents ($\text{Li}_2[\mathbf{1}] - \text{K}_2[\mathbf{1}]$, $\text{Li}_2[\mathbf{2}] - \text{K}_2[\mathbf{2}]$; Scheme 1). In all cases, the reaction with HBpin was already instantaneous at room temperature, as judged by the rapid fading of the intensely colored $\text{M}_2[\text{R}_2\text{-DBA}]$ solutions after addition of the borane ($\text{M}^+ = \text{Li}^+$: red; $\text{M}^+ = \text{Na}^+$, K^+ : green). According to *in situ* NMR spectroscopy, 1 equiv. of HBpin was sufficient to quantitatively convert $\text{Na}_2[\mathbf{1}]$, $\text{K}_2[\mathbf{1}]$, and $\text{K}_2[\mathbf{2}]$ to $\text{Na}_2[\mathbf{3}]$, $\text{K}_2[\mathbf{3}]$, and $\text{K}_2[\mathbf{4}]$, respectively, featuring newly formed B–H and B–Bpin bonds together with two tetracoordinate B centers on their DBA cores (type-I structures; Scheme 1). In contrast, equimolar mixtures of $\text{Li}_2[\mathbf{1}]$ or $\text{Na}_2[\mathbf{2}]$ with HBpin contained $\text{Li}_2[\mathbf{3}]$ or $\text{Na}_2[\mathbf{4}]$ and significant amounts of still unconsumed starting materials. Full conversion of the DBA-dianion salts to the corresponding type-I addition products required the use of excess HBpin. These $[\text{DBA-R}_2]^{2-}$ -reactivity trends are the same as those previously reported for H_2 activation:¹³ (i) M^+ ions with larger charge-radius ratio have a higher tendency to form contact-ion pairs with $[\text{DBA-R}_2]^{2-}$ in solution, which leads to the reactivity order $\text{Li}_2[\text{DBA-R}_2] <$



Scheme 1 Addition of pinacolborane (HBpin) across the two B atoms of 9,10-dihydro-9,10-diboraanthracene (DBA) dianions to form B–B bonds. Depending on the nature of the counter cations, the dianionic type-I structures compete with monoanionic type-II structures, which are formed by transfer of the H^- ligand onto a second equivalent of HBpin. (i) 1–5 equiv. HBpin (see SI), THF- d_8 , room temperature; (ii) prolonged storage in 1,2-dimethoxyethane (DME) during crystallization experiments.



$\text{Na}_2[\text{DBA-R}_2] < \text{K}_2[\text{DBA-R}_2]$.¹¹ (ii) Smaller B-bonded substituents R impede substrate access to $[\text{DBA-R}_2]^{2-}$ less than bulkier groups, resulting in the reactivity order $[\mathbf{2}]^{2-} < [\mathbf{1}]^{2-}$.

The accordingly least reactive $\text{Li}_2[\mathbf{2}]$ is indeed a peculiarity, since a type-I product (putative $\text{Li}_2[\mathbf{4}]$) could not even be detected as an intermediate. Rather, the monoanion salt $\text{Li}[\mathbf{5}]$ was formed without H^- ligand at the DBA moiety (type-II structure; Scheme 1). Again, an excess of HBpin was necessary to enforce quantitative transformation of $\text{Li}_2[\mathbf{2}]$ to $\text{Li}[\mathbf{5}]$, which is accompanied by a formal release of LiH . According to quantum-chemical calculations, this would be an energetically unfavorable process^{20,21} and it is, therefore, reasonable to assume that excess HBpin acts as H^- scavenger. The NMR-spectroscopically observed formation of $[\text{BH}_4]^-$ and B_2pin_3 ²² supports this assumption, since H^- is known to induce a corresponding decay of HBpin.¹⁷ In summary, we propose that $\text{Li}_2[\mathbf{2}]$ and HBpin are in a dynamic addition-elimination equilibrium, which is shifted towards $\text{Li}[\mathbf{5}]$ formation by H^- transfer to HBpin and the subsequent irreversible decomposition of the resulting $[\text{H}_2\text{Bpin}]^-$ adduct. Consistent with this view, addition of LiH to a THF- d_8 solution of $\text{Li}[\mathbf{5}]$ leads back to $\text{Li}_2[\mathbf{2}]$ and HBpin (NMR-spectroscopic control).

The characteristic NMR data of type-I/II structures are exemplarily discussed with reference to the $\text{Na}_2[\mathbf{4}]/\text{Li}[\mathbf{5}]$ couple. The ¹¹B NMR spectrum of $\text{Na}_2[\mathbf{4}]$ contains three signals. Two of them appear in the chemical shift range of tetracoordinate B nuclei [−22.0 ppm (BBpin), −17.4 ppm (d, $^1J_{\text{BH}} = 68$ Hz, BH)] and the third is characteristic of a tricoordinate B center [42.6 ppm (broad, Bpin)].²³ The $\delta(^{11}\text{B})$ values of $\text{Li}[\mathbf{5}]$ prove the presence of one tetra- and two tricoordinate B atoms [−19.7 ppm (BBpin), 41.6 ppm (Bpin), 61.8 ppm (BC₃)]. Accordingly, only the ¹H{¹¹B} NMR spectrum of $\text{Na}_2[\mathbf{4}]$ shows the signal of a B-bonded H atom (2.14 ppm). The ¹H integral values in the spectra of both $\text{Na}_2[\mathbf{4}]$ and $\text{Li}[\mathbf{5}]$ are consistent with the presence of one Bpin substituent in each of these molecules. In line with the proposed average C_s symmetry of $\text{Na}_2[\mathbf{4}]$ and $\text{Li}[\mathbf{5}]$ in solution, their ¹³C{¹H} NMR spectra exhibit only six resonances in the aromatic region. Single crystals of the type-II compound $[\text{Li}(\text{thf})_2][\mathbf{5}]$ were grown by gas-phase diffusion of *n*-hexane into a C₆H₆ solution of thf-solvated $\text{Li}[\mathbf{5}]$ (Fig. 2). Attempts at the crystallization also of type-I compounds gave specimens suitable for X-ray diffraction only from 1,2-dimethoxyethane (DME) solutions of thf-solvated $\text{Na}_2[\mathbf{4}]$ (room temperature, 1–2 d). Since X-ray





Fig. 2 X-ray-crystallographically derived solid-state structures of the solvates $[\text{Li}(\text{thf})_2][\mathbf{5}]$ and $[\text{Na}(\text{thf})_2][\mathbf{5}]$, featuring one B–Bpin bond. C-bonded H atoms are omitted for clarity; the thf ligands are simplified as wireframes; Li: pink, B: green, C: black, O: red, Na: yellow spheres.

analysis revealed that the crystals consisted of $[\text{Na}(\text{thf})_2][\mathbf{5}]$, as opposed to $\text{Na}_2[\mathbf{4}]$ (Fig. 2), it appears that $\text{Na}_2[\mathbf{4}]$ is also susceptible to formal NaH elimination (triggered by residual HBpin under the crystallization conditions). The DBA moiety of $[\text{Na}(\text{thf})_2][\mathbf{5}]$ has one sp^3 -hybridized [B(1)] and one sp^2 -hybridized B atom [B(2); $\Sigma(\angle_{\text{CBC}}) = 360.0^\circ$]. At B(1), the Bpin substituent is attached in an axial position and with a bond length of B(1)–B(3) = 1.731(9) Å.²⁴ The $[\text{Na}(\text{thf})_2]^+$ cation is coordinated by an O atom belonging to Bpin and by the centroid of a phenylene ring, creating a contact-ion pair. The solid-state structure of $[\text{Li}(\text{thf})_2][\mathbf{5}]$ does not merit further discussion, given that it differs from that of $[\text{Na}(\text{thf})_2][\mathbf{5}]$ mainly in details of cation-anion association.²⁰


Reactivities of $\text{M}_2[\text{DBA-R}_2]/\text{HBpin}$ addition products

Hydridoborate ions $[\text{R}_3\text{B-H}]^-$ and negatively charged adducts $[\text{R}_3\text{B-Bpin}]^-$ are widely used sources of H^- and $[\text{Bpin}]^-$ nucleophiles, respectively.^{25,26} Type-I compounds such as $\text{Na}_2[\mathbf{4}]$ provide both functionalities in the same molecule, which raises the question of whether H^- or $[\text{Bpin}]^-$ are preferentially transferred to, e.g., chlorosilanes as archetypal electrophiles. Treatment of freshly prepared $\text{Na}_2[\mathbf{4}]$ with 1 equiv. of Et_3SiCl resulted in selective H^- abstraction to form $\text{Na}[\mathbf{5}]$ and Et_3SiH , showing that H^- wins the competition [cf. abstraction of MH by HBpin during the formation of $\text{M}[\mathbf{5}]$ ($\text{M}^+ = \text{Li}^+, \text{Na}^+$)]. What about the possibility of $[\text{Bpin}]^-$ transfer in the absence of a competing H^- ion? We have found that $\text{Li}[\mathbf{5}]$ is inert to Et_3SiCl , Et_3SiBr , and MeI at room temperature for several days and therefore does not appear to be a $[\text{Bpin}]^-$ donor. To probe if a polarity-inverted reactivity is present, $\text{Li}[\mathbf{5}]$ was next combined with the silanide salt $\text{Li}[\text{SiPh}_3]$, which indeed gave pinB-SiPh_3 in a clean reaction.²⁸ Here, the highly delocalized byproduct $[\mathbf{2}]^{2-}$ turns out to be a good enough leaving group to promote the transfer of a $[\text{Bpin}]^+$ electrophile (Scheme 2). To confirm this remarkable result further, we also prepared the formal hydridoborate anion $\text{Li}[\mathbf{6}]$ through H^- -adduct formation between the Lewis acid $\mathbf{2}$ and the H^- source LiH and then removed H^+ by deprotonation of $\text{Li}[\mathbf{6}]$ with $\text{Li}[\text{C}(\text{SiMe}_3)_3]$ ²⁹ (Scheme 2).²⁰ Again, $[\mathbf{2}]^{2-}$ is liberated, accompanied by the formation of $\text{HC}(\text{SiMe}_3)_3$, which renders $\text{Li}[\mathbf{6}]$ a model system of $\text{Li}[\mathbf{5}]$ in which all reactive parts are stripped down to their absolute essence.³⁰



Scheme 2 Reactions of $\text{Li}[\mathbf{5}]$ and $\text{Li}[\mathbf{6}]$ with an Si-centered nucleophile and a C-centered Brønsted base to demonstrate transfer of $[\text{Bpin}]^+$ and H^+ along with the suitability of $[\mathbf{2}]^{2-}$ as leaving group. (i) THF- d_6 , room temperature, overnight; (ii) THF- d_6 , 115 °C, 35 h.





Scheme 3 a) Substituent-redistribution reaction at $\text{Na}_2[3]$ producing $\text{Na}_2[7]$ with two Bpin bonds. Substituent-redistribution reaction at $\text{Li}_2[9]$ leading to the pair of more symmetric compounds $\text{Li}_2[10]/\text{Li}_2[11]$. b) X-ray-crystallographically derived solid-state structure of the solvate $[\text{Na}(\text{dme})_2]_2[7]$, featuring two B–Bpin bonds. C-bonded H atoms are omitted for clarity; the dme ligands are shown as wireframes; B: green, C: black, O: red, Na: yellow spheres. (i) DME, room temperature, prolonged storage during crystallization experiments; (ii) THF- d_6 , room temperature.

The reactions between $\text{M}_2[1]/\text{M}_2[2]$ and HBpin described up to this point are not only relevant for H–B-bond activation, but also represent B–B-bond formation reactions off the beaten track. The latter aspect gains additional weight because $\text{Na}_2[3]$ (one B–B bond) is prone to $\text{H}^-/[\text{Bpin}]^-$ scrambling, thereby generating a DBA with two B–B bonds: When a DME solution of $\text{Na}_2[3]$ was stored at room temperature, $[\text{Na}(\text{dme})_2]_2[7]$ precipitated in single-crystalline form (Scheme 3).

The dianion $[7]^{2-}$ can be regarded as Lewis pair $\{1 \cdot 2[\text{Bpin}]^-\}$; the necessary byproduct $\text{Na}_2[8]$ ^{11,13} remained in the mother liquor and was detected by NMR spectroscopy after workup.

The ^1H , ^{11}B , and $^{13}\text{C}\{^1\text{H}\}$ NMR spectroscopic characterization of re-dissolved $[\text{Na}(\text{dme})_2]_2[7]$ was in line with an average C_{2v} or C_{2h} symmetry but allowed no conclusion regarding a mutual *cis* or *trans* orientation of the two Bpin ligands. The assignment of $[\text{Na}(\text{dme})_2]_2[7]$ as a C_1 -symmetric *trans* complex ($\cong C_{2h}$ in solution) was finally achieved by X-ray crystallography ($\text{B}(1) - \text{B}(2) = 1.711(3)$, Scheme 3). Note that the attempted synthesis of *cis*- $\text{Na}_2[7]$ from $\text{Na}_2[1]$ and B_2pin_2 failed, because the two compounds do not react with each other. Also the investigation of the transformation $2 \text{Na}_2[3] \rightarrow \text{Na}_2[7] + \text{Na}_2[8]$ was complemented by a model reaction, in which a Me group mimicked the Bpin group of $\text{Na}_2[3]$ (Scheme 3): Addition of MeLi (1 equiv.) to $\text{Li}[6]$ did not lead to an NMR-spectroscopically detectable diadduct $\text{Li}_2[9]$, but rather to an equimolar mixture of the more symmetric $\text{Li}_2[10]$ and $\text{Li}_2[11]$ ¹³ scrambling products,²⁰ the former corresponding to $\text{Na}_2[7]$.

Taken together, a clear picture of the reactivity trends of the DBA substituents emerges from the model studies, which is of immediate importance with respect to the transfer of H or

Bpin fragments onto substrates in the course of hydroboration reactions: a substituent R residing on the tetracoordinate B atom of a $\text{B}(\text{sp}^2), \text{B}(\text{sp}^3)$ -DBA can be removed as an R^+ fragment because the formerly B–R-bonding electron pair subsequently becomes part of a Clar's sextet³¹ within the central B_2C_4 ring. However, in a $\text{B}(\text{sp}^3), \text{B}(\text{sp}^3)$ -DBA, this energetically favorable electron delocalization after R^+ transfer is blocked by the second tetracoordinate B atom, which is why the same R now possesses mainly R^- character.

Hydroboration reactions

The substrates under investigation contained less polar ($\text{C}=\text{C}$) as well as more polar ($\text{C}=\text{N}$, $\text{C}=\text{O}$) double bonds and possessed different steric demands (e.g., $\text{Ph}(\text{H})\text{C}=\text{NPh} < \text{Ph}(\text{H})\text{C}=\text{N}t\text{Bu}$). Hydroborations were regularly performed with a loading of 25 mol% $\text{Li}_2[2]$ to facilitate the NMR-spectroscopic detection of reaction intermediates, side products, and byproducts; in the selected case of $\text{Ph}_2\text{C}=\text{O}$, it was confirmed that a $\text{Li}_2[2]$ loading of 5 mol% is sufficient for full conversion of the starting materials and thus the reactions are truly catalytic in the DBA. To ensure full comparability, we always added the HBpin to a mixture of freshly prepared $\text{Li}_2[2]$ and the unsaturated substrate.³²

Two fundamentally different reaction mechanisms were found to be operative under these conditions.³³ Which of the two comes into play depends decisively on the steric requirements of the substrate. Only in the case of the bulkiest substrate, the imine $\text{Ph}(\text{H})\text{C}=\text{N}t\text{Bu}$, does HBpin activation by $\text{Li}_2[2]$ initiate the transformation; the actual catalytic cycle is a textbook example of "hidden borohydride catalysis".¹⁷ In all other cases, [4+2] cycloaddition of the substrates' double bonds occurs prior to HBpin addition; subsequent hydroboration is then catalyzed by the cycloadducts.

Mechanism 1: hidden borohydride catalysis. NMR spectra recorded on a freshly prepared mixture of $\text{Li}_2[2]$, $\text{Ph}(\text{H})\text{C}=\text{N}t\text{Bu}$, and HBpin showed the characteristic resonances of the above-mentioned HBpin-activation product $\text{Li}[5]$, together with the signals of $\text{Li}[\text{BH}_4]$ and B_2pin_3 . The resonances of the imine were still prominently visible, and the spectrum did not indicate transformation of this starting material. The situation remained unchanged for several hours at room temperature. Yet, heating the sample to 100 °C for 40 h led to 85% conversion of the imine, but it took another 50 h at 100 °C to drive the reaction to completion. The NMR data of the borylamine primary product agreed well with literature data;³⁴ after *in situ* hydrolysis, we observed the resonances of the free amine $\text{Ph}(\text{H})_2\text{C}-\text{N}(\text{H})t\text{Bu}$.¹³ We took the initial appearance of the $[\text{BH}_4]^-$ ion as a warning signal that the present case might be an example of hidden catalysis by the system $\text{BH}_3 \cdot \text{thf}/[\text{BH}_4]^-$.¹⁷ Indeed, preliminary quantum-chemical calculations indicated that such a process might be at play (see below for an in-depth theoretical treatment of the reaction mechanism). To substantiate the assumption of hidden borohydride catalysis further, we adapted a test reaction recommended by Thomas and coworkers¹⁷ and repeated the experiment in the presence of *N,N,N',N'*-tetramethylethylenediamine (TMEDA),




$\text{R}^1\text{C}(\text{E})=\text{R}^2 \xrightarrow[\text{THF-}d_8]{25 \text{ mol\% } \text{Li}_2[2], 1.4 \text{ eq } \text{HBpin}} \text{R}^1\text{C}(\text{E})\text{Bpin}-\text{R}^2$				
substrate	$T / ^\circ\text{C}^{[a]}$	$t / \text{h}^{[a]}$	conversion (NMR) ^[a]	product after aq. workup; yield
	r. t. (100)	4 (35)	100% (29%)	71%
	r. t. (50)	< 0.1 (3)	100% (40%)	64%
	r. t. (100)	21 (18)	100% (0%)	91%
	r. t. (r. t.)	< 0.1 (24)	100% (54%)	[b]

Table 1 Hydroboration of unsaturated substrates proceeding via DBA-cycloadduct catalysis. [a] Numbers in brackets refer to the blind test in the absence of $\text{Li}_2[2]$; [b] Pure samples of $i\text{PrN}=\text{C}(\text{H})-\text{N}(\text{Bpin})\text{Pr}$ were only obtained by sublimation from the reaction mixture, albeit with substantial decrease of the yield.

which is supposed to act as a BH_3 scavenger. Subsequently, we detected the diagnostic signal of $(\text{BH}_3)_2 \cdot \text{tmeda}^{35}$ and noted a significantly slower reaction. All in all, this confirmed our working hypothesis that the hydroboration of $\text{Ph}(\text{H})\text{C}=\text{NtBu}$ by HBpin is only initiated by $\text{Li}_2[2]$, but catalyzed by $\text{BH}_3 \cdot \text{thf}/[\text{BH}_4]^-$.

Mechanism II: DBA-cycloadduct catalysis. Hydroborations of $\text{Ph}(\text{H})\text{C}=\text{NPh}$, $\text{Ph}_2\text{C}=\text{O}$, and $\text{Ph}_2\text{C}=\text{CH}_2$ with HBpin in the presence of $\text{Li}_2[2]$ proceeded quantitatively already at room temperature (Table 1). We found that the key elements of each reaction scenario are the same for all three substrates. It is therefore sufficient to discuss the experimental facts using $\text{Ph}(\text{H})\text{C}=\text{NPh}$ as an example. Imine $\text{Ph}(\text{H})\text{C}=\text{NPh}$ undergoes an instantaneous and quantitative [4+2] cycloaddition reaction with $\text{Li}_2[2]$ to afford the dianionic bicyclo[2.2.2]octadiene derivative $\text{Li}_2[12]$ (Scheme 4). As a distinct difference, the corresponding cycloaddition with the bulkier $\text{Ph}(\text{H})\text{C}=\text{NtBu}$ is very slow at room temperature (and underlies a dynamic



Scheme 4 Crossover experiment to demonstrate that hydroboration of $\text{Ph}(\text{H})\text{C}=\text{NPh}$ with HBpin can be mediated not only by $\text{Li}_2[2]$ (actual catalyst: $\text{Li}_2[12]$), but also by pre-formed $\text{Li}_2[13]$ or $\text{Li}_2[14]$ without generating $\text{Ph}_2(\text{H})\text{C}-\text{OBpin}$ or $\text{H}_3\text{C}-\text{CH}_2\text{Bpin}$ as crossover side product, respectively ($\text{THF-}d_8$, room temperature). Note that the high catalyst loadings were used for the sole purpose of being able to detect any catalyst degradation or trace formation of crossover products by NMR spectroscopy.




Fig. 3 Comparison of Lewis structures and computed charge densities of a) Kinjo's bicyclic catalyst **B** and b) compounds $[14]^{2-}$ and $\text{Li}_2[14]$.

addition-elimination equilibrium at 100°C).¹³ Consequently, under the prevailing reaction conditions, only in the case of $\text{Ph}(\text{H})\text{C}=\text{NtBu}$ is free $\text{Li}_2[2]$ still available for initial HBpin activation, while in the other cases the actual active species must be a different one. Thus, the different steric demands of $\text{Ph}(\text{H})\text{C}=\text{NtBu}$ and, e.g., $\text{Ph}(\text{H})\text{C}=\text{NPh}$ lead to a bifurcation of the reaction mechanism into Mechanism I and a new Mechanism II.

Probable candidate catalysts under Mechanism II would be the respective [4+2] cycloadducts. This assumption is supported by crossover experiments in which the hydroboration of $\text{Ph}(\text{H})\text{C}=\text{NPh}$ was efficiently mediated by pre-formed $\text{Li}_2[13]$ or $\text{Li}_2[14]$, obtained from $\text{Li}_2[2]$ and $\text{Ph}_2\text{C}=\text{O}$ or $\text{H}_2\text{C}=\text{CH}_2$, respectively. According to *in situ* NMR spectroscopy, the reactions furnished exclusively $\text{Ph}(\text{H})_2\text{C}-\text{N}(\text{Bpin})\text{Ph}$ (and no $\text{Ph}_2(\text{H})\text{C}-\text{OBpin}$ or $\text{H}_3\text{C}-\text{CH}_2\text{Bpin}$), leaving $\text{Li}_2[13]$ or $\text{Li}_2[14]$ intact (Scheme 4).

At first glance, the molecular scaffolds of $[12]^{2-}$, $[13]^{2-}$, and $[14]^{2-}$ may appear similar to Kinjo's bicyclic compound **B** (Fig. 3a), which he employed as “electrostatic catalyst” for the hydroboration of various aldehydes and ketones. Here, the key to activation of the $\text{H}-\text{Bpin}$ bond is believed to lie in electrostatic interactions between the negative partial charge on the borane's H atom and positively polarized regions in the binding pocket of **B**.⁷

However, decisive differences between Kinjo's and our catalysts should arise from (i) the presence of electronegative N and O atoms in **B**, which are (largely) absent in $[12]^{2-}$, $[13]^{2-}$, and $[14]^{2-}$, and (ii) the fact that **B** is a neutral compound whereas our catalysts are dianion salts. A comparison of the computed charge densities of **B**, $[14]^{2-}$, and $\text{Li}_2[14]$ confirms this view (Fig. 3b): while positively polarized regions are indeed found in the binding pocket of **B**, they are completely missing in $[14]^{2-}$. If the Li^+ counter cations are included in the




charge-density calculations, positively polarized areas also emerge for $\text{Li}_2[\mathbf{14}]$ but remain largely associated with Li^+ . Although the charge distribution in $\text{Li}_2[\mathbf{14}]$ does not directly correspond to the charge distribution in the binding pocket of **B**, it would nevertheless be conceivable that an interaction between the negatively polarized borane-H atom and one of the Li^+ cations could still activate the H–Bpin bond.³⁶ However, our calculations show that Li^+ and HBpin preferentially interact via an O atom (and not the H atom) of the borane (Fig. S93).^{37,38} Kinjo's mechanism is therefore not applicable to our case.

What is a plausible alternative? Although a dynamic equilibrium $\text{Li}_2[\mathbf{C}] \rightleftharpoons \text{Li}_2[\mathbf{2}] + \text{substrate}$ does not exist at room temperature for the substrates $\text{Ph}(\text{H})\text{C}=\text{NPh}$, $\text{Ph}_2\text{C}=\text{O}$, and $\text{Ph}_2\text{C}=\text{CH}_2$, it is conceivable that HBpin attack induces the reversible cleavage of one E–B(bridgehead) bond of $\text{Li}_2[\mathbf{C}]$ (Scheme 5; E = CH_2 , NPh, O). In other words, there may be a competition between the B atoms of DBA and HBpin for the same E^- donor. Precedence exists in the form of the acetone/ $\text{Na}_2[\mathbf{2}]$ cycloadduct $\text{Na}_2[\mathbf{15}]$, which straightforwardly inserts CO_2 into its O–B bond to afford $\text{Na}_2[\mathbf{16}]$ (Scheme 5).¹² In the putative negatively charged adduct $[\mathbf{D}]^{2-}$, the electronic situation at the H–B bond should be comparable to the case of, e.g., $[\text{BH}_4]^-$. Thus, hydride transfer from $[\mathbf{D}]^{2-}$ to the unsaturated substrate, followed by “[Bpin]⁺” transfer, would yield the respective hydroboration product and regenerate the catalyst $\text{Li}_2[\mathbf{C}]$. Like Mechanism I, the essence of Mechanism II has also been confirmed by quantum-chemical calculations (*vide infra*).

Switch between Mechanisms I and II in the double hydroboration of $i\text{PrN}=\text{C}=\text{N}i\text{Pr}$. The carbodiimide $i\text{PrN}=\text{C}=\text{N}i\text{Pr}$ has two C=N bonds and can therefore be singly or doubly hydroborated. So far, no catalyst has been described to promote both reaction scenarios.^{39–43} We now report that $\text{Li}_2[\mathbf{2}]$ can catalyze the monohydroboration of $i\text{PrN}=\text{C}=\text{N}i\text{Pr}$ to give $i\text{PrN}=\text{C}(\text{H})-\text{N}(\text{Bpin})i\text{Pr}$ (room temperature, instantaneous). The reaction is selective even in the presence of 2.4 equiv. of HBpin and proceeds via Mechanism II (the moderate steric bulk of the heterocumulene allows formation of its [4+2] cycloadduct, which has been fully characterized).²⁰ Upon heating the sample to 100 °C (23 h), a second hydroboration furnishes $i\text{Pr}(\text{pinB})\text{N}-\text{CH}_2-\text{N}(\text{Bpin})i\text{Pr}$. Due to the larger steric bulk of $i\text{PrN}=\text{C}(\text{H})-\text{N}(\text{Bpin})i\text{Pr}$ compared to $i\text{PrN}=\text{C}=\text{N}i\text{Pr}$, Mechanism I is operative in the second step (the $[\text{BH}_4]^-$ ion was detected by *in situ* NMR spectroscopy).

Quantum Chemical Calculations

Technical details. Geometry optimizations and Hessian calculations were performed at the $\omega\text{B97XD}/6\text{-31+G}(\text{d,p})^{44-46}$ level of theory including implicit solvation by the solvent model based on density (SMD).⁴⁷ Our analysis showed that explicit treatment of solvent molecules is crucial to obtain a correct description of entropy contributions along the reaction path as well as reliable kinetic barrier heights. Therefore, the optimal solvent coordination number for each intermediate was



Scheme 5 Conceptual relationship between H–Bpin activation by $\text{Li}_2[\mathbf{C}]$ and CO_2 activation by $\text{Na}_2[\mathbf{15}]$ ($\text{R}_2\text{C}=\text{E} = \text{H}_2\text{C}=\text{CH}_2$, $\text{Ph}_2\text{C}=\text{CH}_2$, $\text{Ph}(\text{H})\text{C}=\text{NPh}$, $\text{Ph}_2\text{C}=\text{O}$)

determined by free energy calculations (for details, see Fig. S94). Unless otherwise denoted, optimized geometries were confirmed to be the desired minimum-energy structures or transition states by vibrational frequency analysis. Single-point calculations were performed at the SMD/ $\omega\text{B97XD}/6\text{-311++G}(\text{d,p})$ level (solvent: THF; $\epsilon = 7.4257$). All free-energy values were calculated for the corresponding experimental temperature and included a concentration correction^{48,49} that accounts for the change in standard states going from gas phase to condensed phase. All calculations were performed in Gaussian 16, Revision A.03.⁵⁰

Computational characterization of hydroboration Mechanism I.

Since well-explored hidden borohydride catalysis plays a prominent role in Mechanism I while our primary interest lies in the characterization of DBA-catalyzed hydroboration reactions, we limited the investigation of Mechanism I to key intermediates and did not calculate kinetic barriers. In order to determine the reaction mechanism, a variety of possible reaction paths was considered (at the experimental temperature of 100 °C).

Since our above-mentioned experimental results exclude direct H^- transfer from $\text{Li}_2[\mathbf{4}]$ to $\text{Ph}(\text{H})\text{C}=\text{NtBu}$, a corresponding reaction mechanism was not considered theoretically. Rather, inspired by experiments of Clark and coworkers,⁵¹ we presumed that HBpin would be able to take up the H^- ion provided by $\text{Li}_2[\mathbf{4}]$, forming $\text{Li}[\text{H}_2\text{Bpin}]$, analogous to Clark's reactive species $[\text{tBuO}(\text{H})\text{Bpin}]^-$. HBpin could thereby act as H^- shuttle to $\text{Ph}(\text{H})\text{C}=\text{NtBu}$. Indeed, this approach led to an overall exergonic reaction ($-24 \text{ kcal mol}^{-1}$) with the free energy of the highest-lying intermediate $\text{Li}[\text{H}_2\text{Bpin}]$ being $+13 \text{ kcal mol}^{-1}$ (Fig. S96).

Despite its thermodynamic feasibility, this mechanism fails to explain the experimentally observed formation of $[\text{BH}_4]^-$ and B_2pin_3 . To take these two species into account, we further investigated their formation from decomposition of $[\text{H}_2\text{Bpin}]^-$. We first calculated the complete reduction of 1 equiv. HBpin by 3 equiv. $\text{Li}_2[\mathbf{4}]$. The energetically costly formation of 1 equiv. $\text{Li}_2[\text{pin}]$ per equivalent of $\text{Li}[\text{BH}_4]$ generated renders this reaction exceedingly endergonic and thermodynamically out of scope ($+36 \text{ kcal mol}^{-1}$, Fig. S95; $\text{H}_2\text{pin} = \text{pinacol}$).



As an alternative approach, we propose a stepwise decomposition of $\text{Li}[\text{H}_2\text{Bpin}]$ via B–H/B–O σ -bond metathesis with 2 equiv. HBpin (Fig. S97). Formation of the final products $\text{Li}[\text{BH}_4]$ and B_2pin_3 is exergonic by $-21 \text{ kcal mol}^{-1}$, likely due to the isodesmic nature of this reaction and to the formation of two stable BO_3 motifs. Once formed, $\text{Li}[\text{BH}_4]$ takes on the role as the actual imine-reducing agent. On this basis, we can now establish a catalytic cycle for hydroboration of $\text{Ph}(\text{H})\text{C}=\text{NtBu}$ by HBpin that is fully consistent with all our experimental results (Fig. 4): In a first step, transition metal-like addition of HBpin to precatalyst $\text{Li}_2[\mathbf{2}]$ affords $\text{Li}_2[\mathbf{4}]$, the starting point of catalytic

Cycle I (Fig. 4a, right). While the newly formed B–B bond persists throughout all subsequent reaction steps, the π -ligand on $\text{Li}_2[\mathbf{4}]$ is transferred to a second HBpin molecule from solution. The free energy of the intermediates $\text{Li}[\mathbf{5}]$ and $\text{Li}[\text{H}_2\text{Bpin}]$ amounts to $+19 \text{ kcal mol}^{-1}$ when referenced to the starting point of the cycle, $\text{Li}_2[\mathbf{4}]$ (Fig. 4b, right). This energy penalty, which is equivalent to the most endergonic step of the reaction sequence, results from the loss of π conjugation in the H^- carrier $\text{Li}[\text{H}_2\text{Bpin}]$. As outlined above, decomposition of $\text{Li}[\text{H}_2\text{Bpin}]$ generates $\text{Li}[\text{BH}_4]$. In a slightly exergonic reaction,




Fig. 4 a) Right: Catalytic Cycle I (hydride shuttle: $\text{Li}[\text{H}_2\text{Bpin}]$). Left: Catalytic Cycle II, hidden catalysis (hydride shuttle: $\text{Li}[\text{BH}_4]$). b) Right: Free energy diagram for catalytic Cycle I. Left: Free energy diagram for hidden catalysis Cycle II including BH_3 , thf/ $\text{Li}[\text{BH}_4]$. Energies are referenced to the starting point of the respective cycle. Level of theory: SMD(solvent=THF)/ ω B97XD/6-311++G(d,p) from optimized structures at SMD(solvent=THF)/ ω B97XD/6-31+G(d,p). Explicit thf molecules are omitted for clarity.



the double bond of Ph(H)C=NtBu inserts into a B–H bond of $[\text{BH}_4]^-$, affording the amide-borane adduct **Int1**. The next step involves a ligand exchange at the negatively charged N atom of **Int1** through nucleophilic attack of **Int1** on HBpin from solution, which releases $\text{BH}_3\cdot\text{thf}$ and forms **Int2**.⁵² **Int2** possesses an activated H–Bpin bond (compare again Clark's intermediate) and thus represents the second highest thermodynamic barrier (+12 kcal mol⁻¹) of Cycle I (Fig. 4b, right). For the last reaction step, there are two possible options, which leads to a bifurcation of the catalytic cycle into Cycles I and II (Fig. 4a). Following Cycle I, Li[5] abstracts the H⁻ ligand from **Int2** in a notably exergonic reaction (–26 kcal mol⁻¹). As a result, the hydroboration product **P** is formed and Cycle I can start anew. The second possibility to produce **P** is H⁻ transfer from **Int2** to $\text{BH}_3\cdot\text{thf}$ (generated in the preceding step), thereby entering Cycle II (left panels of Fig. 4a-b). The driving force to **P** is significantly higher along Cycle II (–38 kcal mol⁻¹) than along Cycle I (–26 kcal mol⁻¹). Moreover, since Cycle II starts from Li[BH₄] rather than Li₂[4], it bypasses the formation of Li[5] and Li[H₂Bpin], the highest-lying intermediate of Cycle I. The steps between the formation of Li[BH₄] and the formation of **Int2** are equivalent in both cycles. Given that $\text{BH}_3\cdot\text{thf}$ has the highest H⁻ affinity of all species along cycles I and II (Table S8), it will take over the role as H⁻ shuttle as soon as it is available, effectively making the DBAs Li₂[4] and Li[5] obsolete. We, therefore, predict that only in the beginning the reaction proceeds with activation of HBpin by Li₂[2]. After a few cycles, the mechanism switches to hidden catalysis, where $\text{BH}_3\cdot\text{thf}$ is the catalytically active species. From this point on, Cycle II outcompetes Cycle I.

Computational characterization of hydroboration Mechanism II. As previously discussed, less sterically demanding substrates than Ph(H)C=NtBu readily form bicyclo[2.2.2]octadiene derivatives by [4+2] cycloaddition to Li₂[2] at room temperature. For the following reasons, the combination of Ph(H)C=NPh and Li₂[14] was chosen as representative model system for our theoretical study: (i) Li₂[14] should be the most challenging candidate to evaluate the feasibility of the key ring-opening step in our proposed catalytic mechanism, because, unlike bridging O or NR units, CH₂ fragments do not carry electron lone pairs as obvious sites of attack for incoming HBpin molecules (see Scheme 5). (ii) Li₂[14] possesses a symmetric scaffold so that only one kind of reactive center has to be considered.

Our proposed catalytic cycle for the Li₂[14]-mediated hydroboration of Ph(H)C=NPh is shown in Fig. 5; the corresponding free energies of the intermediates/transition states along the reaction path are depicted in Fig. 6. Similar to Mechanism I, the first step of Mechanism II is again activation of HBpin, but the actual mode is different: Instead of the previously observed transition metal-like cleavage of the H–Bpin bond, the borane is now nucleophilically attacked by a bridging CH₂ group of Li₂[14].⁵³

As a consequence, the tricyclic scaffold is opened, leading to formation of a tetracoordinate C–(H)Bpin unit and a tricoordinate DBA-B center (**Int3**; Fig. 6).




Fig. 5 Hydroboration Mechanism II for less sterically demanding substrates. The corresponding energy diagram is given in Fig. 6.

Int3 lies about +7 kcal mol⁻¹ above the reactants. The corresponding barrier (**TS1**) amounts to about +25 kcal mol⁻¹. The activated H⁻ ligand of the C–(H)Bpin moiety in **Int3** is subject to intramolecular H⁻ transfer to the tricoordinate DBA-B center. This H⁻ shift restores the π -conjugated BO₂ motif in the Bpin residue and an electron octet on the DBA-B atom to afford the stabilized intermediate **Int4**, which lies only about +1 kcal mol⁻¹ above the reactants. The corresponding barrier (**TS2**) is about +15 kcal mol⁻¹ higher in energy than the reactants, *i.e.* considerably lower than the barrier of the first step. **Int4** then acts as an H⁻ donor to the imine, generating the amide $[\text{Ph(H)}_2\text{C=NPh}]^-$. The barrier of this step is about +3 kcal mol⁻¹ and the resulting **Int5** is –18 kcal mol⁻¹ lower in energy than the reactants.⁵⁴ $[\text{Ph(H)}_2\text{C=NPh}]^-$ forms an adduct with the Bpin residue of **Int5**, affording **Int6** (–13 kcal mol⁻¹). The corresponding barrier lies –1 kcal mol⁻¹ below the reactants and +17 kcal mol⁻¹ above **Int5**. The latter barrier height would have to be overcome if the reaction energy of the **Int4** → **Int5** step is instantaneously dissipated into the solvent. If this is not the case, the barrier associated with **TS4** would in fact be lower. In any case, the barrier of the **Int5** → **Int6** step is at least 8 kcal mol⁻¹ lower than the rate-determining **TS1** (+25 kcal mol⁻¹). The catalytic cycle is completed by the restitution of the tricyclic framework of [14]²⁻ and the simultaneous release of the hydroboration product $\text{Ph(H)}_2\text{C=N(Bpin)Ph}$ (**P**). The barrier of this last step is about +3 kcal mol⁻¹ above the reactants (+16 kcal mol⁻¹ above the previous intermediate) while the exothermicity of the overall reaction is high (–29 kcal mol⁻¹).

As a final remark on key technical details of the calculations, we emphasize that the first step of the mechanism, *i.e.*,






Fig. 6 Free energy diagram of hydroboration Mechanism II. Orange: Transition states and intermediate including counterions and explicit solvent molecules. The **TS2** geometry was obtained via a relaxed scan (Fig. S100) and subsequent restricted optimization of the counterions and explicit THF molecules. **TS3** and **TS4** geometries were obtained via relaxed scans (Fig. S101 and Fig. S102, respectively). Level of theory: SMD(solvent=THF)/ ω B97XD/6-311++G(d,p) from optimized structures at SMD(solvent=THF)/ ω B97XD/6-31+G(d,p).

nucleophilic attack of the C₂ bridge on HBpin, possesses by far the highest barrier (**TS1**; Fig. 6). It is about +10 kcal mol⁻¹ higher than the second highest barrier (**TS2**) with respect to the reactants. Concerning the respective preceding intermediate, the difference is even larger (+25 kcal mol⁻¹ vs. +8 kcal mol⁻¹) since **Int3** lies about +7 kcal mol⁻¹ above the reactants.

Hence, we investigated the influence of the counterions and their first solvation shell on the performance of the catalyst for both steps in more detail. Rather than modeling the effects of the Li⁺ counter cations and explicit THF solvent molecules, we tested the consequences of a mere inclusion of continuum effects by the SMD model. We found the resulting barrier for the trigonal-bipyramidal transition state (**TS1**) to increase by +7 kcal mol⁻¹ to +32 kcal mol⁻¹, which is too high considering the experimental conditions. Furthermore, **TS2** is also destabilized by +4 kcal mol⁻¹ (Fig. S98). This effect can be rationalized by considering coordination of the Li⁺ cation via the O atom of the HBpin molecule. Electron density is thereby withdrawn from the adjacent B center, rendering it more electrophilic and, in turn, facilitating its attack on the electron-rich C₂ bridge of Li₂[14]. Furthermore, the tetracoordinate B center in the resulting **Int3** is also stabilized by a Li⁺...O interaction (Fig. S98). Since all other barriers are significantly lower than **TS1** and **TS2**, we calculated them – as well as the corresponding intermediates – without explicit counterions and THF molecules to reduce calculation cost.

Conclusions

We have successfully expanded the range of [DBA-Me₂]²⁻-catalyzed reactions to include hydroborations with pinacolborane (HBpin; DBA-Me₂ = 9,10-dimethyl-9,10-dihydro-9,10-diboraanthracene). In comparison to analogous hydrogenation reactions, distinct differences become apparent: The general entry step for [DBA-Me₂]²⁻-catalyzed hydrogenations is H–H-bond addition across the two B atoms of the doubly reduced arylborane. Thus, any competing [4+2] cycloaddition reaction between the unsaturated substrate (to be hydrogenated) and the [DBA-Me₂]²⁻ dianion leads to catalyst poisoning. Although H–Bpin-bond activation with [DBA-Me₂]²⁻ is also possible, it only plays a role in the hydroboration of sterically loaded substrates such as Ph(H)C=NtBu and proceeds via HBpin degradation to [BH₄]⁻ and B₂pin₃, providing an example of “hidden borohydride catalysis”. In all other cases where [4+2] cycloaddition to [DBA-Me₂]²⁻ is not hampered by the steric demands of the substrates (*e.g.*, Ph(H)C=NPh), the cycloadducts are not dead ends, but rather constitute the active hydroboration catalysts. Using the hydroboration of Ph(H)C=NPh with HBpin, mediated by the cycloadduct between [DBA-Me₂]²⁻ and H₂C=CH₂, as representative model reaction, we have unveiled the underlying reaction mechanism in a joint experimental and theoretical effort: The 1,2-ethanediyl-bridged tricyclic catalyst can be viewed as intramolecular B–C adduct, in which a Lewis-acidic tricoordinate DBA-B center protects a Lewis-basic [:CH₂–CH₂–B(Me)Ar₂]²⁻ fragment to create a stable resting state. In the presence of HBpin, the tricycle can reversibly open to form



an HBpin-alkyl adduct in which the H–Bpin bond is activated for H[−] transfer to the tricoordinate DBA-B center, from which H[−] migrates further to the polar Ph(H)C=NPh bond. The resulting amide [Ph(H)₂C–NPh][−] takes up [Bpin]⁺ from [pinB–CH₂–CH₂–B(Me)Ar₂][−] to form Ph(H)₂C–N(Bpin)Ph, thereby regenerating the catalyst and completing the catalytic cycle. According to thorough DFT calculations, our mechanistic proposal is energetically feasible under the experimentally applied reaction conditions (room temperature, THF-*d*₈, Li⁺ counter cations). Quantum-chemical calculations by Kinjo and coworkers have led them to explicitly rule out such a scenario for their related neutral bicyclo[2.2.2]octadienes, assembled from 1,3,2,5-diazadiborinines rather than [Me₂-DBA]^{2−} dianions.⁷ Instead, they postulated H–Bpin-bond activation through electrostatic interactions with positively polarized regions of the still intact tricyclic catalyst. This comparison clearly demonstrates the multifaceted character of doubly B-doped bicyclo[2.2.2]octadienes and shows how seemingly subtle modifications of their molecular scaffolds can significantly alter their catalytic properties. In the future, it will be interesting to explore what else this class of compounds can contribute to the field of main group catalysis.

Data availability

The datasets supporting this article have been uploaded as part of the supplementary information.

Author contributions

S. E. P. performed all preparative experiments. C. H. performed all calculations. M. B. performed the X-ray crystal structure analyses. F. F., H.-W. L, B. E., and M. W. supervised the project. The manuscript was written through contributions of all authors.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 G. Erker and D. W. Stephan, *Frustrated Lewis Pairs I & II*, Springer, Heidelberg, 2013.
- 2 A. R. Jupp and D. W. Stephan, *Trends Chem.*, 2019, **1**, 35–48.
- 3 S. E. Prey and M. Wagner, *Adv. Synth. Catal.*, 2021, **363**, 2290–2309.
- 4 Y. Su and R. Kinjo, *Chem. Soc. Rev.*, 2019, **48**, 3613–3659.
- 5 D. Wu, L. Kong, Y. Li, R. Ganguly and R. Kinjo, *Nat. Commun.*, 2015, **6**, 7340.
- 6 D. Wu, R. Ganguly, Y. Li, S. N. Hoo, H. Hirao and R. Kinjo, *Chem. Sci.*, 2015, **6**, 7150–7155.
- 7 D. Wu, R. Wang, Y. Li, R. Ganguly, H. Hirao and R. Kinjo, *Chem*, 2017, **3**, 134–151.
- 8 B. Wang, Y. Li, R. Ganguly, H. Hirao and R. Kinjo, *Nat. Commun.*, 2016, **7**, 11871.
- 9 B. Wang and R. Kinjo, *Tetrahedron*, 2018, **74**, 7273–7276.
- 10 A. Lorbach, M. Bolte, H.-W. Lerner and M. Wagner, *Organometallics*, 2010, **29**, 5762–5765. DOI: 10.1039/D2SC05518J
- 11 E. von Grotthuss, M. Diefenbach, M. Bolte, H.-W. Lerner, M. C. Holthausen and M. Wagner, *Angew. Chem. Int. Ed.*, 2016, **55**, 14067–14071.
- 12 E. von Grotthuss, S. E. Prey, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2018, **57**, 16491–16495.
- 13 E. von Grotthuss, S. E. Prey, M. Bolte, H.-W. Lerner and M. Wagner, *J. Am. Chem. Soc.*, 2019, **141**, 6082–6091.
- 14 E. von Grotthuss, F. Nawa, M. Bolte, H.-W. Lerner and M. Wagner, *Tetrahedron*, 2019, **75**, 26–30.
- 15 Harman et al. have reported a number of remarkable (catalytic) transformations mediated by Au complexes of bidentate diphosphine ligands with DBA backbones: (a) J. W. Taylor, A. McSkimming, M.-E. Moret and W. H. Harman, *Angew. Chem. Int. Ed.*, 2017, **56**, 10413–10417; (b) J. W. Taylor, A. McSkimming, L. A. Essex and W. H. Harman, *Chem. Sci.*, 2019, **10**, 9084–9090; (c) J. W. Taylor and W. H. Harman, *Chem. Commun.*, 2020, **56**, 4480–4483; (d) J. W. Taylor and W. H. Harman, *Chem. Commun.*, 2020, **56**, 13804–13807.
- 16 (a) M. A. Dureen, A. Lough, T. M. Gilbert and D. W. Stephan, *Chem. Commun.*, 2008, 4303–4305; (b) P. Eisenberger, A. M. Bailey and C. M. Crudden, *J. Am. Chem. Soc.*, 2012, **134**, 17384–17387; (c) D. W. Stephan, *Acc. Chem. Res.*, 2015, **48**, 306–316; (d) F.-G. Fontaine, M.-A. Courtemanche, M.-A. Légaré and É. Rochette, *Coord. Chem. Rev.*, 2017, **334**, 124–135; (e) P. Vasko, I. A. Zulkifly, M. Á. Fuentes, Z. Mo, J. Hicks, P. C. J. Kamer and S. Aldridge, *Chem. Eur. J.*, 2018, **24**, 10531–10540; (f) A. Falconnet, M. Magre, B. Maity, L. Cavallo and M. Rueping, *Angew. Chem. Int. Ed.*, 2019, **58**, 17567–17571; (g) Y. Lebedev, I. Polishchuk, B. Maity, M. Dinis Veloso Guerreiro, L. Cavallo and M. Rueping, *J. Am. Chem. Soc.*, 2019, **141**, 19415–19423; (h) D. W. Stephan, *Chem*, 2020, **6**, 1520–1526.
- 17 A. D. Bage, K. Nicholson, T. A. Hunt, T. Langer and S. P. Thomas, *ACS Catal.*, 2020, **10**, 13479–13486.
- 18 K. Wade, *Electron Deficient Compounds*, Springer, New York, 1971.
- 19 I. A. I. Mkhallid, J. H. Barnard, T. B. Marder, J. M. Murphy and J. F. Hartwig, *Chem. Rev.*, 2010, **110**, 890–931.
- 20 See the ESI for further details.
- 21 The free Gibbs energy for the reaction [Li(thf)₂]₂[4] + THF → [Li(thf)₂]₂[5] + [Li(thf)₃]H amounts to +27.4 kcal mol^{−1} at room temperature.
- 22 W. Clegg, A. J. Scott, C. Dai, G. Lesley, T. B. Marder, N. C. Norman and L. J. Farrugia, *Acta Crystallogr. Sect. C: Cryst. Struct. Commun.*, 1996, **52**, 2545–2547.
- 23 The ¹¹B NMR signal of the Bpin group in Na₂[4] (42.6 ppm; THF-*d*₈) may seem unusually downfield shifted, as the resonance of B₂pin₂ appears at 32.6 ppm (own measurement in THF-*d*₈). However, the signals of Bpin groups attached to anionic, tetracoordinate B centers appear in the same range as that of Na₂[4] (cf. K[B₂pin₂-OtBu] δ = 41.1; THF-*d*₈): S. Pietsch, E. C. Neeve, D. C. Apperley, R. Bertermann, F. Mo, D. Qiu, M. S. Cheung, L. Dang, J. Wang, U. Radius, Z. Lin, C. Kleeberg and T. B. Marder, *Chem. Eur. J.*, 2015, **21**, 7082–7099. The calculated ¹¹B NMR shifts of [Na(thf)₂]₂[4] are: δ = 45.8 (Bpin), −18.9 (BH), −23.3 (BBpin) (GIAO method at the SMD(solvent=THF)/ωB97XD/6-311++G(d,p) level).
- 24 Further B–Bpin bond lengths (Å) useful for comparison are (a) B₂pin₂ = 1.7041(15), (b) [K(18-c-6)][B₂pin₂OMe] = 1.753(2); [Me₄N][B₂pin₂F]·THF = 1.736(8), and (c) [Mg(HC((Me)CN(2,6-*i*Pr₂C₆H₃))₂)] [pinB–BPh₃] = 1.718(7): (a) C. Kleeberg, A. G. Crawford, A. S. Batsanov, P. Hodgkinson, D. C. Apperley, M. S. Cheung, Z. Lin and T. B. Marder, *J. Org. Chem.*, 2012, **77**, 785–789; (b) ref. 23; (c) A.-F.



- Pécharman, M. S. Hill, C. L. McMullin and M. F. Mahon, *Angew. Chem. Int. Ed.*, 2017, **56**, 16363–16366.
- 25 E. C. Neeve, S. J. Geier, I. A. I. Mkhali, S. A. Westcott and T. B. Marder, *Chem. Rev.*, 2016, **116**, 9091–9161.
- 26 A. B. Cuenca, R. Shishido, H. Ito and E. Fernández, *Chem. Soc. Rev.*, 2017, **46**, 415–430.
- 27 C. Kleeberg and C. Borner, *Eur. J. Inorg. Chem.*, 2013, **2013**, 2799–2806.
- 28 Li[SiPh₃] was chosen as the nucleophile, because its high steric demand disfavors the formation of a stable adduct with the tricoordinate DBA-B atom in Li[5]. Moreover, silylboranes such as Ph₃Si-Bpin have considerable application potential: (a) T. Ohmura and M. Sugimoto, *Bull. Chem. Soc. Jpn.*, 2009, **82**, 29–49; (b) M. Oestreich, E. Hartmann and M. Mewald, *Chem. Rev.*, 2013, **113**, 402–441; (c) S. Bähr, W. Xue and M. Oestreich, *ACS Catal.*, 2019, **9**, 16–24; (d) C. Moberg, *Synthesis*, 2020, **52**, 3129–3139; (e) K. Hirano and M. Uchiyama, *Adv. Synth. Catal.*, 2021, **363**, 2340–2353; (f) J.-J. Feng, W. Mao, L. Zhang and M. Oestreich, *Chem. Soc. Rev.*, 2021, **50**, 2010–2073.
- 29 G. Becker, H.-M. Hartmann, A. Münch and H. Riffel, *Z. Anorg. Allg. Chemie*, 1985, **530**, 29–42.
- 30 Quantitative deprotonation of Li[6] is also possible using Ph(H)₂C–N(Li)tBu as the base. Given that the amide can be formed through H[−] transfer to Ph(H)C=NtBu, this deprotonation experiment also marks the hitherto missing keystone of the reaction mechanism previously proposed for the catalytic hydrogenation of the imine by Li₂[2]/H₂: In contrast to conventional FLPs, the H₂-activation product Li₂[11] (Scheme 3) delivers first an H[−] and second an H⁺ ion (cf. ref. 13).
- 31 T. Wassmann, A. P. Seitsonen, A. M. Saitta, M. Lazzeri and F. Mauri, *J. Am. Chem. Soc.*, 2010, **132**, 3440–3451.
- 32 For the following reasons, Li₂[2] was chosen despite its somewhat different behavior toward HBpin compared to the other M₂[1]/M₂[2] compounds: (i) The neutral Me-substituted DBA **2** is more convenient to handle than its H-substituted congener **1**. (ii) We have already published the Li⁺ salts of some cycloadducts between [DBA-Me]₂^{2−} and unsaturated substrates, which are the actual catalytically active species and thus the focus of our mechanistic investigations (cf. refs. 12, 13). (iii) Since our DFT calculations include the counter cations, the computational cost for the treatment of Li⁺ is lower than for the treatment of Na⁺/K⁺.
- 33 We initially relied on this protocol because it is most similar to the one applied for our earlier hydrogenation reactions (cf. ref. 13), which we want to use for comparison. Later, it was found that adding the unsaturated substrate first is essential to exclude contributions of the hidden borohydride catalysis as an undesired background reaction.
- 34 X. Wang and X. Xu, *RSC Adv.*, 2021, **11**, 1128–1133.
- 35 A. R. Gatti and T. Wartik, *Inorg. Chem.*, 1966, **5**, 329–330.
- 36 M. Magre, M. Szewczyk and M. Rueping, *Chem. Rev.*, 2022, **122**, 8261–8312.
- 37 Sen et al. identified a comparable Li⁺⋯O coordination as the key interaction in lithium phenolate-catalyzed HBpin additions to carbonyl compounds. The resulting decrease in O=B π donation leads to an increase in the Lewis acidity of the HBpin molecule and in turn facilitates its interaction with the carbonyl-O atom. We adopted such a scenario to Li₂[14] and computed a corresponding reaction barrier of +31 kcal mol^{−1}. Although this value is lower than that of the uncatalyzed reaction by −14 kcal mol^{−1}, it is still higher than **TS1** by +6 kcal mol^{−1}. Since **TS1** represents the highest barrier in our hydroboration Mechanism II (see the ESI for details), also Sen's mechanism can be ruled out for our case: M. K. Bisai, T. Das, K. Vanka and S. S. Sen, *Chem. Commun.*, 2018, **54**, 6843–6846. DOI: 10.1039/D2SC05518J
- 38 Hydroboration of Ph(H)C=NPh with HBpin in THF-d₈ required about the same time, regardless of whether Li₂[14] or Na₂[14] was used as the catalyst (NMR-spectroscopic control). We take this as justification that Li₂[14] is a representative example catalyst to study Mechanism II. It should also be noted that the necessary explicit treatment of the counter cation's coordination sphere is much more complex in the case of the larger Na⁺. Its coordination sphere includes six neighbors while the one of Li⁺ only contains four. Consequently, not only does the size of the system increase, but also a greater variety of coordination polyhedra can be expected.
- 39 C. Weetman, M. S. Hill and M. F. Mahon, *Chem. Eur. J.*, 2016, **22**, 7158–7162.
- 40 D. Mukherjee, S. Shirase, T. P. Spaniol, K. Mashima and J. Okuda, *Chem. Commun.*, 2016, **52**, 13155–13158.
- 41 M. D. Anker, M. Arrowsmith, R. L. Arrowsmith, M. S. Hill and M. F. Mahon, *Inorg. Chem.*, 2017, **56**, 5976–5983.
- 42 A. Ramos, A. Antiñolo, F. Carrillo-Hermosilla, R. Fernández-Galán and D. García-Vivó, *Chem. Commun.*, 2019, **55**, 3073–3076.
- 43 N. Sarkar, S. Bera and S. Nembenna, *J. Org. Chem.*, 2020, **85**, 4999–5009.
- 44 J.-D. Chai and M. Head-Gordon, *Phys. Chem. Chem. Phys.*, 2008, **10**, 6615–6620.
- 45 R. Krishnan, J. S. Binkley, R. Seeger and J. A. Pople, *J. Chem. Phys.*, 1980, **72**, 650–654.
- 46 T. Clark, J. Chandrasekhar, G. W. Spitznagel and P. von Ragué Schleyer, *J. Comput. Chem.*, 1983, **4**, 294–301.
- 47 A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2009, **113**, 6378–6396.
- 48 A. Ben-Naim, *J. Phys. Chem.*, 1978, **82**, 792–803.
- 49 J. R. Pliengo Jr and J. M. Riveros, *Phys. Chem. Chem. Phys.*, 2002, **4**, 1622–1627.
- 50 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Peterson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, Gaussian 16 (Revision A.03), Gaussian Inc., Wallingford, CT, USA, 2016.
- 51 I. P. Query, P. A. Squier, E. M. Larson, N. A. Isley and T. B. Clark, *J. Org. Chem.*, 2011, **76**, 6452–6456.
- 52 Another possible source of Bpin residues is Li[5]. Attack of **Int1** at the Bpin substituent of Li[5] would lead to the immediate formation of **P** and Li₂[2] via [Bpin]⁺ transfer and with release of BH₃·thf (cf. the analogous reaction Li[SiPh₃] + Li[5] → pinB–SiPh₃ + Li₂[2]; Scheme 2). According to our calculations this alternative scenario would lead to a decrease in the driving force of the overall reaction from −14 kcal mol^{−1} to −8 kcal mol^{−1}. On the other hand, formation of **Int2** (+15 kcal mol^{−1}) would be bypassed. However, an interaction between the amide anion of **Int1** and negatively charged [5][−] is electrostatically unfavorable and the reactive B(pin) center of [5][−] is sterically less



accessible than that of HBpin. We therefore assume that the thermodynamically less favorable formation of **Int2** (+15 kcal mol⁻¹) is preferred by kinetics. Since the formation of Li[**5**] and Li[H₂Bpin], which is the energetically most costly step of Cycle I, is not circumvented, a switch to Cycle II will take place in any case.

- 53 An analogous attack of a B-bonded Me residue on HBpin or a B–H/B–C σ -bond metathesis leading directly to **Int4** (Fig. 5) can be ruled out due to significantly higher reaction barriers (Fig. S98, **TSa** and **TSb**).
- 54 H⁻ transfer to the alternative electrophile HBpin would be thermodynamically less favorable by +13 kcal mol⁻¹. In general, HBpin appears to be the weakest H⁻ acceptor of the investigated species (see Table S8).

View Article Online
DOI: 10.1039/D2SC05518J

