



Kent Academic Repository

Tamè, Luigi (2015) *Hand Posture Alters Perceived Finger Numerosity.*
In: UNIVERSITY OF LONDON RESEARCH DAY ACTING AND INTERACTING BODIES, 2 June 2015, London, UK.

Downloaded from

<https://kar.kent.ac.uk/98835/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/Session 3>

This document version

Other

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Hand Posture Alters Perceived Finger Numerosity

Luigi Tamè, Elanah Dransfield, Matthew R Longo

Birkbeck, University of London

Abstract

Patients with posterior parietal lesions commonly fail in identifying their fingers, a condition known as *finger agnosia*. Recent research has shown that healthy people may also perform poorly in certain tasks of finger identification. Here, we investigated whether the representations of finger numerosity is modulated by the spatial relationships between the fingers. We used the 'in between' test, a classical measure of finger agnosia, in which participants estimate the number of unstimulated fingers between two touched fingers. Stimulation consisted of pairs of mechanical tactile stimuli delivered on the back of the second phalanx of the fingers of one hand. Across blocks, the fingers were placed in three postures: (1) with fingers touching each other, (2) fingers separated by one centimetre, or (3) fingers spread to the maximum comfortable splay. Participants judged the number of unstimulated fingers 'in between' the two touches and responded vocally as quickly and accurately as possible. Critically, participants gave larger numerosity estimates when the fingers were positioned far apart compared to when they were close together or touching. Our results demonstrate that increasing the spatial distance between the fingers makes participants experience the fingers as more numerous.

Keywords: TACTILE, FINGERS, BODY STRUCTURAL REPRESENTATION, SPACE