



Kent Academic Repository

Lukova, Andrea, Dunmore, Christopher J., Skinner, Matthew M. and Kivell, Tracy L. (2023) *Trabecular distribution of distal femur in extant apes*. In: PASC-SCPA 2023, 19-20 Jan 2023, Canada. (Unpublished)

Downloaded from

<https://kar.kent.ac.uk/100674/> The University of Kent's Academic Repository KAR

The version of record is available from

This document version

Other

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Preferred format: podium presentation

Trabecular distribution of distal femur in extant apes

Andrea Lukova¹, Christopher J. Dunmore¹, Tracy L. Kivell^{1,2}, Matthew M. Skinner^{1,2}

¹Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK

²Centre for the Exploration of the Deep Human Journey, University of the Witwatersrand, Johannesburg, South Africa

Extant great apes are often used to model aspects of fossil hominin locomotor behaviours. Comparative investigation of trabecular bone, which (re-)models to reflect loads incurred during life, can provide novel insights into the locomotor reconstruction of fossil taxa [1]. Here we analyze the distal femoral epiphysis of [*Homo sapiens*] (N = 26), [*Gorilla gorilla*] (N = 14), [*Pan troglodytes verus*] (N = 15), and [*Pongo*] sp. (N = 9) to determine how variation in trabecular structure reflects differences in locomotor behaviours. Canonical holistic morphometric analysis (cHMA) of relative bone volume fraction (rBV/TV) and degree of anisotropy (DA) is used to infer patterns of joint loading in extant taxa.

A principal component analysis of rBV/TV and DA distributions show clear separation between taxa. Trabecular distribution in humans is consistent with medial (due to the ground reaction forces) and lateral (due to the resistance of the knee adduction moment provided by the quadriceps and gastrocnemius muscles and lateral collateral ligament) loading. Distribution in non-human apes is consistent with primarily medial loading due to the higher knee adduction moment, varus angle and ground reaction forces. Signals of a more extended knee in female gorillas compared to males (or chimpanzees) may reflect a more extended knee position during vertical climbing and higher arboreality in females [3]. Orangutans showed the most homogenous distribution of trabecular structure across both condyles, consistent with more variable knee joint postures. These results provide the comparative context to interpret knee posture and, in turn, locomotor behaviours in fossil hominins.

References: [1] Georgiou, L., Dunmore, C. J., Bardo, A., Buck, L. T., Hublin, J. J., Pahr, D. H., ... & Skinner, M. M. (2020). Evidence for habitual climbing in a Pleistocene hominin in South Africa. *Proceedings of the National Academy of Sciences*, 117(15), 8416-8423. [2] Bachmann, S., Dunmore, C. J., Skinner, M. M., Pahr, D. H. and Synek, A. (2022). A computational framework for canonical holistic morphometric analysis of trabecular bone. *Scientific Reports*, 12, 1-13. [3] Isler, K. (2005). 3D-kinematics of vertical climbing in hominoids. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 126(1), 66-81.

Acknowledgments: For access to specimens we thank the following individuals/institutions: Max Planck Institute for Evolutionary Anthropology (C. Boesch, J-J. Hublin); Museum für Naturkunde - Leibniz Institute for Evolution and Biodiversity Science (F. Mayer, C. Funk); Powell-Cotton Museum (I. Livne); Royal Museum for Central Africa (E. Gilissen); University of Florence (J. Moggi-Cecchi, S. Bortoluzzi); Johann-Friedrich-Blumenback-Institute for Zoology and Anthropology, Georg-August University, Goettingen (B. Grosskopf); Frankfurt Senckenberg Museum (V. Volpato); University of the Witswatersand (L. Berger, B. Zipfel); Science Academy of the Czech Republic (J. Svoboda). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 819960).