



Kent Academic Repository

Gilmer, Jannik, Trageser, Timo, Čaić, Luis, Virovets, Alexander, Bolte, Michael, Lerner, Hans-Wolfram, Fantuzzi, Felipe and Wagner, Matthias (2023) *Catalyst-free diboration and silaboration of alkenes and alkynes using bis(9-heterofluorenyl)s*. *Chemical Science*, 14 (17). pp. 4589-4596. ISSN 2041-6520.

Downloaded from

<https://kar.kent.ac.uk/100715/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1039/D3SC01395B>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: J. Gilmer, T. Trageser, L. ai, A. Virovets, M. Bolte, H. Lerner, F. Fantuzzi and M. Wagner, *Chem. Sci.*, 2023, DOI: 10.1039/D3SC01395B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Catalyst-free diboration and silaboration of alkenes and alkynes using bis(9-heterofluorenyls)

Jannik Gilmer,^{a,b} Timo Trageser,^a Luis Čaić,^a Alexander Virovets,^a Michael Bolte,^a Hans-Wolfram Lerner,^a Felipe Fantuzzi^{*b} and Matthias Wagner^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Diboration and silaboration reactions are prominent tools to introduce valuable functional groups into organic substrates. To date, most diboranes(4) and silyboranes used for this purpose are electronically and/or kinetically stabilized and require activation by a catalyst. We show here that the tetraaryl (μ -hydrido)diborane(4) anion **[3]**⁻ and the silyl (hydrido)borate (**[4]**⁻)/Me₃SiBr system react spontaneously with the archetypal olefin ethylene in the absence of a catalyst. The actual active species in both cases are the valence isoelectronic intermediates [FluB–B(H)Flu]⁻ (**[1]**⁻) and FluB–Si(H)Flu (**2**), which consist of two 9-heterofluorenyl halves that get attached to the 1 and 2 positions of ethylene. At room temperature, **[1]**⁻ is present in a dynamic equilibrium with its isolable isomer **[3]**⁻, while **2** has to be released *in situ* at low temperatures by H⁻ abstraction from **[4]**⁻. Quantum-chemical calculations show qualitatively identical reaction mechanisms for **[1]**⁻ and **2**. Since the reactions start with π coordination of the ethylene molecule to a vacant B(p_z) orbital, the high Lewis-acidity and low steric hindrance of the 9-borafluorenyl fragments are the keys to success. As the reaction proceeds, back-donation from the B–E bond into the ethylene π^* orbital becomes increasingly important (E = B, Si). The scope of the reactions has been extended to *t*Bu(H)C=CH₂ and *t*BuC \equiv CH on the one hand and FluB–Si(Cl)Flu as well as FluB–Si(Cl)Ph₂ on the other.


Introduction

Hydroborations are addition reactions of an H–BR₂ bond to alkenes or alkynes.¹ The resulting organoboranes are versatile precursors for further functionalization or C–C coupling. The ease with which hydroboration reactions proceed depends critically on the Lewis acidity of the hydroborane and, in turn, on the nature of its R substituents: While the dialkyl borane 9-borabicyclo[3.3.1]nonane (9-BBN) smoothly adds to unsaturated organic substrates already at RT,² significantly elevated temperatures are required for hydroborations with the less Lewis acidic catecholborane (HBcat) or pinacolborane (HBpin).^{1c} As a consequence, considerable research efforts have been devoted to the development of efficient hydroboration catalysts, such as [ClRh(PPh₃)₃] (Wilkinson's complex).^{1,3}

Switching from hydroboration to diboration doubles the number of boryl groups that can be introduced in a single step. However, this advantage comes at a price because strongly Lewis acidic diboranes(4) such as B₂Cl₄, which spontaneously adds to ethylene or acetylene,⁴ are difficult to prepare and impractical to handle.⁵ Thus, the most commonly used diboranes(4) today are the commercially available compounds B₂cat₂ and B₂pin₂, which routinely require activation by catalysts

(Scheme 1).⁶ These can be as diverse as noble metal complexes⁷ and plain alkoxide ions.⁸ The latter act through tetracoordination of one B center to form B(sp²)–B(sp³) diboranes with strongly polarized B–B bonds.⁹

Diborations typically provide products containing two identical functionalizable groups that are afterwards difficult to address individually. To remedy this deficiency, silaboration reactions have been developed during which two chemically distinct substituents of largely orthogonal reactivity are introduced – a boryl and a silyl group.¹⁰ Similar to the B–B bond in B₂pin₂, the B–Si bonds of most silyboranes are rather inert and consequently need to be activated by transition-metal complexes or nucleophiles.^{6c,11,12} This is because the vast majority of available silyboranes carry electronically tamed or kinetically protected boryl moieties like Bpin,^{10a} Bcat,¹³ or



Scheme 1 General schemes for hydroboration, diboration, and silaboration reactions on alkenes and alkynes.

^aInstitut für Anorganische und Analytische Chemie, Goethe-Universität Frankfurt, Max-von-Laue-Straße 7, D-60438 Frankfurt am Main, Germany. E-mail: matthias.wagner@chemie.uni-frankfurt.de

^bSchool of Chemistry and Forensic Science, University of Kent, Park Wood Rd, Canterbury CT2 7NH, United Kingdom. E-mail: f.fantuzzi@kent.ac.uk

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



BMe₂.¹⁴ As a result, boron tetracoordination by pristine alkenes or alkynes hardly takes place, which explains why these substrates are unable to activate the B–Si bonds on their own. Can an unprecedented catalyst-free silaboration be achieved by purposefully increasing the Lewis acidity of the boryl residue while decreasing its steric demand? For the following reasons, we relied on our recently reported borafluorenyl derivative **2** to answer this question (Fig. 1a):¹⁵ (i) The B center of **2** is part of a formally antiaromatic borole ring and therefore a particularly good electron acceptor, and (ii) it is exposed to its environment due to the planarized 2,2'-biphenylene ligand. Moreover, the silafluorenyl part of **2** renders it a structural analog of corresponding bis(9-borafluorenyl) anions [**1**][−] (Fig. 1a), which we have already investigated in different contexts in the recent past.¹⁶ The [**1**][−]/**2** pair now offers the unique opportunity to compare diborations and silaborations on the basis of two compounds that could not be any more alike. The results of our combined experimental and theoretical study are reported herein – among them the first spontaneous 1,2-addition of a silylborane to a substrate as simple as ethylene.

Results and discussion

All reactions and NMR measurements were carried out in THF or THF-*d*₈; all DFT calculations were performed at the SMD(THF)/ωB97XD/def2-QZVPP//SMD(THF)/ωB97XD/6-31+G** level of theory. Throughout this work, calculated structures **X**^c differ from the corresponding experimental structures **X** in that all aryl-bonded *t*Bu groups are replaced by





Fig. 1 a) The two valence isoelectronic key intermediates M[**1**] and **2** of the catalyst-free diboration and silaboration reactions, respectively (M = Li, K). M[**1**] is in a dynamic equilibrium with its isolable isomer M[**3**]; **2** forms *in situ* through thf-ligand dissociation from the NMR-detectable adduct **2**(thf) ($\Delta G_{\text{diss}} = 10.5$ kcal mol^{−1}). Carbon atoms marked with “*” bear *t*Bu groups. b) DFT calculations (SMD(THF)/ωB97XD/def2-QZVPP//SMD(THF)/ωB97XD/6-31+G**) predict that [Li(thf)₂][**1**^c] is energetically less favorable by 2.0 kcal mol^{−1} than [Li(thf)₂][**3**^c] ($\Delta G^{\ddagger} < 10$ kcal mol^{−1}). Compound numbers with a superscript “C” indicate calculated structures for which the Ar-bonded *t*Bu groups have been replaced by H atoms. For clarity, schematic representations of the calculated structures are shown; full graphical representations are included in the ESI.

H atoms, in order to reduce CPU time and the number of potential conformers.

Since the immediate, valence isoelectronic starting materials [**1**][−] and **2** are not (yet) available, we first required suitable synthesis equivalents that are either in equilibrium with [**1**][−]/**2** or capable of releasing the actual reactive species *in situ*. An appropriate synthesis equivalent for M[**1**] should be its isolable isomer M[**3**] (M = Li, K; Scheme 2), because M[**3**] behaved as if it were M[**1**] already in previous reactivity studies.^{16d,f} Moreover, quantum-chemical calculations indicate that (i) [Li(thf)₂][**3**^c] is only 2.0 kcal mol^{−1} more stable than [Li(thf)₂][**1**^c] and (ii) the rearrangement [**3**^c] → [**1**^c] is associated with barriers < 10 kcal mol^{−1} (Fig. 1b; more details in Scheme S2).¹⁷ On the BSi side, the thf adduct **2**(thf) of the elusive **2** can straightforwardly be generated from Na[**4**] through H-abstraction with Me₃SiBr at low temperatures (Scheme 2; **2**^c(thf) → **2**^c + THF; $\Delta G_{\text{diss}} = 10.5$ kcal mol^{−1}).^{15,17}

According to our experience, the reactivity of anionic aryl boranes is often strongly influenced by the nature of their counter cations.^{15,18} We have therefore conducted all transformations involving the [**1**][−] anion with both Li[**3**] and K[**3**]. Since in the present context the counter-cation



Scheme 2 Diboration and silaboration of ethylene, starting from K[**3**] or Na[**4**]/Me₃SiBr and furnishing K[**5**] and **6**(thf), respectively. The reactions proceed *via* the valence isoelectronic intermediates [**1**]^{n−} and [**2**]^{n−}. (i) THF-*d*₈, −196 °C to RT, 99% yield; (ii) THF, −78 °C to RT, 56% yield. Carbon atoms marked with “*” bear *t*Bu groups. In the crystal-structure plots, the [K(thf)₂]⁺ counter cation of [**5**][−], C-bonded H atoms, and Ar-bonded *t*Bu groups are omitted for clarity; H: gray, B: green, C: black, O: red, Si: blue.




dependence turned out to be marginal, we will henceforth refer only to the K^+ salts, as these gave the qualitatively best crystallographic analyses (all results obtained with the Li^+ salts are provided in the ESI).

In a first experiment, we assessed the behavior of $K[3]$ and the $Na[4]/Me_3SiBr$ system toward ethylene, the archetypal olefin. In both cases, the reaction was complete within minutes at $T \leq RT$ and furnished the 1,2-bis(9-borafluorenyl)ethane salt $K[5]$ and the 1-(9-silafluorenyl)-2-(9-borafluorenyl)ethane $6(thf)$, respectively (Scheme 2). While the formation of $K[5]$ was quantitative (99% yield), a somewhat lower product selectivity of 90% was determined by NMR in the case of $6(thf)$ (56% yield), which we attribute to issues associated with the *in situ* release of 2 . The 1H NMR spectrum of $K[5]$ shows a signal at 1.47 ppm (4H) assignable to the chemically equivalent protons of the 1,2-ethanediy bridge. A severely broadened ^{13}C resonance at 15.2 ppm is characteristic of B-bonded alkyl-C atoms.¹⁹ Only one $tBuC_6H_3$ signal set is observed in both the 1H and the $^{13}C\{^1H\}$ NMR spectrum, testifying to a high (average) symmetry of $[5]^-$ in solution. The computed ^{13}C chemical shifts of $[5]^-$ deviate on average by < 2 ppm from the experimentally determined values (a significantly larger mean deviation > 4 ppm was determined for the computed ^{13}C shifts of a hypothetical diboration product with a dibenzo[*g,p*]chrysene-type framework; cf. $[3]^-$).¹⁷ The 1,2-bis(9-borafluorenyl)ethane structure of $[K(thf)_2][5]$ was ultimately proven by single-crystal X-ray diffraction (SCXD; Scheme 2). As sole discrepancy, the B...B-bridging H^- ion of the twisted cyclic HB_2C_2 core, which was located in the difference Fourier map and freely refined, was not detectable even in the


$^1H\{^{11}B\}$ NMR spectrum. Also, no doublet splitting was observable in the ^{11}B NMR spectrum, although the resonance clearly appeared in the spectral region of tetracoordinated ^{11}B nuclei (9.3 ppm).¹⁹ To confirm that these spectral peculiarities do not contradict our structural proposal, but are inherent to $K[5]$, we first compared its 1H NMR data with those of the known $FluB-CH_2CH_2-BFlu$ ($BFlu = 2,7$ -di-*t*Bu-9-borafluorenyl), a neutral analog of $[5]^-$ devoid of the B...B bridging H^- ion.²⁰ We found differences in the chemical shift values of corresponding protons of up to 0.5 ppm, but after addition of 1 equiv. $Li[HBEt_3]$ the spectra of the reaction mixture and of $K[5]$ became essentially superimposable. Unlike $K[5]$, the less symmetric $6(thf)$ leads to two different CH_2 and $tBuC_6H_3$ signal sets in the 1H as well as the $^{13}C\{^1H\}$ NMR spectrum. Only one of the two 1,2-ethanediy ^{13}C resonances is broadened. The signal of the Si-bonded H atom appears as a triplet at 4.68 ppm ($1H$, $^3J_{H,H} = 3.5$ Hz) and shows ^{29}Si satellites ($^1J_{H,Si} = 192$ Hz). Its well-resolved shape points against a bonding interaction with the adjacent B atom in solution, even though $B(\mu-H)Si$ structural motifs are known.²¹ The pronounced shielding of the ^{11}B nucleus in $6(thf)$ (13.6 ppm) is therefore likely due to thf coordination. In line with this, 1H NMR spectra ($THF-d_8$) recorded on a sample of $6(thf)$ isolated from $THF-h_8$ revealed the presence of 1 equiv. of $THF-h_8$. The solid-state structure of $6(thf)$ features a terminal Si-H atom, a thf ligand attached to the B atom, and an approximate *s-trans* configuration about the 1,2-ethanediy bridge ($B-C-C-Si = 167^\circ$).

According to quantum-chemical calculations, the reaction mechanisms underlying the formation of $K[5]$ and $6(thf)$ starting



Scheme 3 a) Computed reaction mechanism for the diboration and silaboration of ethylene, starting from $[1^C]^-$ and 2^C and leading to the primary intermediates $[8^C]^-$ and $gauche-6^C$, respectively. b) Conformational changes and subsequent adduct formations furnishing the final products $[5^C]^-$ and $6^C(thf)$. Compound numbers with a superscript "C" indicate calculated structures for which the Ar-bonded *t*Bu groups have been replaced by H atoms. All energies in $kcal\ mol^{-1}$ at the SMD(THF)/ ω B97XD/def2-QZVPP//SMD(THF)/ ω B97XD/6-31+G** level of theory. Prox, dist: proximal, distal with respect to the B...E core.






Scheme 5 a) Diboration and silaboration of $t\text{Bu}(\text{H})\text{C}=\text{CH}_2$, starting from $\text{K}[\mathbf{3}]$ or the chlorosilane $\mathbf{9}(\text{thf})$ and furnishing $\text{K}[\mathbf{11}]/\text{K}[\mathbf{12}]$ (ratio 10:1) and $\mathbf{13}$, respectively. (i) $\text{THF}-d_8$, RT, quant. conv. by NMR. Upon heating, the mixture $\text{K}[\mathbf{11}]/\text{K}[\mathbf{12}]$ is quantitatively converted to the 1,1-diboration product $\text{K}[\mathbf{12}]$. The putative 1,1-silaboration product $\mathbf{14}$ was not observed, but obtained *via* a different route (*cf.* Scheme 5c). b) Solid-state structures of co-crystallized $\{[\mathbf{11}]_{0.73}[\mathbf{12}]_{0.27}\}^-$ and of $\mathbf{13}$. The $[\text{K}(\text{thf})_2]^+$ counter cation of $\{[\mathbf{11}]_{0.73}[\mathbf{12}]_{0.27}\}^-$, C-bonded H atoms, and Ar-bonded $t\text{Bu}$ groups are omitted for clarity; H: gray, B: green, C: black, Si: blue, Cl: yellow. c) Synthesis of $\mathbf{14}$ from $\mathbf{15}$, $t\text{BuC}\equiv\text{CH}$, and $(\text{HBFlu})_2$ *via* a hydrosilylation/hydroboration sequence [$(\text{HBFlu})_2 = 2,7$ -di-*tert*-butyl-9H-9-borafluorene dimer]. Carbon atoms marked with "*" bear $t\text{Bu}$ groups.

found that the diborylethane $\text{K}[\mathbf{11}]$ is formed together with its diborylmethane isomer $\text{K}[\mathbf{12}]$ in a 10:1 ratio (RT, quantitative conversion). Silaboration of $t\text{Bu}(\text{H})\text{C}=\text{CH}_2$ gave exclusively the 1,2-addition product $\mathbf{13}$, with the $t\text{Bu}$ group adjacent to the boryl substituent. Upon heating to 50°C for 5 d, $\text{K}[\mathbf{11}]$

underwent quantitative rearrangement to $\text{K}[\mathbf{12}]$,³⁰ consistent with this, $[\mathbf{12}^-]$ is predicted to be thermodynamically more stable than $[\mathbf{11}^-]$ by 6.0 kcal mol^{-1} . $\mathbf{13}$ remained inert at 50°C , although its 1-boryl-1-silyl-isomer $\mathbf{14}$ (Scheme 5a) should also be thermodynamically favored by 5.5 kcal mol^{-1} (*cf.* $\mathbf{13}^-$ vs. $\mathbf{14}^-$). Since rearrangement cannot be enforced by applying higher temperatures (prolonged heating of $\mathbf{13}$ to 100°C results in decomposition), we alternatively resorted to the successive hydrosilylation/hydroboration of $t\text{BuC}\equiv\text{CH}$ to prove that a stable $\mathbf{14}$ exists in principle (Scheme 5c).³¹ Given the established access routes to $[\mathbf{12}]^-$ and $\mathbf{14}$ *via* hydroboration of 9-vinyl-9-heterofluorenes, it is reasonable to propose that the $[\mathbf{11}^-] \rightarrow [\mathbf{12}]^-$ isomerization can proceed along a dehydroboration/rehydroboration sequence. Assuming an analogous scenario for a putative $\mathbf{13} \rightarrow \mathbf{14}$ rearrangement, either the transition state associated with the dehydroboration step is too high to be overcome at $T < 100^\circ\text{C}$ or additional high-temperature reaction channels open after dehydroboration. As the most obvious distinguishing feature, $[\mathbf{11}]^-$ generates four sets of $t\text{BuC}_6\text{H}_3$ signals and $[\mathbf{12}]^-$ only two, in both the ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum. Due to the stereogenic center in $[\mathbf{11}]^-$, the protons of the 1,2-ethylenediyl group represent an AMX spin system, whereas the $t\text{BuH}_2\text{C}-\text{CH}$ linker of $[\mathbf{12}]^-$ has an A_2X spin system. In $[\mathbf{11}]^-$, both $\text{BC}-\text{CB}$ ^{13}C resonances are severely broadened; in $[\mathbf{12}]^-$, this applies solely to the CB_2 signal. At RT, all NMR signals of the 9-borafluorenyl group of $\mathbf{13}$ are broadened (almost) beyond detection, which we attribute to a coalescence phenomenon induced by rotation of the substituent about the $\text{B}-\text{C}$ bond. Indeed, cooling of the sample to $T = -30^\circ\text{C}$ results in the expected four $t\text{BuC}_6\text{H}_3$ signal sets. The chemical shift values of the two ^{13}C sets corresponding to the 9-borafluorenyl moiety show characteristic deviations from those of $\mathbf{6}(\text{thf})$ and similarities to those of tricoordinate 9-bromo-9-borafluorene,³² indicating the absence of a B-coordinating thf ligand or of a $\text{B}\cdots\text{Si}$ -bridging Cl atom. The coalescence temperature of 9-borafluorenyl rotation decreases from RT to approximately -30°C upon going from $\mathbf{13}$ to $\mathbf{14}$, pointing to the $t\text{Bu}$ group as major origin of rotational hindrance. Thus, the room-temperature ^1H NMR spectrum of $\mathbf{14}$ shows three $t\text{BuC}_6\text{H}_3$ signal sets, one of them with double intensity. Gas-phase diffusion of *n*-hexane into a THF solution of $\text{K}[\mathbf{11}]/\text{K}[\mathbf{12}]$, that had never been exposed to temperatures above RT, resulted in the growth of co-crystals of composition $[\text{K}(\text{thf})_2]\{[\mathbf{11}]_{0.73}[\mathbf{12}]_{0.27}\}^-$. Almost all atoms of both species coincide in the crystal lattice except for their $\{\text{B}_2(\mu-\text{H})(\mu-\text{CH}_2\text{CH}t\text{Bu})\}$ and $\{\text{B}_2(\mu-\text{H})(\mu-\text{CHCH}_2t\text{Bu})\}$ fragments, which overlap partly (shown in blue and red, respectively, in Scheme 5b). From this superposition it becomes obvious that $[\mathbf{12}]^-$ is the sterically less encumbered isomer. The solid-state structure of $\mathbf{13}$ confirms the 1,2-addition mode of the silaboration reaction as well as its proposed regioselectivity. Furthermore, the SCXD structure of free $\mathbf{13}$ bears striking resemblance with the computed intermediate *gauche*- $\mathbf{6}^c$ (Scheme 3), in line with the proposition that the initially formed *gauche*- $\mathbf{6}^c$ is favored over *anti*- $\mathbf{6}^c$ as long as no thf coordination occurs. In view of the fact that the diborations/silaborations with $\text{K}[\mathbf{3}]$ and $\mathbf{9}(\text{thf})$ proceed uncatalyzed and with remarkable ease, the






Scheme 6 Diboration and silaboration of $t\text{BuC}\equiv\text{CH}$, starting from $\text{K}[3]$ or the chlorosilane $9(\text{thf})$ and furnishing $\text{K}[16]$ and 17 , respectively. Carbon atoms marked with "*" bear $t\text{Bu}$ groups. In the crystal-structure plots, the $[\text{K}(\text{thf})_2]^+$ counter cation of $[16]^-$, C-bonded H atoms, and Ar-bonded $t\text{Bu}$ groups are omitted for clarity; H: gray, B: green, C: black, Si: blue, Cl: yellow. (i) $\text{THF}-d_6$, RT, quant. conv. by NMR.

next question was: Can the scope of the two reactions be extended even to alkynes? As a first trial, we treated both starting materials with $t\text{BuC}\equiv\text{CH}$ and observed clean formation of the 1,2-*cis*-addition products $\text{K}[16]$ and 17 , respectively, within minutes at RT (Scheme 6). As in 13 , the $t\text{Bu}$ group of 17 ends up next to the 9-borafluorenyl unit. These structural assignments are based on SCXD and (2D) NMR experiments. The configuration of 17 would be considered "abnormal" in most transition metal-catalyzed silaboration reactions of alkynes $\text{RC}\equiv\text{CH}$, where the silyl group is usually introduced adjacent to the R substituent.³³ There, inverting this "normal" regioselectivity is hard to achieve and requires elaborate catalyst design.³⁴ A notable NMR feature of ditopic organoborane $\text{K}[16]$ is that both the $\text{B}(\mu\text{-H})\text{B}$ and the $\text{C}=\text{CH}$ signals are split into well-resolved doublets in the $^1\text{H}\{^{11}\text{B}\}$ NMR spectrum due to pronounced $^3J_{\text{H,H}}$ coupling of 8.1 Hz.

Finally, we investigated the extent to which our silaboration reaction depends on (i) the sterically undemanding planar framework of the 9-silafluorenyl fragment and (ii) an enhanced Lewis acidity of the Si center due to the compressed endocyclic C–Si–C angle.³⁵ To this end, we formally cut the biphenyl backbone of the 9-silafluorenyl and switched to an $\text{Si}(\text{Cl})\text{Ph}_2$ substituent (*cf.* $18(\text{thf})$; its synthesis is analogous to that of $9(\text{thf})$ and described in the ESI; Scheme 7). We found that the reaction of $18(\text{thf})$ with ethylene and $t\text{BuC}\equiv\text{CH}$ proceeds with similar ease and selectivity as in the case of $9(\text{thf})$ to give the final products $19(\text{thf})$ and 20 (see the ESI for the SCXD results).



Scheme 7 Silaboration of ethylene or $t\text{BuC}\equiv\text{CH}$, starting from diphenylsilane $18(\text{thf})$ and furnishing $19(\text{thf})$ or 20 , respectively. Carbon atoms marked with "*" bear $t\text{Bu}$ groups. (i) exc. ethylene, $\text{THF}-d_6$, $-196\text{ }^\circ\text{C}$ to RT, quant. conv. by NMR; (ii) 1 equiv. $t\text{BuC}\equiv\text{CH}$, $\text{THF}-d_6$, RT, quant. conv. by NMR.

We conclude that neither of the factors (i) and (ii) is critical for the reaction to proceed. In addition to this fundamental insight, an advantage of $18(\text{thf})$ in terms of applications is the easier availability of the starting material $\text{Cl}(\text{H})\text{SiPh}_2$ compared to $\text{Cl}(\text{H})\text{SiFlu}$ (15).¹⁵

Conclusions

We have successfully employed a pair of otherwise isostructural B–B- and B–Si-bonded compounds ($[1]^-$ and 2) to compare diboration and silaboration reactions under essentially identical steric conditions. Irrespective of their different charge, both molecules undergo spontaneous 1,2-addition to $\text{C}=\text{C}$ double and $\text{C}\equiv\text{C}$ triple bonds without the need for a catalyst. Such reactivity is rare for electron-precise diboranes and unprecedented for silyboranes. We attribute it to the high Lewis acidity and the low steric shielding of the installed planar borafluorenyl units. In the case of the substrate $t\text{Bu}(\text{H})\text{C}=\text{CH}_2$, also the 1,1-diboration product $[12]^-$ was observed at RT, albeit to a small degree of less than 10%. Upon heating to $50\text{ }^\circ\text{C}$, the corresponding 1,2-diboration product $[11]^-$ quantitatively rearranges to $[12]^-$. We never encountered an analogous 1,1-silaboration product – not even at elevated temperatures. Using quantum chemical calculations, we developed a mechanistic proposal that is consistent with all experimental facts. The crucial activation step occurs *via* simultaneous (i) coordination of a filled substrate π orbital to the empty $\text{B}(p_z)$ orbital and (ii) back-bonding from the filled B–E σ orbital to the substrate π^* orbital (E = B, Si). Apart from the insights into catalyst-free diboration and silaboration reactions gained from the studies on our diboranes and silyboranes, the products obtained are promising precursors for future syntheses of B,Si-doped PAHs.

Author contributions

J.G. performed the experimental studies, characterized all new compounds, and performed the quantum-chemical calculations. T.T. optimized the synthesis of $\text{M}[5]$ and performed the syntheses of $\text{M}[11]$ as well as $\text{M}[12]$. L.C. assisted with the synthesis of 14 . M.B.



performed the X-ray crystal structure analyses of the compounds **9**(thf), [K(thf)₂][**11**]_{0.73}[**12**]_{0.27}, [Li(thf)₄][**12**], **14**, **20**. A.V. performed the X-ray crystal structure analyses of the compounds [K(thf)₂][**5**], **6**(thf), **13**, [K(thf)₂][**16**], and **17**, and further optimized the structures [K(thf)₂][**11**]_{0.73}[**12**]_{0.27}, [Li(thf)₄][**12**], and **20**. H.W.L., F.F., and M.W. supervised the project. The manuscript was written by J.G. and M.W. and edited by all co-authors.

Data availability

The datasets supporting this article have been uploaded as part of the supplementary information.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

J.G. thanks the German Academic Exchange Service for a research grant. We thank the center for scientific computing (CSC) Frankfurt and the University of Kent for providing HPC resources that contributed to the computational investigation of this work.

Notes and references

- (a) K. Burgess and M. J. Ohlmeyer, *Chem. Rev.*, 1991, **91**, 1179–1191; (b) I. Beletskaya and A. Pelter, *Tetrahedron*, 1997, **53**, 4957–5026; (c) C. C. Chong and R. Kinjo, *ACS Catal.*, 2015, **5**, 3238–3259; (d) S. J. Geier, C. M. Vogels, J. A. Melanson and S. A. Westcott, *Chem. Soc. Rev.*, 2022, **51**, 8877–8922.
- H. C. Brown and J. C. Chen, *J. Org. Chem.*, 1981, **46**, 3978–3988.
- D. Männig and H. Nöth, *Angew. Chem. Int. Ed.*, 1985, **24**, 878–879.
- (a) G. Urry, J. Kerrigan, T. D. Parsons and H. I. Schlesinger, *J. Am. Chem. Soc.*, 1954, **76**, 5299–5301; (b) P. Ceron, A. Finch, F. Frey, J. Kerrigan, T. Parsons, G. Urry and H. I. Schlesinger, *J. Am. Chem. Soc.*, 1959, **81**, 6368–6371; (c) C. Pubill-Ulldemolins, E. Fernández, C. Bo and J. M. Brown, *Org. Biomol. Chem.*, 2015, **13**, 9619–9628.
- Further selected examples of catalyst-free diboration reactions: (a) H. Klusik, C. Pues and A. Berndt, *Z. Naturforsch.*, 1984, **39b**, 1042–1045; (b) S. Luckert, E. Eversheim, U. Englert, T. Wagner and P. Paetzold, *Z. Anorg. Allg. Chem.*, 2001, **627**, 1815–1823; (c) C. Kojima, K.-H. Lee, Z. Lin and M. Yamashita, *J. Am. Chem. Soc.*, 2016, **138**, 6662–6669; (d) J. Böhnke, H. Braunschweig, A. Deißberger, T. Dellermann, R. D. Dewhurst, J. O. C. Jiménez-Halla, S. Kachel, H. Kelch and D. Prieschl, *Chem. Commun.*, 2017, **53**, 12132–12135.
- (a) T. B. Marder and N. C. Norman, *Top. Catal.*, 1998, **5**, 63–73; (b) E. C. Neeve, S. J. Geier, I. A. I. Mkhalid, S. A. Westcott and T. B. Marder, *Chem. Rev.*, 2016, **116**, 9091–9161; (c) A. B. Cuenca, R. Shishido, H. Ito and E. Fernández, *Chem. Soc. Rev.*, 2017, **46**, 415–430; (d) S. Ding, L. Xu and Z. Miao, *Molecules*, 2019, **24**, 1325; (e) L. Tendera, F. Fantuzzi, T. B. Marder and U. Radius, *Chem. Sci.*, 2023, **14**, 2215–2228.
- (a) T. Ishiyama, N. Matsuda, N. Miyaura and A. Suzuki, *J. Am. Chem. Soc.*, 1993, **115**, 11018–11019; (b) R. T. Baker, P. Nguyen, T. B. Marder and S. A. Westcott, *Angew. Chem. Int. Ed.*, 1995, **34**, 1336–1338. DOI: 10.1039/D3SC01395B
- (a) A. Bonet, C. Pubill-Ulldemolins, C. Bo, H. Gulyás and E. Fernández, *Angew. Chem. Int. Ed.*, 2011, **50**, 7158–7161; (b) N. Miralles, J. Cid, A. B. Cuenca, J. J. Carbó and E. Fernández, *Chem. Commun.*, 2015, **51**, 1693–1696.
- R. D. Dewhurst, E. C. Neeve, H. Braunschweig and T. B. Marder, *Chem. Commun.*, 2015, **51**, 9594–9607.
- (a) M. Sugimoto, H. Nakamura and Y. Ito, *Chem. Commun.*, 1996, 2777–2778; (b) M. Sugimoto, H. Nakamura and Y. Ito, *Angew. Chem. Int. Ed.*, 1997, **36**, 2516–2518; (c) S. Onozawa, Y. Hatanaka and M. Tanaka, *Chem. Commun.*, 1997, 1229–1230; (d) J. Carlos Araujo Da Silva, M. Birot, J.-P. Pillot and M. Pétraud, *J. Organomet. Chem.*, 2002, **646**, 179–190; (e) H. Ito, Y. Horita and E. Yamamoto, *Chem. Commun.*, 2012, **48**, 8006–8008; (f) E. Yamamoto, R. Shishido, T. Seki and H. Ito, *Organometallics*, 2017, **36**, 3019–3022; (g) Y. Morimasa, K. Kabasawa, T. Ohmura and M. Sugimoto, *Asian J. Org. Chem.*, 2019, **8**, 1092–1096; (h) Y. Gu, Y. Duan, Y. Shen and R. Martin, *Angew. Chem. Int. Ed.*, 2020, **59**, 2061–2065.
- (a) M. Oestreich, E. Hartmann and M. Mewald, *Chem. Rev.*, 2013, **113**, 402–441; (b) C. Moberg, *Synthesis*, 2020, **52**, 3129–3139; (c) J.-J. Feng, W. Mao, L. Zhang and M. Oestreich, *Chem. Soc. Rev.*, 2021, **50**, 2010–2073.
- Rare exceptions requiring particular substrates are reported in: (a) M. Sugimoto, T. Fukuda, H. Nakamura and Y. Ito, *Organometallics*, 2000, **19**, 719–721; (b) K. Oshima, T. Ohmura and M. Sugimoto, *Chem. Commun.*, 2012, **48**, 8571–8573; (c) M. Kondo, J. Kanazawa, T. Ichikawa, T. Shimokawa, Y. Nagashima, K. Miyamoto and M. Uchiyama, *Angew. Chem. Int. Ed.*, 2020, **59**, 1970–1974.
- M. Sugimoto, T. Matsuda, H. Nakamura and Y. Ito, *Tetrahedron*, 1999, **55**, 8787–8800.
- E. Bonnefon, M. Birot, J. Dunogues, J.-P. Pillot, C. Courseille and F. Taulelle, *Main Group Metal Chemistry*, 1996, **19**, 761–767.
- J. Gilmer, M. Bolte, A. Virovets, H.-W. Lerner, F. Fantuzzi and M. Wagner, *Chem. Eur. J.*, 2023, **29**, e202203119.
- (a) T. Kaese, A. Hübner, M. Bolte, H.-W. Lerner and M. Wagner, *J. Am. Chem. Soc.*, 2016, **138**, 6224–6233; (b) T. Kaese, H. Budy, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2017, **56**, 7546–7550; (c) J. Gilmer, H. Budy, T. Kaese, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2020, **59**, 5621–5625; (d) T. Trageser, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2020, **59**, 7726–7731; (e) H. Budy, J. Gilmer, T. Trageser and M. Wagner, *Eur. J. Inorg. Chem.*, 2020, **2020**, 4148–4162; (f) T. Trageser, D. Bebej, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2021, **60**, 13500–13506.
- See the ESI for more details.
- (a) E. von Grotthuss, S. E. Prey, M. Bolte, H.-W. Lerner and M. Wagner, *J. Am. Chem. Soc.*, 2019, **141**, 6082–6091; (b) H. Budy, T. Kaese, M. Bolte, H.-W. Lerner and M. Wagner, *Angew. Chem. Int. Ed.*, 2021, **60**, 19397–19405.
- H. Nöth and B. Wrackmeyer, *Nuclear Magnetic Resonance Spectroscopy of Boron Compounds*, in *NMR basic Principles and Progress*, ed. P. Diehl, E. Fluck and R. Kosfeld, Springer Verlag, Berlin, Heidelberg, New York, 1978.
- T. Kaese, T. Trageser, H. Budy, M. Bolte, H.-W. Lerner and M. Wagner, *Chem. Sci.*, 2018, **9**, 3881–3891.
- (a) B. Wrackmeyer, W. Milius and O. L. Tok, *Chem. Eur. J.*, 2003, **9**, 4732–4738; (b) H. Gao, R. Müller, E. Irran, H. F. T. Klare, M. Kaupp and M. Oestreich, *Chem. Eur. J.*, 2022, **28**, e202104464.
- In addition to the diboration mechanism outlined in Scheme 3, we also considered alternative mechanisms starting from isomers of [3⁻] other than [1⁻]. All of these have considerably



- higher overall barriers and should therefore contribute less to the overall reaction scenario (see the ESI for further details).
- 23 For steric reasons, this is different to the proposed mechanism for the addition of propylene to the bulkier [pinB-B(OMe)pin]⁻, where initially only one of the two olefinic C atoms coordinates to the anionic diborane (Ref[8a]).
- 24 G. Knizia and J. E. M. N. Klein, *Angew. Chem. Int. Ed.*, 2015, **54**, 5518–5522.
- 25 N. G. Rondan, K. N. Houk and R. A. Moss, *J. Am. Chem. Soc.*, 1980, **102**, 1770–1776.
- 26 S. P. Gnanasekar and E. Arunan, *Aust. J. Chem.*, 2020, **73**, 767–774.
- 27 The NBO calculations show a similar picture: The transition states [TS1^c]/[TS2^c] are mainly stabilized by the interaction of the ethylene π orbital with the p_z orbital of the respective sp²-hybridized borafuorenyl moiety (60.8/43.8 kcal mol⁻¹). An interaction between the B–E σ orbital and the π* orbital of ethylene is small at these stages (8.2/1.9 kcal mol⁻¹). However, the latter becomes increasingly pronounced as the reaction course progresses.
- 28 Ohmura, Suginome, and co-workers reported a transition metal-catalyzed silaboration that also tolerates an SiCl function (Ref[34a]).
- 29 A. Hübner, Z.-W. Qu, U. Englert, M. Bolte, H.-W. Lerner, M. C. Holthausen and M. Wagner, *J. Am. Chem. Soc.*, 2011, **133**, 4596–4609.
- 30 Reaction of Li[3] with 3,3-dimethyl-but-1-ene at RT in C₆D₆ gives Li[12] with 70% selectivity, which precipitates from an *n*-hexane/C₆D₆ solution at –30 °C. Following this procedure, Li[12] can be isolated in higher purity compared to the thermal rearrangement protocol starting from M[11] (M = Li, K).
- 31 We have confirmed that Karstedt's catalyst does not promote the rearrangement **13** → **14**.
- 32 A. Hübner, M. Diefenbach, M. Bolte, H.-W. Lerner, M. C. Holthausen and M. Wagner, *Angew. Chem. Int. Ed.*, 2012, **51**, 12514–12518.
- 33 Assuming that the alkyne insertion into the B–Si bond of **9**(thf) proceeds *via* a transition state analogous to TS2^c (cf. Scheme 3), we anticipate that the observed regioselectivity is due to steric effects: to avoid unfavorable interactions with the "three-dimensional" silyl group, the sterically demanding *t*Bu substituent of the incoming alkyne must be oriented above the planar borafuorene moiety, which ultimately promotes the observed "abnormal" regioselectivity.
- 34 (a) T. Ohmura, K. Oshima, H. Taniguchi and M. Suginome, *J. Am. Chem. Soc.*, 2010, **132**, 12194–12196; (b) T. Ohmura, K. Oshima and M. Suginome, *Angew. Chem. Int. Ed.*, 2011, **50**, 12501–12504; (c) K. Kojima, Y. Nagashima, C. Wang and M. Uchiyama, *ChemPlusChem*, 2019, **84**, 277–280; (d) M. Zhao, C.-C. Shan, Z.-L. Wang, C. Yang, Y. Fu and Y.-H. Xu, *Org. Lett.*, 2019, **21**, 6016–6020.
- 35 (a) A. H. J. F. de Keijzer, F. J. J. de Kanter, M. Schakel, R. F. Schmitz and G. W. Klumpp, *Angew. Chem. Int. Ed.*, 1996, **35**, 1127–1128; (b) S. Deerenberg, M. Schakel, A. H. J. F. de Keijzer, M. Kranenburg, M. Lutz, A. L. Spek and K. Lammertsma, *Chem. Commun.*, 2002, **4**, 348–349.

