



Kent Academic Repository

Adetimehin, Adeyemi Daniel, Mole, Calvin Gerald, Finaughty, Devin A. and Heyns, Marise (2023) *Incidental observation of bone modification by Crematogaster cf. liengmei (Hymenoptera: Formicidae) in Cape Town, South Africa*. Forensic science, medicine, and pathology . ISSN 1556-2891.

Downloaded from

<https://kar.kent.ac.uk/102947/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1007/s12024-023-00714-2>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Incidental observation of bone modification by *Crematogaster* cf. *liengmei* (Hymenoptera: Formicidae) in Cape Town, South Africa

Adeyemi Daniel Adetimehin¹ · Calvin Gerald Mole¹ · Devin Alexander Finaughty^{2,3} · Marise Heyns^{1,4}

Accepted: 1 September 2023
© The Author(s) 2023

Abstract

Different taxa of insects have been reported to modify the skeletal remains of vertebrates during feeding or the creation of their larval and pupal chambers. Anecdotal reports on the modification of skeletal remains by ants exist but are rare. In this paper, a case of modification of bone due to the feeding activity of the ant, *Crematogaster* cf. *liengmei* (Hymenoptera: Formicidae), is reported. In November 2022, a bone was encountered incidentally in the vicinity of the Table Mountain National Park. The bone had been colonized by several individuals of *Crematogaster* cf. *liengmei*. Several ants were observed feeding on and removing bone particles and soft tissue remnants. A closer observation revealed striae/furrows on the epiphyseal ends of the bone. A 60-kg pig cadaver used in a winter decomposition trial in the same area was similarly colonized by this ant species, resulting in soft-tissue modification. Due to the similarity in the striae/furrows observed on the bone and the bitemarks inflicted by the ants on the right ear of the pig cadaver in the vicinity, it was concluded that the striae/furrows seen on the bone were created by the ants. Our observations show that *Crematogaster* cf. *liengmei* is an important taphonomic bio-agent that can contribute to the modification of skeletal remains in terrestrial environments.

Keywords Ants · Forensic entomology · *Crematogaster* cf. *liengmei* · Insect scavenging · Taphonomy

Case report

In November 2022, during routine data collection for an ongoing decomposition and cadaver entomofauna successional study in Table Mountain National Park, a bone was incidentally encountered in the vicinity of the study site (Fig. 1). Upon inspection, it was observed that the bone was colonized by several ants (Fig. 1). A 3-min undisturbed visual observation followed by a short video recording (Online Resource 1) and photography demonstrated active

feeding by the ants on the bone. A closer inspection of the bone revealed several striae/furrows at the epiphyseal ends, suspected to originate from the feeding activities of the ants (Figs. 2 and 3). In addition, a few individual ants were observed removing some bone particles and/or soft tissue remnants away from the bone (Fig. 3).

Crematogaster cf. *liengmei* (Hymenoptera: Formicidae) individuals

The ants were identified as *Crematogaster* cf. *liengmei* with the assistance of a local ant specialist using morphological descriptions in Fisher and Bolton [1]. Although identification of the ants to the species level would have been ideal, this is sometimes impossible due to the lack of morphological descriptions and occurrence of morphological variability in insects such as ants [2, 3]. Thus, the abbreviation “cf.” (*confer*, which means “compare to”) indicates the identification is provisional [4, 5].

The necrophagous behavior of this ant has previously been reported on neonate pig cadavers at the same study site [6]. The morphology of the striae/furrows observed on

✉ Adeyemi Daniel Adetimehin
yemiadetimehin@gmail.com

¹ Division of Forensic Medicine and Toxicology, Department of Pathology, University of Cape Town, Cape Town, South Africa

² Division of Natural Sciences, School of Chemistry and Forensic Science, University of Kent, Canterbury, UK

³ Division of Clinical Anatomy and Biological Anthropology, Department of Human Biology, University of Cape Town, Cape Town, South Africa

⁴ School of Medicine, Faculty of Life and Health Sciences, Ulster University, Derry/Londonderry, UK

Fig. 1 A bone encountered incidentally in the vicinity of the Table Mountain National Park



Fig. 2 *Crematogaster* cf. *liengmei* individuals aggregating and feeding on the fleshy remnants and bone particles on the anterior part of the bone



the bone bears some similarity to the bite marks seen on the external part of the right ear (Figs. 4 and 5) of an adult pig cadaver, a day after colonization by *Crematogaster* cf. *liengmei*. Due to this similarity, we concluded that the striae/furrows on the bone were created by this same ant species.

Discussion

The activities of vertebrate scavengers (e.g., vulture, mongoose, raccoon, domestic dog, feral cat, genet, jackal,

Fig. 3 *Crematogaster* cf. *liengmei* individuals aggregating, feeding on, and removing (yellow arrows) fleshy remnants and bone particles on the posterior end of the bone



Fig. 4 *Crematogaster* cf. *liengmei* individuals aggregating and feeding on the flesh on the external part of the adult pig's right ear (yellow arrow) as early as day 1 after cadaver deployment



porcupine, civet, warthog, and rat amongst many others) on vertebrate remains have been extensively investigated and documented in various parts of the world [7–19]. Scavenging activities have the potential to cause significant modification to bodies exposed in the environment,

including artefactual disarticulation, and scattering of remains. Such activity alters the rate of decomposition and induces bone modification that may be misconstrued as human-induced ante/peri-mortem trauma.

Fig. 5 Bitemarks (yellow arrow) inflicted by *Crematogaster* cf. *liengmei* on the external part of the adult pig's right ear on day 2



Invertebrate insect scavengers such as ants have garnered far less attention. Yet their activities on and around vertebrate remains have forensic implications [6, 20, 21]. Globally, several species of ants have been reported to prey on the immature (eggs, larvae, and pupae) and adult stages of other forensically important insects [20–22]. Additionally, they alter soft tissues causing artefacts and hemorrhage, creating sites for adult blow fly oviposition [6, 20–24]. Finally, they prevent fly landing and egg laying on decomposing vertebrate remains [6, 20–22]. These necrophagous and predatory behaviors by ants have been suggested to alter the decomposition of vertebrate remains and the estimation of the minimum time since death when using entomological evidence [20, 21]. Specifically, several members of the genus *Crematogaster* have been reported on decomposing human and animal remains [20, 21] and have been observed to create skin artefacts and alter entomofaunal interactions with a cadaver as described above [25–27]. While several reports exist on the necrophagous and predatory behaviors of various ant species, to the best of our knowledge, only one paper exists on the impact of ants on the skeletal remains of vertebrates [3].

Other invertebrates, including members of the insect orders Isoptera (termites), Coleoptera (beetles), Lepidoptera (moths), and Hymenoptera (wasps and bees), have been documented to feed on and modify the skeletal remains of vertebrates for nutritional purposes and/or the creation of their larval and pupal chambers [28–35]. However, previous reports on the modification of skeletal remains by ants

have been anecdotal and speculative [29, 31]. Go [3] is the first author to provide empirical evidence on the contribution of ants as taphonomic bio-agents on vertebrate skeletal remains. During analysis of the skeletal remains of an individual recovered from the Manila North Cemetery in the Philippines, several individuals of *Nylanderia* species (Hymenoptera: Formicidae) were observed nesting in the skeletal remains. Furthermore, post-mortem skeletal alteration in the form of tiny holes, scalloped edges, and shallow striae on several parts of the skeleton (i.e., ankles, tibia, fibula) were attributed to the activities (e.g., gnawing) of the ant species.

The observed striae/furrows on the bone in this study are superficially similar in morphology to the bitemarks/furrows inflicted by some vertebrate [see e.g., 16, 36, 37], and invertebrate (i.e., tenebrionid beetles) [31] scavengers on skeletal remains. These striae/furrows can potentially be misinterpreted as vertebrate scavenging, human-inflicted ante/peri-mortem trauma, or physicochemical weathering. Our incidental observation on the activities of *Crematogaster* cf. *liengmei* provides further evidence on the impact of ants on skeletal remains.

Limited information is available on the biology, ecology, and foraging behavior of *Crematogaster* cf. *liengmei* [6]. Generally, members of the genus *Crematogaster* are known to be tree- and ground-dwelling ants with generalized and omnivorous feeding habits [2, 6, 25]. Their generalized feeding habits may explain why they are able to colonize and feed on the soft tissues of the cadaver and

cadaver-associated entomofauna [6]. In addition, they often establish multiple nesting sites in terrestrial environments [2, 6, 25], the proximity of which dictates their ability to immediately colonize vertebrate remains. Above all, the exploitation of vertebrate remains as sources of nutrients by ants, including *Crematogaster* species may be linked to the size and nutritional status of the nests, age and previous foraging experience of the nest inhabitants, and quality of food [6, 25, 38–40].

Congruent with the findings of our previous study [6], we attribute the occurrence of several minute striae/furrows on the epiphyseal ends of the bone to the release of formic acid and other glandular chemicals secreted during the feeding activity of ants. As previously suggested, forensic pathologists, anthropologists, paleobiologists, crime scene investigators, and archaeologists should take cognizance of the presence of ants around skeletal remains of buried and surface decomposing human/animal cadavers. The feeding activity of ants can create striae, furrows, and/or edge gnawings that can mimic vertebrate and invertebrate (e.g., termites and beetles) scavenging, or human-inflicted ante/peri-mortem trauma. It is worth noting in this paper that the morphological state and integrity of the bone prior to its incidental discovery was unknown. Also, the striae/furrows on the bone were observed macroscopically without magnification or microscopic analysis. For these reasons, the possibility of interference by other vertebrate scavengers (e.g., Cape grey mongoose) as documented in Spies et al. [11] and other invertebrate scavengers (e.g., beetles) on the bone prior to our observations cannot be entirely ruled out. Consequently, future field and laboratory-based studies incorporating macroscopic and microscopic analyses will be conducted to provide additional information about the bone modification performed by ants and other invertebrates. This information will be useful in forensic, anthropological, and archaeological investigations.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12024-023-00714-2>.

Acknowledgements We thank the Authorities of the South African National Parks and Table Mountain National Park for granting us access and hosting the winter decomposition and cadaver entomofauna successional study in their premises. We thank Mr. John Morris (Farm Manager: Mariendahl Experimental Farm, Stellenbosch University) for providing the adult pig used in the winter decomposition and cadaver entomofauna successional study. We thank Dr. Jean-Bernard Huchet of the CNRS Muséum National d'Histoire naturelle, Paris, France for his revision and helpful comments on the earlier version of the manuscript and for providing additional bibliographic references. We are grateful to the South African National Research Foundation and University of Cape Town for providing Adeyemi Daniel Adetimehin with the Grantholder Student-Linked Bursary, International Students' Scholarship, JW Jagger Centenary Gift Scholarship, and Postgraduate Research Training Grant for his academic programme. The financial assistance of the National Research Foundation towards this research is hereby acknowledged. Opinions expressed and conclusions arrived

at are those of the authors and are not necessarily to be attributed to the National Research Foundation.

Author contribution Conceptualization: Adeyemi Daniel Adetimehin, Marise Heyns, and Devin Alexander Finaughty; Methodology: Adeyemi Daniel Adetimehin, Marise Heyns, and Devin Alexander Finaughty; Formal analysis, investigation, and visualization: Adeyemi Daniel Adetimehin; Writing—original draft preparation: Adeyemi Daniel Adetimehin; Writing—review and editing: Adeyemi Daniel Adetimehin, Marise Heyns, Devin Alexander Finaughty, and Calvin Gerald Mole; Funding acquisition and resources: Marise Heyns; Project administration: Marise Heyns and Calvin Gerald Mole; Supervision: Marise Heyns, Devin Alexander Finaughty, and Calvin Gerald Mole.

Funding Open access funding provided by University of Cape Town. This study was funded by the South African National Research Foundation (NRF) through a Research Grant awarded to Dr. Marise Heyns (grant number: CSUR116299).

Data availability All data generated or analyzed in relation to this study are included in this published article and its supplementary information file.

Declarations

Ethical approval Ethical approval for the decomposition and cadaver entomofauna successional studies in the Table Mountain National Park was granted by the University of Cape Town, Faculty of Health Sciences Animal Ethics Committee (UCT FHS AEC Reference number: 021_021). Approval to conduct the study in the Table Mountain National Park was also obtained from the authorities of the Table Mountain National Park (Permit number: CRC/2022-2023/024--2019/V1).

Conflict of interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Fisher BL, Bolton B. Ants of Africa and Madagascar, A Guide to the Genera. Berkeley: University of California Press; 2016.
2. Blaimer BB. Taxonomy and natural history of the *Crematogaster* (*Decracema*)-group (Hymenoptera: Formicidae) in Madagascar. *Zootaxa*. 2010;2714:1–39.
3. Go M. A case of human bone modification by ants (Hymenoptera: Formicidae) in the Philippines. *Forensic Anthropology*. 2018;1(2):117–23.
4. Bengtson P. Open nomenclature. *Palaeontology*. 1988;31(1):223–7.

5. Sigovini M, Keppel E, Tagliapietra D. Open Nomenclature in the biodiversity era. *Methods in Ecology and Evolution*. 2016;7(10):1217–25.
6. Adetimehin AD, Mole CG, Finaughty DA, Heyns M. Caught in the act: impact of *Crematogaster cf. liengmei* (Hymenoptera: Formicidae) necrophagous behavior on neonate pigs (*Sus scrofa domestica* L.) in the Western Cape Province of South Africa. *International Journal of Legal Medicine*. 2022; <https://doi.org/10.1007/s00414-022-02835-9>.
7. Reeves NM. Taphonomic effects of vulture scavenging. *Journal of Forensic Sciences*. 2009;54(3):523–8.
8. Smith JK. Raccoon scavenging and the taphonomic effects on early human decomposition and PMI estimation. Dissertation: University of Tennessee; 2015.
9. Steadman DW, Daurtartas A, Kenyherc MW, Jantz LM, Mundorff A, Vidoli GM. Differential scavenging among pig, rabbit, and human subjects. *Journal of Forensic Sciences*. 2018;63(4):1684–91.
10. Spies MJ, Gibbon VE, Finaughty DA. Forensic taphonomy: vertebrate scavenging in the temperate southwestern Cape. *South Africa Forensic Science International*. 2018;290:62–9.
11. Spies MJ, Finaughty DA, Gibbon VE. Forensic taphonomy: scavenger-induced scattering patterns in the temperate southwestern Cape, South Africa—a first look. *Forensic Science International*. 2018;290:29–35.
12. Spies MJ, Finaughty DA, Friedling LJ, Gibbon VE. The effect of clothing on decomposition and vertebrate scavengers in cooler months of the temperate southwestern Cape. *South Africa Forensic Science International*. 2020;309:110197.
13. Dibner H, Valdez CM, Carter DO. An experiment to characterize the decomposer community associated with carcasses (*Sus scrofa domestica*) on Oahu. *Hawaii Journal of Forensic Sciences*. 2019;64(5):1412–20.
14. Garcia S, Smith A, Baigent C, Connor M. The scavenging patterns of feral cats on human remains in an outdoor setting. *Journal of Forensic Sciences*. 2020;65(3):948–52.
15. Keyes CA, Myburgh J, Brits D. Animal scavenging on pig cadavers in the Lowveld of South Africa. *Forensic Science International*. 2021;327:110969.
16. Keyes CA, Myburgh J, Brits D. Scavenger activity in a peri-urban agricultural setting in the Highveld of South Africa. *International Journal of Legal Medicine*. 2021;135:979–91.
17. Keyes CA, Myburgh J, Brits D. Identifying forensically relevant urban scavengers in Johannesburg. *South Africa Science & Justice*. 2022;62(3):399–409.
18. Flint CA, Sawyer SJ, Rhinesmith-Carranza J, Tomberlin JK. Rodent scavenging of pig remains potentially increases oviposition sites for primary colonizers. *Journal of Forensic Sciences*. 2022;67(4):1728–33.
19. Indra L, Errickson D, Young A, Löscher S. Uncovering forensic taphonomic agents: animal scavenging in the European context. *Biology*. 2022;11(4):601.
20. Campobasso CP, Marchetti D, Introna F, Colonna MF. Postmortem artifacts made by ants and the effect of ant activity on decomposition rates. *American Journal of Forensic Medicine and Pathology*. 2009;30:84–7.
21. Eubank MD, Lin C, Tarone AM. The role of ants in vertebrate carrion decomposition. *Food Webs*. 2019;18: e00109.
22. Paula MC, Morishita GM, Cavarson CH, Gonçalves CR, Tavares PRA, Mendonça A, Suárez YR, Antonialli-Junior WF. Action of ants on vertebrate carcasses and blow flies (Calliphoridae). *Journal of Medical Entomology*. 2016;53(6):1283–91.
23. Ventura F, Gallo M, De Stefano F. Postmortem skin damage due to ants. *American Journal of Forensic Medicine and Pathology*. 2010;31:120–1.
24. Meyer F, Monroe MD, Williams HN, Goddard J. *Solenopsis invicta x richteri* (Hymenoptera: Formicidae) necrophagous behavior causes post-mortem lesions in pigs which serve as oviposition sites for Diptera. *Forensic Science International: Reports*. 2020;2:100067.
25. Longino GT. The *Crematogaster* (Hymenoptera, Formicidae, Myrmecinae) of Costa Rica. *Zootaxa*. 2003;151:1–150.
26. Bonacci T, Brandmayr TZ, Brandmayr P, Vercillo V, Porcelli F. Successional patterns of the insect fauna on a pig carcass in southern Italy and the role of *Crematogaster scutellaris* (Hymenoptera, Formicidae) as a carrion invader. *Entomological Science*. 2011;14(2):125–32.
27. Bugelli V, Forni D, Bassi LA, Di Paolo M, Marra D, Lenzi S, Toni C, Giusiani M, Domenici R, Gherardi M, Vanin S. Forensic entomology and the estimation of the minimum time since death in indoor cases. *Journal of Forensic Sciences*. 2015;60(2):525–31.
28. Pittoni E. Necropoli di Pill'e Matta Quartucciu (Cagliari, Sardinia): wild bee and solitary wasp activity and bone diagenetic factors. *International Journal of Osteoarchaeology*. 2009;19(3):386–96.
29. Backwell LR, Parkinson AH, Roberts EM, d'Errico F, Huchet J-B. Criteria for identifying bone modification by termites in the fossil record. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 2012;337:72–87.
30. Backwell L, Huchet J-B, Harrison JDG, D'errico F. Invertebrate modification of bone. In: Pokines JT, Symes SA, L'Abbe EN, editors. *Manual of Forensic Taphonomy*. 2nd ed. New York: CRC Press; 2022. p. 631–66.
31. Holden AR, Harris JM, Timm RM. Paleocological and taphonomic implications of insect-damaged Pleistocene vertebrate remains from Rancho La Brea, southern California. *PLoS ONE*. 2013;8(7): e67119.
32. Huchet J-B, Le Mort F, Rabinovich R, Blau S, Coqueugniot H, Arensburg B. Identification of dermestid pupal chambers on Southern Levant human bones: inference for reconstruction of Middle Bronze Age mortuary practices. *Journal of Archaeological Science*. 2013;40(10):3793–803.
33. Huchet J-B. Approche ichnologique et taphonomique des altérations ostéolytiques dues aux insectes en contexte archéologique. In: Denys C, Patou-Mathis M, editors. *Manuel de Taphonomie*. Actes Sud-Errance (éd), MNHN & CNRS; 2014a. pp. 185–207.
34. Huchet J-B. L'archéontologie funéraire. In: Charabidze D, Goselin M, editors. *Insectes, cadavres et scènes de crime, Principes et applications de l'entomologie médico-légale*. Editions De Boeck; 2014b. pp. 201–224.
35. Viero A, Montisci M, Pelletti G, Vanin S. Crime scene and body alterations caused by arthropods: implications in death investigation. *International Journal of Legal Medicine*. 2019;133:307–16.
36. Moraitis K, Spiliopoulou C. Forensic implications of carnivore scavenging on human remains recovered from outdoor locations in Greece. *Journal of Forensic and Legal Medicine*. 2010;17(6):298–303.
37. Young A, Marquez-Grant N, Stillman R, Smith MJ, Korstiens AH. An investigation of red fox (*Vulpes vulpes*) and Eurasian badger (*Meles meles*) scavenging, scattering, and removal of deer remains: forensic implications and applications. *Journal of Forensic Sciences*. 2015;60:S39–55.
38. Carroll CR, Janzen DH. Ecology of foraging by ants. *Annual Review of Ecology and Systematics*. 1973;4:231–57.
39. Traniello JFA. Foraging strategies of ants. *Annual Review of Entomology*. 1989;34:19–210.
40. Nooten SS, Chan KH, Schultheiss P, Bogar TA, Guénard B. Ant body size mediate functional performance and species interactions in carrion decomposer communities. *Functional Ecology*. 2022;00:1–13.

This work is not currently under consideration or published in another journal.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.