



Kent Academic Repository

Edrich, Elizabeth, Duvenage, Lucian and Gourlay, Campbell W. (2024) *Alternative Oxidase – Aid or obstacle to combat the rise of fungal pathogens?* *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1865 (2). ISSN 0005-2728.

Downloaded from

<https://kar.kent.ac.uk/104571/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1016/j.bbabi.2024.149031>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Alternative Oxidase – Aid or obstacle to combat the rise of fungal pathogens?

Elizabeth S.M. Edrich^a, Lucian Duvenage^b, Campbell W. Gourlay^{a,*}

^a Kent Fungal Group, School of Biosciences, University of Kent, Kent CT2 9HY, UK

^b CMM AFRICA Medical Mycology Research Unit, Department of Pathology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

ARTICLE INFO

Keywords:

Alternative Oxidase
Mitochondria
Fungi
Plants
Pathogen

ABSTRACT

Fungal pathogens present a growing threat to both humans and global health security alike. Increasing evidence of antifungal resistance in fungal populations that infect both humans and plant species has increased reliance on combination therapies and shown the need for new antifungal therapeutic targets to be investigated. Here, we review the roles of mitochondria and fungal respiration in pathogenesis and discuss the role of the Alternative Oxidase enzyme (Aox) in both human fungal pathogens and phytopathogens. Increasing evidence exists for Aox within mechanisms that underpin fungal virulence. Aox also plays important roles in adaptability that may prove useful within dual targeted fungal-specific therapeutic approaches. As improved fungal specific mitochondrial and Aox inhibitors are under development we may see this as an emerging target for future approaches to tackling the growing challenge of fungal infection.

1. Introduction

Fungal pathogens are responsible for over one billion human infections and over 1.6 million deaths annually [1–4], as well as a third of all global crop failures [1,5]. However, despite the threat that fungal strains present to animal species and the threat to food security worldwide, they remain under-researched [3] [6].

Human fungal pathogens can cause superficial, sub cutaneous or systemic infections, which in the case of immunocompromised individuals can be associated with high mortality. The outcome for patients is also strongly correlated with the speed and accuracy of diagnosis, a rise in antifungal resistance and socioeconomic factors that restrict treatment ability [7–9]. Antifungal resistance is likely to become a major issue, driven by factors such as a limited number of identified cellular targets for antifungal development, the over-use of agricultural fungicides and the emergence of more fungal pathogens in an ever-warming climate [10,11].

There is therefore the need to conduct research into new antifungal targets and strategies to prevent fungal infection. Mitochondria could prove to be a useful target against plant fungal pathogens [12] and growing evidence, driven by an increase in our understanding of respiratory chain physiology, suggests that inhibitors may also be developed to tackle human infection [13,14].

2. Fungal respiration

As with many eukaryotes, fungal pathogens possess a well conserved classical Electron Transport Chain (ETC) which is used to generate a proton motive force (PMF) that can drive ATP synthesis, and which is important for the numerous processes that mitochondrial function supports. Forward electron transfer through the respiratory complexes (FET) provides the thermodynamically favoured reaction that is coupled to oxidative phosphorylation and ATP production. However reverse electron transfer (RET) can occur, whereby electrons flow backwards through Complex I. RET is thought to be induced by reduction of the Ubiquinone pool (UQP) to between 40 and 60 %, requiring a high PMF, a large thermodynamic driving force, and a high ΔpH [15,16].

Sites for mitochondrial Reactive Oxygen Species (ROS) production, namely superoxide (O_2^-) generation, include Complex I and Complex III, particularly sites I_F , I_Q and III_{QO} [15]. Site I_Q is responsible for most of the superoxide production during RET, whereby electrons are forced into Complex I through a high QH_2 / Q ratio and high PMF [15]. Superoxides can be generated by both FET and RET [15], whereby production and removal depend on substrate availability (like succinate oxidation, which drives RET [17]) QH_2/Q ratio, rate of oxygen consumption, and mitochondrial dysfunction or inhibition [15]. The multifactorial nature of ROS generation and management requires

* Corresponding author.

E-mail address: C.W.Gourlay@kent.ac.uk (C.W. Gourlay).

<https://doi.org/10.1016/j.bbabio.2024.149031>

Received 1 August 2023; Received in revised form 16 November 2023; Accepted 3 January 2024

Available online 7 January 2024

0005-2728/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

continual homeostasis, as ROS are essential within diverse cellular processes in both host and pathogen [18], but highly inflated ROS levels have been attributed to mitochondrial disease [19]. Induction of ROS production has also been implicated in the mode of action of several antifungals [20,21] suggesting a role for mitochondria and/or redox homeostasis as a target in current therapeutic approaches. However, understanding the full signalling mechanisms of ROS within fungal pathogenesis represents a growing field of study.

Intuitively the conservation of the highly conserved classical ETC provides a barrier to its development as an attractive target for human fungal pathogens. However, fungal specific differences that are essential for function have been reported, such as alternative NADH dehydrogenases, which may bypass Complex I activity [22,23]. One key difference in the respiratory chain of many human fungal pathogens is the presence of a cyanide-insensitive alternative oxidase (Aox), which is not found in mammalian mitochondria. Aox branches from the main respiratory chain at the level of the UQP (Fig. 1) and has a catalytic di-iron centre orientated towards the mitochondrial matrix. Interestingly, this membrane-bound oxidase is non-proton motive, and therefore does not have a significant role in ATP production, but rather oxidises ubiquinol and reduces oxygen to water, bypassing the ETC prior to proton translocation by complexes III and IV (Fig. 1). However, there is evidence for ATP generation through the Aox/Complex I pathway in *Botrytis cinerea* [22,24], and inhibition of Aox in *Gaeumannomyces graminis* leads to a decreased rate of ATP synthesis [25], indicating that the

extent of Aox involvement in ATP generation may be species specific.

While Aox is not found in mammals, it is highly conserved amongst pathogenic fungi. Some fungal species like *Aspergillus niger*, *Aspergillus flavus* and *Candida albicans* have multiple isoforms [26] [27], there are notable exceptions, for example in *Candida glabrata* and *Pneumocystis jiroveci*, which, like *Saccharomyces cerevisiae*, do not contain a multi-subunit Complex I or Aox. It may be the case that Crabtree positive yeasts, that primarily use aerobic fermentation for increased growth rates under high glucose availability [28], do not require an Aox for mitochondrial homeostasis, as growth is predominantly supported by fermentation rather than respiration. Crabtree negative yeast, that utilise respiration for energy generation such as *Candida albicans* and *Cryptococcus neoformans*, have retained Aox and maintain metabolic flexibility by utilising both alternative and glucose carbon sources. This flexibility may contribute to host colonisation and virulence in the nutrient-scarce host [29–32], whereby presence of Aox in fungal pathogens may also assist in the maintenance of mitochondrial function upon immune challenge, the regulation of ROS production, and pathogenesis [33–35].

Interestingly, fungi have been shown to utilise multiple respiratory pathways, for example *Aspergillus nidulans* can alternate between both classical and alternative respiratory pathways to generate sterigmatocystin, a precursor to Aflatoxin B₁ [26,36]. A third ‘parallel’ respiratory chain (PAR), in *Candida albicans* and *Candida parapsilosis* has been proposed [37–40], which may contribute up to 10 % of total




Fig. 1. Schematic of the Electron Transport Chain (ETC) found in fungal pathogens.

Both a classic ETC and an alternative respiratory chain are found in most fungi. Where known, crystal structures of the ETC components are shown [Complex I, Protein Data Bank (PDB) 3M9S [172], Complex II PDB 3VR8 [173], Trypanosomal Aox PDB 3VV9 [174], Complex III PDB 1KY0 [175], Complex IV PDB 8DH6 (to be published), ATP Synthase PDB 1QO1 [176]]. Aox branches from the main respiratory chain at the level of the UQP. This pathway produces little ATP, but instead dissipates energy as heat and bypassing proton transfer through downstream Complexes III and IV. Known inhibitors of each ETC component are listed in red. Created with BioRender.com. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

respiration capacity. The ability to utilise alternate routes of respiration suggests that these yeasts may have evolved within an environment that presents regular and significant challenges to electron transport. Alternatively, it may be that alternate respiration supports signalling and metabolic changes that are required for adaptability within changing environments. This may include factors produced by competing or co-colonising microbes that damage or inhibit growth [41], such as phenazine production by *Pseudomonas aeruginosa* [42], or detoxification of nitric oxide (NO) released by host cells during infection [43,44]. This could also involve nutritional challenges that limit access to key metabolites, co-factors that support respiration or environmental challenges such as temperature variation or oxygen availability [41,45].

3. The role of mitochondria in fungal pathogenicity

An increasing interest in fungal respiration has highlighted several mitochondrial roles which are thought to contribute to human fungal pathogenicity [46], although some of the signalling mechanisms behind this involvement are still unclear. Respiration deficiency leads to diminished virulence in mice intravenously infected with *Candida albicans* [47], and respiratory inhibition with sodium nitroprusside (SNP), an NO donor, and salicylhydroxamic acid (SHAM), an Aox inhibitor, reduces *Candida albicans* cell viability and increases phagocytic uptake by macrophages [37]. Increased phagocytosis has also been seen after treatment of *Candida albicans* with the Complex III inhibitor Antimycin A, thought to be attributable to β -glucan exposure [48]. However, pre-treatment of the cells with SHAM and SNP prior to murine infection exhibited increased virulence through transcriptional changes and cell wall remodelling, presenting a higher renal fungal burden, and increased immune infiltrate than untreated cells [37]. This data therefore indicates that mitochondrial signalling mechanisms need to be investigated further.

Active infection and dissemination inside the host is presumed to be energetically demanding for pathogenic yeasts, especially for switches to hyphal growth forms for *Candida albicans* via the Ras1/cAMP/PKA signalling pathway [49,50]. Other instances of increased respiratory demand for ATP in *Candida albicans* include escape from macrophage engulfment through catabolism of amino acids for morphogenesis [51]. This increased ATP demand could be attributed to the action of ATP-driven pumps belonging to the ATP Binding Cassette family (ABC). ABC transporters are profound influx-efflux pumps that rely on ATP hydrolysis to transport a variety of molecules, including sterols, metabolites, and drugs, proving crucial for pathogenic activity of *Magnaporthe grisea* [52,53], and are thought to contribute to the multi-drug resistant phenotype seen in *Candida auris* [54]. As well as heavy reliance on ATP production, Vacuolar ATPases (V-ATPases) generate a pH gradient through ATP hydrolysis which drives secondary transporters to maintain cellular ion homeostasis. This process requires Ergosterol for optimal V-ATPase function and is sensitive to combination treatment with both Fluconazole and Amiodarone in the *C. albicans* murine candidiasis model [55]. These investigations prove that ATP is crucial for a variety of processes that underpin fungal pathogenesis, although this could be in combination with other mitochondrial signalling pathways.

The maintenance of mitochondrial morphology for stress tolerance and virulence is highlighted in *Cryptococcus neoformans*, whereby fully functioning inherited mitochondria from the *MATa* parent are critical for growth under fluctuating temperature, low oxygen availability and iron regulation [56–60]. Defects occurring in Complex I, II or IV can impair conidiation and sexual development of the yeasts *Neurospora crassa* and *Podospora anserina*, even when alternative NADH dehydrogenases are present and respiration is maintained by Aox [61–69]. Interestingly, a mutation in the *Cryptococcus neoformans* NADH promoter region increased production of melanin, Glucuronoxylomannan (GXM) release, and ATP, virulence enhancing traits which occurred through serial passage in the *Galleria mellonella* model [70]. These

studies suggest that while conservation of mitochondrial components and morphology is seen across multiple taxa, mitochondrial plasticity may have a role in enhancing virulence and host evasion.

Mitochondrial morphology is thought to have importance in the ER-mitochondria encounter structure (ERMES), which has been highlighted for cell fitness, immune evasion, and virulence in both *Candida albicans* and *Aspergillus fumigatus* alike [71]. Mitochondrial contact sites with both the ER and peroxisomes have been thought to contribute to lipid homeostasis through shuttling of tricarboxylic-acid (TCA) cycle intermediates like citrate from peroxisomes to the mitochondria, although details of metabolite transfer and regulation of contact sites is still unclear. Interestingly, although direct lipid transit pathways are yet to be elucidated, a recent study by Enkler *et. al* [72] suggested that Arf1 couples fatty acid β -oxidation to mitochondrial ATP synthesis and can regulate mitochondrial fission and fusion [72]. Adequate regulation of mitochondrial fission and fusion mechanisms is important in *Cryptococcus neoformans*, whereby mitochondrial fusion defects lead to increased ETC inhibitor sensitivity and loss of virulence in a murine model [57], and mitochondrial fragmentation seen in *Aspergillus fumigatus* is seen during human granulocyte killing as a response to oxidative stress [73]. The function of mitochondria in fungal pathogenesis is multi-factorial, with ATP production, organelle signalling and morphology underpinning virulence mechanisms. However, deeper investigations into virulence signalling pathways involving mitochondria should be explored.

4. Aox function in plant fungal pathogens

The ETC of plants has been extensively studied [74], and Aox activity has been documented in both plants and phytopathogenic fungi alike. However, evidence shows that the ETC of plants differs to that of other eukaryotes due to the number of subunits found for each mitochondrial complex. For example, the Complex I of plants has nearly 50 different subunits [75], and studies of *Pichia stipitis* and *Neurospora crassa* show that fungal Aox differs from Aox in plants in that it occurs as a monomer and is not induced by α -keto acids such as pyruvate [76]. The disparity between plant and fungal complex subunits, including additional proteins found in plant complexes [77,78] is thought to assist in antifungal therapies that target phytopathogenic respiration.

In plant fungal pathogens such as *Moniliophthora perniciosa* and *Sclerotinia sclerotiorum*, Aox is reported to be more active during the mycelial growth phase, suggesting that the metabolic control provided by alternative respiration is a crucial factor in morphogenesis [79,80]. Interestingly, the activation of Aox for fungal growth has also been thought to contribute to mycotoxin production by food-colonising fungi, such as Aflatoxins produced by *Aspergillus flavus* [81–83]. It is interesting to note that Aox is activated and upregulated in *Solanum lycopersicum*, *Arabidopsis thaliana* and *Nicotiana attenuata* in response to bacterial and viral attack, mainly for oxidative and nitrosative stress management [84–88], although the roles of Aox in stress signalling during fungal infection of other plant species requires further research.

For other plant pathogens, such as *Botrytis cinerea*, *Ustilago maydis* and *Magnaporthe grisea*, Aox is required for active resistance to Quinone Outside Inhibitor (Q_oI) fungicides such as the strobilurins, Azoxystrobin and Pyraclostrobin [89–92]. This class of fungicides inhibit mitochondrial respiration through binding to Cytochrome *bc*₁, blocking the movement of electrons at the quinone outer binding site [93], although field resistance is becoming an increasing problem, such as in *Mycosphaerella fijiensis* and *Mycosphaerella musicola* infections of bananas [94] and *Pyrenophora tritici-repentis* infections of Argentinian wheat [95]. Increasing Q_oI resistance has been attributed to the presence of an Aox in *Mycosphaerella graminicola* and *Aspergillus flavus* [26,96], whereby Aox provides an alternative route for electron transport away from the target site of the Q_oI fungicides to maintain electron flux. In the presence of the strobilurin Azoxystrobin, *Fusarium graminearum* upregulated transcription of Aox and rapidly increased oxygen uptake [97]. To address the

emerging antifungal resistance in agricultural practices, research into new succinate dehydrogenase inhibitors (SDHIs), such as Carboxin has begun, whereby fungal respiration is inhibited through blockage of the ubiquinone binding sites of Complex II. SDHIs have been rapidly gaining interest due to their broad, high antifungal activity [98], however phytopathogenic sensitivity to SDHIs is slowly shifting [99] and evidence suggests that SDHI site-specific inhibition may give rise to resistance if not monitored correctly [100]. This may, in part, be due to the presence of fungal Aox at the level of the UQP to provide an alternative respiratory pathway under this inhibition. The potential benefits of Aox inhibitors in the agrochemical industry has been recently reviewed [81].

Although Aox has proposed functions for virulence in plant fungal pathogens, not all phytopathogens have a predicted sequence, such as the *Puccinia* species responsible for wheat rust disease and *Melampsora lini* which causes flax rust (Fig. 2). Interestingly, while resistance mechanisms of wheat and flax towards these pathogens has been documented [101,102] there are few noted instances of antifungal resistance for these pathogens themselves, which, in conjunction with these other studies, indicates that Aox may have a role in phytopathogenic resistance to antifungal drugs. While much needs to be investigated in plant-pathogen interactions, given that both plants and phytopathogenic fungi can induce Aox independently for multi-factorial stress relief, one cannot rule out the possibility that both fungi and plants may use Aox in within the environmental niche of an active phytopathogenic infection. Investigations into the role of Aox on both the host and pathogen sides of an infection could provide an insight into tackling antifungal resistance impacting food security.

5. Aox function in human fungal pathogens

Most human fungal pathogens possess at least one Aox (Fig. 3). The importance of Aox in morphogenesis and resistance to oxidative stress

from the human host has been demonstrated in several fungal pathogens. Aox1 from *Cryptococcus neoformans* was shown to be induced at 37 °C and was reported to play a role in virulence in the murine inhalation model [103] and in *Paracoccidioides brasiliensis*, Aox is upregulated in response to oxidative stress and for the mycelial-to-yeast transformation, a crucial step in paracoccidioidomycosis [104,105]. *Aspergillus flavus* and *Aspergillus fumigatus* have multiple isoforms of Aox, whereby the isoform AoxA was found to attribute resistance to oxidative stress and macrophage killing [26,106,107]. Aox is also upregulated in *Candida albicans* and *Candida auris* in response to oxidative stress conditions [108–110], plays a role in hyphal growth and biofilm formation [111,112] and deletion of Aox in *Candida albicans* leads to increased Fluconazole susceptibility [113]. A recent study assessed *Candida albicans* respiratory capacity when exposed to SNP in combination with the known Aox inhibitor salicylhydroxamic acid (SHAM). *Candida albicans* treated with this combination displayed a rapid transition to hyphal growth upon relief from inhibition and *Candida albicans* treated with both SNP and SHAM also displayed a decrease in caspofungin resistance [37]. This indicated that Aox has a significant role in the hyphal switching phenotype in *Candida albicans*, and that transcription of a second alternative oxidase, Aox2, is induced in the presence of ETC inhibitors to buffer respiratory stress and increase alternative respiration capacity [37,114]. Interestingly, deletion of Aox2 also leads to decreased virulence of *Candida albicans* in the murine model through increased immune recognition [115]. However, some reports suggest that Aox1 is dispensable for virulence in *Candida albicans* [116] and *Aspergillus fumigatus* [106], this may be due to differences between experimental approaches or strain backgrounds and remains a point to be clarified.

Interestingly, both *Candida glabrata* and *Pneumocystis jiroveci* do not have a predicted Aox sequence (Fig. 3), however evidence of respiratory inhibition of *Pneumocystis jiroveci* with SHAM has been reported,




Fig. 2. Key plant fungal pathogens.

Schematic illustrating key plant fungal pathogens. Pathogenic species without a known or predicted Aox sequence in the UniProt database [177] are highlighted in red. Created with BioRender.com. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. Key human fungal pathogens.

Schematic illustrating key human fungal pathogens. Pathogenic species without a known or predicted Aox sequence in the UniProt database [177] are highlighted in red. Created with BioRender.com. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

suggesting an Aox may be present [117,118]. While incidences of *Candida glabrata* based Candidiasis are increasing, the genetic similarity of *Candida glabrata* to *Saccharomyces cerevisiae* and differences to *Candida albicans* indicate that both species must have evolved different routes to pathogenesis, independently of Aox [119]. Unfortunately, little is known about the metabolic requirements of *Pneumocystis jiroveci* for infection, although it has been postulated that *Pneumocystis* pneumonia is based on reactivation of a latent infection, using the host as an environmental reservoir to facilitate human-human transmission [120,121] which also appears Aox independent. The importance of Aox in virulence itself appears pathogen specific, as parasitic *Trypanosoma brucei* alternative oxidase (TAO) has proven to be essential for respiration in the bloodstream form of the parasite [122,123], with intense therapeutic interest providing the only known published crystal structure of Aox [124] and evidence of TAO inhibition with ascofuranone [125]. However, while interesting links have been made for Aox involvement in parasite infection, this falls out of the scope of this review.

Although human-fungi interactions are widely investigated, direct signalling roles for Aox in fungal pathogenesis and host dissemination are still unclear. Because of its ability to maintain respiration in the presence of classical ETC inhibitors, Aox in human fungal pathogens could be proposed to relieve respiratory stresses induced by the host during infection, such as NO. In humans, NO is produced during host infection by phagocytes by nitric oxide synthase iNOS (or NOS2) as part of the arsenal of oxidants that can inhibit or kill invading pathogens [126]. While no direct links between human NO production and fungal Aox during pathogenesis have been recorded, studies on plant mitochondria show that while NO inhibits cytochrome c oxidase, Aox remains uninhibited [127]. The evidence presented suggests that Aox does play a role in fungal pathogenesis, although most likely as an indirect

mediator of oxidative stresses induced by the host and maintenance of fungal morphology in the host environment, rather than a direct virulence component. Perhaps for human fungal pathogens, Aox activity should be seen as a latent, adaptive mechanism which activates under selective pressure for metabolic homeostasis. This would provide a background benefit to fungal pathogens in immune evasion but would be dispensable as an active component of virulence mechanisms required for dissemination in the host. However, further investigations into Aox activation and function within different human fungal pathogens is required to substantiate these postulations.

6. Problems and potential for targeting respiration in human fungal pathogens

Current antifungal therapies either inhibit biosynthetic machinery for ergosterols (e.g., azoles), nucleic acids (antimetabolites e.g., Flucytosine) or disrupt the cell membrane or wall (e.g., polyenes or echinocandins). While conservation of mitochondrial machinery between both animals and fungi presents a problem for suitable antifungal treatments, current inhibitors of the fungal ETC, highlighted in Fig. 1, include both natural and synthetic compounds. Rotenone and Metformin are known inhibitors of Complex I, investigated for their efficacy against cancer and antifungal properties against *Candida albicans* and *Aspergillus niger* [128–132], although Metformin is classically used to treat Type II Diabetes [133]. Complex II is prone to inhibition by 3-nitropropionic acid (3-NPA) or thenoyltrifluoroacetone (TTFA), although new studies have highlighted Gracillin, a natural steroidal saponin as a potential anti-tumour drug [134,135]. Numerous Complex III inhibitors have been identified, of which Strobilurins such as Azoxystrobin are rapidly gaining interest [83,136–138]. For example, combination treatment of

Complex III and Aox inhibition was found to enhance sensitivity to caspofungin in *C. parapsilosis* [139]. However, while Complex III inhibition is attractive for antifungal activity [13], inhibitors of this kind can often bind to both pathogenic and host complexes alike, so different applications for this inhibitor class are being investigated, such as cancer therapeutics [138,140–142]. While inhibition of yeast ATP synthase is possible using Bedaquiline [143] and Oligomycin [144], the strong conservation between fungal and mammalian isoforms of the mitochondrial ATP synthase renders this enzyme as an antifungal target obsolete, although it does provide a promising antibiotic target [145].

Prolonged reliance on treatment guidelines has presented increasing numbers of resistance in clinical cases [146–148]. Resistance patterns have begun to emerge in various fungal species, and reliance on combination therapies of azoles, polyenes, echinocandins and anti-metabolites mean that monotherapies are declining in potency against rapidly adapting fungal pathogens. For example, mutations in the ergosterol pathway and sterol biosynthesis of *Candida tropicalis*, *Candida albicans* and *Candida lusitanae* leads to resistance to azoles and Amphotericin B as monotherapies [149–154] and interestingly this mutation in *Candida lusitanae* *ERG3* is also induced via micafungin monotherapy, which then provides cross resistance to multiple antifungal classes [155]. Treatments are now more aimed towards combination therapy to target invasive fungal infections such as the use of Caspofungins with other antifungal classes to treat Candidiasis [156], combinations of multiple triazole types such as PC945 and voriconazole for treatment of *Aspergillus fumigatus* based Aspergillosis [157] and recommended treatment for Cryptococcosis includes a combination of Amphotericin B and Flucytosine, followed by Fluconazole in a maintenance program that can last up to a year [158,159], although antifungal hetero-resistance and cross-tolerance has been documented [160].

The increasing amount of antifungal resistance and the reliance on combination therapies means focus is shifting towards compounds that can re-sensitise resistant fungal pathogens to current antifungals, rather than identify novel antifungal compounds. Interestingly, a group of Indazole derivatives have been identified to convert azoles from fungistatic back to fungicidal for *Candida albicans* infections through inhibition of the cytochrome *bc*₁ complex [161]. One compound, Inz-5, enhanced the ability of macrophages to contain *C. albicans*, and Inz-1 showed selection for yeast complex *bc*₁ over human *bc*₁, which offsets the current issues other Complex III inhibitors face in host treatment. Other synergisms are beginning to emerge, such as Tetrandrine which increased the antifungal activity of Fluconazole in the murine Candidiasis model [162] and even a combination of Fluconazole with small molecule ENOblock [163] or SHAM [164] showed synergism against *Candida albicans*. With the indication that research is moving away from monotherapeutic antifungals, inhibition of Aox could be effective in combination therapies with existing antifungal drugs such as Echinocandins and Azoles which are currently circulated for the treatment of *Candidiasis* and *Cryptococcosis*, but not as a monotherapy alone. Another postulation includes the use of known natural stressors for pathogens such as NO, which can be produced by SNP, in combination with Aox inhibitors. While using NO stress in combination with Aox inhibition is an attractive proposal, research into efficacy against fungal pathogens in vitro and in vivo is required. Research into fungal respiration machinery, especially the role of Aox, and its links to virulence, remain understudied.

In summary, due to the connection of mitochondria to pathogenesis, cell wall regulation and lipid metabolism, fungal-specific respiratory inhibitors may prove to be effective against pathogens either in isolation or in combination with current antifungals. However, the conservation of the respiratory machinery in eukaryotes and the robust and adaptive nature of fungal respiration is a challenge for drug development, so investigation into compounds that can re-sensitise drug resistant fungal pathogens to existing therapies may also provide relief from infection. Characterisation of fungal-specific respiratory chain components are needed, together with a deeper understanding of the roles of those

already characterised, such as Aox.

7. Proposed roles for Aox

Studies so far suggest that Aox does not have a direct role in fungal virulence but does have a role in maintenance of oxidative stress mechanisms, ROS production, and even Complex I driven respiration in *Botrytis cinerea* [24,165]. It could be postulated that Aox, having close links and established electron transfer from the UQP, and yet no dominant role in ATP synthesis, could act as a switch between FET and RET in response to environmental cues for both ROS production and scavenging. Interestingly, this concept is supported by an investigation in murine mitochondria using xenotypic expression of Aox from *Ciona intestinalis* [166]. Aox is known to oxidise ubiquinol and reduces oxygen to water, bypassing the ETC prior to proton translocation by complexes III and IV, producing heat as a byproduct. It would make sense, therefore, if Aox acted in FET to stop electron leak through I_Q or III_{QO} by oxidation of quinone which could reduce ROS production in stress conditions induced by host dissemination. More interestingly, as RET induction requires an unfavourable thermodynamic force and high QH₂/Q ratio, one could speculate that Aox could contribute to RET induction itself through generation and release of heat energy as the RET driving force for electron movement into Complex I and maintenance of the reduced quinone state. While induction of RET and induction of ROS production seems counterproductive to fungal pathogens, certain morphological developments induce high ROS, such as capsule development in *Cryptococcus neoformans* [167] and *Aspergillus niger* and *P. penicillium* spore germination [168]. This, in conjunction with studies that show Aox as sensitive to pH changes [169] and other postulations highlighting Aox as important in ROS homeostasis [26] [170] [171] supports the idea that Aox could act a FET/RET switch to provide ROS for energy demanding virulence mechanisms in fungal pathogens or defend against host-generated ROS through induction of FET, independently of ATP production. Experiments investigating ROS production, FET/RET initiation and thermodynamics in relation to Aox activity should be considered.

8. Conclusion

While fungal pathogens present a growing threat to both human health and food security, research into antifungal therapies are still neglected. Here, we investigated the role of the electron transport chain in fungal virulence and antifungal resistance, including the Aox pathway. Interestingly, while the direct role of Aox in pathogenic virulence remains unclear, studies have showed a potential homeostatic role for metabolism under both biotic and abiotic stresses. This stress tolerance mechanism is thought to contribute to pathogenic survival in the host and contribute to current antifungal resistance through control of ROS and NO accumulation, which is uncoupled from ATP production. To address the issues faced by antifungal resistance, application of an Aox-specific antifungal may re-sensitise resistant fungal pathogens to drugs such as Fluconazole, although research into combination therapies and Aox-specific inhibitors still needs to be pursued.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

References

- [1] M.C. Fisher, N.A.R. Gow, S.J. Gurr, Tackling emerging fungal threats to animal health, food security and ecosystem resilience, *Philos. Trans. R. Soc. B* 371 (1709) (2016) 20160332.
- [2] B. Brown, D. Denning, G. Gow, L. Levitz, N. Netea, W. White, Hidden killers: human fungal infections, *Sci. Transl. Med.* 4 (165) (2012) 165rv13.
- [3] F. Almeida, M.L. Rodrigues, C. Coelho, The still underestimated problem of fungal diseases worldwide, *Front. Microbiol.* 10 (2019).
- [4] C. Fisher Matthew, J. Gurr Sarah, A. Cuomo Christina, S. Bleher David, J. Hailing, H. Stukenbrock Eva, et al., Threats posed by the fungal kingdom to humans, wildlife, and agriculture, *mBio* 11 (3) (2020), <https://doi.org/10.1128/mbio.00449-20>.
- [5] S. Savary, A. Ficke, J. Aubertot, C. Hollier, Crop losses due to diseases and their implications for global food production losses and food security, *Food Secur.* 4 (4) (2012) 519–537.
- [6] M.L. Rodrigues, J.D. Nosanchuk, Fungal diseases as neglected pathogens: a wake-up call to public health officials, *PLoS Negl. Trop. Dis.* 14 (2) (2020) e0007964.
- [7] L.E. Cowen, D. Sanglard, S.J. Howard, P.D. Rogers, D.S. Perlin, Mechanisms of antifungal drug resistance, *Cold Spring Harb. Perspect. Med.* 5 (7) (2014 Nov 10) a019752.
- [8] B. Havliczkova, V.A. Czaika, M. Friedrich, Epidemiological trends in skin mycoses worldwide, *Mycoses* 51 (Suppl. 4) (2008 Sep) 2–15.
- [9] J. Baker, D.W. Denning, The SSS revolution in fungal diagnostics: speed, simplicity and sensitivity, *Br. Med. Bull.* (2023) ldad011.
- [10] B. Ball, M. Langille, J. Geddes-McAlister, Fun(gi)omics: advanced and diverse technologies to explore emerging fungal pathogens and define mechanisms of antifungal resistance, *mBio* 11 (5) (2020), <https://doi.org/10.1128/mbio.01020-20>.
- [11] J.A. Lucas, N.J. Hawkins, B.A. Fraaije, Chapter two - the evolution of fungicide resistance, *Adv. Appl. Microbiol.* 90 (2015) 29–92.
- [12] R. Medina, M.E.E. Franco, L.C. Bartel, V. Martinez Alcántara, M.C.N. Saparrat, P. A. Balatti, Fungal Mitogenomes: Relevant Features to Planning Plant Disease Management, *Frontiers in Microbiology*, 2020, p. 11.
- [13] L. Duvenage, C.A. Munro, C.W. Gourlay, The potential of respiration inhibition as a new approach to combat human fungal pathogens, *Curr. Genet.* 65 (6) (2019) 1347–1353.
- [14] Q. Yulin, W. Jinxin, L. Quanzhen, H. Bing, Recent progress in research on mitochondrion-targeted antifungal drugs: a review, *Antimicrob. Agents Chemother.* 67 (6) (2023) 3.
- [15] E.T. Gibbs, C.A. Lerner, M.A. Watson, H. Wong, A.A. Gerencser, M.D. Brand, Site IQ in mitochondrial complex I generates S1QEL-sensitive superoxide/hydrogen peroxide in both the reverse and forward reactions, *Biochem. J.* 480 (5) (2023 Mar 15) 363–384.
- [16] E.L. Robb, A.R. Hall, T.A. Prime, S. Eaton, M. Szibor, C. Viscomi, et al., Control of mitochondrial superoxide production by reverse electron transport at complex I, *J. Biol. Chem.* 293 (25) (2018) 9869–9879.
- [17] R. Roca, W. Whitworth, P. Prag, M. Murphy, R. Ramakrishnan, Tumor necrosis factor induces pathogenic mitochondrial ROS in tuberculosis through reverse electron transport, *Science* 376 (6600) (2022) eabh2841.
- [18] A. Warris, E.R. Ballou, Oxidative responses and fungal infection biology, *Semin. Cell Dev. Biol.* 89 (2019) 34–46.
- [19] I.G. Kirkinetzos, C.T. Moraes, Reactive oxygen species and mitochondrial diseases, *Semin. Cell Dev. Biol.* 12 (6) (2001) 449–457.
- [20] N. Delattin, B.P.A. Cammue, K. Thevissen, Reactive oxygen species-inducing antifungal agents and their activity against fungal biofilms, *Future Med. Chem.* 6 (1) (2014) 77–90.
- [21] E. Shekhova, O. Kniemeyer, A.A. Brakhage, Induction of mitochondrial reactive oxygen species production by Itraconazole, Terbinafine, and Amphotericin B as a mode of action against *Aspergillus fumigatus*, *Antimicrob. Agents Chemother.* 61 (11) (2017 Oct 24) e00978 (17, Print 2017 Nov).
- [22] T. Joseph-Horne, D.W. Hollomon, P.M. Wood, Fungal respiration: a fusion of standard and alternative components, *Biochim. Biophys. Acta (BBA) - Bioenergetics* 1504 (2) (2001) 179–195.
- [23] T. Joseph-Horne, D.W. Hollomon, Functional diversity within the mitochondrial electron transport chain of plant pathogenic fungi, *Pest Manag. Sci.* 56 (1) (2000) 24–30.
- [24] H. Tamura, A. Mizutani, H. Yukioka, N. Miki, K. Ohba, M. Masuko, Effect of the methoxyiminoacetamide fungicide, SSF129, on respiratory activity in *Botrytis cinerea*, *Pestic. Sci.* 55 (7) (1999) 681–686.
- [25] T. Joseph-Horne, P.M. Wood, C.K. Wood, A.L. Moore, J. Headrick, D. Hollomon, Characterization of a Split respiratory pathway in the wheat “take-all” fungus, *Gaeumannomyces graminis var. tritici* *, *J. Biol. Chem.* 273 (18) (1998) 11127–11133.
- [26] F. Tian, S.Y. Lee, S.Y. Woo, H.S. Chun, Alternative oxidase: a potential target for controlling aflatoxin contamination and propagation of *Aspergillus flavus*, *Front. Microbiol.* (2020) 11.
- [27] W. Huh, S. Kang, Characterization of the gene family encoding alternative oxidase from *Candida albicans*, *Biochem. J.* 356 (2) (2001) 595–604.
- [28] C. Malina, R. Yu, J. Björkeröth, E.J. Kerkhoven, J. Nielsen, Adaptations in metabolism and protein translation give rise to the Crabtree effect in yeast, *Proc. Natl. Acad. Sci.* 118 (51) (2021) e2112836118.
- [29] D.S. Childers, I. Raziunaite, G. Mol Avelar, J. Mackie, S. Budge, D. Stead, et al., The rewiring of ubiquitination targets in a pathogenic yeast promotes metabolic flexibility, host colonization and virulence, *PLoS Pathog.* 12 (4) (2016) e1005566.
- [30] A. Burgain, É. Pic, L. Markey, F. Tebbji, C.A. Kumamoto, A. Sellam, A novel genetic circuitry governing hypoxic metabolic flexibility, commensalism and virulence in the fungal pathogen *Candida albicans*, *PLoS Pathog.* 15 (12) (2019) e1007823.
- [31] C. Malina, R. Yu, J. Björkeröth, E.J. Kerkhoven, J. Nielsen, Adaptations in metabolism and protein translation give rise to the Crabtree effect in yeast, *Proc. Natl. Acad. Sci.* 118 (51) (2021) e2112836118.
- [32] T. Bouklas, L. Masone, B.C. Fries, Differences in Sirtuin regulation in response to calorie restriction in *Cryptococcus neoformans*, *J. Fungi* 4 (1) (2018) 26.
- [33] VdP Martins, T.M. Dinamarco, C. Curti, S.A. Uyemura, Classical and alternative components of the mitochondrial respiratory chain in pathogenic fungi as potential therapeutic targets, *J. Bioenerg. Biomembr.* 43 (1) (2011) 81–88.
- [34] A.L. Moore, T. Shiba, L. Young, S. Harada, K. Kita, K. Ito, Unraveling the heater: new insights into the structure of the alternative oxidase, *Annu. Rev. Plant Biol.* 64 (1) (2013) 637–663.
- [35] G.C. Vanlerberghe, L. McIntosh, ALTERNATIVE OXIDASE: from gene to function, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48 (1) (1997) 703–734.
- [36] Á.P. Molnár, Z. Németh, E. Fekete, M. Flippin, N.P. Keller, L. Karaffa, Analysis of the relationship between alternative respiration and sterigmatocystin formation in *Aspergillus nidulans*, *Toxins* 10 (4) (2018) 168.
- [37] L. Duvenage, A. Walker Louise, B. Aleksandra, A. Johnston Simon, M. MacCallum Donna, A. Munro Carol, et al., Inhibition of classical and alternative modes of respiration in *Candida albicans* leads to cell wall remodeling and increased macrophage recognition, *mBio* 10 (1) (2019), <https://doi.org/10.1128/mbio.02535-18>.
- [38] G. Milani, W. Jarmuszkiewicz, C.M. Sluse-Goffart, A.Z. Schreiber, A.E. Vercesi, F. E. Sluse, Respiratory chain network in mitochondria of *Candida parapsilosis*: ADP/O appraisal of the multiple electron pathways, *FEBS Lett.* 508 (2) (2001) 231–235.
- [39] H. Guedouari, R. Gergondey, A. Bourdais, O. Vanparis, A.L. Bulteau, J. M. Camadro, et al., Changes in glutathione-dependent redox status and mitochondrial energetic strategies are part of the adaptive response during the filamentation process in *Candida albicans*, *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1842 (9) (2014) 1855–1869.
- [40] E.J. Helmerhorst, M. Stan, M.P. Murphy, F. Sherman, F.G. Oppenheim, The concomitant expression and availability of conventional and alternative, cyanide-insensitive, respiratory pathways in *Candida albicans*, *Mitochondrion* 5 (3) (2005) 200–211.
- [41] J. Baishya, C.A. Wakeman, Selective pressures during chronic infection drive microbial competition and cooperation, *npj Biofilms Microbiom.* 5 (1) (2019) 16.
- [42] D.K. Morales, N. Grahl, C. Okegbe, L.E.P. Dietrich, N.J. Jacobs, D.A. Hogan, Control of *Candida albicans* metabolism and biofilm formation by *Pseudomonas aeruginosa* phenazines, *mBio* 4 (1) (2013 Jan 29) 526.
- [43] L.R. Hoffman, A.R. Richardson, L.S. Houston, H.D. Kulasekara, W. Martens-Habbena, M. Klausen, et al., Nutrient availability as a mechanism for selection of antibiotic tolerant *Pseudomonas aeruginosa* within the CF airway, *PLoS Pathog.* 6 (1) (2010) e1000712.
- [44] Y. Zhao, J. Lim, J. Xu, J. Yu, W. Zheng, Nitric oxide as a developmental and metabolic signal in filamentous fungi, *Mol. Microbiol.* 113 (5) (2020) 872–882.
- [45] A. Pradhan, Q. Ma, L.J. de Assis, I. Leaves, D.E. Larcombe, A.V. Rodriguez Rondon, et al., Anticipatory stress responses and immune evasion in fungal pathogens, *Trends Microbiol.* 29 (5) (2021) 416–427.
- [46] B. Black, C. Lee, L.C. Horianopoulos, W.H. Jung, J.W. Kronstad, Respiring to infect: emerging links between mitochondria, the electron transport chain, and fungal pathogenesis, *PLoS Pathog.* 17 (7) (2021) e1009661.
- [47] S. Aoki, S. Ito-Kuwa, Y. Nakamura, T. Masuhara, Comparative pathogenicity of a wild-type strain and respiratory mutants of *Candida albicans* in mice, *Zentralblatt für Bakteriologie* 273 (3) (1990) 332–343.
- [48] C. Shuna, L. Minghui, Y.A. Hassan Rabeay, Heintz-Buschart Anna, W. Junsong, B. Ursula, Inhibition of respiration of *Candida albicans* by small molecules increases phagocytosis efficacy by macrophages, *mSphere* 5 (2) (2020), <https://doi.org/10.1128/msphere.00016-20>.
- [49] N. Grahl, E.G. Demers, A.K. Lindsay, C.E. Harty, S.D. Willger, A.E. Piispanen, et al., Mitochondrial activity and Cyr1 are key regulators of Ras1 activation of *C. Albicans* virulence pathways, *PLoS Pathog.* 11 (8) (2015) e1005133.
- [50] C. Lin, Y. Chen, Conserved and divergent functions of the cAMP/PKA signaling pathway in *Candida albicans* and *Candida tropicalis*, *J. Fungi* 4 (2) (2018).
- [51] F.G.S. Silao, M. Ward, K. Ryman, A. Wallström, B. Brinddefalk, K. Udekwi, et al., Mitochondrial proline catabolism activates Ras1/cAMP/PKA-induced filamentation in *Candida albicans*, *PLoS Genet.* 15 (2) (2019) e1007976.
- [52] A. Gupta, B.B. Chattoo, Functional analysis of a novel ABC transporter ABC4 from *Magnaporthe grisea*, *FEMS Microbiol. Lett.* 278 (1) (2008) 22–28.
- [53] M. Urban, T. Bhargava, J.E. Hamer, An ATP-driven efflux pump is a novel pathogenicity factor in rice blast disease, *EMBO J.* 18 (3) (1999) 512–521.
- [54] M. Wasi, N.K. Khandelwal, A.J. Moorhouse, R. Nair, P. Vishwakarma, G. Bravo Ruiz, et al., ABC transporter genes show upregulated expression in drug-resistant clinical isolates of *Candida auris*: a genome-wide characterization of ATP-Binding Cassette (ABC) transporter genes, *Front. Microbiol.* 10 (2019).
- [55] Y. Zhang, S. Gamarra, G. Garcia-Effron, S. Park, D.S. Perlin, R. Rao, Requirement for Ergosterol in V-ATPase function underlies antifungal activity of azole drugs, *PLoS Pathog.* 6 (6) (2010) e1000939.
- [56] Z. Yan, J. Xu, Mitochondria are inherited from the MATa parent in crosses of the basidiomycete fungus *Cryptococcus neoformans*, *Genetics* 163 (4) (2003) 1315–1325.

- [57] L. Chang Andrew, Tamara L. Doering, Maintenance of mitochondrial morphology in *Cryptococcus neoformans* is critical for stress resistance and virulence, *mBio* 9 (6) (2018), <https://doi.org/10.1128/mbio.01375-18>.
- [58] L.C. Horianopoulos, J.W. Kronstad, Connecting iron regulation and mitochondrial function in *Cryptococcus neoformans*, *Curr. Opin. Microbiol.* 52 (2019) 7–13.
- [59] S.S. Ingavale, Y.C. Chang, H. Lee, C.M. McClelland, M.L. Leong, K. Kwon-Chung, Importance of mitochondria in survival of *Cryptococcus neoformans* under low oxygen conditions and tolerance to cobalt chloride, *PLoS Pathog.* 4 (9) (2008) e1000155.
- [60] S. Giles Steven, Batinić-Haberle Ines, R. Perfect John, Gary M. Cox, *Cryptococcus neoformans* mitochondrial superoxide dismutase: an essential link between antioxidant function and high-temperature growth, *Eukaryot. Cell* 4 (1) (2005) 46–54.
- [61] M. Duarte, R. Sousa, A. Videira, Inactivation of genes encoding subunits of the peripheral and membrane arms of neurospora mitochondrial complex I and effects on enzyme assembly, *Genetics* 139 (3) (1995) 1211–1221.
- [62] M. Duarte, A. Videira, Respiratory chain complex I is essential for sexual development in *Neurospora* and binding of iron sulfur clusters are required for enzyme assembly, *Genetics* 156 (2) (2000) 607–615.
- [63] M. Duarte, N. Mota, L. Pinto, A. Videira, Inactivation of the gene coding for the 30.4-kDa subunit of respiratory chain NADH dehydrogenase: is the enzyme essential for *Neurospora*? *Mol. Gen. Genet.* MGG 257 (3) (1998) 368–375.
- [64] T.A.A. Harkness, R.A. Rothery, J.H. Weiner, S. Werner, J.E. Azevedo, A. Videira, et al., Disruption of the gene encoding the 78-kilodalton subunit of the peripheral arm of complex I in *Neurospora crassa* by repeat induced point mutation (RIP), *Curr. Genet.* 27 (4) (1995) 339–350.
- [65] R. Navarro-Espíndola, F. Suaste-Olmos, L. Peraza-Reyes, Dynamic regulation of peroxisomes and mitochondria during fungal development, *J. Fungi* 6 (4) (2020).
- [66] M. Isabel, A. Dencher Norbert, V. Arnaldo, K. Frank, Supramolecular Organization of the Respiratory Chain in *Neurospora crassa* mitochondria, *Eukaryot. Cell* 6 (12) (2007) 2391–2405.
- [67] F. Krause, C.Q. Scheckhuber, A. Werner, S. Rexroth, N.H. Reifschneider, N. A. Dencher, et al., Supramolecular organization of cytochrome c oxidase- and alternative oxidase-dependent respiratory chains in the filamentous fungus *Podospora anserina**, *J. Biol. Chem.* 279 (25) (2004) 26453–26461.
- [68] M.F.P.M. Maas, C.H. Sellem, F. Krause, N.A. Dencher, A. Sainsard-Chanet, Molecular gene therapy: overexpression of the alternative NADH dehydrogenase NDI1 restores overall physiology in a fungal model of respiratory complex I deficiency, *J. Mol. Biol.* 399 (1) (2010) 31–40.
- [69] M.S. Chae, F.E. Nargang, Investigation of regulatory factors required for alternative oxidase production in *Neurospora crassa*, *Physiol. Plant.* 137 (4) (2009) 407–418.
- [70] M. Merryman, J. Crigler, R. Seipelt-Thiemann, E. McClelland, A mutation in *C. neoformans* mitochondrial NADH dehydrogenase results in increased virulence in mice, *Virulence* 11 (1) (2020) 1366–1378.
- [71] B. Geißel, M. Penka, M. Neubauer, J. Wagener, The ER-mitochondria encounter structure contributes to hyphal growth, mitochondrial morphology and virulence of the pathogenic mold *Aspergillus fumigatus*, *Int. J. Med. Microbiol.* 307 (1) (2017) 37–43.
- [72] L. Enkler, V. Szentgyörgyi, M. Pennauer, C. Prescianotto-Baschong, I. Riezman, A. Wiesyk, et al., Arf1 coordinates fatty acid metabolism and mitochondrial homeostasis, *Nat. Cell Biol.* 25 (2023) 1157–1172.
- [73] D. Ruf, V. Brantl, J. Wagener, Mitochondrial fragmentation in *aspergillus fumigatus* as early marker of granulocyte killing activity, *Front. Cell. Infect. Microbiol.* (2018) 8.
- [74] P. Schertl, H. Braun, Respiratory electron transfer pathways in plant mitochondria, *Front. Plant Sci.* (2014) 5.
- [75] H. Braun, S. Binder, A. Brennicke, H. Eubel, A.R. Fernie, I. Finkemeier, et al., The life of plant mitochondrial complex I, *Mitochondrion* 19 (2014) 295–313.
- [76] A. Umbach, J. Siedow, The cyanide-resistant alternative oxidases from the Fungi *Pichia stipitis* and *Neurospora crassa* are monomeric and lack regulatory features of the plant enzyme, *Arch. Biochem. Biophys.* 378 (2000) 234–245.
- [77] S. Huang, A.H. Millar, Succinate dehydrogenase: the complex roles of a simple enzyme, *Curr. Opin. Plant Biol.* 16 (3) (2013) 344–349.
- [78] A.H. Millar, H. Eubel, L. Jänsch, V. Kruff, J.L. Heazlewood, H. Braun, Mitochondrial cytochrome c oxidase and succinate dehydrogenase complexes contain plant specific subunits, *Plant Mol. Biol.* 56 (1) (2004 Sep) 77–90.
- [79] D.P.T. Thomazella, P.J.P.L. Teixeira, H.C. Oliveira, E.E. Saviani, J. Rincones, I. M. Toni, et al., The hemibiotrophic cacao pathogen *Moniliophthora perniciosa* depends on a mitochondrial alternative oxidase for biotrophic development, *New Phytol.* 194 (4) (2012) 1025–1034.
- [80] T. Xu, Y. Wang, W. Liang, F. Yao, Y. Li, D. Li, et al., Involvement of alternative oxidase in the regulation of sensitivity of *Sclerotinia sclerotiorum* to the fungicides azoxystrobin and procymidone, *J. Microbiol.* 51 (3) (2013) 352–358.
- [81] M. Szibor, C. Schenkl, M.R.O. Barsottini, L. Young, A.L. Moore, Targeting the alternative oxidase (AOX) for human health and food security, a pharmaceutical and agrochemical target or a rescue mechanism? *Biochem. J.* 479 (12) (2022 Jun 30) 1337–1359.
- [82] A.K. Pandey, M.K. Samota, A. Kumar, A.S. Silva, N.K. Dubey, Fungal mycotoxins in food commodities: present status and future concerns, *Front. Sustain. Food Syst.* 7 (2023) 2571–2581.
- [83] F. Tian, S.Y. Lee, S.Y. Woo, H.Y. Choi, S.B. Park, H.S. Chun, Effect of plant-based compounds on the antifungal and antifungal efficiency of strobilurins against *Aspergillus flavus*, *J. Hazard. Mater.* 415 (2021) 125663.
- [84] Fu Li-Jun, K. Shi, M. Gu, Yan-Hong Zhou, De-Kun Dong, Wu-Sheng Liang, et al., Systemic induction and role of mitochondrial alternative oxidase and nitric oxide in a compatible Tomato–Tobacco mosaic virus interaction, *MPMI* 23 (1) (2010) 39–48.
- [85] E. Keunen, M. Jozefczak, T. Remans, J. Vangronsveld, A. Cuyppers, Alternative respiration as a primary defence during cadmium-induced mitochondrial oxidative challenge in *Arabidopsis thaliana*, *Environ. Exp. Bot.* 91 (2013) 63–73.
- [86] G.G.K. Oh, B.M. O’Leary, S. Signorelli, A.H. Millar, Alternative oxidase (AOX) 1a and 1d limit proline-induced oxidative stress and aid salinity recovery in *Arabidopsis*, *Plant Physiol.* 188 (3) (2022 Mar 4) 1521–1536.
- [87] B.H. Simons, F.F. Millenaar, L. Mulder, L.C. Van Loon, H. Lambers, Enhanced expression and activation of the alternative oxidase during infection of *Arabidopsis* with *Pseudomonas syringae* pv *tomato1*, *Plant Physiol.* 120 (2) (1999) 529–538.
- [88] L. Zhang, Y. Oh, H. Li, I.T. Baldwin, I. Galis, Alternative oxidase in resistance to biotic stresses: *Nicotiana attenuata* AOX contributes to resistance to a pathogen and a piercing-sucking insect but not *Manduca sexta* larvae, *Plant Physiol.* 160 (3) (2012 Nov) 1453–1467.
- [89] Cruz Avila-Adame, Wolfram Köller, Disruption of the alternative oxidase gene in *Magnaporthe grisea* and its impact on host infection, *MPMI* 15 (5) (2002) 493–500.
- [90] E. Sierra-Campos, I. Velázquez, D. Matuz-Mares, A. Villavicencio-Quejjeiro, J. P. Pardo, Functional properties of the *Ustilago maydis* alternative oxidase under oxidative stress conditions, *Mitochondrion* 9 (2) (2009) 96–102.
- [91] C.A. Cárdenas-Monroy, T. Pohlmann, G. Piñón-Zárate, G. Mata-Ortega, G. Guerra, M. Feldbrügge, et al., The mitochondrial alternative oxidase Aox1 is needed to cope with respiratory stress but dispensable for pathogenic development in *Ustilago maydis*, *PLoS One* 12 (3) (2017) e0173389.
- [92] Z. Lin, J. Wu, P.A. Jamieson, C. Zhang, Alternative oxidase is involved in the pathogenicity, development, and oxygen stress response of *Botrytis cinerea*, *Phytopathology*® 109 (10) (2019) 1679–1688.
- [93] D. Fernández-Ortuño, J. Tores, A. Vicente, A. Pérez-García, Mechanisms of resistance to QoI fungicides in phytopathogenic fungi, *Int. Microbiol.* 11 (2008) 1–9.
- [94] T.Y.K. Oliveira, T.C. Silva, S.I. Moreira, F.S. Christiano, M.C.G. Gasparoto, B. A. Fraaije, et al., Evidence of resistance to QoI fungicides in contemporary populations of *Mycosphaerella fijiensis*, *M. musicola* and *M. thailandica* from banana plantations in Southeastern Brazil, *Agronomy* 12 (12) (2022) 2952.
- [95] F.J. Sautua, M.A. Carmona, Detection and characterization of QoI resistance in *Pyrenophora tritici-repentis* populations causing tan spot of wheat in Argentina, *Plant Pathol.* 70 (9) (2021) 2125–2136.
- [96] M. Miguez, C. Reeve, P.M. Wood, D.W. Hollomon, Alternative oxidase reduces the sensitivity of *Mycosphaerella graminicola* to QoI fungicides, *Pest Manag. Sci.* 60 (1) (2004) 3–7.
- [97] I. Kaneko, H. Ishii, Effect of azoxystrobin on activities of antioxidant enzymes and alternative oxidase in wheat head blight pathogens *Fusarium graminearum* and *Microdochium nivale*, *J. Gen. Plant Pathol.* 75 (5) (2009) 388–398.
- [98] S. Li, X. Li, H. Zhang, Z. Wang, H. Xu, The research progress in and perspective of potential fungicides: succinate dehydrogenase inhibitors, *Bioorg. Med. Chem.* 50 (2021) 116476.
- [99] W. Chen, L. Wei, W. Zhao, B. Wang, H. Zheng, P. Zhang, et al., Resistance risk assessment for a novel succinate dehydrogenase inhibitor pydiflumetofen in *Fusarium asiaticum*, *Pest Manag. Sci.* 77 (1) (2021 Jan) 538–547.
- [100] H.F. Avenot, T.J. Michailides, Progress in understanding molecular mechanisms and evolution of resistance to succinate dehydrogenase inhibiting (SDHI) fungicides in phytopathogenic fungi, *Crop Prot.* 29 (7) (2010) 643–651.
- [101] A. Karelov, N. Kozub, O. Sozinova, Y. Pirkko, I. Sozinov, A. Yemets, et al., Wheat genes associated with different types of resistance against stem rust (*Puccinia graminis* Pers.), *Pathogens* 11 (10) (2022).
- [102] C.A. Cullis, Disease resistance genes in flax, in: C.A. Cullis (Ed.), *Genetics and Genomics of Linum Cham*, Springer International Publishing, 2019, pp. 215–225.
- [103] A. Shamima, C. McDade Henry, M. Gorlach Jenifer, H. Garrett, M. Cox Gary, John R. Perfect, Role of alternative oxidase gene in pathogenesis of *Cryptococcus neoformans*, *Infect. Immun.* 71 (10) (2003) 5794–5802.
- [104] P. Martins Vicente, M. Dinamarco Taisa, M. Soriani Frederico, G. Tudella Valéria, C. Oliveira Sergio, H. Goldman Gustavo, et al., Involvement of an alternative oxidase in oxidative stress and mycelium-to-yeast differentiation in *Paracoccidioides brasiliensis*, *Eukaryot. Cell* 10 (2) (2011) 237–248.
- [105] O. Hernández Ruiz, A. Gonzalez, A.J. Almeida, D. Tamayo, A.M. Garcia, A. Restrepo, et al., Alternative oxidase mediates pathogen resistance in *Paracoccidioides brasiliensis* infection, *Crop Sci.* 51 (10) (2011) e1353.
- [106] N. Grah, T.M. Dinamarco, S.D. Willger, G.H. Goldman, R.A.S. Cramer, *Aspergillus fumigatus* mitochondrial electron transport chain mediates oxidative stress homeostasis, hypoxia responses and fungal pathogenesis, *Mol. Microbiol.* 84 (2) (2012) 383–399.
- [107] T. Magnani, F.M. Soriani, VdP Martins, AcDF Policarpo, C.A. Sorgi, L.H. Faccioli, et al., Silencing of mitochondrial alternative oxidase gene of *Aspergillus fumigatus* enhances reactive oxygen species production and killing of the fungus by macrophages, *J. Bioenerg. Biomembr.* 40 (6) (2008) 631–636.
- [108] N. Del-Saz, M. Ribas-Carbo, A.E. McDonald, H. Lambers, A.R. Fernie, I. Florez-Sarasa, An *in vivo* perspective of the role(s) of the alternative oxidase pathway, *Trends Plant Sci.* 23 (3) (2018) 206–219.
- [109] A.E. McDonald, D.V. Gospodaryov, Alternative NAD(P)H dehydrogenase and alternative oxidase: proposed physiological roles in animals, *Mitochondrion* 45 (2019) 7–17.
- [110] A.C. Copesey, M.R.O. Barsottini, B. May, F. Xu, M.S. Albury, L. Young, et al., Kinetic characterisation and inhibitor sensitivity of *Candida albicans* and *Candida auris* recombinant AOX expressed in a self-assembled proteoliposome system, *Sci. Rep.* 11 (1) (2021) 14748.

- [111] L. Duvenage, L.A. Walker, A. Bojarczuk, S.A. Johnston, D.M. McCallum, C. A. Munro, et al., Alternative oxidase induction protects *Candida albicans* from respiratory stress and promotes hyphal growth, *bioRxiv* (2018) 405670.
- [112] T. Wang, X. Xie, K. Li, Y. Deng, H. Chen, Alternative oxidase promotes biofilm formation of *Candida albicans*, *Curr. Med. Sci.* 38 (3) (2018) 443–448.
- [113] L. Yan, M. Li, Y. Cao, P. Gao, Y. Cao, Y. Wang, et al., The alternative oxidase of *Candida albicans* causes reduced fluconazole susceptibility, *J. Antimicrob. Chemother.* 64 (4) (2009) 764–773.
- [114] Z. Liu, P. Basso, S. Hossain, S.D. Liston, N. Robbins, L. Whitesell, et al., Multifactor transcriptional control of alternative oxidase induction integrates diverse environmental inputs to enable fungal virulence, *Nat. Commun.* 14 (1) (2023) 4528.
- [115] Z. Liu, P. Basso, S. Hossain, S.D. Liston, N. Robbins, L. Whitesell, et al., Multifactor transcriptional control of alternative oxidase induction integrates diverse environmental inputs to enable fungal virulence, *Nat. Commun.* 14 (1) (2023 Jul 27) 4528-w.
- [116] W. Huh, S. Kang, Characterization of the gene family encoding alternative oxidase from *Candida albicans*, *Biochem. J.* 356 (2) (2001) 595–604.
- [117] I. Ittarat, W. Asawamahsakda, M.S. Bartlett, J.W. Smith, S.R. Meshnick, Effects of atovaquone and other inhibitors on *Pneumocystis carinii* dihydroorotate dehydrogenase, *Antimicrob. Agents Chemother.* 39 (2) (1995 Feb) 325–328.
- [118] S. Merali, D. Vargas, M. Franklin, A.B.J. Clarkson, S-adenosylmethionine and *Pneumocystis carinii*, *J. Biol. Chem.* 275 (20) (2000 May 19) 14958–14963.
- [119] T. Galadón, L. Carreté, The birth of a deadly yeast: tracing the evolutionary emergence of virulence traits in *Candida glabrata*, *FEMS Yeast Res.* 16(2):foV110 (2016).
- [120] A. Apostolopoulou, J.A. Fishman, The pathogenesis and diagnosis of pneumocystis jiroveci pneumonia, *J. Fungi* 8 (11) (2022).
- [121] P. Badiee, A. Rezapour, A. Abbasian, H.R. Foroutan, H. Jafarian, Prevalence of colonization and mitochondrial large subunit rRNA mutation of *Pneumocystis jiroveci* among Iranian children, *Iran. J. Microbiol.* 8 (5) (2016 Oct) 326–330.
- [122] A.B.C. Clarkson, J. Bienen EJ, Pollakis G, Grady RW, Respiration of bloodstream forms of the parasite *Trypanosoma brucei brucei* is dependent on a plant-like alternative oxidase *, *J. Biol. Chem.* 264 (30) (1989) 17770–17776.
- [123] M. Chaudhuri, R.D. Ott, G.C. Hill, Trypanosome alternative oxidase: from molecule to function, *Trends Parasitol.* 22 (10) (2006) 484–491.
- [124] T. Shiba, Y. Kido, K. Sakamoto, D.K. Inaoka, C. Tsuge, R. Tatsumi, et al., Structure of the trypanosome cyanide-insensitive alternative oxidase, *Proc. Natl. Acad. Sci. U. S. A.* 110 (12) (2013 Mar 19) 4580–4585.
- [125] S.K. Menzies, L.B. Tulloch, G.J. Florence, T.K. Smith, The trypanosome alternative oxidase: a potential drug target? *Parasitology* 145 (2) (2018) 175–183.
- [126] K.D. Goughenour, J. Zhao, J. Xu, Z.P. Zhao, A. Ganguly, C.M. Freeman, et al., Murine inducible nitric oxide synthase expression is essential for antifungal defenses in kidneys during disseminated *Cryptococcus neoformans* infection, *J. Immunol.* 207 (8) (2021 Oct 15) 2096–2106.
- [127] A.H. Millar, D.A. Day, Nitric oxide inhibits the cytochrome oxidase but not the alternative oxidase of plant mitochondria, *FEBS Lett.* 398 (2) (1996) 155–158.
- [128] Z. Bai, L.M. Harvey, B. McNeil, Physiological responses of chemostat cultures of *Aspergillus niger* (B1-D) to simulated and actual oxidative stress, *Biotechnol. Bioeng.* 82 (6) (2003) 691–701.
- [129] E.J. Kot, V.L. Olson, L.J. Rolewicz, D.O. McClary, An alternate respiratory pathway in *Candida albicans*, *Antonie Van Leeuwenhoek* 42 (1) (1976) 33–48.
- [130] F. Nasrin, Study of antimicrobial and antioxidant potentiality of anti-diabetic drug metformin, *Int. J. Pharm. Drug Anal.* 2 (3) (2014) 220–224.
- [131] Mascaraque V, Navas C, Hernaéz ML, Gil C, Molero G2. Proteomic study of the effect of metformin on *C. albicans*. *Access Microbiol.*;3(12):po0078.
- [132] Jaiswal S. Meherunisa, V. Seth, Study of metformin effect on antimicrobial property, *Int. Arch. BioMed. Clin. Res.* 4 (3) (2018).
- [133] I. Pernicova, M. Korbonits, Metformin—mode of action and clinical implications for diabetes and cancer, *Nat. Rev. Endocrinol.* 10 (3) (2014) 143–156.
- [134] W. Liu, Y. Wang, J. Chen, Z. Lin, M. Lin, X. Lin, et al., Beneficial effects of Gracillin from *Rhizoma Paridis* against gastric carcinoma via the potential TIPE2-mediated induction of endogenous apoptosis and inhibition of migration in BGC823 cells, *Front. Pharmacol.* 12 (2021).
- [135] Y. Li, H. Liu, X. Liu, B. Xiao, M. Zhang, Y. Luo, et al., Gracillin shows potential efficacy against non-small cell lung cancer through inhibiting the mTOR pathway, *Front. Oncol.* (2022) 12.
- [136] X. Zhang, H. Liu, Y. Gao, H. Wang, B. Guo, J. Li, Synthesis and antifungal activities of new type β -Methoxyacrylate-based Strobilurin analogues, *Chin. J. Chem.* 30 (7) (2012) 1517–1524.
- [137] A. Gao, Y. Fu, K. Zhang, M. Zhang, H. Jiang, L. Fan, et al., Azoxystrobin, a mitochondrial complex III Qo site inhibitor, exerts beneficial metabolic effects in vivo and in vitro, *Biochim. Biophys. Acta Gen. Subj.* 1840 (7) (2014) 2212–2221.
- [138] K. Bhattacharya, A.K. Bag, R. Tripathi, S.K. Samanta, B.C. Pal, C. Shaha, et al., Mahanine, a novel mitochondrial complex-III inhibitor induces G0/G1 arrest through redox alteration-mediated DNA damage response and regresses glioblastoma multiforme, *Am. J. Cancer Res.* 4 (6) (2014 Nov 19) 629–647.
- [139] C. Georgios, E. Lewis Russell, Dimitrios P. Kontoyiannis, Inhibition of *Candida parapsilosis* mitochondrial respiratory pathways enhances susceptibility to Caspofungin, *Antimicrob. Agents Chemother.* 50 (2) (2006) 744–747.
- [140] H. Chen, L. Li, Y. Lu, Y. Shen, M. Zhang, L. Ge, et al., Azoxystrobin reduces oral carcinogenesis by suppressing mitochondrial complex III activity and inducing apoptosis, *Cancer Manag. Res.* 12 (2020) 11573–11583.
- [141] S. Takahashi, T. Shinomiya, Y. Nagahara, Azoxystrobin induces apoptosis and cell cycle arrest in human leukemia cells independent of p53 expression, *Anticancer Res.* 42 (3) (2022) 1307.
- [142] A. Kapur, P. Mehta, A.D. Simmons, S.S. Erickson, G. Mehta, S.P. Palecek, et al., Atovaquone: an inhibitor of oxidative phosphorylation as studied in gynecologic cancers, *Cancers* 14 (9) (2022).
- [143] M. Luo, W. Zhou, H. Patel, A.P. Srivastava, J. Szymersky, M.M. Bonar, et al., Bedaquiline inhibits the yeast and human mitochondrial ATP synthases, *Commun. Biol.* 3 (1) (2020) 452.
- [144] J. Szymersky, D. Osowski, D.E. Walters, D.M. Mueller, Oligomycin frames a common drug-binding site in the ATP synthase, *Proc. Natl. Acad. Sci.* 109 (35) (2012) 13961–13965.
- [145] R. Mackie, N. Al-Bakkar, M. Kfoury, R. Roufayel, J. Sabatier, Z. Fajloun, Inhibitors of ATP synthase as new antibacterial candidates, *Antibiotics* 12 (4) (2023) 650.
- [146] J.A. Hendrickson, C. Hu, S.L. Aitken, N. Beyda, Antifungal resistance: a concerning trend for the present and future, *Curr. Infect. Dis. Rep.* 21 (12) (2019) 47.
- [147] K. Shannon, I. Gabriel, Q. Monica, S. Karen, O. Belinda, A. Greenko Jane, et al., Antifungal resistance trends of *Candida auris* clinical isolates in New York and New Jersey from 2016 to 2020, *Antimicrob. Agents Chemother.* 66 (3) (2022) 2242.
- [148] S.S. Gonçalves, A.C.R. Souza, A. Chowdhary, J.F. Meis, A.L. Colombo, Epidemiology and molecular mechanisms of antifungal resistance in *Candida* and *Aspergillus*, *Mycoses* 59 (4) (2016) 198–219.
- [149] J.H. Rex, C.R. Cooper, W.G. Merz, J.N. Galgiani, E.J. Anaissie, Detection of amphotericin B-resistant *Candida* isolates in a broth-based system, *Antimicrob. Agents Chemother.* 39 (4) (1995) 906–909.
- [150] V. Patrick, Larcher Gérald, Bergès Thierry, R. Gilles, C. Dominique, Bouchara Jean-Philippe, Mechanisms of azole resistance in a clinical isolate of *Candida tropicalis*, *Antimicrob. Agents Chemother.* 49 (11) (2005) 4608–4615.
- [151] A. Forastiero, A.C. Mesa-Arango, A. Alastruey-Izquierdo, L. Alcazar-Fuoli, L. Bernal-Martinez, T. Pelaez, et al., *Candida tropicalis* antifungal cross-resistance is related to different azole target (Erg11p) modifications, *Antimicrob. Agents Chemother.* 57 (10) (2013) 4769–4781.
- [152] S.L. Kelly, D.C. Lamb, D.E. Kelly, N.J. Manning, J. Loeffler, H. Hebart, et al., Resistance to fluconazole and cross-resistance to amphotericin B in *Candida albicans* from AIDS patients caused by defective sterol $\Delta 5,6$ -desaturation, *FEBS Lett.* 400 (1) (1997) 80–82.
- [153] S.G. Whaley, E.L. Berkow, J.M. Rybak, A.T. Nishimoto, K.S. Barker, P.D. Rogers, Azole antifungal resistance in *Candida albicans* and emerging non-albicans *Candida* species, *Front. Microbiol.* 7 (2017) 2173.
- [154] Y. Young Laura, M. Hull Christina, H. Joseph, Disruption of ergosterol biosynthesis confers resistance to amphotericin B in *Candida lusitanae*, *Antimicrob. Agents Chemother.* 47 (9) (2003) 2717–2724.
- [155] Scott Nancy E, Edwin ES, Kline Susan E, Anna S. Rapid evolution of multidrug resistance in a *Candida lusitanae* infection during micafungin monotherapy. *Antimicrob. Agents Chemother.* 2023;0(0):543.
- [156] S. Su, H. Yan, L. Min, H. Wang, X. Chen, J. Shi, et al., The antifungal activity of caspofungin in combination with antifungals or non-antifungals against *Candida* species in vitro and in clinical therapy, *Expert Rev. Anti-Infect. Ther.* 20 (2) (2022) 161–178.
- [157] T. Colley, G. Sehra, L. Daly, G. Kimura, T. Nakaoki, Y. Nishimoto, et al., Antifungal synergy of a topical triazole, PC945, with a systemic triazole against respiratory *Aspergillus fumigatus* infection, *Sci. Rep.* 9 (1) (2019) 9482.
- [158] J.R. Perfect, W.E. Dismukes, F. Dromer, D.L. Goldman, J.R. Graybill, R.J. Hamill, et al., Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Diseases Society of America, *Clin. Infect. Dis.* 50 (3) (2010) 291–322.
- [159] K.R. Iyer, N.M. Revie, C. Fu, N. Robbins, L.E. Cowen, Treatment strategies for cryptococcal infection: challenges, advances and future outlook, *Nat. Rev. Microbiol.* 19 (7) (2021) 454–466.
- [160] Y. Feng, G. Vladimir, L. Hui, Z. Cheng, G. Lu, B. Judith, et al., Adaptation to fluconazole via aneuploidy enables cross-adaptation to amphotericin B and Flucytosine in *Cryptococcus neoformans*, *Microbiol. Spectr.* 9 (2) (2021) 723.
- [161] B.M. Vincent, J. Langlois, R. Srinivas, A.K. Lancaster, R. Scherz-Shouval, L. Whitesell, et al., A fungal-selective cytochrome *bc₁* inhibitor impairs virulence and prevents the evolution of drug resistance, *Cell Chem. Biol.* 23 (8) (2016) 978–991.
- [162] J. Shi, S. Li, A. Gao, K. Zhu, H. Zhang, Tetrandrine enhances the antifungal activity of fluconazole in a murine model of disseminated candidiasis, *Phytomedicine* 46 (2018) 21–31.
- [163] L. Li, T. Zhang, J. Xu, J. Wu, Y. Wang, X. Qiu, et al., The synergism of the small molecule ENOblock and fluconazole against fluconazole-resistant *Candida albicans*, *Front. Microbiol.* 10 (2019) 2071.
- [164] L. Yan, M. Li, Y. Cao, P. Gao, Y. Cao, Y. Wang, et al., The alternative oxidase of *Candida albicans* causes reduced fluconazole susceptibility, *J. Antimicrob. Chemother.* 64 (4) (2009) 764–773.
- [165] Q. Li, Z. Bai, A. O'Donnell, L.M. Harvey, P.A. Hoskisson, B. McNeil, Oxidative stress in fungal fermentation processes: the roles of alternative respiration, *Biotechnol. Lett.* 33 (3) (2011) 457–467.
- [166] M. Szibor, T. Gainutdinov, E. Fernandez-Vizcarra, E. Dufour, Z. Gizatullina, G. Debska-Vielhaber, et al., Bioenergetic consequences from xenotopic expression of a tunicate AOX in mouse mitochondria: switch from RET and ROS to FET, *Biochim. Biophys. Acta Bioenerg.* 1861 (2) (2020 Feb 1) 148137.
- [167] N. Trevijano-Contador, S.A. Rossi, E. Alves, S. Landin-Ferreiro, O. Zaragoza, Capsule enlargement in *Cryptococcus neoformans* is dependent on mitochondrial activity, *Front. Microbiol.* 8 (2017) 1423.

- [168] R. Cao, L. Tan, K. Li, Q. Wan, G. Wu, J. Wang, et al., The germination of fungal spores in water and enhanced their resistance to chlor(am)ine: characteristics and mechanisms, *Chem. Eng. J.* 454 (2023) 140184.
- [169] W. Jarmuszkiewicz, L. Hryniewiecka, F.E. Sluse, The effect of pH on the alternative oxidase activity in isolated *Acanthamoeba castellanii* mitochondria, *J. Bioenerg. Biomembr.* 34 (3) (2002) 221–226.
- [170] G.C. Vanlerberghe, Alternative oxidase: a mitochondrial respiratory pathway to maintain metabolic and signaling homeostasis during abiotic and biotic stress in plants, *Int. J. Mol. Sci.* 14 (4) (2013) 6847.
- [171] R. El-Khoury, K.K. Kemppainen, E. Dufour, M. Szibor, H.T. Jacobs, P. Rustin, Engineering the alternative oxidase gene to better understand and counteract mitochondrial defects: state of the art and perspectives, *Br. J. Pharmacol.* 171 (8) (2014) 2243–2249.
- [172] R.G. Efremov, R. Baradaran, L.A. Sazanov, The architecture of respiratory complex I, *Nature* 465 (7297) (2010) 441–445.
- [173] H. Shimizu, A. Osanai, K. Sakamoto, D.K. Inaoka, T. Shiba, S. Harada, et al., Crystal structure of mitochondrial quinol–fumarate reductase from the parasitic nematode *Ascaris suum*, *J. Biochem.* 151 (6) (2012) 589–592.
- [174] T. Shiba, Y. Kido, K. Sakamoto, D.K. Inaoka, C. Tsuge, R. Tatsumi, et al., Structure of the trypanosome cyanide-insensitive alternative oxidase, *Proc. Natl. Acad. Sci.* 110 (12) (2013) 4580–4585.
- [175] C. Lange, C. Hunte, Crystal structure of the yeast cytochrome bc₁ complex with its bound substrate cytochrome c, *Proc. Natl. Acad. Sci.* 99 (5) (2002) 2800–2805.
- [176] S. Stock, L. Leslie, W. Walker, Molecular architecture of the rotary motor in ATP synthase, *Science* 286 (5445) (1999) 1700–1705.
- [177] U.C. The, UniProt: the universal protein knowledgebase in 2021, *Nucleic Acids Res.* 49 (2021) D480–D489.