



Kent Academic Repository

White, Lewis C. and Hydon, Peter E. (2024) *Moving frames: difference and differential-diff Lagrangians. Symmetry, Integrability and Geometry: Methods and Applications*, 20 . ISSN 1815-0659.

Downloaded from

<https://kar.kent.ac.uk/104622/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.3842/SIGMA.2024.006>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY-SA (Attribution-ShareAlike)

Additional information

For the purpose of open access, the author(s) has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Moving Frames: Difference and Differential-Difference Lagrangians

Lewis C. WHITE and Peter E. HYDON

*School of Mathematics, Statistics and Actuarial Science, University of Kent,
Canterbury, Kent, CT2 7NF, UK*

E-mail: lcwhite29@gmail.com, P.E.Hydon@kent.ac.uk

Received September 19, 2023, in final form January 09, 2024; Published online January 15, 2024

<https://doi.org/10.3842/SIGMA.2024.006>

Abstract. This paper develops moving frame theory for partial difference equations and for differential-difference equations with one continuous independent variable. In each case, the theory is applied to the invariant calculus of variations and the equivariant formulation of the conservation laws arising from Noether's theorem. The differential-difference theory is not merely an amalgam of the differential and difference theories, but has additional features that reflect the need for the group action to preserve the prolongation structure. Projectable moving frames are introduced; these cause the invariant derivative operator to commute with shifts in the discrete variables. Examples include a Toda-type equation and a method of lines semi-discretization of the nonlinear Schrödinger equation.

Key words: moving frames; difference equations; differential-difference equations; variational calculus; Noether's theorem

2020 Mathematics Subject Classification: 39A14; 58D19; 47E07

1 Introduction

The modern formulation of moving frames introduced by Fels and Olver [2, 3] is a powerful tool in the analysis of partial differential equations (PDEs). It enables one to reduce a given system of PDEs to an invariant system by factoring out Lie symmetry group orbits (locally, at least). If the symmetries are extraneous to the problem of interest (for instance, projective symmetries in computer vision [20]), the moving frame provides a major simplification. For a clear, straightforward introduction to moving frames, see Mansfield's text [12].

Many PDE systems of interest are Euler–Lagrange equations with a Lie group of variational symmetries. For such systems, moving frames are most effectively applied by invariantizing the Lagrangian functional directly [10]. With this approach, Noether's theorem has an elegant formulation in terms of the adjoint action of the Lie group on a set of invariants [4, 5, 6].

Difference equations have discrete independent variables, so any Lie symmetries act only on the dependent variables. Commonly, the action varies with the discrete variables. Moving frames can be adapted to an equation on a finite set of points by using a finite-dimensional product space. Finite difference approximation of a given differential equation requires consistency as the points coalesce. This constraint led to the introduction of multi-space [19], which has been used to construct (highly accurate) invariant approximations of ordinary differential equations (ODEs) [9]. More generally, a discrete moving frame attaches a finite-dimensional product space to each base point, without imposing coalescence or any other structure in advance [14]. The discrete moving frame construction works for any number of independent variables and has been used to generalize multi-space to higher dimensions [17].

This paper is a contribution to the Special Issue on Symmetry, Invariants, and their Applications in honor of Peter J. Olver. The full collection is available at <https://www.emis.de/journals/SIGMA/Olver.html>




Figure 1. (a) The total space for scalar OΔEs is $\mathcal{T} = \mathbb{Z} \times \mathbb{R}$; (b) the total space for scalar PΔEs with two independent variables is $\mathcal{T} = \mathbb{Z}^2 \times \mathbb{R}$.

Difference equations have an intrinsic structure that arises from their mesh point labels. We restrict attention to the most common case, an m -dimensional logically rectangular mesh. The labels $n^i \in \mathbb{Z}$, $i = 1, \dots, m$, can be regarded as the independent variables. Each label belongs to an ordered set, so it is helpful to incorporate this ordering into the moving frame definition. This has been achieved for ordinary difference equations (OΔEs), resulting in the invariant variational calculus and equivariant Noether’s theorem [15, 16]. The current paper extends this approach to partial difference equations (PΔEs) and to differential-difference equations (DΔEs) with one continuous independent variable.

Section 2 summarizes the building-blocks of the PΔE theory, from which we develop difference moving frames (see Section 3), the invariant calculus of variations (see Section 4), and the equivariant formulation of Noether’s conservation laws (see Section 5). Even for scalar PΔEs, it is necessary to use several generating invariants and to take the relations between these into account. The more dependent and independent variables there are, the more complex these relations can become. For clarity, we illustrate the general theory with fairly straightforward examples. In particular, we use a Toda-type equation as a running example to show the various aspects of the theory.

There is one aspect of the PΔE theory that is simpler than its counterpart for PDEs: the Lie group action on the independent variables is trivial. This is not necessarily the case for DΔEs, although there are constraints on the group action, as discussed in Section 6. These constraints suggest the idea of a projectable moving frame, which is a major simplification. Section 7 outlines the invariant variational calculus and Noether conservation laws for DΔEs, emphasizing those aspects of the theory that do not follow immediately from the PΔE theory. Section 8 presents some examples, including a method of lines semi-discretization of the nonlinear Schrödinger equation. Concluding remarks are given in Section 9.

2 A brief summary of the building-blocks

2.1 Difference prolongation space

A given differential equation can be represented as a variety within an appropriate jet space (see [18]). A difference equation has discrete independent variables, so to use moving frames, one must represent the equation as a variety within an appropriate continuous space. Such spaces are subspaces of the difference prolongation space [21], which is described briefly in this section.

Consider a difference equation with independent variables $\mathbf{n} := (n^1, n^2, \dots, n^m) \in \mathbb{Z}^m$ and dependent variables $\mathbf{u} := (u^1, u^2, \dots, u^q) \in \mathbb{R}^q$. These variables are coordinates on the *total space* $\mathbb{Z}^m \times \mathbb{R}^q$; a solution of the PΔE is a graph on the total space. Figure 1 illustrates the total spaces for a scalar OΔE and PΔE. For simplicity, we assume that the equation holds for

all $\mathbf{n} \in \mathbb{Z}^m$; our results apply *mutatis mutandis* to difference equations on any product lattice (see [7]). We also assume that all functions are smooth in their continuous arguments.

The total space is mapped to itself by horizontal translations

$$\mathbf{T}_{\mathbf{I}}: \mathbb{Z}^m \times \mathbb{R}^q \rightarrow \mathbb{Z}^m \times \mathbb{R}^q, \quad \mathbf{T}_{\mathbf{I}}: (\mathbf{n}, \mathbf{u}) \mapsto (\mathbf{n} + \mathbf{I}, \mathbf{u}),$$

where $\mathbf{I} \in \mathbb{Z}^m$ is a multi-index. Over each \mathbf{n} , one can construct the prolongation space $P(\mathbb{R}^q)$, which is an infinite-dimensional Cartesian product space with coordinates $u_{\mathbf{J}}^{\alpha} \in \mathbb{R}$, where $\mathbf{J} \in \mathbb{Z}^m$. Let $P_{\mathbf{n}}(\mathbb{R}^q)$ denote the prolongation space over a given \mathbf{n} ; this is a continuous fibre over the fixed base point \mathbf{n} . Every graph on the total space defines a point in $P_{\mathbf{n}}(\mathbb{R}^q)$, with the coordinate $u_{\mathbf{J}}^{\alpha}$ taking the value of u^{α} given by the graph at $\mathbf{n} + \mathbf{J}$.

Horizontal translation extends naturally to the total prolongation space $\mathbb{Z}^m \times P(\mathbb{R}^q)$ as follows:

$$\mathbf{T}_{\mathbf{I}}: (\mathbf{n}, (u_{\mathbf{J}}^{\alpha})) \mapsto (\mathbf{n} + \mathbf{I}, (u_{\mathbf{J}}^{\alpha})).$$

Suppose that f is a function on $\mathbb{Z}^m \times P(\mathbb{R}^q)$; its restriction to $P_{\mathbf{n}}(\mathbb{R}^q)$ is denoted by

$$f_{\mathbf{n}}((u_{\mathbf{J}}^{\alpha})) = f(\mathbf{n}, (u_{\mathbf{J}}^{\alpha})).$$

The pullback $\mathbf{T}_{\mathbf{I}}^*$ of $f_{\mathbf{n}+\mathbf{I}}((u_{\mathbf{J}}^{\alpha}))$ to $P_{\mathbf{n}}(\mathbb{R}^q)$ is

$$\mathbf{T}_{\mathbf{I}}^* f_{\mathbf{n}+\mathbf{I}}((u_{\mathbf{J}}^{\alpha})) = f(\mathbf{n} + \mathbf{I}, (u_{\mathbf{J}+\mathbf{I}}^{\alpha})).$$

This can be represented on the prolongation space over \mathbf{n} as a mapping $\mathbf{S}_{\mathbf{I}}$, called the *shift* by \mathbf{I} , which acts on smooth functions $f \in C^{\infty}(P_{\mathbf{n}}(\mathbb{R}^q))$ as follows:

$$\mathbf{S}_{\mathbf{I}} f(\mathbf{n}, (u_{\mathbf{J}}^{\alpha})) = f(\mathbf{n} + \mathbf{I}, (u_{\mathbf{J}+\mathbf{I}}^{\alpha})).$$

To summarize, each shift operator represents the action of the pullback on functions as follows: $\mathbf{S}_{\mathbf{I}} f_{\mathbf{n}} := \mathbf{T}_{\mathbf{I}}^* f_{\mathbf{n}+\mathbf{I}}$. Although $\mathbf{S}_{\mathbf{I}}$ represents a translation, it does not change the fibre. By using the shift operators, one can represent a given PΔE as a variety on $P_{\mathbf{n}}(\mathbb{R}^q)$. We do this from here on, using the Einstein summation convention to denote sums over all variables other than \mathbf{n} , as far as possible. For simplicity, we omit the multi-index subscript $\mathbf{0}$ on variables, except where this may cause confusion. In particular, u^{α} denotes $u_{\mathbf{0}}^{\alpha}$ henceforth.

Remark 2.1. For a given PΔE, one can restrict attention to a finite-dimensional subspace of the prolongation space, provided that this includes all $u_{\mathbf{J}}^{\alpha}$ for which \mathbf{J} lies within the stencil of the PΔE with respect to u^{α} (see [7]), together with any other relevant $u_{\mathbf{J}}^{\alpha}$ for the problem being considered. However, for generality, we use the full prolongation space over \mathbf{n} .

2.2 The difference variational calculus

This section summarizes the difference variational calculus from the formal viewpoint introduced by Kupershmidt [11]. It closely resembles the formal differential variational calculus described in [18]. Summation by parts replaces integration by parts, and a difference version of the divergence is used, which we now describe.

Each shift operator $\mathbf{S}_{\mathbf{J}}$, where $\mathbf{J} = (j^1, \dots, j^m)$, may be written as a product of unit shift operators, $\mathbf{S}_i := \mathbf{S}_{\mathbf{1}_i}$, and their inverses. Here $\mathbf{1}_i$ is the multi-index whose only non-zero entry is the i^{th} one, which is 1. Thus \mathbf{S}_i is the forward shift in the n^i -direction. By the composition rule for translations, $\mathbf{S}_i \mathbf{S}_j = \mathbf{S}_j \mathbf{S}_i$ and $\mathbf{S}_{\mathbf{J}} = \mathbf{S}_1^{j_1} \cdots \mathbf{S}_m^{j_m}$; consequently, $(\mathbf{S}_{\mathbf{J}})^{-1} = \mathbf{S}_{-\mathbf{J}}$. The identity operator, $\text{id} := \mathbf{S}_{\mathbf{0}}$, maps every function to itself, and the forward difference operator in the direction n^k is $\mathbf{D}_{n^k} = \mathbf{S}_k - \text{id}$. A *difference divergence* is an expression of the form $\text{Div}(F) = \mathbf{D}_{n^k} F^k$ for some $F := (F^1, \dots, F^m)$. It is straightforward to write a given expression of the form $(\mathbf{S}_{\mathbf{J}} - \text{id})f$, where f is a function, as a difference divergence; however, the resulting functions F^k are not unique if $m > 1$ and may be messy.

Definition 2.2. A *conservation law* of a given system of PΔEs is a difference divergence expression, $\mathcal{C} = \text{Div}(F)$, such that $\mathcal{C} = 0$ on all solutions of the system.

A linear difference operator on $P_{\mathbf{n}}(\mathbb{R}^q)$ is an operator of the form $\mathcal{H} = h^{\mathbf{J}}S_{\mathbf{J}}$, where each $h^{\mathbf{J}}$ is a function. The formal adjoint of \mathcal{H} is the operator \mathcal{H}^\dagger defined by

$$f\mathcal{H}g - (\mathcal{H}^\dagger f)g \in \text{im}(\text{Div})$$

for all functions f, g . Explicitly, $\mathcal{H}^\dagger f = S_{-\mathbf{J}}(h^{\mathbf{J}}f)$, because

$$fh^{\mathbf{J}}S_{\mathbf{J}}g - (S_{-\mathbf{J}}(h^{\mathbf{J}}f))g = (S_{\mathbf{J}} - \text{id})\{(S_{-\mathbf{J}}(h^{\mathbf{J}}f))g\}.$$

A special case is the very useful *summation by parts* formula

$$f(S_{\mathbf{J}}g) = (S_{-\mathbf{J}}f)g + (S_{\mathbf{J}} - \text{id})\{(S_{-\mathbf{J}}f)g\}.$$

The basic variational problem is to find the extrema of a given functional

$$\mathcal{L}[\mathbf{u}] = \sum_{\mathbf{n}} L(\mathbf{n}, [\mathbf{u}]),$$

where $[\mathbf{u}]$ represents finitely many shifts of the dependent variables. Extrema are found by requiring that

$$\left\{ \frac{d}{d\epsilon} \mathcal{L}[\mathbf{u} + \epsilon \mathbf{w}] \right\} \Big|_{\epsilon=0} = 0,$$

for all $\mathbf{w}: \mathbb{Z}^m \rightarrow \mathbb{R}^q$ that vanish sufficiently rapidly as any independent variable approaches infinity. Using summation by parts,

$$\frac{d}{d\epsilon} \Big|_{\epsilon=0} \mathcal{L}(\mathbf{n}, [\mathbf{u} + \epsilon \mathbf{w}]) = \left(S_{\mathbf{J}} w^\alpha \frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \right) = w^\alpha E_{u^\alpha}(L) + \text{Div}(A_{\mathbf{u}}(\mathbf{n}, [\mathbf{u}], [\mathbf{w}])),$$

where

$$E_{u^\alpha} = S_{-\mathbf{J}} \frac{\partial}{\partial u_{\mathbf{J}}^\alpha}$$

is the difference Euler–Lagrange operator with respect to u^α , and

$$\text{Div}(A_{\mathbf{u}}(\mathbf{n}, [\mathbf{u}], [\mathbf{w}])) = \sum_{\mathbf{J}} (S_{\mathbf{J}} - \text{id}) \left(w^\alpha S_{-\mathbf{J}} \frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \right).$$

As \mathbf{w} is arbitrary and the sum of $\text{Div}(A_{\mathbf{u}}(\mathbf{n}, [\mathbf{u}], [\mathbf{w}]))$ over \mathbf{n} is zero (by Stokes' theorem), the extrema satisfy the following system of Euler–Lagrange (difference) equations,

$$E_{u^\alpha}(L) = S_{-\mathbf{J}} \left(\frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \right) = 0.$$

Remark 2.3. If the dependent variables are regarded as depending smoothly on a continuous parameter t as well as \mathbf{n} , the same result is achieved by using

$$\frac{d}{dt} \Big|_{(u^\alpha)' = w^\alpha} L[\mathbf{u}] = 0,$$

where $(u^\alpha)' = du^\alpha/dt$. This approach is used later to derive the Lie group invariant version of the Euler–Lagrange equations.

2.3 Variational point symmetries and Noether's theorem

We now outline some relevant facts about Lie point symmetries, with application to variational calculus (see [7, 18] for further details). Let G be an R -dimensional Lie group parametrized by $\varepsilon = (\varepsilon^1, \dots, \varepsilon^R) \in \mathbb{R}^R$ in some neighbourhood of the identity, e . For now, we restrict attention to such a neighbourhood, in which the elements of G are $\Gamma(\varepsilon)$, where Γ depends smoothly on ε , with $\Gamma(\mathbf{0}) = e$. Locally, the left action of G on the coordinates $\mathbf{u} = (u^1, \dots, u^q)$ is denoted by $\hat{\mathbf{u}} = \Gamma(\varepsilon) \cdot \mathbf{u}$. The R -dimensional Lie algebra \mathcal{X} of infinitesimal generators has a basis

$$\mathbf{v}_r = Q_r^\alpha(\mathbf{n}, \mathbf{u}) \partial_{u^\alpha}, \quad r = 1, \dots, R, \quad \text{where} \quad Q_r^\alpha = \left. \frac{\partial \hat{u}^\alpha}{\partial \varepsilon^r} \right|_{\varepsilon=\mathbf{0}},$$

so every infinitesimal generator of a one-parameter (local) Lie subgroup of point transformations is of the form $\mathbf{v} = Q^\alpha(\mathbf{n}, \mathbf{u}) \partial_{u^\alpha}$, where $Q^\alpha = c^r Q_r^\alpha$ for some real constants c^r . The q -tuple $\mathbf{Q} = (Q^1, \dots, Q^R)$ is the *characteristic* of the Lie subgroup whose infinitesimal generator is \mathbf{v} .

Each infinitesimal generator is a tangent vector field on the total space. It is represented on the prolongation space $P_{\mathbf{n}}(\mathbb{R}^q)$ by the prolonged vector field

$$\text{pr } \mathbf{v} = (\text{S}_{\mathbf{J}} Q^\alpha) \frac{\partial}{\partial u_{\mathbf{J}}^\alpha}.$$

From here on, we refer to $\text{pr } \mathbf{v}$ simply as \mathbf{v} , because it will always be clear whether the generator is acting on the total space or the prolongation space over \mathbf{n} . Note that \mathbf{n} is invariant under the Lie group action, as are the shift operators $\text{S}_{\mathbf{J}}$.

Denote the left action of a general group element $g \in G$ (not necessarily in the neighbourhood of the identity) on the total space by $\tilde{\mathbf{u}} = g \cdot \mathbf{u}$ and define

$$\tilde{\mathbf{v}}_r = Q_r^\alpha(\mathbf{n}, \tilde{\mathbf{u}}) \partial_{\tilde{u}^\alpha}, \quad r = 1, \dots, R.$$

The adjoint representation of g on \mathcal{X} can be expressed as a matrix, $\mathcal{A}d(g) = (a_r^s(g))$, whose components are determined from the following relations (see [15]):

$$\tilde{\mathbf{v}}_r = a_r^s(g) \tilde{\mathbf{v}}_s, \quad r = 1, \dots, R. \quad (2.1)$$

By regarding the infinitesimal generators as differential operators and applying the left-hand side of the identity (2.1) to each \tilde{u}^α in turn, one obtains

$$\left(\frac{\partial \tilde{u}^\alpha}{\partial u^\beta} \right) Q_r^\beta(\mathbf{n}, \mathbf{u}) = \mathbf{v}_r(\tilde{u}^\alpha) = Q_s^\alpha(\mathbf{n}, \tilde{\mathbf{u}}) a_r^s(g), \quad (2.2)$$

where the matrix $(\partial \tilde{u}^\alpha / \partial u^\beta)$ is the Jacobian matrix of the transformation $g: \mathbf{u} \rightarrow \tilde{\mathbf{u}}$. Prolonging this result to each $u_{\mathbf{J}}^\alpha$ gives the identities

$$\left(\frac{\partial \tilde{u}_{\mathbf{J}}^\alpha}{\partial u_{\mathbf{J}}^\beta} \right) Q_r^\beta(\mathbf{n} + \mathbf{J}, \mathbf{u}_{\mathbf{J}}) = Q_s^\alpha(\mathbf{n} + \mathbf{J}, \tilde{\mathbf{u}}_{\mathbf{J}}) a_r^s(g).$$

Definition 2.4. The point transformations generated by \mathbf{v} are *variational symmetries* of the Lagrangian $L(\mathbf{n}, [\mathbf{u}])$ if there exist functions $B^i(\mathbf{n}, [\mathbf{u}])$ such that

$$\mathbf{v}(L) := (\text{S}_{\mathbf{J}} Q^\alpha) \frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} = D_{n^i} B^i. \quad (2.3)$$

The Lagrangian is invariant under the symmetries generated by \mathbf{v} if $B^i = 0$ for all i . Summing (2.3) by parts leads to Noether's theorem for PΔEs (see [7]).




Figure 2. Moving frame defined by a cross-section; $\mathcal{O}(z)$ denotes the group orbit through z .

Theorem 2.5 (difference Noether's theorem). *Suppose that a Lagrangian L has a variational symmetry with characteristic $\mathbf{Q} \neq \mathbf{0}$. Then the system of Euler–Lagrange equations has the following conservation law:*

$$-D_n B^i + \sum_{\mathbf{J}} (S_{\mathbf{J}} - \text{id}) \left(Q^\alpha S_{-\mathbf{J}} \frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \right) = 0. \quad (2.4)$$

From here on, we restrict attention to Lie groups of variational point symmetries that leave the Lagrangian invariant, so $B^i = 0$, $i = 1, \dots, m$.

2.4 Moving frames

We now outline some basics of moving frames on an arbitrary manifold (see [2, 3, 12] for further details). Let M be a smooth manifold. Suppose that a Lie group, G , of point transformations has a smooth left action on M that is free and regular¹ in a neighbourhood $\mathcal{U} \subset M$ of a point $z \in M$. Freeness and regularity are necessary and sufficient to guarantee the existence of a cross-section \mathcal{K} that is transverse to the group orbits that foliate \mathcal{U} , as shown in Figure 2. Moreover, each orbit intersects \mathcal{K} at a unique point.

If a group action is not free and regular, it can be made so by replacing M by a Cartesian product space M^N of sufficiently high dimension and using the induced product action (see Boutin [1]). Henceforth, we assume that M is of sufficiently high dimension that the action is free and regular.

Definition 2.6 (moving frame). Given a smooth Lie group action $G \times M \rightarrow M$, a moving frame is a smooth equivariant map $\rho: \mathcal{U} \subset M \rightarrow G$. Here \mathcal{U} is called the domain of the frame.

Given a left action, $g \cdot z$, a left equivariant map satisfies $\rho(g \cdot z) = g\rho(z)$ and a right equivariant map satisfies $\rho(g \cdot z) = \rho(z)g^{-1}$. The frame is called left or right accordingly. The inverse of a right frame is a left frame.

To find a right frame for an R -dimensional Lie group, G , write the cross-section \mathcal{K} as a system of equations $\psi_r(z) = 0$, $r = 1, \dots, R$. Then solve the *normalization equations*,

$$\psi_r(g \cdot z) = 0, \quad r = 1, \dots, R, \quad (2.5)$$

to obtain the unique group element $g = \rho(z)$ that maps z to the intersection of its group orbit with \mathcal{K} (see Figure 2). Both $\rho(g \cdot z)$ and $\rho(z)g^{-1}$ satisfy the equation $\psi_r(\rho(g \cdot z) \cdot (g \cdot z)) = 0$, so by uniqueness, the solution is a right frame.

¹The action is free if the only group element $g \in G$ that fixes every point in the neighbourhood is the identity. The action is regular if the orbits form a regular foliation.

An important part of the method of moving frames is to choose a cross-section \mathcal{K} that makes computations as simple as possible. It is usually easiest to choose a cross-section on which R of the coordinates on \mathcal{U} are constant. The normalization equations are then expressed as

$$g \cdot z_1 = c_1, \quad g \cdot z_2 = c_2, \quad \dots, \quad g \cdot z_R = c_R,$$

where z_r are coordinates and c_r are fixed constants. We will use this approach in our examples.

Given a left action $G \times M \rightarrow M$ and a right frame ρ , let

$$\iota(z) = \rho(z) \cdot z = g \cdot z|_{g=\rho(z)}.$$

The components of $\iota(z)$ are invariant under the Lie group action, because

$$\iota(g \cdot z) = \rho(g \cdot z) \cdot (g \cdot z) = \rho(z) \cdot g^{-1}g \cdot z = \rho(z) \cdot z = \iota(z).$$

The non-constant components of $\iota(z)$ are called the *normalized invariants*. Note that $\rho(\rho(z) \cdot z)$ is the identity element of G , so $\iota(\iota(z)) = \iota(z)$. In other words, the operator ι projects z to its invariant components. This operator extends to functions $f(z)$, as follows: $\iota(f(z)) = f(\iota(z))$. Note that $\iota(\iota(f(z))) = \iota(f(z))$, so ι projects out the invariant component of $f(z)$; hence, ι is called the *invariantization operator*. Indeed, if $F(z)$ is any invariant, the following *replacement rule* applies:

$$F(z) = F(\iota(z)). \tag{2.6}$$

Consequently, the set of all invariants is generated by the normalized invariants $\iota(z)$.

2.5 Discrete moving frames

The discrete moving frame developed by Mansfield, Marí Beffa and Wang [14] and Marí Beffa and Mansfield [17] can be thought of as a moving frame adapted to discrete base points. The Lie group action on M is extended to the diagonal (left) action on the Cartesian product manifold $\mathcal{M} = M^N$:

$$g \cdot (z_1, z_2, \dots, z_N) \mapsto (g \cdot z_1, g \cdot z_2, \dots, g \cdot z_N).$$

No assumptions are made about any relationship between the elements z_1, \dots, z_N .

Definition 2.7 (discrete moving frames). Let G^N denote the Cartesian product of N copies of the group G . A map

$$\rho: M^N \rightarrow G^N, \quad \rho(z) = (\rho_1(z), \dots, \rho_N(z))$$

is a right discrete moving frame if

$$\rho_k(g \cdot z) = \rho_k(z)g^{-1}, \quad k = 1, \dots, N,$$

and a left discrete moving frame if

$$\rho_k(g \cdot z) = g\rho_k(z), \quad k = 1, \dots, N.$$

The frame is right (resp. left) equivariant under the action of the Lie group.

A set of normalization equations yields a corresponding right discrete moving frame. The component ρ_k is the unique element of G that takes z to the cross-section \mathcal{K}_k . The sequence of moving frames with a nontrivial intersection of domains (ρ_k) which makes up the discrete moving frame is, locally, uniquely determined by the cross-section $\mathcal{K} = (\mathcal{K}_1, \dots, \mathcal{K}_N)$ to the group orbit through z (see [14] for more details). Again, no assumptions are made about any relationship between the components of the cross-section.

The invariants of the right (discrete) frame are $I_{k,j} := \rho_k(z) \cdot z_j$. If M is q -dimensional, each z_j has components z_j^1, \dots, z_j^q . So the components of $I_{k,j}$ are

$$I_{k,j}^\alpha := \rho_k(z) \cdot z_j^\alpha, \quad \alpha = 1, \dots, q.$$

The discrete moving frame applies to a wide variety of discrete domains. We now show how it can be adapted to the difference prolongation space for PΔEs, yielding the difference moving frame.

3 Difference moving frames

In view of Remark 2.1, one can restrict attention to a finite prolongation space \mathcal{M} of arbitrarily high dimension. This enables one to treat a difference moving frame as a particular type of discrete moving frame, so that the key definitions and theorems for discrete moving frames in [14, 15, 17] apply. The main distinction is that for the difference frame, there is a relation between the cross-sections and frames on the different fibres: they must be consistent with the pullback to any particular fibre.

3.1 Difference frames and invariants

Let \mathcal{K} and $\rho([\mathbf{u}])$ denote the cross-section and frame on \mathbf{n} , respectively. The cross-section on \mathbf{n} , denoted \mathcal{K} , is replicated for all the other base points $\mathbf{n} + \mathbf{J}$ if and only if the cross-section over $\mathbf{n} + \mathbf{J}$ is represented on \mathcal{M} by $S_{\mathbf{J}}\mathcal{K}$. Consequently, $\rho_{\mathbf{J}}([\mathbf{u}]) = S_{\mathbf{J}}\rho([\mathbf{u}])$; this constraint can be extended to the infinite-dimensional prolongation space $P_{\mathbf{n}}(\mathbb{R}^q)$. From here on we use ρ (resp. $\rho_{\mathbf{J}}$) as shorthand for $\rho_0([\mathbf{u}])$ (resp. $\rho_{\mathbf{J}}([\mathbf{u}])$).

Definition 3.1. A difference moving frame is a discrete moving frame such that \mathcal{M} is a prolongation space and the cross-section over $\mathbf{n} + \mathbf{J}$ is represented on \mathcal{M} by $S_{\mathbf{J}}\mathcal{K}$.

The normalized invariants for difference moving frames are

$$I_{\mathbf{K},\mathbf{J}} := \rho_{\mathbf{K}} \cdot \mathbf{u}_{\mathbf{J}} = (S_{\mathbf{K}}\rho) \cdot (S_{\mathbf{J}}\mathbf{u}).$$

By definition, $S_i I_{\mathbf{K},\mathbf{J}} = I_{\mathbf{K}+1_i, \mathbf{J}+1_i}$. Hence, every invariant $I_{\mathbf{K},\mathbf{J}}$ can be expressed as a shift of

$$I_{\mathbf{K}-\mathbf{J},0} = (S_{\mathbf{K}-\mathbf{J}}\rho) \cdot \mathbf{u}. \tag{3.1}$$

Definition 3.2 (discrete Maurer–Cartan invariants). Given a right discrete moving frame $\rho([\mathbf{u}])$ (which commonly is expressed as a matrix), the right discrete *Maurer–Cartan invariants* are the components of

$$K_{(i)} = (S_i \rho) \rho^{-1} = \iota(S_i \rho), \quad i = 1, \dots, m,$$

together with their shifts $S_{\mathbf{J}}K_{(i)}$. Invariance of $K_{(i)}$ follows from equivariance of the frame.

Definition 3.3. A set of invariants is a *generating set* for the algebra of functions of difference invariants if any difference invariant in the algebra can be written as a function of elements of the generating set and their shifts.

The invariantization of a multiply-shifted frame is obtained by concatenating the matrices of Maurer–Cartan invariants and their shifts. For instance,

$$\iota(S_i S_j \rho) = (S_i S_j \rho) \rho^{-1} = (S_j K_{(i)}) K_{(j)}.$$

Consequently, (3.1) can be written as $I_{\mathbf{K}-\mathbf{J},\mathbf{0}} = \{(S_{\mathbf{K}-\mathbf{J}} \rho)^{-1}\} I_{\mathbf{0},\mathbf{0}}$, where the term in braces is a concatenation of the Maurer–Cartan invariants and their shifts. This establishes the following result.

Proposition 3.4. *Given a right difference moving frame $\rho([\mathbf{u}])$, the set of all invariants is generated by $I_{\mathbf{0},\mathbf{0}} = \rho \cdot \mathbf{u}$ and the set of components of $K_{(j)} = (S_j \rho) \rho^{-1}$, $j = 1, \dots, m$.*

Definition 3.5 (syzygy). A syzygy on a set of invariants is a relation between the invariants that expresses functional dependency.

In other words, a syzygy on a set of invariants is a function of the invariants that becomes an identity when it is expressed in terms of the underlying variables $[\mathbf{u}]$. In general, there are syzygies between the invariants in Proposition 3.4, which lead to useful recurrence relations. Such relations enable all of these invariants to be expressed in terms of a small² generating set of invariants, κ^β , and their shifts, $\kappa_{\mathbf{J}}^\beta := S_{\mathbf{J}} \kappa^\beta$.

Example 3.6. As a running example to illustrate the theory, we use the Lagrangian

$$L = \ln \left| \frac{u_{1,0} - u_{0,1}}{u_{1,1} - u_{0,0}} \right|, \quad (3.2)$$

where $\mathbf{n} = (n^1, n^2)$ and $u_{i,j}$ represents the value of u at $(n^1 + i, n^2 + j)$. The Euler–Lagrange equation is

$$E_u(L) = \frac{1}{u_{1,1} - u_{0,0}} - \frac{1}{u_{-1,1} - u_{0,0}} - \frac{1}{u_{1,-1} - u_{0,0}} + \frac{1}{u_{-1,-1} - u_{0,0}} = 0, \quad (3.3)$$

which is a Toda-type equation that is satisfied by all solutions of the autonomous dpKdV equation and the cross-ratio equation (see [7] for details). This equation is partitioned into two independent components, with $n^1 + n^2$ being either odd or even.

The Lagrangian (3.2) has a six-parameter Lie group of variational symmetries, whose infinitesimal generators are linear combinations of

$$\begin{aligned} \mathbf{v}_1 &= \partial_{u_{0,0}}, & \mathbf{v}_2 &= u_{0,0} \partial_{u_{0,0}}, & \mathbf{v}_3 &= u_{0,0}^2 \partial_{u_{0,0}}, \\ \mathbf{v}_4 &= (-1)^{n^1+n^2} \partial_{u_{0,0}}, & \mathbf{v}_5 &= (-1)^{n^1+n^2} u_{0,0} \partial_{u_{0,0}}, & \mathbf{v}_6 &= (-1)^{n^1+n^2} u_{0,0}^2 \partial_{u_{0,0}}, \end{aligned}$$

where we use ∂_z as shorthand for $\partial/\partial z$ from here on. The corresponding characteristics are

$$\begin{aligned} Q_1 &= 1, & Q_2 &= u_{0,0}, & Q_3 &= u_{0,0}^2, & Q_4 &= (-1)^{n^1+n^2}, \\ Q_5 &= (-1)^{n^1+n^2} u_{0,0}, & Q_6 &= (-1)^{n^1+n^2} u_{0,0}^2, \end{aligned}$$

and the prolonged generators are

$$\text{pr } \mathbf{v}_r = (S_1^i S_2^j Q_r) \partial_{u_{i,j}}.$$

²The number of generating invariants depends on the number of dependent and independent variables, as well as on the group G and the normalization that is used. Finding a relation between these quantities is an open problem.

Only the symmetries generated by linear combinations of \mathbf{v}_1 , \mathbf{v}_2 and \mathbf{v}_4 leave the Lagrangian invariant. (The other generators produce divergence terms, although these can be absorbed without changing the Euler–Lagrange equation, by adding a divergence to the Lagrangian.)

Consider the two-parameter Lie group action generated by \mathbf{v}_1 and \mathbf{v}_2 ; this is

$$g: u_{i,j} \mapsto \tilde{u}_{i,j} = bu_{i,j} + a,$$

which is defined for every $a \in \mathbb{R}$ and $b \in \mathbb{R}^+$. For the half-space $\mathcal{U} = \{\mathcal{M}: u_{1,1} > u_{0,0}\}$, we choose the normalization equations³ (2.5) to be

$$\tilde{u}_{0,0} = 0, \quad \tilde{u}_{1,1} = 1.$$

Then the frame ρ is the group element with the parameters

$$a = \frac{-u_{0,0}}{u_{1,1} - u_{0,0}}, \quad b = \frac{1}{u_{1,1} - u_{0,0}}.$$

For this frame, the invariantization of $u_{i,j}$ is

$$\iota(u_{i,j}) = (\tilde{u}_{i,j})|_{g=\rho} = \frac{u_{i,j} - u_{0,0}}{u_{1,1} - u_{0,0}}.$$

The basic Maurer–Cartan invariants are

$$\kappa = \iota(u_{1,0}) = \frac{u_{1,0} - u_{0,0}}{u_{1,1} - u_{0,0}}, \quad \lambda = \iota(u_{0,1}) = \frac{u_{0,1} - u_{0,0}}{u_{1,1} - u_{0,0}}.$$

These two invariants generate all invariants $\iota(u_{i,j})$.

As an example, we show how to find $\iota(u_{2,1})$ in terms of these generating invariants and their shifts. First shift λ to involve $u_{2,1}$ and lower shifts of κ and λ :

$$\lambda_{1,0} = \frac{u_{1,1} - u_{1,0}}{u_{2,1} - u_{1,0}}.$$

As shifts of invariants are invariant, the replacement rule yields

$$\lambda_{1,0} = \frac{\iota(u_{1,1}) - \iota(u_{1,0})}{\iota(u_{2,1}) - \iota(u_{1,0})} = \frac{1 - \kappa}{\iota(u_{2,1}) - \kappa}.$$

Rearranging,

$$\iota(u_{2,1}) = \kappa + \frac{1 - \kappa}{\lambda_{1,0}}.$$

Similarly, the replacement rule gives

$$S_1 \iota(u_{i,j}) = \frac{u_{i+1,j} - u_{1,0}}{u_{2,1} - u_{1,0}} = \frac{\iota(u_{i+1,j}) - \kappa}{\iota(u_{2,1}) - \kappa},$$

which leads to the first of two general recurrence relations:

$$\iota(u_{i+1,j}) = \kappa + \left(\frac{1 - \kappa}{\lambda_{1,0}} \right) S_1 \iota(u_{i,j}), \quad (3.4)$$

$$\iota(u_{i,j+1}) = \lambda + \left(\frac{1 - \lambda}{\kappa_{0,1}} \right) S_2 \iota(u_{i,j}). \quad (3.5)$$

Note that the invariantization operator ι does not commute with the shift operators.

The invariant $\iota(u_{2,2})$ can be calculated from the identity $\iota(u_{1,1}) = 1$ in two ways: either use (3.4) first, then (3.5), or vice versa. This leads to the following syzygy between the generating invariants and their shifts:

$$\frac{(\lambda - 1)(\kappa_{0,1} - 1)}{\kappa_{0,1}\lambda_{1,1}} = \frac{(\kappa - 1)(\lambda_{1,0} - 1)}{\lambda_{1,0}\kappa_{1,1}}. \quad (3.6)$$

The syzygy (3.6) arises because two dependent variables (κ, λ) are used instead of one (u) .

³For the other half-space, with $u_{1,1} < u_{0,0}$, an appropriate normalization is $\tilde{u}_{0,0} = 0$ and $\tilde{u}_{1,1} = -1$.

3.2 Differential invariants and syzygies

From Remark 2.3, one can derive the Euler–Lagrange equations by regarding the variables $u_{\mathbf{J}}^{\alpha}$ as depending smoothly on a continuous parameter t . The same is true in the context of invariants, if one stipulates that

- t is invariant under the Lie group action;
- every shift commutes with differentiation with respect to t (as is required for differential-difference equations [21]).

This approach was used in Mansfield et al. [15] to invariantize the Euler–Lagrange equations for OΔEs. We now extend it to PΔEs.

As t is invariant, the Lie group action (for point transformations) extends to the first-order jet space of \mathcal{M} as follows:

$$g \cdot \frac{du_{\mathbf{J}}^{\alpha}}{dt} = \frac{d(g \cdot u_{\mathbf{J}}^{\alpha})}{dt} = \frac{\partial(g \cdot u_{\mathbf{J}}^{\alpha})}{\partial u_{\mathbf{J}}^{\delta}} \frac{du_{\mathbf{J}}^{\delta}}{dt}.$$

As the action is free and regular on \mathcal{M} , it will remain so on the jet space and we may use the same frame to find the first-order differential invariants. Let

$$\sigma^{\alpha} := \iota \left(\frac{du^{\alpha}}{dt} \right) = \frac{\partial(g \cdot u^{\alpha})}{\partial u^{\delta}} \Big|_{g=\rho} \frac{du^{\delta}}{dt}, \quad \alpha = 1, \dots, q. \quad (3.7)$$

The Jacobian matrix of any Lie group transformation on \mathcal{M} is necessarily non-singular, so (3.7) can be inverted, as follows:

$$\frac{du^{\delta}}{dt} = \theta_{\alpha}^{\delta}([\mathbf{u}])\sigma^{\alpha}.$$

Here the coefficients $\theta_{\alpha}^{\delta}([\mathbf{u}])$ are the components of the inverse of the Jacobian matrix, evaluated on the moving frame. Consequently, for each \mathbf{J} ,

$$\iota \left(\frac{du_{\mathbf{J}}^{\delta}}{dt} \right) = \iota(\mathbf{S}_{\mathbf{J}}(\theta_{\alpha}^{\delta}([\mathbf{u}])\sigma^{\alpha})) = \iota(\mathbf{S}_{\mathbf{J}}\theta_{\alpha}^{\delta}([\mathbf{u}]))\mathbf{S}_{\mathbf{J}}\sigma^{\alpha}. \quad (3.8)$$

So all of the first-order differential invariants can be written in terms of the generating invariants κ^{β} , the generating differential invariants σ^{α} , and their shifts. In particular,

$$\sigma^{\delta} = \iota \left(\frac{du^{\delta}}{dt} \right) = \iota(\theta_{\alpha}^{\delta}([\mathbf{u}]))\sigma^{\alpha},$$

so the invariantization of the matrix $(\theta_{\alpha}^{\delta}([\mathbf{u}]))$ is the identity matrix.

To calculate the invariantized Euler–Lagrange equations, it is necessary to determine the differential syzygies

$$\frac{d\kappa^{\beta}}{dt} = \iota \left(\frac{\partial \kappa^{\beta}}{\partial u_{\mathbf{J}}^{\alpha}} \right) \iota \left(\frac{du_{\mathbf{J}}^{\alpha}}{dt} \right) = \mathcal{H}_{\alpha}^{\beta} \sigma^{\alpha}.$$

The terms $\mathcal{H}_{\alpha}^{\beta}$ are linear difference operators whose coefficients are functions of κ^{β} and their shifts.

Example 3.7 (Example 3.6 cont.). We now find the differential invariants for the running example. The action of the group on the derivative $u'_{i,j} = du_{i,j}/dt$ is

$$g \cdot u'_{i,j} = \frac{\partial(g \cdot u_{i,j})}{\partial u_{i,j}} u'_{i,j} = b u'_{i,j}.$$

Therefore, the first-order differential invariants are

$$\iota(u'_{i,j}) = \frac{u'_{i,j}}{u_{1,1} - u_{0,0}}.$$

The generating differential invariant is

$$\sigma = \frac{u'_{0,0}}{u_{1,1} - u_{0,0}},$$

so (3.8) amounts to

$$\iota(u'_{i,j}) = \{\iota(u_{i+1,j+1}) - \iota(u_{i,j})\} S_1^i S_2^j \sigma.$$

For instance,

$$\iota(u'_{0,1}) = \{\iota(u_{1,2}) - \iota(u_{0,1})\} S_2 \sigma = \frac{1 - \lambda}{\kappa_{0,1}} S_2 \sigma.$$

The derivatives of the generating invariants are

$$\begin{aligned} \frac{d\kappa}{dt} &= \frac{u'_{1,0} - u'_{0,0}}{u_{1,1} - u_{0,0}} - \frac{(u_{1,0} - u_{0,0})(u'_{1,1} - u'_{0,0})}{(u_{1,1} - u_{0,0})^2}, \\ \frac{d\lambda}{dt} &= \frac{u'_{0,1} - u'_{0,0}}{u_{1,1} - u_{0,0}} - \frac{(u_{0,1} - u_{0,0})(u'_{1,1} - u'_{0,0})}{(u_{1,1} - u_{0,0})^2}. \end{aligned} \quad (3.9)$$

Invariantizing (3.9), using the replacement rule (2.6), we obtain

$$\begin{aligned} \frac{d\kappa}{dt} &= \iota(u'_{1,0}) - \kappa \iota(u'_{1,1}) + (\kappa - 1) \iota(u'_{0,0}), \\ \frac{d\lambda}{dt} &= \iota(u'_{0,1}) - \lambda \iota(u'_{1,1}) + (\lambda - 1) \iota(u'_{0,0}). \end{aligned} \quad (3.10)$$

Replacing each $\iota(u'_{i,j})$ in (3.10) by its expression in terms of σ gives

$$\frac{d\kappa}{dt} = \mathcal{H}_\kappa \sigma, \quad \frac{d\lambda}{dt} = \mathcal{H}_\lambda \sigma,$$

where

$$\begin{aligned} \mathcal{H}_\kappa &= \frac{1 - \kappa}{\lambda_{1,0}} S_1 - \frac{\kappa(\kappa - 1)(\lambda_{1,0} - 1)}{\lambda_{1,0} \kappa_{1,1}} S_1 S_2 + (\kappa - 1) \text{id}, \\ \mathcal{H}_\lambda &= \frac{1 - \lambda}{\kappa_{0,1}} S_2 - \frac{\lambda(\kappa - 1)(\lambda_{1,0} - 1)}{\lambda_{1,0} \kappa_{1,1}} S_1 S_2 + (\lambda - 1) \text{id}. \end{aligned} \quad (3.11)$$

4 The invariant formulation of the Euler–Lagrange equations

Here we show how to calculate the Euler–Lagrange equations, in terms of invariants, for a given invariant difference Lagrangian. Any such Lagrangian, $L(\mathbf{n}, [\mathbf{u}])$, can be written in terms of the generating invariants κ^β and their shifts $\kappa_{\mathbf{J}}^\beta = S_{\mathbf{J}}\kappa^\beta$:

$$L(\mathbf{n}, [\mathbf{u}]) = L^\kappa(\mathbf{n}, [\kappa]). \quad (4.1)$$

The key result is the following proposition, which generalizes OΔE invariantization [15] to PΔEs.

Proposition 4.1 (invariant Euler–Lagrange equations). *Suppose that the Lagrangian $L(\mathbf{n}, [\mathbf{u}])$ is invariant under an R -parameter Lie group of point transformations, so that (4.1) holds. Given the differential syzygies, $d\kappa^\beta/dt = \mathcal{H}_\alpha^\beta \sigma^\alpha$, the following identity holds:*

$$E_{u^\alpha}(L) \frac{du^\alpha}{dt} = ((\mathcal{H}_\alpha^\beta)^\dagger E_{\kappa^\beta}(L^\kappa)) \sigma^\alpha, \quad (4.2)$$

where

$$E_{\kappa^\beta} = S_{-\mathbf{J}} \frac{\partial}{\partial \kappa_{\mathbf{J}}^\beta}$$

is the difference Euler operator with respect to κ^β . The invariantization of the original system of Euler–Lagrange equations is

$$\iota(E_{u^\alpha}(L)) = (\mathcal{H}_\alpha^\beta)^\dagger E_{\kappa^\beta}(L^\kappa) = 0, \quad \alpha = 1, \dots, q. \quad (4.3)$$

Proof. In the original coordinates,

$$\frac{dL}{dt} = \frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \frac{du_{\mathbf{J}}^\alpha}{dt} = E_{u^\alpha}(L) \frac{du^\alpha}{dt} + \text{Div}(A_{\mathbf{u}}); \quad (4.4)$$

here summation by parts has produced the difference divergence

$$\text{Div}(A_{\mathbf{u}}) = \sum_{\mathbf{J}} (S_{\mathbf{J}} - \text{id}) \left(S_{-\mathbf{J}} \left(\frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \right) (u^\alpha)' \right) =: D_{n^i} (A_\alpha^i(\mathbf{n}, [\mathbf{u}]) (u^\alpha)'), \quad (4.5)$$

where each A_α^i is a linear difference operator. In the invariant coordinates,

$$\begin{aligned} \frac{dL^\kappa}{dt} &= \frac{\partial L^\kappa}{\partial \kappa_{\mathbf{K}}^\beta} \frac{d\kappa_{\mathbf{K}}^\beta}{dt} = E_{\kappa^\beta}(L^\kappa) \frac{d\kappa^\beta}{dt} + \text{Div}(A_{\kappa}) = E_{\kappa^\beta}(L^\kappa) \mathcal{H}_\alpha^\beta \sigma^\alpha + \text{Div}(A_{\kappa}) \\ &= \{(\mathcal{H}_\alpha^\beta)^\dagger (E_{\kappa^\beta}(L^\kappa))\} \sigma^\alpha + \text{Div}(A_{\mathcal{H}} + A_{\kappa}). \end{aligned} \quad (4.6)$$

Here, there are two contributions to the difference divergence:

$$\begin{aligned} \text{Div}(A_{\kappa}) &= \sum_{\mathbf{K}} (S_{\mathbf{K}} - \text{id}) \left(S_{-\mathbf{K}} \left(\frac{\partial L}{\partial \kappa_{\mathbf{K}}^\beta} \right) (\kappa^\beta)' \right) =: D_{n^i} (F_\beta^i(\mathbf{n}, [\kappa]) (\kappa^\beta)'), \\ \text{Div}(A_{\mathcal{H}}) &= E_{\kappa^\beta}(L^\kappa) \mathcal{H}_\alpha^\beta \sigma^\alpha - \{(\mathcal{H}_\alpha^\beta)^\dagger (E_{\kappa^\beta}(L^\kappa))\} \sigma^\alpha =: D_{n^i} (H_\alpha^i(\mathbf{n}, [\kappa]) \sigma^\alpha), \end{aligned}$$

where F_β^i and H_α^i are linear difference operators. The difference between (4.4) and (4.6), summed over \mathbf{n} to annihilate the divergence terms, is

$$0 = \sum_{\mathbf{n}} \left(\frac{dL}{dt} - \frac{dL^\kappa}{dt} \right) = \sum_{\mathbf{n}} (E_{u^\alpha}(L) (u^\alpha)' - (\mathcal{H}_\alpha^\beta)^\dagger (E_{\kappa^\beta}(L^\kappa)) \sigma^\alpha).$$

Note that the invariants at a particular \mathbf{n} are invariantized by the frame at that \mathbf{n} . Each $u^\alpha(\mathbf{n}, t)$ has arbitrary (smooth) dependence on t ; there is no link between $u^\alpha(\mathbf{n}, t)$ and $u^\alpha(\mathbf{m}, t)$ for $\mathbf{n} \neq \mathbf{m}$. Therefore, (4.2) holds for each \mathbf{n} , and so

$$\iota(\mathbf{E}_{u^\alpha}(\mathbf{L}))\sigma^\alpha = (\mathcal{H}_\alpha^\beta)^\dagger(\mathbf{E}_{\kappa^\beta}(L^\kappa))\sigma^\alpha.$$

Equation (4.3) follows from the independence of the differential invariants σ^α . \blacksquare

Corollary 4.2. *Under the conditions of Proposition 4.1, using the notation in its proof,*

$$\text{Div}(A_{\mathbf{u}}) = \text{Div}(A_{\mathcal{H}} + A_{\kappa}).$$

Proof. Compare (4.4) and (4.6), taking (4.2) into account. \blacksquare

Example 4.3 (Example 3.6 cont.). The Lagrangian (3.2) is written in terms of the generating invariants $\kappa = \iota(u_{1,0})$ and $\lambda = \iota(u_{0,1})$ as

$$L^\kappa = \ln |\kappa - \lambda|.$$

Applying the Euler operators with respect to κ and λ gives

$$\mathbf{E}_\kappa(L^\kappa) = \frac{1}{\kappa - \lambda}, \quad \mathbf{E}_\lambda(L^\kappa) = \frac{-1}{\kappa - \lambda}.$$

From (3.11),

$$\begin{aligned} \mathcal{H}_\kappa^\dagger &= \frac{1 - \kappa_{-1,0}}{\lambda} \mathbf{S}_1^{-1} - \frac{\kappa_{-1,-1}(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa \lambda_{0,-1}} \mathbf{S}_1^{-1} \mathbf{S}_2^{-1} + (\kappa - 1)\text{id}, \\ \mathcal{H}_\lambda^\dagger &= \frac{1 - \lambda_{0,-1}}{\kappa} \mathbf{S}_2^{-1} - \frac{\lambda_{-1,-1}(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa \lambda_{0,-1}} \mathbf{S}_1^{-1} \mathbf{S}_2^{-1} + (\lambda - 1)\text{id}. \end{aligned}$$

By Proposition 4.1, the invariant Euler–Lagrange equation for the running example is

$$\begin{aligned} 0 &= \mathcal{H}_\kappa^\dagger \mathbf{E}_\kappa(L^\kappa) + \mathcal{H}_\lambda^\dagger \mathbf{E}_\lambda(L^\kappa) \\ &= \frac{1 - \kappa_{-1,0}}{\lambda(\kappa_{-1,0} - \lambda_{-1,0})} - \frac{1 - \lambda_{0,-1}}{\kappa(\kappa_{0,-1} - \lambda_{0,-1})} - \frac{(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa \lambda_{0,-1}} + 1. \end{aligned}$$

This is the invariantization of the Toda-type equation (3.3) given by the chosen normalization, which has the advantage that L^κ depends only on unshifted invariants.

5 Conservation laws

For ODEs [4, 5, 12], PDEs [6] and OΔEs [15, 16], it has been shown that the conservation laws associated with an R -parameter Lie group of variational point symmetries are equivariant and can be written in terms of the invariants and the moving frame. For PΔEs, we use the same reasoning as in the papers above to show that the R conservation laws can be written in the equivariant form

$$D_{n^i} \{V_l^i(\mathbf{n}, [\boldsymbol{\kappa}]) a_r^l(\rho)\} = 0, \quad r = 1, \dots, R,$$

where $a_r^l(\rho)$ are the components of the adjoint representation of ρ and each $V_l^i(\mathbf{n}, [\boldsymbol{\kappa}])$ is invariant.

For a Lagrangian $L(\mathbf{n}, [\mathbf{u}])$ that is invariant under the one-parameter group generated by \mathbf{v}_r , the corresponding conservation law (2.4) given by Noether's theorem is

$$\sum_{\mathbf{J}} (\mathbf{S}_{\mathbf{J}} - \text{id}) \left(Q_r^\alpha(\mathbf{n}, [\mathbf{u}]) \mathbf{S}_{-\mathbf{J}} \frac{\partial L}{\partial u_{\mathbf{J}}^\alpha} \right) = 0. \quad (5.1)$$

The left-hand side of (5.1) is almost the same as $\text{Div } A_{\mathbf{u}}$ in (4.5), with the exception that $(u^\alpha)'$ in $A_{\mathbf{u}}$ is replaced by the characteristic component $Q_r^\alpha(\mathbf{n}, [\mathbf{u}])$. This replacement can be achieved by substituting the group parameter ε^r for t and evaluating the result at $\varepsilon^r = 0$.

By Corollary 4.2, the conservation law $\text{Div}(A_{\mathbf{u}}) = 0$ amounts to

$$\text{Div}(A_{\boldsymbol{\kappa}} + A_{\mathcal{H}}) = D_{n^i} (F_\beta^i(\mathbf{n}, [\boldsymbol{\kappa}]) (\boldsymbol{\kappa}^\beta)') + H_\alpha^i(\mathbf{n}, [\boldsymbol{\kappa}]) \sigma^\alpha = 0.$$

If t is the group parameter ε^r then $(\boldsymbol{\kappa}^\beta)' = 0$, because each $\boldsymbol{\kappa}^\beta$ is invariant. Consequently, the conservation law given by Noether's theorem is

$$D_{n^i} (H_\alpha^i(\mathbf{n}, [\boldsymbol{\kappa}]) \sigma^\alpha) = 0, \tag{5.2}$$

where σ^α is the invariantization of the tangent vector to the group generated by \mathbf{v}_r , evaluated at $\varepsilon = 0$.

Proposition 5.1. *Suppose that the conditions of Proposition 4.1 hold. If the linear difference operators in (5.2) are*

$$H_\alpha^i(\mathbf{n}, [\boldsymbol{\kappa}]) = \mathcal{C}_\alpha^{i, \mathbf{J}}(\mathbf{n}, [\boldsymbol{\kappa}]) \mathbf{S}_{\mathbf{J}},$$

then Noether's theorem gives the R conservation laws,

$$D_{n^i} \{ \mathcal{C}_\alpha^{i, \mathbf{J}}(\mathbf{n}, [\boldsymbol{\kappa}]) \mathbf{S}_{\mathbf{J}} \{ \iota(Q_s^\alpha(\mathbf{n}, \mathbf{u})) a_r^s(\rho) \} \} = 0, \quad r = 1, \dots, R.$$

Proof. This proof is a slimmed-down analogue of its counterpart for OΔEs (see [15]). We replace t by the parameter ε^r and let $\widehat{\mathbf{u}}(\mathbf{n}, \mathbf{u}, \varepsilon^r)$ be the orbit of the one-parameter local Lie group generated by \mathbf{v}_r . Let \widehat{u}_r^α denote $d\widehat{u}^\alpha/d\varepsilon^r$ and note that $\widehat{\mathbf{u}}(\mathbf{n}, \mathbf{u}, 0) = \mathbf{u}$. By the chain rule,

$$\sigma_r^\alpha := \iota(\widehat{u}_r^\alpha) = \left\{ \frac{\partial(g \cdot \widehat{u}^\alpha)}{\partial \widehat{u}^\beta} \widehat{u}_r^\beta \right\} \Big|_{g=\rho}.$$

In particular, from (2.2),

$$\sigma_r^\alpha \Big|_{\varepsilon^r=0} = \left\{ \frac{\partial(g \cdot u^\alpha)}{\partial u^\beta} Q_r^\beta(\mathbf{n}, \mathbf{u}) \right\} \Big|_{g=\rho} = \{ Q_s^\alpha(\mathbf{n}, g \cdot \mathbf{u}) a_r^s(g) \} \Big|_{g=\rho} = \iota(Q_s^\alpha(\mathbf{n}, \mathbf{u})) a_r^s(\rho).$$

Substitute $\sigma_r^\alpha \Big|_{\varepsilon^r=0}$ for σ^α in (5.2) to complete the proof. ■

By the prolongation formula, the conservation laws amount to

$$D_{n^i} \{ \mathcal{C}_\alpha^{i, \mathbf{J}}(\mathbf{n}, [\boldsymbol{\kappa}]) (\mathbf{S}_{\mathbf{J}} \iota(Q_s^\alpha)) a_r^s(\rho_{\mathbf{J}}) \} = 0.$$

The adjoint representation is a Lie group representation, so

$$a_r^s(\rho_{\mathbf{J}}) = a_l^s(\rho_{\mathbf{J}} \rho^{-1}) a_r^l(\rho) = a_l^s(\iota(\rho_{\mathbf{J}})) a_r^l(\rho).$$

This leads to the following corollary.

Corollary 5.2. *The conservation laws for a difference frame may be written in the form*

$$D_{n^i} \{ V_l^i(\mathbf{n}, [\boldsymbol{\kappa}]) a_r^l(\rho) \} = 0,$$

where

$$V_l^i(\mathbf{n}, [\boldsymbol{\kappa}]) = \mathcal{C}_\alpha^{i, \mathbf{J}}(\mathbf{n}, [\boldsymbol{\kappa}]) (\mathbf{S}_{\mathbf{J}} \iota(Q_s^\alpha)) a_l^s(\iota(\rho_{\mathbf{J}})).$$

Example 5.3 (Example 3.6 cont.). For the running example, the equivariant conservation laws are obtained as follows:

$$\begin{aligned}
\text{Div}(A_{\mathcal{H}}) &= E_{\kappa}(L^{\kappa})\mathcal{H}_{\kappa}\sigma + E_{\lambda}(L^{\kappa})\mathcal{H}_{\lambda}\sigma - \{(\mathcal{H}_{\kappa})^{\dagger}E_{\kappa}(L^{\kappa}) + (\mathcal{H}_{\lambda})^{\dagger}E_{\lambda}(L^{\kappa})\}\sigma \\
&= (S_1 - \text{id}) \left(\frac{1 - \kappa_{-1,0}}{\lambda(\kappa_{-1,0} - \lambda_{-1,0})} \sigma \right) + (S_2 - \text{id}) \left(\frac{\lambda_{0,-1} - 1}{\kappa(\kappa_{0,-1} - \lambda_{0,-1})} \sigma \right) \\
&\quad + (S_1 S_2 - \text{id}) \left(-\frac{(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa\lambda_{0,-1}} \sigma \right) \\
&= D_{n^1} \left\{ \frac{1 - \kappa_{-1,0}}{\lambda(\kappa_{-1,0} - \lambda_{-1,0})} \sigma - \frac{(\kappa_{-1,0} - 1)(\lambda - 1)}{\kappa_{0,1}\lambda} S_2 \sigma \right\} \\
&\quad + D_{n^2} \left\{ \frac{\lambda_{0,-1} - 1}{\kappa(\kappa_{0,-1} - \lambda_{0,-1})} \sigma - \frac{(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa\lambda_{0,-1}} \sigma \right\}.
\end{aligned}$$

The invariantized (unprolonged) infinitesimals are $\iota(Q_1) = 1$ and $\iota(Q_2) = 0$, so

$$\iota(Q_s(\mathbf{n}, \mathbf{u}))a_r^s(\rho) = a_r^1(\rho).$$

The adjoint action of $g: u \mapsto \tilde{u} = bu + a$ on the infinitesimal generators gives $\mathbf{v}_1 = b\tilde{\mathbf{v}}_1$ and $\mathbf{v}_2 = -a\tilde{\mathbf{v}}_1 + \tilde{\mathbf{v}}_2$, so the components of the adjoint matrix are

$$a_1^1 = b, \quad a_1^2 = 0, \quad a_2^1 = -a, \quad a_2^2 = 1.$$

On the frame ρ (at (m, n)), these components are

$$a_1^1(\rho) = \frac{1}{u_{1,1} - u_{0,0}}, \quad a_1^2(\rho) = 0, \quad a_2^1(\rho) = \frac{u_{0,0}}{u_{1,1} - u_{0,0}}, \quad a_2^2(\rho) = 1. \quad (5.3)$$

Applying S_2 to (5.3) gives the components on the frame at $(m, n + 1)$. In particular,

$$\begin{aligned}
a_1^1(S_2\rho) &= \frac{1}{u_{1,2} - u_{0,1}} = \frac{\kappa_{0,1}}{1 - \lambda} a_1^1(\rho), \\
a_2^1(S_2\rho) &= \frac{u_{0,1}}{u_{1,2} - u_{0,1}} = \frac{\kappa_{0,1}}{1 - \lambda} a_2^1(\rho) + \frac{\lambda\kappa_{0,1}}{1 - \lambda} a_2^2(\rho).
\end{aligned}$$

Consequently, in terms of the frame (5.3), the equivariant versions of the conservation laws arising from Noether's theorem are

$$\begin{aligned}
0 &= D_{n^1} \left\{ \left(\frac{1 - \kappa_{-1,0}}{\lambda(\kappa_{-1,0} - \lambda_{-1,0})} + \frac{\kappa_{-1,0} - 1}{\lambda} \right) a_1^1(\rho) \right\} \\
&\quad + D_{n^2} \left\{ \left(\frac{\lambda_{0,-1} - 1}{\kappa(\kappa_{0,-1} - \lambda_{0,-1})} - \frac{(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa\lambda_{0,-1}} \right) a_1^1(\rho) \right\}, \\
0 &= D_{n^1} \left\{ \left(\frac{1 - \kappa_{-1,0}}{\lambda(\kappa_{-1,0} - \lambda_{-1,0})} + \frac{\kappa_{-1,0} - 1}{\lambda} \right) a_2^1(\rho) + (\kappa_{-1,0} - 1)a_2^2(\rho) \right\} \\
&\quad + D_{n^2} \left\{ \left(\frac{\lambda_{0,-1} - 1}{\kappa(\kappa_{0,-1} - \lambda_{0,-1})} - \frac{(\kappa_{-1,-1} - 1)(\lambda_{0,-1} - 1)}{\kappa\lambda_{0,-1}} \right) a_2^1(\rho) \right\}.
\end{aligned}$$

6 Differential-difference structure

Differential-difference moving frames are set in a continuous space that embodies prolongation with respect to both derivatives and shifts (see [21]). Just as for PΔEs, the discrete independent variables, $\mathbf{n} = (n^1, \dots, n^m)$, and the corresponding shift operators are invariant. In general, the




Figure 3. A graph on (ordered) slices in the total space $\mathcal{T} = \mathbb{R} \times \mathbb{Z} \times \mathbb{R}$.

invariant derivative operators do not commute with one another or with the shift operators, a problem that can be resolved with substantial technical machinery.⁴ However, such machinery is not needed for systems of DΔEs for $\mathbf{u} = (u^1, \dots, u^q) \in \mathbb{R}^q$ with just one continuous independent variable, x , provided that the group action on x is sufficiently simple. For consistency with our presentation of difference moving frames, we restrict attention to such systems and actions.

The total space, $\mathcal{T} = \mathbb{R} \times \mathbb{Z}^m \times \mathbb{R}^q$, consists of a continuous *slice*, $\mathcal{T}_{\mathbf{n}} = \mathbb{R} \times \{\mathbf{n}\} \times \mathbb{R}^q$, over each $\mathbf{n} \in \mathbb{Z}^m$. Consequently, each graph on \mathcal{T} defined by $\mathbf{u} = \mathbf{f}(x, \mathbf{n})$ restricts to a graph on every slice, as illustrated in Figure 3. We consider only graphs that are smooth on each slice, which can be prolonged by differentiation as many times as needed.⁵ The differential prolongation structure on the slice over any \mathbf{n} is embodied by the infinite jet space, $J^\infty(\mathcal{T}_{\mathbf{n}})$. This space has vertical coordinates $u_{j;\mathbf{0}}^\alpha$, where j denotes the number of derivatives with respect to x . In particular, $u_{0;\mathbf{0}}^\alpha$ represents u^α at \mathbf{n} .

The differential prolongation structure over any fixed \mathbf{n} is replicated over all other points $\mathbf{n} + \mathbf{K}$. Consequently, one can use difference prolongation to construct the (differential-difference) *prolongation space* over \mathbf{n} , denoted $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$. This space has coordinates $(x, (u_{j;\mathbf{K}}^\alpha))$, where $u_{j;\mathbf{K}}^\alpha$ represents the value of $u_{j;\mathbf{0}}^\alpha$ on the jet space over $\mathbf{n} + \mathbf{K}$.

Just as for difference equations, the horizontal translation $\mathbf{T}_{\mathbf{I}}$ maps \mathbf{n} to $\mathbf{n} + \mathbf{I}$ without changing any other coordinates. The pullback of

$$\mathbf{T}_{\mathbf{I}}: P(J^\infty(\mathcal{T}_{\mathbf{n}})) \rightarrow P(J^\infty(\mathcal{T}_{\mathbf{n}+\mathbf{I}}))$$

to $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$ is the shift operator $\mathbf{S}_{\mathbf{I}}$, which can be regarded as a product of unit forward shifts S_i , $i = 1, \dots, m$, and their inverses. This acts on functions $f \in C^\infty(P(J^\infty(\mathcal{T}_{\mathbf{n}})))$ as follows:

$$\mathbf{S}_{\mathbf{I}}f(x, \mathbf{n}, (u_{j;\mathbf{K}}^\alpha)) = f(x, \mathbf{n} + \mathbf{I}, (u_{j;\mathbf{K}+\mathbf{I}}^\alpha)).$$

All shift operators $\mathbf{S}_{\mathbf{K}}$ commute with one another and with the total derivative operator,

$$D = \frac{\partial}{\partial x} + u_{j+1;k} \frac{\partial}{\partial u_{j;k}}.$$

For notational consistency, we define $D_{(j)} := D^j$, so the vertical coordinates on $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$ may be written as $u_{j;\mathbf{K}} = \mathbf{S}_{\mathbf{K}}D_{(j)}u_{0;\mathbf{0}}^\alpha$.

The differential-difference divergence of a given $(1+m)$ -tuple of functions in $C^\infty(P(J^\infty(\mathcal{T}_{\mathbf{n}})))$, $A = (A^0; A^1, \dots, A^m)$, is

$$\text{Div}(A) := DA^0 + D_{n^i}A^i,$$

where $D_{n^i} = S_i - \text{id}$ is the i^{th} forward difference operator. Note: we sum over i from 1 to m only. A linear differential-difference operator on the prolongation space $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$ is an operator of

⁴Moving frames for differential Euler–Lagrange equations can be expressed naturally in terms of the variational bicomplex [10]; for a constructive approach that lends itself to symbolic computation, see [6].

⁵If graphs are only locally smooth, restrict attention to neighbourhoods in which they are smooth.

the form $\mathcal{H} = h^{j;\mathbf{K}} \mathbf{S}_{\mathbf{K}} D_{(j)}$, for given functions $h^{j;\mathbf{K}}$. The formal adjoint of \mathcal{H} is the unique operator \mathcal{H}^\dagger such that

$$f\mathcal{H}g - (\mathcal{H}^\dagger f)g \in \text{im}(\text{Div}).$$

Using the standard identities

$$\mathbf{S}_{\mathbf{K}}^\dagger = \mathbf{S}_{-\mathbf{K}}, \quad D_{(j)}^\dagger = (-D)_{(j)} := (-1)^j D^j,$$

one obtains

$$\mathcal{H}^\dagger f = (-D)_{(j)} \mathbf{S}_{-\mathbf{K}}(h^{j;\mathbf{K}} f).$$

Definition 6.1. A *conservation law* of a given system of DΔEs is a differential-difference divergence expression, $\mathcal{C} = \text{Div}(A)$, such that $\mathcal{C} = 0$ on all solutions of the system.

A given DΔE defines a variety in the continuous space $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$, on which it is possible to construct moving frames that respect both the differential and difference structures. Every Lie group, G , of point transformations of the total space, whose prolongation to $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$ preserves these structures, consists of *projectable* diffeomorphisms on each slice:

$$g: \mathcal{T} \rightarrow \mathcal{T}, \quad g: (x, \mathbf{n}, \mathbf{u}) \mapsto (g \cdot x, g \cdot \mathbf{n}, g \cdot \mathbf{u}) := (\tilde{x}(x), \mathbf{n}, \tilde{\mathbf{u}}(x, \mathbf{n}, \mathbf{u})).$$

The projectability condition arises because mappings for which \tilde{x} depends on \mathbf{n} or \mathbf{u} are incompatible with the prolongation structure (see [21] for details). This is a key distinction between DΔEs and PDEs, for which Lie point transformations do not have to be projectable. For each \mathbf{n} , the mapping g is a diffeomorphism, and therefore the Jacobian determinants,

$$J_x := \frac{d\tilde{x}}{dx}, \quad J_{\mathbf{u}} := \det \left(\frac{\partial \tilde{u}^\alpha}{\partial u^\beta} \right),$$

are nonzero. The prolongation conditions give the action of g on $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$ (recursively):

$$g \cdot u_{j+1;\mathbf{0}}^\alpha = \tilde{u}_{j+1;\mathbf{0}}^\alpha := \frac{D\tilde{u}_{j;\mathbf{0}}^\alpha}{D\tilde{x}}, \quad g \cdot u_{j;\mathbf{K}}^\alpha = \tilde{u}_{j;\mathbf{K}}^\alpha := \mathbf{S}_{\mathbf{K}} \tilde{u}_{j;\mathbf{0}}^\alpha.$$

A tilde $\tilde{}$ over a function or operator denotes that x and each $u_{j;\mathbf{K}}^\alpha$ are replaced by \tilde{x} and $\tilde{u}_{j;\mathbf{K}}^\alpha$ respectively.

From here on, we consider R -parameter Lie groups of (projectable) point transformations, whose generators are of the form

$$\mathbf{v}_r = \xi_r(x) \partial_x + \eta_r^\alpha(x, \mathbf{n}, \mathbf{u}) \partial_{u^\alpha}, \quad r = 1, \dots, R.$$

The adjoint action of g on the Lie algebra spanned by $\mathbf{v}_1, \dots, \mathbf{v}_R$ satisfies the identities

$$\mathbf{v}_r = a_r^s(g) \tilde{\mathbf{v}}_s, \quad r = 1, \dots, R. \tag{6.1}$$

The characteristic corresponding to \mathbf{v}_r is $\mathbf{Q}_r = (Q_r^1, \dots, Q_r^q)$, where

$$Q_r^\alpha = \eta_r^\alpha(x, \mathbf{n}, \mathbf{u}) - \xi_r(x) u_{1;\mathbf{0}}^\alpha.$$

This enables the differential-difference prolongation of \mathbf{v}_r to be written as

$$\text{pr } \mathbf{v}_r = \xi_r D + X_r, \quad \text{where } X_r := (\mathbf{S}_{\mathbf{K}} D_{(j)} Q_r^\alpha) \partial_{u_{j;\mathbf{K}}^\alpha}. \tag{6.2}$$

The operator X_r is the characteristic form of the generator; it acts only on the vertical coordinates of $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$. From here on, we write $\text{pr } \mathbf{v}_r$ simply as \mathbf{v}_r ; generators are assumed to be prolonged wherever this is needed. Note that (6.1) holds equally for the prolonged generators.

Lemma 6.2. *Let G be a Lie group of projectable transformations of the total space \mathcal{T} , and let $g \in G$. Using the above notation, the following identities hold:*

$$\xi_r D = a_r^s(g) \tilde{\xi}_s \tilde{D}; \quad (6.3)$$

$$X_r = a_r^s(g) \tilde{X}_s; \quad (6.4)$$

$$J_x \xi_r = \tilde{\xi}_s a_r^s(g); \quad (6.5)$$

$$\left(\frac{\partial \tilde{u}^\alpha}{\partial u^\beta} \right) Q_r^\beta = \tilde{Q}_s^\alpha a_r^s(g). \quad (6.6)$$

Proof. The chain rule and (6.1) give

$$\xi_r D = \xi_r \frac{d\tilde{x}}{dx} \tilde{D} = \mathbf{v}_r(\tilde{x}) \tilde{D} = a_r^s(g) \tilde{\mathbf{v}}_s(\tilde{x}) \tilde{D} = a_r^s(g) \tilde{\xi}_s \tilde{D},$$

which proves (6.3). To obtain (6.4), substitute (6.2) into the prolongation of (6.1) and take (6.3) into account. Then apply (6.3) to \tilde{x} and (6.4) to \tilde{u}^α to prove (6.5) and (6.6). ■

The construction of a differential-difference moving frame is essentially the same as that of a difference moving frame. Suppose that the Lie group G acts smoothly, freely and regularly on $\mathcal{M} \subset P(J^\infty(\mathcal{T}_{\mathbf{n}}))$, a finite prolongation space over \mathbf{n} whose coordinates include all relevant variables, including at least one derivative $u_{j;\mathbf{K}}^\alpha$, $j \geq 1$. Let the cross-section and frame on \mathcal{M} be \mathcal{K} and ρ , respectively. Such a moving frame uses the \mathbf{K}^{th} translate of \mathcal{K} at every other base point $\mathbf{n} + \mathbf{K}$; the cross-section and frame at $\mathbf{n} + \mathbf{K}$ are represented on \mathcal{M} by $S_{\mathbf{K}}\mathcal{K}$ and $\rho_{0;\mathbf{K}} = S_{\mathbf{K}}\rho$, respectively. This construction extends immediately to $P(J^\infty(\mathcal{T}_{\mathbf{n}}))$, because \mathcal{M} has arbitrary dimension.

As usual, the invariantization (denoted by ι) of a function, operator, etc., is obtained by evaluating the transformed quantity on the frame ρ . The freedom to choose a cross-section leads to the possibility that $\iota(x)$ may depend on one or more of the variables $u_{j;\mathbf{K}}^\alpha$, because $\tilde{x} = g \cdot x$ depends on x and the group parameters.

Lemma 6.3. *The invariantized total derivative, $\mathcal{D} = \iota(D)$, commutes with all (invariant) shift operators $S_{\mathbf{K}}$ if and only if $\mathcal{J} = J_x|_{g=\rho}$ is a function of x alone.*

Proof. Evaluating the identity $D = J_x \tilde{D}$ on ρ gives $D = \mathcal{J}\mathcal{D}$. Each S_i commutes with D , so

$$[\mathcal{D}, S_i] = \mathcal{J}^{-1} D S_i - (S_i \mathcal{J}^{-1}) S_i D = \mathcal{J}^{-1} [D, S_i] - (D_{n^i} \mathcal{J}^{-1}) S_i D = -(D_{n^i} \mathcal{J}^{-1}) S_i D.$$

The right-hand side is zero if and only if $S_i \mathcal{J} = \mathcal{J}$, which requires that \mathcal{J} is independent of n^i and $[\mathbf{u}]$. So a necessary and sufficient condition for every shift operator to commute with \mathcal{D} is that \mathcal{J} depends on neither \mathbf{n} nor $[\mathbf{u}]$. ■

Every group parameter that occurs in J_x also occurs in \tilde{x} , which may also have one further parameter associated with translations in x . So a sufficient condition for \mathcal{D} to commute with all shifts is that $\iota(x)$ depends on x alone.⁶ Indeed, even if \tilde{x} includes translations, the requirement for the parameters in J_x to depend on x alone (on ρ) ensures that there exist normalizations for which $\iota(x)$ depends on x only: simply set $\iota(x)$ to be constant. These observations motivate the following general definition.

⁶The only exception to this condition being necessary occurs when G includes translations in x . As an example, consider $(\tilde{x}, \tilde{u}) = (x + a, e^a u + b)$. The unusual normalization $\iota(u) = 0$, $\iota(u_{1;0}) = 1$ gives $a = -\ln(u_{1;0})$, so $\iota(x) = x - \ln(u_{1;0})$, but \mathcal{D} commutes with all shifts because $\mathcal{J} = 1$. A normalization for which $\iota(x)$ depends on x only is $\iota(x) = 0$, $\iota(u) = 0$.

Definition 6.4. Given a prolongation space⁷ \mathcal{P} on which the Lie group G acts smoothly, freely and regularly, let \mathcal{B} denote the space coordinatized by the continuous independent variables, $\mathbf{x} = (x^1, \dots, x^p)$. A moving frame, ρ , is *projectable* if $\iota(\mathbf{x}) = \rho \cdot \mathbf{x}$ is a function of \mathbf{x} alone.

If G consists of projectable transformations, then \mathcal{B} is invariant. Thus, one can restrict attention to the action of G on \mathcal{B} , ignoring those elements of G that fix every $\mathbf{x} \in \mathcal{B}$. Let $G_{\mathcal{B}}: \mathcal{B} \rightarrow \mathcal{B}$ denote the resulting Lie group of restricted transformations.

Lemma 6.5. *Let G be a Lie group of projectable transformations that acts smoothly, freely and regularly on a prolongation space \mathcal{P} . Then there exists a projectable moving frame on \mathcal{P} if and only if $G_{\mathcal{B}}$ acts freely and regularly on \mathcal{B} .*

Proof. Suppose that $G_{\mathcal{B}}$ acts freely and regularly on \mathcal{B} ; smoothness is inherited from G . A moving frame $\rho_{\mathcal{B}}$ on \mathcal{B} can be constructed by imposing normalization conditions. This determines the group parameters that occur in $G_{\mathcal{B}}$ and ensures that $\rho_{\mathcal{B}} \cdot \mathbf{x}$ depends on \mathbf{x} alone. Then $\rho_{\mathcal{B}}$ can be extended to a moving frame ρ on \mathcal{P} by choosing a normalization that determines the remaining parameters in G . Conversely, if there exists a projectable moving frame ρ on \mathcal{P} , its restriction to \mathcal{B} is a moving frame, whose existence requires $G_{\mathcal{B}}$ to act freely and regularly. ■

In particular, if x is the only continuous independent variable, a projectable moving frame exists only if $G_{\mathcal{B}}$ depends on at most one parameter, for otherwise the restricted action is not free. The moving frame is projectable if either x is invariant, or \tilde{x} depends on just one group parameter that is determined by a normalization equation of the form $\iota(x) = \text{const}$. Either of these conditions ensures that \mathcal{D} commutes with $\mathbf{S}_{\mathbf{K}}$. We restrict attention to projectable moving frames for the remainder of this paper.

7 Differential-difference variational calculus

The D Δ E variational calculus in terms of invariants is derived in much the same way as its counterpart for P Δ Es, so we present the basic method concisely to avoid too much repetition. However, there are some important differences that stem from the group action on the continuous independent variable x ; we describe these in detail. Here and henceforth, square brackets around an expression denotes the expression and finitely many of its derivatives and shifts.

Let G be an R -parameter Lie group of variational point symmetries for a given Lagrangian functional,

$$\mathcal{L} = \sum_{\mathbf{n}} \int \mathbf{L}(x, \mathbf{n}, [\mathbf{u}]) dx,$$

that leave the one-form $\mathbf{L}(x, \mathbf{n}, [\mathbf{u}]) dx$ invariant under the group action. Then

$$\mathbf{L} dx = \tilde{\mathbf{L}} d\tilde{x} = \tilde{\mathbf{L}} J_x dx. \tag{7.1}$$

In particular, invariance under the transformations generated by \mathbf{v}_r amounts to the condition

$$\mathbf{v}_r(\mathbf{L}) + \mathbf{L} D\xi_r = 0$$

(see [18] for details). A useful equivalent form of this condition is

$$X_r(\mathbf{L}) + D(\mathbf{L}\xi_r) = 0, \tag{7.2}$$

where X_r is the generator in characteristic form (whose action on x and dx is trivial).

⁷This definition includes jet spaces for differential equations, as projectability is also useful in this context.

The Euler–Lagrange equations are

$$E_{u^\alpha}(\mathbf{L}) := \mathbf{S}_{-\mathbf{K}}(-D)_{(j)} \left(\frac{\partial \mathbf{L}}{\partial u_{j;\mathbf{K}}^\alpha} \right) = 0, \quad \alpha = 1, \dots, m.$$

Just as for PΔEs, these equations can be obtained by allowing $[\mathbf{u}]$ (but *not* x) to depend smoothly on a continuous invariant parameter $t \in \mathbb{R}$. Then, using $'$ to denote the derivative with respect to t ,

$$\frac{d}{dt}(\mathbf{L}dx) = \frac{\partial \mathbf{L}}{\partial u_{j;k}^\alpha} (u_{j;k}^\alpha)' dx = E_{u^\alpha}(\mathbf{L})(u_{0;0}^\alpha)' dx + \text{Div}(\mathbf{A}_{\mathbf{u}})dx, \quad (7.3)$$

where $\text{Div}(\mathbf{A}_{\mathbf{u}})$ consists of the divergence terms arising from the summation and integration by parts. Note that $X_r(\mathbf{L})$ is obtained from $d\mathbf{L}/dt$ by setting $[(u_{0;0}^\alpha)' = Q_r^\alpha]$. So, from (7.2) and (7.3), the Noether conservation law corresponding to invariance under \mathbf{v}_r is

$$0 = -Q_r^\alpha E_{u^\alpha}(\mathbf{L}) = \text{Div}(\mathbf{A}_{\mathbf{u}})|_{[(u_{0;0}^\alpha)' = Q_r^\alpha]} - X_r(\mathbf{L}) = \text{Div}(\mathbf{A}_{\mathbf{u}})|_{[(u_{0;0}^\alpha)' = Q_r^\alpha]} + D(\mathbf{L}\xi_r). \quad (7.4)$$

Evaluating (7.1) on the frame gives

$$\mathbf{L}dx = \iota(\mathbf{L})\iota(dx) =: L^\kappa(\iota(x), \mathbf{n}, [\kappa])\iota(dx),$$

where κ denotes the generating invariants that depend on $[\mathbf{u}]$ (and possibly also on x). Explicitly, $\iota(dx) = \mathcal{J}dx$, so $L^\kappa = \mathbf{L}\mathcal{J}^{-1}$. As the frame is projectable, \mathcal{J} depends only on x ; in particular, if x is invariant, $\mathcal{J} = 1$.

To obtain the invariantized Euler–Lagrange equations, differentiate the one-form $L^\kappa\iota(dx)$ with respect to t . Each κ^β depends on t through its dependence on $[\mathbf{u}]$, giving rise to the syzygies $d\kappa^\beta/dt = \mathcal{H}_\alpha^\beta\sigma^\alpha$, where $\sigma^\alpha = \iota((u_{0;0}^\alpha)')$ and each \mathcal{H}_α^β is an invariant differential-difference operator. Specifically,

$$\mathcal{H}_\alpha^\beta = \iota \left(\frac{\partial \kappa^\beta}{\partial u_{j;\mathbf{K}}^\alpha} \right) \mathcal{D}_{(j)} \mathbf{S}_{\mathbf{K}},$$

where $\mathcal{D} = \mathcal{J}^{-1}D$ is the invariantized total derivative and $\mathcal{D}_{(j)} := \mathcal{D}^j$. Note that \mathcal{D} commutes with each $\mathbf{S}_{\mathbf{K}}$; thus the invariant derivatives and shifts of κ^β can be written as $\kappa_{j;\mathbf{K}}^\beta := \mathcal{D}_{(j)} \mathbf{S}_{\mathbf{K}} \kappa^\beta$. Integration by parts is straightforward: given two functions, f and g ,

$$f(\mathcal{D}g)\iota(dx) = f(Dg)dx = \{D(fg) - (Df)g\}dx = \{\mathcal{D}(fg) - (\mathcal{D}f)g\}\iota(dx).$$

As $\mathcal{D}(fg)\iota(dx)$ is a divergence, define the adjoint of \mathcal{D} relative to $\iota(dx)$ to be $\mathcal{D}^\dagger = -\mathcal{D}$. (The bold dagger \dagger distinguishes this adjoint from the standard adjoint, \dagger .) As the frame is projectable, $\mathbf{S}_{\mathbf{K}}^\dagger = \mathbf{S}_{\mathbf{K}}^\dagger = \mathbf{S}_{-\mathbf{K}}$. Therefore, using the notation $(-\mathcal{D})_{(j)} := (-1)^j \mathcal{D}_{(j)}$, we obtain

$$\begin{aligned} \frac{d}{dt}(L^\kappa\iota(dx)t) &= \frac{\partial L^\kappa}{\partial \kappa_{j;\mathbf{K}}^\beta} \frac{d\kappa_{j;\mathbf{K}}^\beta}{dt} \iota(dx) \\ &= \left((-\mathcal{D})_{(j)} \mathbf{S}_{-\mathbf{K}} \frac{\partial L^\kappa}{\partial \kappa_{j;\mathbf{K}}^\beta} \right) \frac{d\kappa^\beta}{dt} \iota(dx) + \text{Div}(\mathbf{A}_{\kappa})\iota(dx) \\ &= E_{\kappa^\beta}(L^\kappa)(\mathcal{H}_\alpha^\beta\sigma^\alpha)\iota(dx) + \text{Div}(\mathbf{A}_{\kappa})\iota(dx) \\ &= \{(\mathcal{H}_\alpha^\beta)^\dagger(E_{\kappa^\beta}(L^\kappa))\}\sigma^\alpha\iota(dx) + \text{Div}(\mathbf{A}_{\mathcal{H}} + \mathbf{A}_{\kappa})\iota(dx). \end{aligned} \quad (7.5)$$

The one-forms $\text{Div}(A_{\kappa})\iota(dx)$ and $\text{Div}(A_{\mathcal{H}})\iota(dx)$ are defined by the above; Div has the same form as Div , but with D replaced by \mathcal{D} . The following identity is useful:

$$\text{Div}(A^0; A^1, \dots, A^m)\iota(dx) = \text{Div}(A^0; \mathcal{J}A^1, \dots, \mathcal{J}A^m)dx. \quad (7.6)$$

Similarly to PΔEs, the i^{th} component of A_{κ} (resp. $A_{\mathcal{H}}$) is of the form $F_{\beta}^i(\kappa^{\beta})'$ (resp. $H_{\alpha}^i\sigma^{\alpha}$); here F_{β}^i and H_{α}^i are invariant differential-difference operators.

Proposition 7.1. *Suppose that the Lagrangian one-form $L(\mathbf{n}, x, [\mathbf{u}])dx$ is invariant under an R -parameter Lie group of point transformations. In the above notation,*

$$E_{u^{\alpha}}(L)(u_{0;0}^{\alpha})'dx = ((\mathcal{H}_{\alpha}^{\beta})^{\dagger}E_{\kappa^{\beta}}(L^{\kappa}))\sigma^{\alpha}\iota(dx), \quad (7.7)$$

so the invariantized Euler–Lagrange DΔEs are

$$\iota(E_{u^{\alpha}}(L)) = (\mathcal{H}_{\alpha}^{\beta})^{\dagger}E_{\kappa^{\beta}}(L^{\kappa}) = 0, \quad \alpha = 1, \dots, q. \quad (7.8)$$

Furthermore,

$$\text{Div}(A_{\mathbf{u}})dx = \text{Div}(A_{\mathcal{H}} + A_{\kappa})\iota(dx). \quad (7.9)$$

Proof. The proof is essentially the same as for PΔEs. In view of (7.6),

$$0 = \sum_{\mathbf{n}} \int \frac{d}{dt} (Ldx - L^{\kappa}\iota(dx)) = \sum_{\mathbf{n}} \int E_{u^{\alpha}}(L)(u_{0;0}^{\alpha})'dx - \{(\mathcal{H}_{\alpha}^{\beta})^{\dagger}(E_{\kappa^{\beta}}(L^{\kappa}))\}\sigma^{\alpha}\iota(dx).$$

This holds for arbitrary functions $(u_{0;0}^{\alpha})'$, which are independent at each base point \mathbf{n} . Therefore, (7.7) follows, from which (7.8) is obtained by invariantizing and using the independence of the functions σ^{α} . Equation (7.9) is derived by equating the right-hand sides of (7.3) and (7.5), taking (7.7) into account. \blacksquare

In the original variables, there are two types of contribution to each Noether conservation law (7.4) for which $\xi_r \neq 0$. The first type arises from integration and summation by parts, so can be treated in much the same way as for PΔEs. However, unlike the PΔE case, A_{κ} cannot be neglected. Each fundamental invariant κ^{β} satisfies $\mathbf{v}_r(\kappa^{\beta}) = 0$. However, only the vertical variables $[\mathbf{u}]$ depend on t , so the characteristic form of the generator is used. If $t = \varepsilon^r$, then $(\kappa^{\beta})'$, evaluated at $\varepsilon^r = 0$, reduces to

$$X_r(\kappa^{\beta}) = \mathbf{v}_r(\kappa^{\beta}) - \xi_r D(\kappa^{\beta}) = -\xi_r D(\kappa^{\beta}) = -\mathcal{D}(\kappa^{\beta})\iota(\xi_s)a_r^s(\rho). \quad (7.10)$$

The last equality arises from (6.3), evaluated on the moving frame ρ .

The counterpart of the remaining term in (7.4), namely $D(L\xi_r)$, is derived as follows. From (6.5) and (7.1),

$$L\xi_r = LJ_x^{-1}\tilde{\xi}_s a_r^s(g) = \tilde{L}\tilde{\xi}_s a_r^s(g),$$

for all $g \in G$; on the moving frame, this amounts to

$$L\xi_r = L^{\kappa}\iota(\xi_s)a_r^s(\rho).$$

Consequently,

$$D(L\xi_r)dx = \mathcal{D}(L^{\kappa}\iota(\xi_s)a_r^s(\rho))\iota(dx). \quad (7.11)$$

Combining all of the above, we obtain the following formulation of the Noether conservation laws.

Proposition 7.2. *Suppose that the conditions of Proposition 7.1 hold. Then Noether's theorem gives the R conservation laws*

$$\text{Div}(A_r)\iota(dx) = 0, \quad r = 1, \dots, R, \quad (7.12)$$

whose components are

$$A_r^0 = H_\alpha^0 \{ \iota(Q_s^\alpha) a_r^s(\rho) \} - F_\beta^0 \{ \mathcal{D}(\kappa^\beta) \iota(\xi_s) a_r^s(\rho) \} + L^\kappa \iota(\xi_s) a_r^s(\rho), \quad (7.13)$$

$$A_r^i = H_\alpha^i \{ \iota(Q_s^\alpha) a_r^s(\rho) \} - F_\beta^i \{ \mathcal{D}(\kappa^\beta) \iota(\xi_s) a_r^s(\rho) \}, \quad i = 1, \dots, m. \quad (7.14)$$

Proof. For each r in turn, replace t by ε^r and u^α by

$$\widehat{u}^\alpha = u^\alpha + \varepsilon^r Q_r^\alpha(\mathbf{n}, x, [\mathbf{u}]) + \mathcal{O}((\varepsilon^r)^2)$$

(prolonged as necessary), and evaluate the results at $\varepsilon^r = 0$. Using the same reasoning as for PDEs,

$$\sigma_r^\alpha \Big|_{\varepsilon^r=0} = \left\{ \frac{\partial(g \cdot u^\alpha)}{\partial u^\beta} Q_r^\beta \right\} \Big|_{g=\rho} = \{ \widetilde{Q}_s^\alpha a_r^s(g) \} \Big|_{g=\rho} = \iota(Q_s^\alpha) a_r^s(\rho),$$

which is substituted for σ^α in $A_{\mathcal{H}}$. The proof is completed by replacing $(\kappa^\beta)'$ in A_κ by the right-hand side of (7.10), and adding the remaining term (7.11). ■

Corollary 7.3. *Each component of the conservation laws (7.12) is equivariant with respect to the moving frame ρ , because there exist functions f_s^i of the invariants such that*

$$A_r^i = f_s^i(\mathbf{n}, \iota(x), [\kappa]) a_r^s(\rho), \quad i = 0, \dots, m, \quad r = 1, \dots, R.$$

Proof. The invariant differential-difference operators in (7.13) and (7.14) act linearly on products of invariants and adjoint components $a_r^s(\rho)$. Consequently, every term in A_r^i is of the form $\phi_l^i a_r^l(\rho_{j;\mathbf{K}})$, where $\rho_{j;\mathbf{K}} = \mathcal{D}_{(j)} \mathbf{S}_{\mathbf{K}} \rho$ and ϕ_l^i is invariant. The invariantization of $\rho_{1;0}$ (in a matrix representation) is the curvature matrix, $\rho_{1;0} \rho^{-1}$ (see [6, 12]). By applying powers of \mathcal{D} to the curvature matrix and eliminating derivatives of order $1, \dots, j-1$, one finds that the term in braces in the identity below is invariant:

$$\rho_{j;0} = \{ \rho_{j;0} \rho^{-1} \} \rho.$$

Shifting this, and using our earlier result that $\rho_{0;\mathbf{K}} \rho^{-1}$ is invariant for a difference moving frame, shows that

$$\rho_{j;\mathbf{K}} = \{ \rho_{j;\mathbf{K}} \rho_{0;\mathbf{K}}^{-1} \} \{ \rho_{0;\mathbf{K}} \rho^{-1} \} \rho = \{ \rho_{j;\mathbf{K}} \rho^{-1} \} \rho$$

is a product of invariants (in braces) and ρ . The adjoint matrices $(a_r^l(g))$ constitute a Lie group representation, so

$$\phi_l^i a_r^l(\rho_{j;\mathbf{K}}) = \{ \phi_l^i a_s^l(\rho_{j;\mathbf{K}} \rho^{-1}) \} a_r^s(\rho),$$

which is in the required form. ■

8 Examples

Example 8.1. To illustrate the calculations in a simple context (without any particular application), consider the Lagrangian one-form

$$Ldx = \frac{(u_{1;0})^2}{u_{0;1} - u_{0;0}} dx, \quad (8.1)$$

whose Euler–Lagrange equation is

$$E_u(L) = \left(\frac{-2u_{2;0}}{u_{0;1} - u_{0;0}} + \frac{2u_{1;0}u_{1;1} - (u_{1;0})^2}{(u_{0;1} - u_{0;0})^2} - \frac{(u_{1;-1})^2}{(u_{0;0} - u_{0;-1})^2} \right) = 0. \quad (8.2)$$

The one-form (8.1) is invariant under the two-parameter Lie group of point transformations

$$g: (x, n, u) \mapsto (\tilde{x}, n, \tilde{u}) = (bx, n, bu + a).$$

The infinitesimal generators are linear combinations of $\mathbf{v}_1 = \partial_u$ and $\mathbf{v}_2 = x\partial_x + u\partial_u$, which yield $(\xi_1, Q_1) = (0, 1)$ and $(\xi_2, Q_2) = (x, u_{0;0} - xu_{1;0})$. Then (7.4) gives the following conservation laws (expressed as one-forms for comparison):

$$\begin{aligned} -Q_1 E_u(L) dx &= D \left(\frac{2u_{1;0}}{u_{0;1} - u_{0;0}} \right) dx + D_n \left(\frac{-(u_{1;-1})^2}{(u_{0;0} - u_{0;-1})^2} \right) dx = 0, \\ -Q_2 E_u(L) dx &= D \left(\frac{u_{1;0}(2u_{0;0} - xu_{1;0})}{u_{0;1} - u_{0;0}} \right) dx + D_n \left(\frac{(u_{1;-1})^2(xu_{1;0} - u_{0;0})}{(u_{0;0} - u_{0;-1})^2} \right) dx = 0. \end{aligned} \quad (8.3)$$

Note that (8.3) includes the term $D(L\xi_2)dx$.

Reflecting the identities $\mathbf{v}_1 = b\tilde{\mathbf{v}}$ and $\mathbf{v}_2 = -a\tilde{\mathbf{v}}_1 + \tilde{\mathbf{v}}_2$, the adjoint representation is given by

$$a_1^1(g) = b, \quad a_1^2(g) = 0, \quad a_2^1(g) = -a, \quad a_2^2(g) = 1.$$

The normalization $\iota(x) = 1$, $\iota(u_{0;0}) = 0$ gives a projectable moving frame ρ , on which

$$a = \frac{-u_{0;0}}{x}, \quad b = \frac{1}{x}.$$

Therefore, $\iota(dx) = x^{-1}dx$ and the invariantized total derivative operator is $\mathcal{D} = xD$. On the moving frame, the adjoint representation has components

$$a_1^1(\rho) = \frac{1}{x}, \quad a_1^2(\rho) = 0, \quad a_2^1(\rho) = \frac{u_{0;0}}{x}, \quad a_2^2(\rho) = 1.$$

A generating set of invariants is

$$\kappa^1 = \iota(u_{1;0}) = u_{1;0}, \quad \kappa^2 = \iota(u_{0;1}) = \frac{u_{0;1} - u_{0;0}}{x},$$

which satisfy the syzygy

$$\kappa_{0;1}^1 = \iota(u_{1;1}) = \kappa^1 + \kappa_{1;0}^2 + \kappa^2. \quad (8.4)$$

The Lagrangian one-form amounts to

$$L^\kappa \iota(dx) = \frac{(\kappa^1)^2}{\kappa^2} \iota(dx),$$

and so

$$E_{\kappa^1}(L^\kappa) = \frac{2\kappa^1}{\kappa^2}, \quad E_{\kappa^2}(L^\kappa) = -\left(\frac{\kappa^1}{\kappa^2}\right)^2.$$

In terms of $\sigma = \iota(u'_{0;0}) = x^{-1}u'_{0;0}$, the t -derivatives of the generating invariants are $(\kappa^\beta)' = \mathcal{H}^\beta \sigma$, where

$$\mathcal{H}^1 = \mathcal{D} + \text{id}, \quad \mathcal{H}^2 = \mathcal{S} - \text{id}.$$

Consequently,

$$\begin{aligned} \frac{dL^\kappa}{dt} \iota(dx) &= \left(\frac{2\kappa^1}{\kappa^2} (\kappa^1)' - \left(\frac{\kappa^1}{\kappa^2} \right)^2 (\kappa^2)' \right) \iota(dx) = E_{\kappa^\beta}(L^\kappa) (\kappa^\beta)' \iota(dx) \\ &= E_{\kappa^\beta}(L^\kappa) (\mathcal{H}^\beta \sigma) \iota(dx) = ((\mathcal{H}^\beta)^\dagger E_{\kappa^\beta}(L^\kappa)) \sigma \iota(dx) + \text{Div}(A_{\mathcal{H}}) \iota(dx), \end{aligned}$$

where

$$\text{Div}(A_{\mathcal{H}}) = \mathcal{D}(E_{\kappa^1}(L^\kappa)\sigma) + D_n(\{\mathcal{S}^{-1}E_{\kappa^2}(L^\kappa)\}\sigma).$$

In this example, L^κ does not involve derivatives or shifts of the generating invariants, and hence $\text{Div}(A_{\mathcal{H}}) = 0$. The invariantized Euler–Lagrange equation is

$$\begin{aligned} 0 &= (\mathcal{H}^\beta)^\dagger E_{\kappa^\beta}(L^\kappa) = (-\mathcal{D} + \text{id}) \left(\frac{2\kappa^1}{\kappa^2} \right) + (\mathcal{S}^{-1} - \text{id}) \left(-\left(\frac{\kappa^1}{\kappa^2} \right)^2 \right) \\ &= \frac{2(\kappa^1 - \kappa_{1;0}^1)}{\kappa^2} + \frac{\kappa^1(\kappa^1 + 2\kappa_{1;0}^2)}{(\kappa^2)^2} - \left(\frac{\kappa_{0;-1}^1}{\kappa_{0;-1}^2} \right)^2. \end{aligned}$$

For comparison, one can invariantize (8.2) directly, using (8.4) and

$$\iota(u_{2;0}) = \kappa_{1;0}^1, \quad \iota(u_{1;-1}) = \kappa_{0;-1}^1, \quad \iota(u_{0;-1}) = -\kappa_{0;-1}^2.$$

From Proposition 7.2, the conservation laws given by Noether's theorem amount to

$$\begin{aligned} \mathcal{D}\{(E_{\kappa^1}(L^\kappa)\iota(Q_s) + L^\kappa\iota(\xi_s))a_r^s(\rho)\} \iota(dx) \\ + D_n\{\{\mathcal{S}^{-1}E_{\kappa^2}(L^\kappa)\}\iota(Q_s)a_r^s(\rho)\} \iota(dx) = 0. \end{aligned} \tag{8.5}$$

Substituting

$$\iota(Q_s)a_1^s(\rho) = \frac{1}{x}, \quad \iota(Q_s)a_2^s(\rho) = \frac{u_{0;0}}{x} - \kappa^1, \quad \iota(\xi_s)a_1^s(\rho) = 0, \quad \iota(\xi_s)a_2^s(\rho) = 1,$$

into (8.5) gives the conservation laws

$$\begin{aligned} 0 &= \mathcal{D} \left\{ \frac{2\kappa^1}{x\kappa^2} \right\} \iota(dx) + D_n \left\{ -\frac{1}{x} \left(\frac{\kappa_{0;-1}^1}{\kappa_{0;-1}^2} \right)^2 \right\} \iota(dx), \\ 0 &= \mathcal{D} \left\{ \frac{2\kappa^1}{\kappa^2} \left(\frac{u_{0;0}}{x} - \kappa^1 \right) + \frac{(\kappa^1)^2}{\kappa^2} \right\} \iota(dx) + D_n \left\{ \left(\frac{\kappa_{0;-1}^1}{\kappa_{0;-1}^2} \right)^2 \left(\kappa^1 - \frac{u_{0;0}}{x} \right) \right\} \iota(dx) \\ &= \mathcal{D} \left\{ \frac{\kappa^1}{\kappa^2} \left(\frac{2u_{0;0}}{x} - \kappa^1 \right) \right\} \iota(dx) + D_n \left\{ \left(\frac{\kappa_{0;-1}^1}{\kappa_{0;-1}^2} \right)^2 \left(\kappa^1 - \frac{u_{0;0}}{x} \right) \right\} \iota(dx). \end{aligned}$$

Example 8.2. Method of lines semi-discretizations are a common source of DΔEs with just one continuous independent variable. The nonlinear Schrödinger (NLS) equation for a field with real and imaginary parts u and v respectively has the following (non-integrable) semi-discretization, with uniform step length h :

$$\begin{aligned} -v_{1;0} + u_{0;0}(u_{0;0}^2 + v_{0;0}^2) + h^{-2}(u_{0;-1} - 2u_{0;0} + u_{0;1}) &= 0, \\ u_{1;0} + v_{0;0}(u_{0;0}^2 + v_{0;0}^2) + h^{-2}(v_{0;-1} - 2v_{0;0} + v_{0;1}) &= 0. \end{aligned}$$

These are the Euler–Lagrange equations corresponding to the Lagrangian one-form

$$\begin{aligned} \text{L}dx = & \left\{ \frac{1}{2}(v_{0;0}u_{1;0} - u_{0;0}v_{1;0}) + \frac{1}{4}(u_{0;0}^2 + v_{0;0}^2)^2 \right. \\ & \left. - \frac{1}{2}h^{-2}((u_{0;1} - u_{0;0})^2 + (v_{0;1} - v_{0;0})^2) \right\} dx, \end{aligned}$$

which is invariant under the two-parameter abelian Lie group of point transformations

$$g: (x, n, u, v) \mapsto (\tilde{x}, \tilde{n}, \tilde{u}, \tilde{v}) = (x + a, n, u \cos b + v \sin b, -u \sin b + v \cos b).$$

The infinitesimal generators are $\mathbf{v}_1 = \partial_x$ and $\mathbf{v}_2 = v\partial_u - u\partial_v$, so (using variable names rather than indices, for clarity)

$$(\xi_1, Q_1^u, Q_1^v) = (1, -u_{1;0}, -v_{1;0}), \quad (\xi_2, Q_2^u, Q_2^v) = (0, v_{0;0}, -u_{0;0}).$$

As the Lie group is abelian, the adjoint representation is the identity, so $a_r^s(g) = \delta_r^s$ for all g .

We now choose the normalization $\iota(x) = 0$, $\iota(v_{0;0}) = 0$, temporarily restricting attention to $u_{0;0} > 0$. (Other normalizations can be used for the remaining coordinate patches.) This gives the frame ρ defined by

$$a|_\rho = -x, \quad b|_\rho = \tan^{-1} \left(\frac{v_{0;0}}{u_{0;0}} \right).$$

In the calculations that follow, $\cos b$ and $\sin b$ (but not b) are evaluated on the frame, so we use

$$a|_\rho = -x, \quad \cos b|_\rho = \frac{u_{0;0}}{\sqrt{u_{0;0}^2 + v_{0;0}^2}}, \quad \sin b|_\rho = \frac{v_{0;0}}{\sqrt{u_{0;0}^2 + v_{0;0}^2}},$$

which extends to other coordinate patches. Note that $\iota(dx) = dx$, and so the invariantized total derivative is $\mathcal{D} = D$. The invariants are generated by

$$\kappa^1 = \iota(u_{0;0}) = \sqrt{u_{0;0}^2 + v_{0;0}^2}, \quad \kappa^2 = u_{0;0}v_{1;0} - v_{0;0}u_{1;0}, \quad \kappa^3 = u_{0;0}u_{0;1} + v_{0;0}v_{0;1}.$$

To see this, note that all derivatives can be obtained from

$$u_{1;0} = \frac{u_{0;0}\kappa^1\kappa_{1;0}^1 - v_{0;0}\kappa^2}{(\kappa^1)^2}, \quad v_{1;0} = \frac{u_{0;0}\kappa^2 + v_{0;0}\kappa^1\kappa_{1;0}^1}{(\kappa^1)^2},$$

and that under the constraint $\iota(v_{0;1}) \geq 0$, all shifts can be obtained from

$$u_{0;1} = \frac{u_{0;0}\kappa^3 - v_{0;0}\phi}{(\kappa^1)^2}, \quad v_{0;1} = \frac{u_{0;0}\phi + v_{0;0}\kappa^3}{(\kappa^1)^2},$$

where

$$\phi = \{(\kappa^1\kappa_{0;1}^1)^2 - (\kappa^3)^2\}^{1/2}.$$

We adopt this constraint for definiteness; if it is not satisfied, replace ϕ by $-\phi$ throughout. By calculating $u_{1;1}$ (or $v_{1;1}$), one obtains the syzygy

$$\frac{\kappa^3 \kappa_{1;1}^1}{\kappa_{0;1}^1} - \kappa_{1;0}^3 + \frac{\kappa_{1;0}^1}{\kappa^1} + \frac{\phi \kappa^2}{(\kappa^1)^2} - \frac{\phi \kappa_{0;1}^2}{(\kappa_{0;1}^1)^2} = 0.$$

In terms of the generating parametric derivatives,

$$\sigma^u = \iota(u'_{0;0}) = \frac{u_{0;0}u'_{0;0} + v_{0;0}v'_{0;0}}{\kappa^1}, \quad \sigma^v = \iota(v'_{0;0}) = \frac{u_{0;0}v'_{0;0} - v_{0;0}u'_{0;0}}{\kappa^1},$$

the derivatives of the generating invariants are

$$\begin{aligned} (\kappa^1)' &= \sigma^u, & (\kappa^2)' &= \frac{2\kappa^2}{\kappa^1} \sigma^u + (\kappa^1 D - \kappa_{1;0}^1) \sigma^v, \\ (\kappa^3)' &= \left(\frac{\kappa^3}{\kappa^1} + \frac{\kappa^3}{\kappa_{0;1}^1} S \right) \sigma^u + \left(\frac{\phi}{\kappa^1} - \frac{\phi}{\kappa_{0;1}^1} S \right) \sigma^v. \end{aligned}$$

As $\iota(dx) = dx$ and $\mathcal{D} = D$, the invariant Euler–Lagrange equations and conservation laws can be calculated directly from

$$L^\kappa = -\frac{1}{2}\kappa^2 + \frac{1}{4}(\kappa^1)^4 - \frac{1}{2}h^{-2}((\kappa_{0;1}^1)^2 - 2\kappa^3 + (\kappa^1)^2).$$

Differentiating this, we obtain

$$\begin{aligned} \frac{d}{dt} L^\kappa &= -\frac{1}{2}(\kappa^2)' + (\kappa^1)^3 (\kappa^1)' - h^{-2}(\kappa_{0;1}^1 (\kappa_{0;1}^1)' - (\kappa^3)' + \kappa^1 (\kappa^1)') \\ &= \{(\kappa^1)^3 - 2h^{-2}\kappa^1\} (\kappa^1)' - \frac{1}{2}(\kappa^2)' + h^{-2}(\kappa^3)' + \underbrace{D_n(-h^{-2}\kappa^1 (\kappa^1)')}_{\text{Div} A_\kappa} \\ &= \left\{ (\kappa^1)^3 - \frac{2\kappa^1}{h^2} - \frac{\kappa^2}{\kappa^1} + \frac{\kappa^3}{h^2\kappa^1} + \frac{\kappa^3}{h^2\kappa_{0;1}^1} S \right\} \sigma^u \\ &\quad + \left\{ -\frac{1}{2}\kappa^1 D + \frac{1}{2}\kappa_{1;0}^1 + \frac{\phi}{h^2\kappa^1} - \frac{\phi}{h^2\kappa_{0;1}^1} S \right\} \sigma^v + D_n(-h^{-2}\kappa^1 (\kappa^1)') \\ &= \left\{ (\kappa^1)^3 - \frac{2\kappa^1}{h^2} - \frac{\kappa^2}{\kappa^1} + \frac{\kappa^3}{h^2\kappa^1} + \frac{\kappa_{0,-1}^3}{h^2\kappa^1} \right\} \sigma^u + \left\{ \kappa_{1;0}^1 + \frac{\phi}{h^2\kappa^1} - \frac{S^{-1}\phi}{h^2\kappa^1} \right\} \sigma^v \\ &\quad + D \left(-\frac{1}{2}\kappa^1 \sigma^v \right) + D_n \left(\frac{\kappa_{0,-1}^3}{h^2\kappa^1} \sigma^u - \frac{S^{-1}\phi}{h^2\kappa^1} \sigma^v \right) + D_n(-h^{-2}\kappa^1 (\kappa^1)'). \end{aligned} \tag{8.6}$$

Consequently, the invariantized Euler–Lagrange equations are

$$\begin{aligned} \iota(E_u(L)) &= (\kappa^1)^3 - \frac{2\kappa^1}{h^2} - \frac{\kappa^2}{\kappa^1} + \frac{\kappa^3}{h^2\kappa^1} + \frac{\kappa_{0,-1}^3}{h^2\kappa^1} = 0, \\ \iota(E_v(L)) &= \kappa_{1;0}^1 + \frac{\phi}{h^2\kappa^1} - \frac{S^{-1}\phi}{h^2\kappa^1} = 0. \end{aligned}$$

In this example, the conservation laws have contributions from both A_κ (see (8.6)) and $A_{\mathcal{H}}$. Using $a_r^s(\rho) = \delta_r^s$ and

$$\begin{aligned} \iota(Q_1^u) &= -\kappa_{1;0}^1, & \iota(Q_1^v) &= -\kappa^2/\kappa^1, & \iota(Q_2^u) &= 0, \\ \iota(Q_2^v) &= -\kappa^1, & \iota(\xi_1) &= 1, & \iota(\xi_2) &= 0, \end{aligned}$$

we obtain the following conservation laws from Proposition 7.2:

$$\begin{aligned}
0 &= D \left\{ -\frac{1}{2} \kappa^1 \iota(Q_1^v) + \iota(\xi_1) L^\kappa \right\} + D_n \left\{ \frac{\kappa_{0,-1}^3}{h^2 \kappa^1} \iota(Q_1^u) - \frac{S^{-1} \phi}{h^2 \kappa^1} \iota(Q_1^v) - \frac{\kappa^1}{h^2} (-\iota(\xi_1) \kappa_{1;0}^1) \right\} \\
&= D \left\{ \frac{1}{4} (\kappa^1)^4 - \frac{(\kappa_{0;1}^1)^2}{2h^2} + \frac{\kappa^3}{h^2} - \frac{(\kappa^1)^2}{2h^2} \right\} + D_n \left\{ -\frac{\kappa_{1;0}^1 \kappa_{0,-1}^3}{h^2 \kappa^1} + \frac{\kappa^2 S^{-1} \phi}{h^2 (\kappa^1)^2} + \frac{\kappa^1 \kappa_{1;0}^1}{h^2} \right\}, \\
0 &= D \left\{ -\frac{1}{2} \kappa^1 \iota(Q_2^v) \right\} + D_n \left\{ \frac{\kappa_{0,-1}^3}{h^2 \kappa^1} \iota(Q_2^u) - \frac{S^{-1} \phi}{h^2 \kappa^1} \iota(Q_2^v) \right\} \\
&= D \left\{ \frac{1}{2} (\kappa^1)^2 \right\} + D_n \left\{ \frac{S^{-1} \phi}{h^2} \right\}.
\end{aligned}$$

9 Concluding remarks

For PΔEs and DΔEs, the prolongation space over a fixed base point \mathbf{n} provides a continuous setting in which moving frames can be used. Difference moving frames respect the ordering of each discrete independent variable and the arbitrariness of the base point. For variational problems, we have shown how to calculate the invariantized Euler–Lagrange equations and equivariant Noether conservation laws directly from an invariant Lagrangian, L^κ .

We have treated the coordinates n^i on the lattice of independent variables, \mathbb{Z}^m , as being fixed by the Lie group of transformations. For a formulation that allows discrete symmetries of the lattice, it would be necessary to replace the Lagrangian L^κ by the Lagrangian m -form, $L^\kappa \text{vol}$, where vol denotes the difference volume form (see [8, 13]). This adds complexity, but little extra insight. However, when each n^i is fixed, only the coefficients of difference forms are transformed by the group action, so one can use moving frames as we have done, without the additional machinery of difference forms.

Our treatment of DΔEs has been restricted to a single continuous independent variable and a projectable moving frame, enabling the shift and invariant differential operators to commute. More generally, let p be the number of continuous independent variables. If $p > 1$, the complexity increases, because the invariant differential operators do not necessarily commute with one another. However, if the moving frame is projectable, all shift operators commute with the invariant derivatives, enabling existing results from PDE theory to be used for the differential part of the calculations. The requirement for the group action to be projectable is not sufficient to guarantee the existence of a projectable moving frame. Nevertheless, projectable moving frames are relevant to many DΔEs of interest, including some well-known integrable systems and method of lines semi-discretizations of PDEs.

Acknowledgments

We thank Professor Elizabeth Mansfield, whose strong advocacy of moving frames, insight and encouragement have led to this project. The work was partially supported by EPSRC grant number EP/R513246/1.

References

- [1] Boutin M., On orbit dimensions under a simultaneous Lie group action on n copies of a manifold, *J. Lie Theory* **12** (2002), 191–203, [arXiv:math-ph/0009021](#).
- [2] Fels M., Olver P.J., Moving coframes. I. A practical algorithm, *Acta Appl. Math.* **51** (1998), 161–213.
- [3] Fels M., Olver P.J., Moving coframes. II. Regularization and theoretical foundations, *Acta Appl. Math.* **55** (1999), 127–208.

-
- [4] Gonçalves T.M.N., Mansfield E.L., On moving frames and Noether's conservation laws, *Stud. Appl. Math.* **128** (2012), 1–29, [arXiv:1006.4660](#).
 - [5] Gonçalves T.M.N., Mansfield E.L., Moving frames and conservation laws for Euclidean invariant Lagrangians, *Stud. Appl. Math.* **130** (2013), 134–166, [arXiv:1106.3964](#).
 - [6] Gonçalves T.M.N., Mansfield E.L., Moving frames and Noether's conservation laws – the general case, *Forum Math. Sigma* **4** (2016), e29, 55 pages, [arXiv:1306.0847](#).
 - [7] Hydon P.E., Difference equations by differential equation methods, *Cambridge Monogr. Appl. Comput. Math.*, Vol. 27, [Cambridge University Press](#), Cambridge, 2014.
 - [8] Hydon P.E., Mansfield E.L., A variational complex for difference equations, *Found. Comput. Math.* **4** (2004), 187–217.
 - [9] Kim P., Olver P.J., Geometric integration via multi-space, *Regul. Chaotic Dyn.* **9** (2004), 213–226.
 - [10] Kogan I.A., Olver P.J., Invariant Euler–Lagrange equations and the invariant variational bicomplex, *Acta Appl. Math.* **76** (2003), 137–193.
 - [11] Kuperschmidt B.A., Discrete Lax equations and differential-difference calculus, *Astérisque* **123** (1985), 212 pages.
 - [12] Mansfield E.L., A practical guide to the invariant calculus, *Cambridge Monogr. Appl. Comput. Math.*, Vol. 26, [Cambridge University Press](#), Cambridge, 2010.
 - [13] Mansfield E.L., Hydon P.E., Difference forms, *Found. Comput. Math.* **8** (2008), 427–467.
 - [14] Mansfield E.L., Marí Beffa G., Wang J.P., Discrete moving frames and discrete integrable systems, *Found. Comput. Math.* **13** (2013), 545–582, [arXiv:1212.5299](#).
 - [15] Mansfield E.L., Rojo-Echeburúa A., Hydon P.E., Peng L., Moving frames and Noether's finite difference conservation laws I, *Trans. Math. Appl.* **3** (2019), tnz004, 47 pages, [arXiv:1804.00317](#).
 - [16] Mansfield E.L., Rojo-Echeburúa A., Moving frames and Noether's finite difference conservation laws II, *Trans. Math. Appl.* **3** (2019), tnz005, 26 pages, [arXiv:1808.03606](#).
 - [17] Marí Beffa G., Mansfield E.L., Discrete moving frames on lattice varieties and lattice-based multispaces, *Found. Comput. Math.* **18** (2018), 181–247.
 - [18] Olver P.J., Applications of Lie groups to differential equations, 2nd ed., *Grad. Texts in Math.*, Vol. 107, [Springer](#), New York, 1993.
 - [19] Olver P.J., Geometric foundations of numerical algorithms and symmetry, *Appl. Algebra Engrg. Comm. Comput.*, Vol. 11, 2001, 417–436.
 - [20] Olver P.J., Sapiro G., Tannenbaum A., Differential invariant signatures and flows in computer vision: A symmetry group approach, in *Geometry-Driven Diffusion in Computer Vision*, *Comput. Imaging Vis.*, Vol. 1, Editor B.M. ter Haar Romeny, [Kluwer Academic Publishers](#), Dordrecht, 1994, 255–306.
 - [21] Peng L., Hydon P.E., Transformations, symmetries and Noether theorems for differential-difference equations, *Proc. A.* **478** (2022), 20210944, 17 pages, [arXiv:2112.06030](#).