



Kent Academic Repository

Kim, Yesol, Koo, Bon-Chul, Pyo, Tae-Soo, Froebrich, Dirk, Jeong, Woong-Seob, Lee, Jae-Joon, Lee, Yong-Hyun, Lee, Ho-Gyu, Kim, Hyun-Jeong and Varricatt, Watson P. (2024) *Extended ionized fe objects in the UWIFE survey*. *Monthly Notices of the Royal Astronomical Society*, 528 (3). pp. 4657-4700. ISSN 1365-2966.

Downloaded from

<https://kar.kent.ac.uk/104822/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1093/mnras/stae295>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.











Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Extended ionized Fe objects in the UWIFE survey

Yesol Kim ^{1,2*}, Bon-Chul Koo ¹, Tae-Soo Pyo ^{3,4}, Dirk Froebrich ⁵, Woong-Seob Jeong ^{2,6},
Jae-Joon Lee ², Yong-Hyun Lee ^{1,7}, Ho-Gyu Lee ², Hyun-Jeong Kim ^{2,8} and Watson P. Varricatt ⁹

¹Department of Physics and Astronomy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

²Korea Astronomy and Space Science Institute, 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Republic of Korea

³Subaru Telescope, National Astronomical Observatory of Japan (NAOJ), National Institutes of Natural Sciences (NINS), 650 North A'ohōku Place, Hilo, HI 96720, USA

⁴School of Mathematical and Physical Science, SOKENDAI (The Graduate University for Advanced Studies), Hayama, Kanagawa 240-0193, Japan

⁵Centre for Astrophysics and Planetary Science, University of Kent, Canterbury CT2 7NH, UK

⁶Department of Astronomy and Space Science, University of Science and Technology, 217 Gajeong-ro Yuseong-gu, Daejeon 34113, Republic of Korea

⁷Samsung SDS, Olympic-ro 35-gil 125, Seoul, Republic of Korea

⁸Department of Earth Science Education, Kongju National University, 56 Gongjudaehak-ro, Gongju-si, Chungcheongnam-do 32588, Republic of Korea

⁹UKIRT Observatory, University of Hawaii, Institute for Astronomy, 640 North A'ohōku Place, Hilo, HI 96720, USA

Accepted 2024 January 22. Received 2024 January 20; in original form 2023 July 6

ABSTRACT

We explore systematically the shocked gas in the first Galactic quadrant of the Milky Way using the United Kingdom Infrared Telescope (UKIRT) Wide-field Infrared Survey for Fe⁺ (UWIFE). The UWIFE survey is the first imaging survey of the Milky Way in the [Fe II] 1.644 μm emission line and covers the Galactic plane in the first Galactic quadrant (7° < l < 62°; |b| ≲ 1°5). We identify 204 extended ionized Fe objects (IFOs) using a combination of a manual and automatic search. Most of the IFOs are detected for the first time in the [Fe II] 1.644 μm line. We present a catalogue of the measured sizes and fluxes of the IFOs and searched for their counterparts by performing positional cross-matching with known sources. We found that IFOs are associated with supernova remnants (25), young stellar objects (100), H II regions (33), planetary nebulae (17), and luminous blue variables (4). The statistical and morphological properties are discussed for each of these.

Key words: catalogues – surveys – circumstellar matter – ISM: kinematics and dynamics – infrared: ISM.

1 INTRODUCTION

Understanding the role of shocks is essential for comprehensively studying the ecology of the Milky Way, since they transfer mass and kinetic energy into the interstellar medium (ISM), provide heavy elements for future star formation by destroying dust grains, and regulate star formation. Shock waves are one of the principal mechanisms of the interaction between stars and the ISM in galaxies, thereby playing an important role in the evolution of the Galaxy. Among the most powerful shock-driving sources are outflows and jets from young stellar objects (YSOs), stellar winds from massive OB stars, and supernova (SN) explosions. To understand the physics of the interactions as well as the nature of the shock-driving sources, observations of emission lines from the shocks are essential.

The [Fe II] a⁴D_{7/2} → a⁴F_{9/2} 1.644 μm transition results in one of the brightest emission lines in near-infrared (NIR). It originates from one of the 16 levels of Fe⁺ that have a low excitation energy. Therefore they are easily excited in shocked gas, resulting in many lines, particularly in NIR. This emission line is thought to be bright in shock-excited gas; one suggested reason is that due to far-ultraviolet (FUV) radiation from the shock front, the Fe atom

is in the form of Fe⁺ over extended regions (McKee, Chernoff & Hollenbach 1984; Hollenbach, Chernoff & McKee 1989; Oliva, Moorwood & Danziger 1989; Koo, Raymond & Kim 2016). In contrast, in photoionized regions, Fe atoms are predominantly at higher ionization states, except when the ionizing radiation is hard enough that it can penetrate further into the interstellar cloud (Koo et al. 2016). Therefore, [Fe II] emission lines from shocked gas are stronger than those from photoionized regions; for example, [Fe II] 1.257 μm/Pa β is over 0.1 in supernova remnants (SNRs) compared to 0.01–0.03 in Orion (Koo & Lee 2015; Mouri, Kawara & Taniguchi 2000). Furthermore, the Fe abundance can be enhanced by shocks owing to grain destruction, making the [Fe II] lines stronger (Koo 2014; Greenhouse et al. 1991; Mouri et al. 2000, and references therein). These characteristics of [Fe II] make its lines extremely useful for studying interstellar shocks (e.g. Dinerstein 1995; Nisini 2008).

For example, the 1.644 μm emission line as a tracer of shocked atomic gas enables us to study shocked regions in jets/outflows of YSOs (Nisini et al. 2002; Caratti o Garatti et al. 2006; Takami et al. 2006; Pyo et al. 2006, 2009; Oh et al. 2016), planetary nebulae (PNe, Welch et al. 1999; Smith, Balick & Gehrz 2005), SNRs (Koo et al. 2007; Lee et al. 2009, 2013), and nebulae of luminous blue variables (LBVs, Smith 2002). Since NIR [Fe II] lines suffer less extinction than widely used optical emission lines such as Hα, [S II], and [O III],

* E-mail: [yskim916@gmail.com](mailto:ykim916@gmail.com)

the NIR lines can give us information on deeply embedded regions inaccessible by optical lines.

Lee et al. (2014) conducted an unbiased [Fe II] 1.644 μm narrow-band imaging survey, which is called the United Kingdom Infrared Telescope (UKIRT) Wide-field Infrared Survey for Fe⁺ (UWIFE). The survey area ($7^\circ < l < 62^\circ$; $|b| \lesssim 1.5^\circ$) is located in the first Galactic quadrant. This survey is the first unbiased, high-resolution [Fe II] survey of the Milky Way. It therefore enables us to discover more [Fe II]-emitting sources and conduct a statistically meaningful investigation of Galactic [Fe II] line sources. Alongside [Fe II]-emitting Galactic SNR to study similar to [Fe II] line objects in nearby galaxies, the survey is expected to systematically detect low-brightness [Fe II] line sources from other kinds of [Fe II] emitters. Therefore, it enables us to assess the level of contribution of each [Fe II]-emitting population. Further spectroscopic studies of new [Fe II] sources found in UWIFE can be used to derive critical densities in the range of $\sim 10^4 - 10^5 \text{ cm}^{-3}$ and temperatures up to 10^4 K (Pesenti et al. 2003), filling the gap in density between [S II] $\lambda 6731 \sim 10^4 \text{ cm}^{-3}$ and [O I] $\lambda 6300 \sim 10^6 \text{ cm}^{-3}$ (Osterbrock 1989). With other [Fe II] lines and emission lines such as [Fe II] 1.533 μm , density diagnostics of $\sim 10^2 - 10^5 \text{ cm}^{-3}$ can be measured and line ratio diagrams with other [Fe II] lines (Pesenti et al. 2003) can help us understand the new parameter range.

Shinn et al. (2014) systematically searched for outflows from ultracompact H II regions (UCHIIs), inferred [Fe II] outflow mass-loss rates, and discussed the traveltime of the [Fe II] outflows using the UWIFE data. The statistical [Fe II] line study of Galactic SNRs in UWIFE and the UKIRT Wide-field Infrared Survey for H₂ (UWISH2, Froebrich et al. 2011) survey revealed a detection rate of 24 per cent for both surveys and suggested a relatively higher coincidence with mixed-morphology and/or radio-bright SNRs (Lee et al. 2019).

A comprehensive catalogue of UWIFE sources will give an opportunity to compare shocked [Fe II] line objects with other tracers in previous large-scale Galactic plane surveys (GPSs). Particularly, the UWIFE survey area is fully covered with the complementary survey, UWISH2 (Froebrich et al. 2011), which was carried out using UKIRT and the Wide-Field Camera (WFCAM, Casali et al. 2007). The catalogue of extended H₂-emitting sources identified in UWISH2 (Froebrich et al. 2015) will be useful for the comparison of shocked molecular gas with higher excitation atomic gas. Also, the Isaac Newton Telescope Photometric H α Survey of the Northern Galactic Plane (IPHAS, Drew et al. 2005) and the UWISH2 survey can provide a chance to compare different outflow/shock tracers. Surveys tracing continuum sources in embedded regions such as the UKIRT Infrared Deep Sky Survey (UKIDSS) GPS (Lucas et al. 2008) in the NIR, the Galactic Legacy Infrared Midplane Survey Extraordinaire (GLIMPSE; Churchwell et al. 2009), the Multiband Imaging Photometer for *Spitzer* GPS (MIPSGAL; Carey et al. 2009) in the mid-infrared (MIR), and the *Herschel* infrared GPS (Molinari et al. 2010) in the far-infrared (FIR) were published.

Furthermore, the source catalogue of various kinds of objects, namely, the catalogue of UCHIIs from the Co-Ordinated Radio ‘N’ Infrared Survey for High-mass star formation (CORNISH, Hoare et al. 2012) and the catalogue of Extended Green Objects (EGO, Cyganowski et al. 2008) can be good candidates to compare with [Fe II] sources, as well as emission-line source catalogues, viz., catalogues of H α emission-line sources from IPHAS (Witham et al. 2008), and Molecular Hydrogen emission-line Object (MHO, Davis et al. 2010). In accordance with these catalogues and aims, we designate [Fe II] 1.644 μm emission-line sources as ionized Fe objects (IFOs) and compile the first comprehensive catalogue of Galactic extended IFOs. The catalogue includes basic physical

properties of IFOs, such as coordinates (l, b), size, position angle (PA), and flux. Information about possible counterparts and their distance is also included.

The organization of the paper is as follows. In Section 2, we describe the data reduction, the source identification, the photometry of the detected sources, and the procedure for searching for counterparts or exciting sources of IFOs. In Section 3, we first present the catalogue of IFOs. The catalogue contains the sizes and fluxes of IFOs as well as their counterparts. The IFOs are classified by their counterpart types. We then explore the statistics of the physical properties and the distribution of IFOs. In Section 4, we discuss the results of the individual types of IFOs. In Section 5, we summarize our paper.

2 DATA AND SOURCE CATALOGUE

2.1 UWIFE survey data

We have used the UWIFE survey data to search for extended IFOs in the Galactic plane (GP). The UWIFE survey was carried out using WFCAM at UKIRT in 2012 and 2013 (Lee et al. 2014). The [Fe II] narrow-band filter was used, having a central wavelength of 1.644 μm and an effective bandwidth of 0.026 μm . The WFCAM hosts four Rockwell Hawaii-II HgCdTe 2 k \times 2 k arrays, each covering 13.65 arcmin \times 13.65 arcmin in area at a pixel scale of 0.4 arcsec. Four pointings of the telescope covered a contiguous area of 0.75 deg² (designated as ‘tile’, following the WFCAM terminology). Each pointing was composed of a set of dithered and microstepped observations, fully sampling the point spread function in good seeing conditions (< 0.8 arcsec). The total integration time per pixel was 720 s. The final [Fe II] images have a nominal 5σ detection limit of 18.7 mag for point sources, with a median seeing of 0.83 arcsec. For extended diffuse sources, the corresponding surface brightness limit is $8.1 \times 10^{-20} \text{ W m}^{-2} \text{ arcsec}^{-2}$.

Lee et al. (2014) also produced continuum-subtracted [Fe II] images (hereafter [Fe II]-H images) by using the *H*-band images from the GPS. The continuum subtraction was carried out in two steps, that is, point-like continuum sources were first removed in both [Fe II] and *H*-band images, and then the point source removed *H*-band images were subtracted from the point source removed [Fe II] images to remove extended continuum sources. The details of the observation and data processing procedure can be found in Lee et al. (2014).

All [Fe II] and [Fe II]-H images from UWIFE are available at the UWIFE web page.¹ The images consist of 220 tiles, where a single tile is a square of 54 arcmin \times 54 arcmin in equatorial coordinates. The tiles are arranged as 55 stripes of four consecutive tiles at constant declination along the GP, covering a region within the First Galactic Quadrant of $7^\circ < l < 62^\circ$; $|b| \lesssim 1.5^\circ$ (see fig. 1 of Lee et al. 2014). On the web page, the UWISH2 H₂ and GPS *JHK*-band images are also available.

2.2 Source identification

In this study, we first aimed to identify IFOs in the continuum-subtracted images (hereafter, [Fe II]-H). We identified most of the IFOs through visual inspection and added several faint IFOs by mean of an automatic source identification, which uses the same algorithm as UWISH2 (Froebrich et al. 2015).

¹<http://gems0.kasi.re.kr/uwif/>

We focused on the extended sources in this study. Visual inspections were carried out twice for the whole survey area. We searched for all possible emission features and, for each feature, we defined an ellipse approximately surrounding the emitting area by eye. Then, the central coordinates, radii, and PA of the ellipses are measured and listed in Table 1. All IFO candidates identified in the [Fe II]-H image were double-checked in both unsubtracted [Fe II] and GPS *H*-band images to confirm whether they were a real source or not. As the UWIFE and GPS observations were separated by several years, variable sources were seen as emission or absorption in the [Fe II]-H images. In particular, since artefacts with a negative digital number (DN) in GPS *H*-band resemble real sources in the [Fe II]-H image, we checked the position of all IFO candidates in the corresponding *H*-band data.

In addition, there are various kinds of artefacts in the [Fe II]-H images. Representative artefacts are: the residuals of bright stars, ghosts, crosstalks, cross-strips after star subtraction, and the diffraction pattern of bright stars (see Appendix A1, e.g.). Residuals of high proper-motion stars were also left in the [Fe II]-H images. We also excluded the features hampered by the artefacts from bright stars. The sources that show diffuse structures in both [Fe II] and *H* which are significantly brighter in [Fe II] compared to the GPS *H* band, or the sources with a low probability of being scattered emission from dust seen in the GPS *H* band, were selected as real sources.

Using the [Fe II]-H images, we conducted an unbiased automatic detection with the code used for identifying MHOs in UWISH2 (Froebich et al. 2015) to benefit from its objectiveness. We adjusted the code to fit the specifications of UWIFE data: (1) remove small-scale features (residual of star subtraction), determine the large-scale background level from a 40 arcsec scale median filter, and calculate its noise value. (2) Draw contours at the 1σ level in ds^2 and identify the isolated contours as ‘regions’. The level was determined empirically to include faint emission of IFOs. The low (1σ) level produces contours around the remaining point sources and noise peaks, but those ‘false’ regions are removed by a minimum size limit in the next stage. (3) Remove contours that are too small ($<4 \text{ arcsec}^2$) or near the image borders. (4) To avoid mistakenly identifying star residual as IFOs, remove contours smaller than 35 arcsec^2 if they are located within 3 arcsec to the Two Micron All-Sky Survey (2MASS) *H*-band stars brighter than 15 mag. This procedure was conducted for all UWIFE tiles except for tile 003, 080, and 196 due to the late release of the *H*-band data in the GPS survey.

All sources identified by the automatic detection were cross-checked by visual inspection. We first examined whether the identified source from the code is an image artefact or not. Appendix A1 shows some examples of the artefacts, including residuals of detector crosstalk and diffraction patterns from saturated stars. These non-astronomical sources can be easily distinguished by comparing them in the [Fe II] and *H*-band images and were removed from the catalogue. We also rejected point-like sources (e.g. high proper-motion stars, variables, [Fe II]-emitting stars, etc.) that are not considered in this paper. Note that the visual identification treats a group of clumpy structures as a single object (e.g. shells of SNR). On the other hand, the code identifies the substructures separately. We fitted each automatically identified IFO with an ellipse and derived the geometrical parameters of the semimajor and semiminor axes, and PA. This process added 14 IFOs, and the complete catalogue is

presented in Table 1, which also provides their coordinates, sizes, fluxes, and counterparts.

2.3 Photometry

We conducted photometry of the IFOs in the [Fe II]-H images. Since our targets have an extended structure, we adopted aperture photometry. In the [Fe II]-H data, artefacts often have higher digital counts than IFOs. Therefore, masking artefacts is a crucial process. We masked the identified artefacts to prevent large uncertainties in the aperture photometry. The residuals of point sources (stars) brighter than 14th magnitude in the *H* band (based on the 2MASS point source catalogue, Skrutskie et al. 2006) were also masked. The size of the masking area was 6 arcsec in diameter, which is large enough to cover general residual patterns. When instrumental artefacts such as electronic crosstalk or diffraction patterns intruded on the aperture, we manually masked them to prevent any contamination.

In order to derive the total flux (F_{tot}) of the identified IFOs in a scientific unit (W m^{-2}), we used the following equation:

$$F_{\text{tot}} = F_0 \cdot \frac{\text{DN}}{t_{\text{exp}}} \cdot 10^{-0.4 m_{\text{zpt}}},$$

F_0 is the in-band flux of Vega falling in the [Fe II] filter ($3.27 \times 10^{-11} \text{ W m}^{-2}$, Lee et al. 2019), whereas t_{exp} and m_{zpt} are the net exposure time (60 s) and the zero-point magnitude of each image, respectively (‘EXPTIME’ and ‘MAGZPT’ in the image header). DN is the total DN falling in the aperture corrected for the sky background. This local background of each source was estimated from a sky annulus with an inner and outer radii of 1.2 and 1.5 times the aperture. We took the mode of the sky values to further avoid the effect of any possible artefacts. The uncertainty of the flux is estimated considering the photometric calibration error from the uncertainty of the zero-point magnitude of $\sim 0.06 \text{ mag}$, which corresponds to ~ 6 per cent of the total flux (Lee et al. 2019). The contribution of Poisson noise from aperture photometry and sky subtraction is negligible. The former is less than one-fifth, and the latter is less than one-tenth of absolute calibration uncertainty.

2.4 Search for associated exciting sources

We have searched for the possible driving source(s) of IFOs via positional cross-matching with previously known sources: SNRs, HII regions, compact (CHII) and UCHII regions, LBVs, PNe, and YSOs. IFOs associated with these sources are classified as SNR-IFO, HII-IFO, CHII-IFO, LBV-IFO, PN-IFO, and YSO-IFO, respectively. The rest of the IFOs are classified as ‘unknown-IFO’. In the following, we describe the processes and catalogues employed for the search for the individual exciting source types.

SNRs have complex and filamentary structures often with a large spatial extent. Thus, a careful identification and the separation of genuine SNR-origin from mere superposition was required. We first selected IFOs located within the boundary of known SNRs, using the central positions and sizes of SNRs in the Galactic SNR catalogue of Green (2019). We then referred to the references in the catalogue and also the SIMBAD Astronomical Database³ for the multiwavelength morphology of SNRs for the confirmation of the association. If an IFO shows a coherent structure occupying a similar extent and/or its morphology implies a spatial correlation with the SNRs, we categorized it as an SNR-IFO. We also checked the area in SIMBAD

²<https://sites.google.com/cfa.harvard.edu/saoimageds9>

³<http://simbad.harvard.edu/simbad/>

Table 1. Catalogue of identified IFOs.

IFO no.	UWIFE designation	l	b	r_1	r_2	PA	Area	F_{tot}	Counterpart
		[deg]	[deg]	[arcsec]		[deg]	[arcsec ²]	[10 ⁻¹⁷ W m ⁻²]	
IFO 001 ^a	J180136.927 – 224228.03	7.19190	+ 0.06028	9.3	2.9	40	84.7	5.88	–
IFO 002	J180210.565 – 214326.69	8.11119	+ 0.43376	2.3	2.0	0	14.4	0.89	–
IFO 003 ^a	J180212.398 – 223720.49	7.33358	–0.01604	2.6	1.8	157	14.7	2.25	YSO
IFO 004	J180219.380 – 213350.56	8.26721	+ 0.48309	5.4	4.2	120	71.2	4.27	YSO
IFO 005	J180511.698 – 195040.56	10.09481	+ 0.74344	18.0	7.2	60	407.1	3.15	PN
IFO 006	J180514.644 – 195027.77	10.10356	+ 0.73511	16.0	7.3	60	366.9	6.04	PN
IFO 007 ^b	J180627.378 – 213227.20	8.75903	–0.34293	1400.0	1400.0	0	6157521.8	–	SNR
IFO 008	J180640.397 – 220136.63	8.35934	–0.62390	4.5	4.2	20	59.3	13.80	PN
IFO 009	J180640.585 – 220126.95	8.36205	–0.62322	6.1	4.9	25	93.9	6.10	PN
IFO 010	J180732.633 – 202606.06	9.84849	–0.02588	3.4	3.4	0	36.3	0.91	YSO
IFO 011	J180916.373 – 201852.96	10.15033	–0.32179	68.0	39.0	150	8331.5	72.10	H II
IFO 012	J180925.793 – 201934.10	10.15813	–0.35954	67.3	62.5	0	13214.3	166.00	H II
IFO 013	J181050.844 – 205738.76	9.76257	–0.95642	4.2	3.4	160	44.8	1.59	YSO
IFO 014	J181051.015 – 205748.39	9.76056	–0.95829	5.9	3.8	170	70.4	2.81	YSO
IFO 015	J181129.632 – 192515.52	11.18511	–0.34766	131.4	131.1	0	54118.7	1090.00	SNR
IFO 016	J181312.424 – 164111.54	13.77907	+ 0.60864	3.5	2.8	90	30.7	0.94	YSO
IFO 017	J181322.096 – 174758.22	12.82064	+ 0.04152	1.3	1.3	0	5.3	0.35	–
IFO 018	J181407.762 – 185101.73	11.98433	–0.62012	3.3	2.1	110	21.7	0.64	YSO
IFO 019	J181408.073 – 185058.01	11.98582	–0.62071	1.7	1.5	60	8.0	0.44	YSO
IFO 020	J181413.148 – 175528.03	12.80784	–0.19604	8.3	7.4	90	192.9	4.79	YSO
IFO 021	J181415.121 – 175557.61	12.80436	–0.20684	10.7	8.3	120	279.0	1.57	YSO
IFO 022	J181419.929 – 175616.05	12.80898	–0.22602	67.6	42.7	160	9068.2	69.50	YSO
IFO 023	J181422.928 – 182508.51	12.39193	–0.46647	3.0	2.7	60	25.4	1.95	YSO
IFO 024	J181434.822 – 164514.38	13.87718	+ 0.28764	8.0	4.0	160	100.3	3.18	UCHII
IFO 025 ^b	J181436.683 – 164507.70	13.88234	+ 0.28200	24.0	20.0	0	1507.9	6.60	UCHII
IFO 026 ^b	J181437.731 – 164526.78	13.87970	+ 0.27580	13.0	8.0	60	326.7	9.40	UCHII
IFO 027	J181521.196 – 160255.94	14.58529	+ 0.46128	4.5	3.9	160	55.1	7.16	PN
IFO 028	J181627.693 – 163553.67	12.45471	–0.99317	7.6	4.3	50	102.6	0.30	–
IFO 029	J181658.050 – 162710.24	14.41440	–0.07172	3.8	3.5	90	41.7	4.66	YSO
IFO 030	J181724.551 – 172216.03	13.65638	–0.60071	4.0	3.1	40	38.9	2.01	YSO
IFO 031	J181750.609 – 120805.80	18.31683	+ 1.78972	1.7	1.4	90	7.4	0.43	YSO
IFO 032	J181750.953 – 120802.39	18.31835	+ 1.78895	2.7	2.5	0	21.2	2.11	YSO
IFO 033	J181758.445 – 120724.48	18.34208	+ 1.76705	2.3	2.0	90	14.45	0.98	YSO
IFO 034	J181828.251 – 165525.05	14.17066	–0.61193	2.0	1.3	0	8.1	0.54	YSO
IFO 035	J181828.590 – 165523.72	14.17162	–0.61295	3.1	2.4	120	23.3	4.89	YSO
IFO 036	J181837.058 – 134248.28	17.01527	+ 0.87651	6.5	4.1	100	83.7	1.51	YSO
IFO 037	J181839.539 – 134237.23	17.02271	+ 0.86910	2.8	2.5	120	21.9	0.83	YSO
IFO 038	J181845.167 – 150257.21	15.85388	+ 0.21557	6.4	5.1	130	102.5	7.34	SNR*
IFO 039	J181847.449 – 135022.70	16.92393	+ 0.77973	1.2	1.2	0	4.5	0.23	YSO
IFO 040	J181849.365 – 134952.55	16.93498	+ 0.77687	2.0	1.5	120	9.4	0.49	YSO
IFO 041	J181855.428 – 135145.51	16.91893	+ 0.74041	10.2	5.6	30	179.4	16.80	HH
IFO 042	J181858.301 – 135236.39	16.91199	+ 0.72350	10.0	6.0	150	188.4	2.65	HH
IFO 043	J181858.835 – 135252.81	16.90897	+ 0.71943	4.0	3.0	140	37.6	0.23	HH
IFO 044	J181901.895 – 135346.30	16.90173	+ 0.70150	15.0	10.0	140	471.2	2.52	HH
IFO 045	J181905.871 – 134522.91	17.03256	+ 0.75343	25.1	19.0	90	1498.2	4.14	YSO
IFO 046 ^a	J181914.708 – 164949.13	14.34049	–0.73101	1.6	1.3	30	6.5	0.33	YSO
IFO 047	J181917.916 – 164355.78	14.43321	–0.69611	4.0	3.0	90	37.6	2.24	YSO
IFO 048	J181922.591 – 134114.45	17.12557	+ 0.72624	7.1	3.0	30	66.9	0.70	YSO
IFO 049	J181925.259 – 134542.71	17.06480	+ 0.68168	5.4	4.0	0	67.8	1.01	YSO
IFO 050	J181927.118 – 151211.16	15.79925	–0.00624	3.5	2.3	30	25.2	8.84	PN
IFO 051 ^b	J182019.871 – 161031.33	15.04081	–0.65134	2.2	2.0	0	13.8	1.33	YSO
IFO 052 ^b	J182020.767 – 161018.45	15.04566	–0.65282	2.8	2.0	0	17.5	4.05	YSO
IFO 053 ^b	J182021.725 – 161015.05	15.04831	–0.65575	2.3	1.7	30	12.2	1.05	YSO
IFO 054 ^b	J182024.436 – 161126.80	15.03583	–0.67472	18.7	9.4	140	552.2	5.11	HCHII
IFO 055 ^b	J182028.170 – 161245.10	15.02369	–0.69815	250.0	100.0	130	78539.8	1257.00	H II
IFO 056	J182032.784 – 160124.98	15.19905	–0.62538	7.6	3.9	0	93.1	0.63	YSO
IFO 057	J182034.306 – 160158.97	15.19359	–0.63521	8.0	2.0	120	50.2	1.12	YSO
IFO 058	J182035.196 – 161942.63	14.93464	–0.77759	330.0	120.0	160	124407.0	473.00	H II
IFO 059	J182035.224 – 140436.84	16.92057	+ 0.28355	1.9	1.8	90	10.7	0.85	YSO
IFO 060	J182035.656 – 140409.72	16.92803	+ 0.28556	5.8	5.1	30	92.9	7.82	YSO
IFO 061	J182036.014 – 140344.82	16.93481	+ 0.28754	3.3	3.3	0	34.2	0.54	YSO
IFO 062 ^b	J182037.224 – 160828.36	15.10369	–0.69649	341.8	197.3	115	211860.0	875.00	H II
IFO 063	J182049.312 – 140353.86	16.95793	+ 0.23896	2.6	2.4	30	19.6	0.60	YSO

Table 1 – continued

IFO no.	UWIFE designation	l [deg]	b [deg]	r_1 [arcsec]	r_2 [arcsec]	PA [deg]	Area [arcsec ²]	F_{tot} [10 ⁻¹⁷ W m ⁻²]	Counterpart
IFO 064	J182056.997 – 161934.88	14.97762	–0.85347	100.0	50.0	160	15707.9	58.27	H II
IFO 065 ^b	J182119.587 – 162224.78	14.97848	–0.95541	15.0	13.5	140	636.2	81.16	LBV
IFO 066	J182121.701 – 160424.16	15.24737	–0.82163	18.2	16.2	0	926.2	8.96	H II
IFO 067	J182134.867 – 161209.80	15.15799	–0.92897	25.5	8.1	135	648.8	11.40	H II
IFO 068	J182228.548 – 171548.32	14.32188	–1.61616	3.5	2.3	30	25.2	1.07	–
IFO 069	J182432.827 – 130950.81	18.17823	–0.13740	101.0	56.0	70	17768.8	21.50	SNR
IFO 070	J182448.037 – 131345.15	18.14955	–0.22238	6.0	5.0	90	94.2	1.38	SNR
IFO 071	J182459.449 – 131552.08	18.14002	–0.27977	130.0	90.0	60	36756.6	110.10	UCHII
IFO 072 ^a	J182548.520 – 130629.85	18.37107	–0.38288	7.0	4.5	120	98.9	82.00	YSO
IFO 073	J182619.105 – 101318.67	20.98261	+ 0.85302	19.9	7.4	120	462.6	6.64	PN
IFO 074	J182656.992 – 113210.92	19.89167	+ 0.10337	4.2	3.4	60	44.8	4.51	YSO
IFO 075	J182851.018 – 124415.55	19.04409	–0.86640	67.4	27.4	15	5801.7	14.10	SNR
IFO 076	J182852.671 – 124311.10	19.06306	–0.86404	16.4	5.5	0	283.3	2.55	SNR
IFO 077 ^a	J182859.486 – 115026.04	19.85474	–0.48056	6.1	2.3	80	44.0	3.45	YSO
IFO 078	J182919.659 – 124153.90	19.13295	–0.95127	132.0	89.4	140	37073.3	192.00	SNR
IFO 079	J182930.563 – 131350.84	18.68160	–1.23735	8.7	3.6	70	98.3	1.89	SNR
IFO 080 ^b	J183314.281 – 100831.20	21.84283	–0.61852	500.0	200.0	40	314159.2	627.00	SNR
IFO 081 ^b	J183328.975 – 110726.68	20.99907	–1.12478	11.8	6.4	25	237.2	49.30	PN
IFO 082 ^a	J183330.115 – 050050.30	26.43603	+ 1.69464	6.8	3.5	165	74.7	6.73	UCHII
IFO 083 ^a	J183330.673 – 050110.94	26.42214	+ 1.68483	15.4	12.0	114	580.5	150.00	UCHII
IFO 084	J183331.327 – 103257.93	21.51352	–0.86841	35.0	17.0	50	1869.2	6.59	SNR
IFO 085	J183333.430 – 103402.86	21.50149	–0.88437	44.8	42.1	90	5925.2	59.70	SNR
IFO 086	J183404.384 – 071820.28	24.45479	+ 0.50637	60.0	30.0	160	5654.8	9.42	UCHII
IFO 087	J183408.045 – 071801.82	24.46632	+ 0.49530	12.7	7.6	30	303.2	1.72	UCHII
IFO 088	J183420.390 – 084722.27	23.16829	–0.23606	600.0	400.0	10	753982.2	584.00	SNR
IFO 089	J183425.284 – 075448.33	23.95514	+ 0.14969	11.0	9.3	20	321.3	11.20	UCHII
IFO 090	J183426.772 – 075428.56	23.96285	+ 0.14677	1.1	1.1	0	3.8	0.32	UCHII
IFO 091	J183541.856 – 072203.61	24.58525	+ 0.12015	15.0	8.0	140	376.9	3.06	YSO
IFO 092	J183648.912 – 071850.94	24.76014	–0.10123	2.3	1.9	20	13.7	2.13	YSO
IFO 093	J183716.440 – 032958.39	28.20160	+ 1.54899	4.0	2.3	10	28.9	0.39	SNR
IFO 094	J183720.713 – 064200.69	25.36590	+ 0.06398	2.0	1.7	30	10.6	1.18	YSO
IFO 095	J183730.398 – 061412.49	25.79595	+ 0.24115	8.0	8.0	0	201.0	3.93	PN/UCHII
IFO 096 ^b	J183740.829 – 061452.41	25.80594	+ 0.19768	180.0	180.0	0	101787.6	221.00	–
IFO 097	J183813.600 – 064815.32	25.37390	–0.17819	25.2	8.0	45	633.3	25.60	UCHII
IFO 098 ^b	J183907.168 – 043230.84	27.48618	+ 0.66204	500.0	400.0	20	628318.5	167.00	SNR
IFO 099	J183909.562 – 071927.89	25.01779	–0.62238	4.0	3.0	0	37.6	2.05	YSO
IFO 100	J183911.798 – 072019.31	25.00933	–0.63714	3.4	2.8	90	29.9	2.56	YSO
IFO 101	J183913.302 – 072057.12	25.00284	–0.64748	1.8	1.5	130	8.4	1.12	YSO
IFO 102	J183931.338 – 054409.74	26.47082	+ 0.02555	3.3	2.3	80	23.8	3.07	LBV
IFO 103	J183931.437 – 054414.64	26.46980	+ 0.02456	2.4	1.4	0	10.5	1.27	LBV
IFO 104	J183950.426 – 043037.60	27.59648	+ 0.51675	4.5	2.0	0	28.2	0.20	YSO
IFO 105	J184120.333 – 045606.47	27.38989	–0.00960	150.0	120.0	70	56548.6	274.00	SNR
IFO 106	J184358.299 – 035306.11	28.62385	–0.11304	216.2	170.1	0	115534.0	197.00	SNR
IFO 107	J184414.391 – 041754.32	28.28667	–0.36139	7.0	3.6	165	79.1	2.70	UCHII
IFO 108	J184422.810 – 041734.78	28.30748	–0.39003	3.9	2.8	90	34.3	0.57	YSO
IFO 109	J184501.647 – 001716.48	31.94472	+ 1.29493	2.6	2.1	90	17.1	0.37	–
IFO 110	J184515.462 – 031604.01	29.31952	–0.11656	4.3	3.5	120	47.2	0.85	–
IFO 111	J184559.282 – 024502.58	29.86281	–0.04271	7.3	3.7	135	84.8	5.47	YSO
IFO 112 ^a	J184829.526 – 021003.18	30.81541	–0.25716	2.9	1.5	150	13.6	0.52	PN
IFO 113	J184927.068 – 005638.37	31.86530	+ 0.01156	240.0	210.0	120	158336.2	3423.55	SNR
IFO 114 ^{ab}	J184933.121 – 003810.21	32.59821	+ 0.35877	8.9	6.3	156	176.1	6.46	PN/H II
IFO 115	J184955.670 – 010153.39	31.84176	–0.13438	1.9	0.9	130	5.3	0.25	YSO
IFO 116 ^{ab}	J185026.138 + 012739.08	33.81379	+ 0.73328	5.5	2.1	68	36.2	0.83	–
IFO 117 ^b	J185125.777 – 000930.42	32.78995	–0.07040	740.0	530.0	170	1232132.7	114.00	SNR
IFO 118	J185128.102 + 002840.15	33.36065	+ 0.21110	8.6	8.0	170	216.1	1.02	HH
IFO 119	J185140.619 + 002850.89	33.38708	+ 0.16604	7.0	3.5	5	70.6	34.60	HH
IFO 120	J185141.136 + 002900.71	33.39049	+ 0.16537	1.4	1.2	90	5.2	0.24	HH
IFO 121	J185144.114 + 002911.43	33.39880	+ 0.15568	7.0	4.0	90	87.9	1.64	HH
IFO 122	J185249.992 + 022802.16	35.28688	+ 0.81455	1.4	0.9	90	3.9	0.56	YSO
IFO 123	J185251.980 + 022804.91	35.29134	+ 0.80753	1.2	1.0	90	3.7	0.74	YSO
IFO 124	J185353.538 + 015714.24	34.95065	+ 0.34503	3.9	2.8	60	34.3	1.87	YSO
IFO 125	J185516.571 + 030512.10	36.11639	+ 0.55408	8.0	4.8	160	120.6	9.03	YSO
IFO 126	J185521.357 + 030154.61	36.07665	+ 0.51134	1.5	1.4	0	6.5	0.42	YSO

Table 1 – *continued*

IFO no.	UWIFE designation	l	b	r_1	r_2	PA	Area	F_{tot}	Counterpart
		[deg]	[deg]	[arcsec]	[arcsec]	[deg]	[arcsec ²]	[10 ⁻¹⁷ W m ⁻²]	
IFO 127 ^a	J185534.205 + 021908.12	35.31833	+ 0.06246	6.3	4.2	175	83.1	1.46	UCHII
IFO 128	J185602.316 + 012139.70	34.66768	-0.40273	1120.0	860.0	150	3025982.1	3942.00	SNR
IFO 129 ^b	J185737.231 + 020350.02	35.47351	-0.43365	11.0	11.0	0	380.1	6.54	PN
IFO 130 ^b	J185738.014 + 020332.58	35.47069	-0.43876	11.0	11.0	0	380.1	7.11	PN
IFO 131	J185808.531 + 010048.26	34.59832	-1.02921	4.5	4.5	0	63.6	3.22	YSO
IFO 132	J185810.531 + 013656.88	35.13815	-0.76162	17.5	14.7	0	808.1	24.60	CHII
IFO 133	J185905.114 + 004833.51	34.52432	-1.33211	2.1	1.4	0	9.2	1.60	-
IFO 134	J185910.539 + 014013.31	35.30084	-0.95905	5.0	3.1	130	48.7	1.56	YSO
IFO 135	J185923.204 + 010413.33	34.79106	-1.27998	2.8	2.8	0	24.6	0.39	-
IFO 136	J190003.046 + 055926.63	39.24352	+ 0.82130	26.6	13.0	170	1086.3	3.29	YSO
IFO 137	J190347.070 + 050946.79	38.93288	-0.38355	35.0	18.0	100	1979.2	8.31	UCHII
IFO 138	J190403.633 + 050753.38	38.93638	-0.45909	5.2	4.4	30	71.8	2.74	UCHII
IFO 139 ^b	J190404.180 + 052703.51	39.22135	-0.31459	197.3	150.1	0	93037.4	618.00	SNR
IFO 140	J190540.231 + 074634.49	41.46973	+ 0.39939	550.0	450.0	100	777544.2	1421.45	SNR
IFO 141	J190659.919 + 052253.12	39.49429	-0.99412	5.4	5.2	90	88.2	12.20	YSO
IFO 142	J190731.328 + 052333.18	39.56407	-1.10471	5.5	4.4	130	76.0	1.17	-
IFO 143	J190734.278 + 070829.06	41.12259	-0.31090	170.0	120.0	115	64088.4	1691.00	SNR
IFO 144	J190813.552 + 052757.00	39.70977	-1.22650	1.6	1.4	90	7.0	0.23	YSO
IFO 145	J190816.446 + 052726.79	39.70784	-1.24102	3.7	1.6	125	18.5	0.40	YSO
IFO 146	J190816.782 + 052506.10	39.67378	-1.26024	3.9	3.6	0	44.1	1.69	YSO
IFO 147	J191106.846 + 090604.55	43.26595	-0.18486	160.0	150.0	120	75398.2	4739.00	SNR
IFO 148 ^a	J191327.650 + 105334.62	45.12149	+ 0.13279	11.2	8.5	137	299.0	14.80	UCHII
IFO 149 ^a	J191327.754 + 105413.66	45.13129	+ 0.13745	3.2	2.4	105	24.1	0.48	UCHII
IFO 150	J191530.963 + 132747.36	47.63089	+ 0.88215	1.7	1.4	120	7.4	0.29	-
IFO 151	J192026.005 + 111955.24	46.30789	-1.17527	5.6	3.9	120	68.6	2.25	YSO
IFO 152	J192029.411 + 111942.04	46.31118	-1.18928	1.3	1.1	90	4.5	0.27	YSO
IFO 153	J192029.485 + 110159.44	46.05061	-1.32806	7.5	5.3	90	124.8	8.71	HH
IFO 154	J192054.201 + 143031.29	49.16624	+ 0.21581	1.6	1.6	0	8.0	0.75	YSO
IFO 155	J192113.714 + 105232.92	45.99659	-1.56168	4.2	2.8	110	36.9	2.34	HH
IFO 156	J192127.938 + 154426.63	50.31742	+ 0.67543	4.5	2.6	30	36.7	1.58	UCHII
IFO 157	J192142.900 + 155351.18	50.48401	+ 0.69629	60.0	20.0	90	3769.9	1.55	PN
IFO 158	J192309.835 + 142912.63	49.40475	-0.27709	40.0	15.0	10	1884.9	8.92	H II
IFO 159 ^b	J192255.023 + 140745.93	49.06144	-0.39309	1200.0	1000.0	160	3769911.3	1647.09	H II
IFO 160 ^b	J192401.145 + 140105.48	49.08966	-0.68118	580.0	430.0	30	783513.2	582.37	SNR
IFO 161	J192348.822 + 143137.35	49.51449	-0.39670	350.0	210.0	115	230907.0	182.47	H II
IFO 162	J192348.169 + 143641.50	49.58771	-0.35441	50.0	30.0	25	4712.3	45.50	LBV
IFO 163	J192354.032 + 143548.00	49.58825	-0.38096	45.0	35.0	0	4948.0	5.76	YSO
IFO 164	J192451.838 + 155729.06	50.89493	+ 0.05763	1.5	1.5	0	7.0	0.24	PN
IFO 165	J192516.759 + 144625.72	49.89974	-0.59204	2.3	1.9	0	13.7	0.72	YSO
IFO 166	J192529.675 + 151646.36	50.36959	-0.39785	1.1	0.9	0	3.1	0.20	-
IFO 167	J192531.202 + 151603.90	50.36214	-0.40884	2.1	1.3	0	8.5	0.79	-
IFO 168	J192531.399 + 151556.79	50.36075	-0.41049	0.8	0.8	0	2.0	0.13	-
IFO 169	J192532.882 + 151538.18	50.35903	-0.41819	1.5	1.3	40	6.1	0.50	-
IFO 170	J192533.417 + 151616.62	50.36947	-0.41501	1.1	1.1	0	3.8	0.26	-
IFO 171	J192534.199 + 151612.08	50.36983	-0.41838	1.2	0.8	90	3.0	0.23	-
IFO 172	J192534.538 + 151632.38	50.37544	-0.41690	1.6	1.4	90	7.0	0.59	-
IFO 173 ^a	J192540.546 + 163305.18	51.50973	+ 0.16761	7.0	5.8	16	127.5	11.50	PN
IFO 174	J192547.157 + 145145.84	50.03612	-0.65762	4.0	2.5	140	31.4	4.22	YSO
IFO 175	J192557.625 + 150231.65	50.21401	-0.60951	1.8	1.6	90	9.0	0.36	YSO
IFO 176	J192557.848 + 150243.23	50.21727	-0.60877	10.0	5.5	0	172.2	1.98	YSO
IFO 177	J192852.403 + 171458.61	52.48804	-0.17205	3.3	2.8	90	29.0	0.92	YSO
IFO 178	J192918.342 + 175615.42	53.14142	+ 0.06679	15.0	10.0	140	471.2	4.95	YSO
IFO 179	J192918.796 + 175723.68	53.15891	+ 0.07429	3.7	1.9	130	22.0	0.31	YSO
IFO 180	J192920.127 + 175716.54	53.15971	+ 0.06871	25.0	8.0	80	628.3	4.41	YSO
IFO 181	J192920.506 + 175458.14	53.12668	+ 0.04898	7.0	5.0	110	109.9	1.11	YSO
IFO 182	J192922.491 + 174442.54	52.98034	-0.03983	4.6	3.5	0	50.5	0.57	-
IFO 183	J192931.617 + 175951.30	53.21927	+ 0.04934	4.4	3.0	90	41.4	1.08	YSO
IFO 184	J192931.871 + 180058.11	53.23604	+ 0.05734	1.7	1.4	90	7.4	0.25	YSO
IFO 185	J192932.874 + 180106.35	53.23994	+ 0.05495	7.0	4.0	45	87.9	1.06	YSO
IFO 186 ^b	J193001.921 + 175455.44	53.20473	-0.09547	5.4	3.3	20	55.9	3.15	YSO
IFO 187	J193120.744 + 192014.92	54.60141	+ 0.31496	1.5	1.3	90	6.1	0.35	-
IFO 188	J193323.546 + 195647.07	55.36730	+ 0.18676	20.0	18.0	40	1130.9	5.22	post-AGBc/YSO
IFO 189	J193831.665 + 202519.19	56.36978	-0.63373	7.0	4.3	150	94.5	10.50	YSO
IFO 190	J193914.355 + 224021.52	58.41141	+ 0.32789	7.0	6.0	40	131.9	3.75	YSO

Table 1 – continued

IFO no.	UWIFE designation	l	b	r_1	r_2	PA	Area	F_{tot}	Counterpart
		[deg]	[deg]	[arcsec]	[arcsec]	[deg]	[arcsec ²]	[10 ⁻¹⁷ W m ⁻²]	
IFO 191	J194014.058 + 232652.51	59.19889	+ 0.51050	5.5	4.2	80	72.5	1.57	–
IFO 192	J194103.922 + 220340.80	58.08778	–0.34083	3.2	1.7	0	17.1	0.62	YSO
IFO 193	J194127.149 + 222739.58	58.47940	–0.22095	14.2	6.8	130	303.3	14.70	YSO
IFO 194	J194241.016 + 225417.72	59.00574	–0.24738	6.5	3.0	60	61.2	2.76	YSO
IFO 195	J194244.693 + 232250.36	59.42558	–0.02322	3.0	3.0	0	28.2	0.65	HH
IFO 196	J194256.665 + 232435.17	59.47362	–0.04848	29.0	18.5	40	1685.4	5.20	HH
IFO 197	J194306.295 + 231810.63	59.39926	–0.13356	1.9	1.3	90	7.7	0.52	–
IFO 198	J194310.286 + 234358.03	59.77970	+ 0.06707	4.9	2.9	60	44.6	2.83	YSO
IFO 199	J194310.930 + 234402.64	59.78203	+ 0.06557	5.3	2.5	90	41.6	2.23	YSO
IFO 200	J194320.930 + 232952.89	59.59633	–0.08502	10.5	8.0	150	263.9	1.34	HH
IFO 201	J194610.902 + 221559.08	58.85575	–1.26581	7.1	2.5	120	55.7	0.96	–
IFO 202	J194620.335 + 243520.73	60.88253	–0.13043	14.0	12.9	80	567.3	17.20	UCHII
IFO 203	J194621.675 + 243516.78	60.88413	–0.13538	1.3	1.3	0	5.3	0.45	YSO
IFO 204	J194646.921 + 251241.33	61.47104	+ 0.09568	40.5	36.5	140	4644.0	111.00	UCHII/H II

Notes.^aIFOs marked with ‘a’ are identified only by an automatic detection method.

^bNote on the individual sources. IFO 7: due to the complexity of the region, flux is not provided. IFO 25–26: the flux of the superposed part is allocated only to IFO 26. IFO 51–55: IFO 51–54 are located inside IFO 55. IFO 62, 98: the flux is derived for a partial region free from severe artefacts. IFO 65, 81, 114, 116, 186: missing flux due to 2MASS-bright star mask. IFO 80, 96, 117: contaminated by an instrumental artefact. The pixels with $\text{DN} > \pm 3\sigma$ are masked for the flux measurement. IFO 129–130: contaminated by an instrumental artefact. The superposed part is excluded from the flux measurement. IFO 139: contaminated by an instrumental artefact. The pixels with $\text{DN} < -2\sigma$ are masked for the flux measurement. IFO 159–160: the flux of the superposed part is allocated only to IFO 160. Note that there is an astrometry problem with certain continuum-subtracted [Fe II] images, where IFO 4, 8, 9, 73, 114, 155, 165, and 186 are located. Therefore, we determined the central positions of the IFOs based on the UKIDSS NIR image.

*IFO 38 is located within the SNR G15.9 + 0.2 domain but highly confined to a southwestern region (see fig. 1 of Sasaki et al. 2018 for an X-ray image of the SNR). Since there is no other possible counterpart in the SIMBAD query and the X-ray emission is coincident, we concluded that the SNR origin cannot be ruled out.

for a possible superposition of unrelated, superposed sources such as PNe along the same line of sight. An IFO without noticeable morphological correlation but positionally coincident with evident SNR emission was categorized as an SNR-IFO (e.g. IFO 38).

Diffuse H II regions also occupy a large spatial area and have complex morphology, so that a SIMBAD/VizieR query by IFO coordinate with an arcminute radius often returns various kinds of incidental sources such as subfilaments of H II regions, jets/outflows from neighbouring YSOs, and merely superposed sources along the line of sight. Therefore, keeping in mind that proximity alone does not necessarily guarantee a true correlation, a morphological correlation was also taken into account for identifying H II regions as an exciting source. If needed, a query with a larger angular scale was conducted to locate the diffuse H II region. We then compared the morphology of IFOs with that of H II regions obtained from high-resolution radio images (GPS; White, Becker & Helfand 2005, NewGPS; Helfand et al. 2006, and the H I, OH, recombination line survey of the Milky Way, THOR continuum; Beuther et al. 2016) and/or data sets from large-scale multiwavelength studies (Povich et al. 2009; Roshi, Churchwell & Anderson 2017; Fujita et al. 2021). We also used small-scale surveys and targeted studies (see Section 4.3). The IFOs with a positive correlation have been categorized as HII-IFOs. However, since the [Fe II] line emission from an H II region is inherently faint, morphological correlation with radio is occasionally hard to confirm. On the basis of this possibility, a few extended and faint IFOs have also been regarded as HII-IFOs although they do not have a clear morphological relationship with an H II region (see comments in Section 4.3).

We further explored whether IFOs are associated with CHII, UCHII, or HCHII regions by querying VizieR within an arcminute-scale radius. Two comprehensive lists of UCHII regions were selected for the VizieR positional matching: the CORNISH UCHII region catalogue (Kalcheva et al. 2018), which is appropriate for the

comparison with the UWIFE survey data in terms of comparable high resolution ($1''.5$) and spatial coverage ($10^\circ < l < 65^\circ$, $|b| < 1^\circ$), and the catalogue presented by Bronfman, Nyman & May (1996) which is a large-scale compilation of *Infrared Astronomical Satellite* (IRAS) FIR colour-selected UCHII regions with higher Galactic latitude coverage ($|b| < 2^\circ$). The IFO positions were subsequently searched in SIMBAD to refer to targeted studies. We compared the [Fe II] line morphologies with available radio continuum images (see Section 4.2). When the IFOs show morphological correlation with radio structures or delineate the boundary of radio structures, they are classified as CHII-IFOs. IFOs having counterparts supposedly earlier or at a lower mass evolutionary stage of an UCHII region (e.g. HCHII region, UCHII precursor, ultracompact embedded cluster, UCEC, which was suggested as a lower mass class of UCHII) are also included in this category (see Section 4.2).

In order to identify IFOs associated with LBVs, the SIMBAD query was conducted with a radius criterion of 10 arcmin. But we noted that the list of LBVs and LBV candidates (hereafter cLBVs) has not been fully incorporated in SIMBAD, so we also used the catalogue of (c)LBVs compiled by Nazé, Rauw & Hutsemékers (2012) which lists the coordinates of 68 (c)LBVs. As far as we know, this is the most comprehensive catalogue of (c)LBVs. For example, Weis & Bomans (2020), in their review article of LBVs, presented a catalogue, but it has a smaller number of (c)LBVs than Nazé et al. (2012), that is, 47 versus 68. In the catalogue of Nazé et al. (2012), 22 LBVs (including 19 candidates) are located inside the UWIFE area. There was also an [Fe II] survey of nine LBVs by Smith (2002). Among the nine LBVs, only one was located within the UWIFE survey area and it has been identified in our survey, too.

For PN-IFOs, the SIMBAD query was used with a radius criterion of 10 arcmin. We additionally compared the morphology of IFOs with multiwavelength data from references in SIMBAD. In order to incorporate recently discovered PNe and PN candidates that

have not been updated in SIMBAD, we made use of the following databases and catalogues. We used the Hong Kong/AAO/Strasbourg H α (HASH) PN database which lists multiwavelength data of newly found ~ 3500 PNe and PN candidates (Parker, Bojčić & Frew 2016). The database includes three large-scale catalogues of Galactic PNe; the Strasbourg-ESO catalogue, the catalogue of Galactic PNe version 2000, and the Macquarie/AAO/Strasbourg H α (MASH) catalogues, together with 159 new PNe from the related IPHAS survey and ~ 400 from the literature. A large number of unpublished, new PN candidates are accessible in this database, which are mostly (1) older, redder, and have lower surface brightness or (2) are more remote and small-scaled, faint PNe (Parker et al. 2016). When the counterpart is not a well-known source and is identified only in the HASH database, we checked the association using the references provided. There are PNe only detectable in NIR, so the recent study of PNe based on the UWISH2 data (Gledhill et al. 2018) was also checked for possible counterparts. This study complements faint or small-scale PNe previously undiscovered.

For the remaining IFOs, we made use of several large-scale catalogues of YSOs alongside catalogues for specific regions or targets. The large-scale survey of YSOs in four evolutionary stages (i.e. quiescent, YSO, protostellar, and massive star-forming stages, Urquhart et al. 2018) was used to find YSO-IFOs, keeping in mind the survey resolution (30 arcsec). The Infrared Array Camera red-source catalogue was also used in the same manner (Robitaille et al. 2008) to locate YSO-candidate counterparts. When there was a positive match, we subsequently displayed their positions on the [Fe II]-H images with H $_2$ contours of UWISH2 data to confirm their association. H $_2$ images are useful since H $_2$ emission is usually more easily excited, forming a series of knots between an IFO and the YSO that drives an H $_2$ outflow. When the positional match and morphological information could not pinpoint an obvious YSO counterpart, we listed up to two YSOs. Also used are small-scale survey catalogues to benefit from a deeper searches for YSOs. Kim, Koo & Davis (2015) conducted a detailed survey of YSO candidates in the infrared dark cloud (IRDC) G53.2 region and investigated their evolutionary stages. Povich & Whitney (2010) investigated the M17 region where we have identified many IFOs, and the study provided the evolutionary stages of the YSOs. Ragan, Bergin & Gutermuth (2009) covered multiple IRDC regions in the UWIFE survey area and a YSO class with MIR colour and distance information was provided. Other small-scale catalogues of YSOs available in Vizier were also used when available (Section 4.1).

Since Herbig–Haro (HH) objects are often bright in [Fe II] emission, we attempted to locate the [Fe II]-emitting HH objects separately from YSO-IFOs. 454 Galactic HH objects have been compiled by Reipurth (2000), who continuously updated the SIMBAD database to include newly found HH objects. We retrieved all HH objects in SIMBAD, up to HH 1213, which includes 3140 substructures (e.g. HH 250A and 250B). First, we search for YSO-IFOs and unknown-IFOs within a radius criterion of 10 arcmin for a given HH object. When there was a match, we looked for a possible association of the IFO with HH object structures via multiwavelength images (mainly H α from IPHAS, Witham et al. 2008). For example, IFO 195 which is associated with the parsec-scale HH 803 has a very compact, small-scale structure. It was originally categorized as an unknown-IFO since we could not find any associated source just based on positional proximity. However, when we plot the IPHAS H α and UWISH2 molecular hydrogen emission contours together, we could associate IFO 195 with the southwestern tip of the series of aligned structures of HH 803 in H α and H $_2$

emission. Fig. 1 shows the representative IFOs with respect to each counterpart.

3 RESULTS

3.1 Catalogue of IFOs and their statistical properties

The full catalogue of extended IFOs is presented in Table 1 and the description of each column of the catalogue is as follows:

Column 1. IFO identifier: designations of IFOs by a catalogue number in ascending order. When the IFO is identified only by a source detection algorithm, we marked them with an ‘a’ after its designation.

Column 2. IFO conventional designation: IFO full-name derived from Right Ascension and Declination (J2000) of the source centre. It follows the 2MASS convention for the naming, that is,

IFO JHH:MM:SS.SSS \pm DD:MM:SS.SS.

Columns 3 and 4. Galactic longitude (l) and latitude (b): the centre position of the source, in units of degree, in Galactic coordinates. For automatically identified IFOs, we adopt the geometric centre of the polygon by two-dimensional Gaussian fitting of an ellipse.

Columns 5 and 6. Semimajor axis (r_1) and semiminor axis (r_2): maximum semimajor and minor angular radius of the IFO in units of arcseconds.

Column 7. Position angle (PA): the angle of the semimajor axis of an ellipse, in a counterclockwise direction, from north to east in units of degree.

Column 8. Area: an area of an ellipse determined by the semimajor and semiminor axes (columns 5 and 6), in square arcseconds.

Column 9. F_{tot} : total flux derived from summing up all flux inside an ellipse drawn from columns 5 and 6. See the photometry section Section 2.3 for details.

Column 10. Counterpart : classification of the IFO indicating the most probable known object as follows: YSO-IFO – outflows or jets from an YSO or YSO candidate, HII-IFO – any outflows surrounding emission originated from the H II region, subdivided into HCHII, UCHII, CHII, and diffuse H II region, SNR-IFO – emission originates in SNR, PN-IFO – emission associated with PN/PN candidates, further classified into PN, PNc, and post-asymptotic giant branch (AGB), LBV-IFO – nebula structure around an LBV or LBV candidate, and unknown-IFO – multiple corresponding known object candidates or no possible known source in the vicinity.

Our IFO catalogue contains 204 sources identified from 219 tiles, which is about 180 deg 2 in total. This number corresponds to an average surface density of ~ 1.1 IFOs per deg 2 in the first quadrant of the GP. This number should be regarded as a lower limit since our source identification methods were conservative. In general, the results of the manual and automatic search by the source detection algorithm were in good agreement. The 14 sources found only by the source detection algorithm, are marked with an ‘a’ after the IFO number in Table 1. They were either very faint or resembled artefacts. The majority of catalogue sources are new discoveries of [Fe II] emission, and represent an order of magnitude increase in the number of extended [Fe II] sources in the first Galactic quadrant.

Table 2 presents basic statistics of IFOs for each counterpart type. We identified 100 YSO-IFO (87 YSOs and 13 HHs), 33 HII-IFO (22 CHII and 11 HII), 25 SNR-IFO, 17 PN-IFO, 4 LBV-IFO, and 25 IFOs without counterparts. Note that if a counterpart source has two distinct [Fe II] structures, they are counted as two separated IFOs which share a common counterpart (e.g. IFO 85 and 86 are from SNR G21.5–0.9 and are counted as 2 SNR-IFOs). Also, one SNR-IFO




Figure 1. Continuum-subtracted [Fe II] and NIR three-colour images of IFOs with various origins: (a) YSO outflow: IRAS 18177 – 1405; (b) CHII region: G35.2S; (c) diffuse H II region: GAL 10.2 – 0.3; (d) LBV nebula: [KW 97] 37–17; (e) PN: IRAS 19234 + 1627; and (f) SNR: W49B. Grey-scale images in the upper rows are UWIFE [Fe II]-H images; Colour-composite images in the lower rows are R/G/B = *KHJ*-band images from the UKIDSS GPS survey. The units of the UWIFE [Fe II]-H images are DN, with the darker colour denoting a higher DN. The UWIFE images of the panels (a) IRAS 18177 – 1405 and (d) [KW97] 37–17 are smoothed with a two-pixel Gaussian. In all images, north is at the top, and east to the left side. Note the following artefacts: panel (a) IRAS 18177 – 1405: diffraction spike from southwest to northeast; (b) G35.2S: crosstalk on the northwest edge of the source; diffraction spikes and an airy disc at the south; (e) IRAS 19234 + 1627: dead pixels on the north and southwestern part at the boundary of the source; and (f) W49B: masked bright stars.

(IFO 7) is included in the number statistics in Table 2 but not used for flux statistics.

In total, 65 per cent of identified IFOs are related to star formation (49 per cent YSO- and 16 per cent HII-IFO), and 22 per cent are associated with evolved objects with 12 per cent of IFOs remaining as of unknown origin. Among them, YSO-IFO is the most frequent population showing [Fe II] emission. However, they account for only 1.6 per cent of the total [Fe II] flux. On the contrary, SNR-IFOs contribute 76 per cent of the total [Fe II] flux, though represent only 12 per cent of the IFOs by number. On average, the SNR-IFOs are 191 times brighter than the YSO-IFOs. The total flux of PN and

LBV-type IFOs is similar, contributing 1 per cent of the total [Fe II] flux, albeit the number of PN-IFOs is four times larger. In order to understand the surface brightness of each type, the size and structure of the [Fe II] sources should be taken into account. In the next section, we will compare each counterpart’s characteristics in more detail.

3.2 Flux and size distribution

In Fig. 2(a), we present the flux distribution of the IFOs. The flux distributions of the individual IFO types are shown in different colours. As mentioned above, some of the IFOs share the same

Table 2. Statistics of IFOs.

	<i>N</i>	Flux _{total}	Flux _{min}	Flux _{max}	Flux _{mean}	$F_{\text{mean}}^{\text{sb}}$
YSO	100	4.3 (−15)	2.0 (−18)	8.2 (−16)	4.3 (−17)	6.6 (−19)
CHII	22	5.2 (−15)	3.2 (−18)	1.5 (−15)	2.4 (−16)	4.0 (−19)
HII	11	4.8 (−14)	8.9 (−17)	1.6 (−14)	4.3 (−15)	0.7 (−19)
PN	17	1.4 (−15)	2.4 (−18)	4.9 (−16)	8.5 (−17)	7.6 (−19)
SNR	25	2.0 (−13)	3.9 (−18)	4.7 (−14)	8.2 (−15)	1.3 (−19)
LBV	4	1.3 (−15)	1.3 (−17)	8.1 (−16)	3.3 (−16)	9.7 (−19)
Unknown	25	2.4 (−15)	1.3 (−18)	2.2 (−15)	9.7 (−17)	5.0 (−19)
Total/mean	204	2.6 (−13)	1.6 (−17)	4.7 (−14)	1.3 (−15)	5.3 (−19)

Notes. *N*: number of IFOs in each type. Flux units are in W m^{-2} . $F_{\text{mean}}^{\text{sb}}$: mean surface brightness of each type (flux divided by area) in $\text{W m}^{-2} \text{arcsec}^{-2}$. Note that one SNR-type (IFO 7) was not used for statistics of fluxes.

exciting/driving source (e.g. 8 of 25 SNR-IFOs and 6 of 17 PN-IFOs). Bearing this in mind, we see that SNR-IFOs and HII-IFOs are bright with F_{tot} as large as $\sim 10^{-14} \text{ W m}^{-2}$, while YSO, PN, and LBV-IFOs are much fainter, with a two-orders of magnitude smaller maximum F_{tot} . YSO and PN-IFOs appear in a similar flux range, but the majority of PN-IFOs are brighter than YSO-IFOs. The unknown-IFOs are generally much fainter than the other types of IFOs.

Fig. 2(b) shows the semimajor axis distribution of IFOs. IFOs appear in a wide range of sizes, from very compact, arcsecond-scale knots to large-scale objects up to ~ 47 arcmin in size. The distribution with respect to types is similar to that of the flux distribution, for example, SNR-IFOs and HII-IFOs are large and bright, while YSO-IFOs and PN-IFOs are small and faint. The radius range (< 10 arcsec) of unknown-IFOs is similar to that of YSO-IFOs except for a few outliers. Although there are some exceptions and scatter, the overall fluxes and sizes seem to be proportional to each other. Especially for HII-IFO, the correlation coefficient of flux and size is 0.87. When divided into CHII and HII region subtypes, it is 0.52 and 0.83, respectively. The correlation coefficient of unknown IFOs is 0.99. In contrast, the coefficient for SNR-IFOs is only 0.39.

Fig. 2(c) presents the surface brightness distribution of IFOs. Unlike the flux and size distributions, the surface brightness distribution of each type shows slightly stratified distributions. Small IFOs appear to have a higher surface brightness in general, that is, YSO-IFOs, PN-IFOs, and unknown-IFOs have higher surface brightness than HII-IFOs and SNR-IFOs. The reason for this might be due to the low surface filling factor of [Fe II]-emitting regions in the latter sources. For example, the IFO with the lowest surface brightness is SNR-IFO 117 (Kes 78). This SNR has a large size and the [Fe II] emission is patchy, apparent only around the northern and southern caps with a marginally detectable limb. For such sources, the true surface brightness of the [Fe II]-emitting regions could be much greater. In Table 1, we made a note for IFOs with small surface filling factors.

3.3 Spatial distribution

Fig. 3 shows the distribution of IFOs in Galactic longitude and latitude. One can notice the Galactic longitude distribution is clustered albeit the sky coverage is more or less homogeneous. The most outstanding overdensities are seen at $l \sim 16^\circ$ and $\sim 51^\circ$. At $l \sim 16^\circ$, the dominant populations are YSO- and HII-IFOs, while at $l \sim 51^\circ$, they are unknown- and YSO-IFOs. Including other clustered IFOs in longitude, the dominant populations responsible for these peak distributions are YSO-IFOs, followed by HII- and unknown-IFOs. A detailed description of the individual peak regions will be presented

later in this section. Note that there are also voids free of IFOs at $40^\circ \lesssim l \lesssim 50^\circ$.

The distribution of the whole population of IFOs in Galactic latitude shows a Gaussian-like distribution. The distribution yields an average latitude at $b = -0^\circ 12$ and standard deviation $\sigma = 0^\circ 65$. Some concentrations of YSO-IFOs are found at $b \sim -0^\circ 7$, 0° , and $0^\circ 8$. The average latitude of YSO-IFOs is $-0^\circ 08 \pm 0^\circ 67$. The centroids of the HII- and SNR-IFO distributions are also below the GP with an average latitude of $b = -0^\circ 09 \pm 0^\circ 63$ and $-0^\circ 27 \pm 0^\circ 58$, respectively. The average latitude of unknown-IFOs is also less than zero, that is, $b = -0^\circ 25 \pm 0^\circ 73$. For comparison, the average latitude of PN-IFOs is $b = 0^\circ 05 \pm 0^\circ 57$. A similar trend has been observed in the UWISH2 survey; the average latitude of the jets and photodissociation regions (PDRs) was $-0^\circ 18 \pm 0^\circ 01$ and $-0^\circ 17 \pm 0^\circ 01$, while that of the PN group was $-0^\circ 01 \pm 0^\circ 01$ toward the Galactic mid-plane (Froeblich et al. 2015). The distribution of IFOs (excluding PN-IFOs) being slightly shifted to the negative latitude might be related to the ‘bone’ structure in the first Galactic quadrant. The bone structure refers to highly elongated, dense giant molecular filaments that are the most probable tracer of spiral arm structure (Zucker, Battersby & Goodman 2018). It is also worth noting the scarcity of IFOs at $0^\circ 9 < b < 1^\circ 5$. The number of LBV-IFOs is too small for their distribution to have any statistical meaning.

Fig. 4 shows the two-dimensional distribution of IFOs in Galactic longitude and latitude along with their flux distributions. Several IFOs in the same system (e.g. jet and counter-jet of an HH object) are shown as concentric circles, as in many cases they are only a few arcseconds away from each other. On average, all populations show clustered distributions with some differences from each other, though the survey coverage is homogeneous. As well as the inhomogeneous distribution of IFOs, all populations except LBV- and unknown-IFOs have more sources toward the Galactic centre ($l \lesssim 30^\circ$). About half of unknown-IFOs are located close to those of YSOs. In addition to the similar physical properties of YSO- and unknown-IFO shown in Fig. 2, we suggest that at least half of the unknown-IFOs might originate from activities involved in YSOs.

The region relatively devoid of IFOs in the one-dimensional longitude and latitude distribution (Fig. 3) turned out to form a large-scale two-dimensional region; IFOs hardly exist toward $l \gtrsim 30^\circ$, $b \gtrsim 0^\circ 9$, and $35^\circ \lesssim l \lesssim 50^\circ$ near the Galactic mid-plane. This might reflect spiral arm structures and the sightline toward them, where we are seeing a shorter sightline toward the Galactic bar at $l \lesssim 30^\circ$. Above this Galactic longitude we are seeing the local arm branching from Perseus Arm and Sagittarius-Scutum Arm (line of sight tangential to $l \sim 45^\circ$) at a greater distance.




Figure 2. (a) F_{tot} distribution of IFOs. Note that the flux of a large-scale IFO 7 is excluded in this figure. (b) Semimajor axis r_1 distribution of IFOs. The semimajor axis of automatically identified IFOs is the best estimate of the coordinate, semimajor, and minor axes from the best-fitting ellipse from IDL procedure 2dgaussfit. (c) Surface brightness distribution of IFOs. IFOs are shown in accordance with their counterparts: YSO, H II region, PN, SNR, LBV, and unknown-IFOs.

We have identified some areas rich in IFOs (upper panel of Fig. 4), where in particular YSO and HII-IFO are ma-

ior causes of overdensity. The respective regions are as follows.




Figure 3. The spatial distribution of IFOs in Galactic longitude and latitude. The top panel shows the distribution of IFOs in Galactic longitude. The bottom panel shows the distribution of IFOs in Galactic latitude. IFOs are shown in accordance with their counterparts: YSO, H II region, PN, SNR, LBV, and unknown-IFOs.

(i) $l \sim 10^{\circ}2$, $b \sim -0^{\circ}3$: this region is coincident with the H II region G10.2 – 0.3, one of the three H II regions in the giant star-forming region W 31. The H II region is known to be very young (~ 0.6 Myr). At least four O stars are residing in it, where the brightest star W 31–1 showed permitted Fe II at $1.6878 \mu\text{m}$ and brackett lines in the NIR spectrum. In the H - and K -band spectra ($\lambda/\Delta\lambda \approx 3000$) the [Fe II] $1.644 \mu\text{m}$ emission line was not detected (Blum, Daminieli & Conti 2001).

(ii) $l \sim 12^{\circ}8$, $b \sim -0^{\circ}2$: this region matches with [MDF2011b] c11, which encircles the O4-6 (super-)giant no. 23 (Messineo et al. 2015). This region is immediately east of the embedded protocluster W 33 Main which is located inside the massive star-forming complex W 33. The K_s -band spectroscopy of no. 23 showed that the extinction of the region is $A_K = 1.20 \pm 0.03$ mag and the luminosity class is III-I. The Oe star no. 22 is located between W 33 Main and no. 23, with line identifications of Fe II $2.0895 \mu\text{m}$ and H_2 , an extinction of $A_K = 2.87 \pm 0.07$ mag.

(iii) $l \sim 15^{\circ}1$, $b \sim -0^{\circ}7$: this overdensity is coincident with one of the most massive star-forming regions, M 17. About a hundred O- and B-type stars are responsible for the emission and the system is quite young (< 1 Myr, Hanson, Howarth & Conti 1997). Bautista &

Pradhan (1998) reported the detection of multiple iron species, including at $1.644 \mu\text{m}$.

(iv) $l \sim 16^{\circ}9$, $b \sim +0^{\circ}8$: multiple compact IFOs are connected to the apex of pillars of creation located in M 16, an active star-forming region. At the tip of the apex, there are protostars in the pillar’s EGGs (‘Evaporating Gaseous Globules’), which are not yet hot enough to emit X-rays. Therefore, the IFOs in M 16 might be tracing some of the youngest protostars.

(v) $l \sim 25^{\circ}4$, $b \sim -0^{\circ}2$: the region corresponds to W 42, an obscured giant H II region. The closest nearby source is [BCD2000] W 42 1, an O5.5 star (Blum, Conti & Daminieli 2000). There are several point-like sources that might be true [Fe II] sources or mere variables.

(vi) $l \sim 30^{\circ}7$, $b \sim -0^{\circ}0$: the IFO is close to one of the closest starburst regions, W 43. This giant H II region has a central open cluster with massive stars.

(vii) $l \sim 49^{\circ}1$, $b \sim -0^{\circ}6$: multiple HII- and YSO-IFOs are located in the vicinity of W 51, which is one of the most massive giant molecular clouds that is optically obscured. All large-scale representative structures, namely W 51 Main, IRS 1, and IRS 2, are bright in the [Fe II] $1.644 \mu\text{m}$ line. Each structure shows a distinct




Figure 4. Two-dimensional distribution of IFOs. The top panel shows the spatial distribution of IFOs having counterparts in the H II region, YSO, and unknown categories. Each circle represents an IFO, and the size of each circle is proportional to its logarithmic F_{tot} (in order of 10^{-17} , 10^{-16} , and 10^{-15} W m^{-2}). Star-forming regions whose positions match those of IFOs in the distribution are labelled. Due to clustered IFOs, many circles overlap. The bottom panel shows the spatial distribution of IFOs with counterparts of PNe, SNRs, and LBVs. Note that the flux of IFO 7 (i.e. SNR G8.7–0.1) is not provided, therefore excluded here.

star-forming phase as follows: W 51 Main – several UCHIIs are located. IRS 1 – evolved H II region with a size of ~ 1 pc. IRS 2 – went through recent star formation, and an $\sim \text{O3}$ star and a massive YSO were found (Barbosa et al. 2008). An LBV-IFO is also coincident with the region, which is a high-mass evolved star (P Cygni supergiant) with evidence for chemical enrichment (Clark et al. 2009).

(viii) $l \sim 53^\circ 2$, $b \sim +0^\circ 0$: multiple YSO-IFOs coincide with an IRDC G53.2, which was formerly catalogued as three IRDCs in the *Midcourse Space Experiment* (MSX) dark cloud (MSXDC) catalogue (Simon et al. 2006). The three IRDCs, viz. MSXDC G053.11+00.05, MSXDC G053.25+00.04, and MSXDC G053.31 + 00.00 harbour hundreds of YSO and YSO candidates, some of them in the vicinity of IFOs.

(ix) $l \sim 59^\circ 4$, $b \sim -0^\circ 2$: the IFOs are located in the central part of SH 2–87, a complex massive star-forming region. The three

submillimetre clumps, SMM 1, SMM 2, and SMM 3 constitute this H II nebula. These three clumps are at separate evolutionary stages (Xue & Wu 2008), and two HII-IFOs were found in the vicinity of the hottest and most massive star-forming clump, SMM 1.

4 DISCUSSION

4.1 Outflows/jets from young stellar objects

Outflows/jets of YSO are composed of ejected and circumstellar swept-up material, and are recognized as an important signpost of recent star-forming activity. This phenomenon plays a key role in conventional disc accretion-outflow theories, the outflow being responsible for the removal of angular momentum and kinetic energy

of accreting material that enables accreting material to overcome the centrifugal force and collapses to form a star (see theories of disc-wind; Pudritz & Norman 1983, X-wind; Shu et al. 1994, and observational studies; Ellerbroek et al. 2013 for reference).

Thanks to the development of IR instruments, previously undiscovered, highly obscured outflows have begun to be found in the NIR. The optical HH objects and their IR counterparts basically refer to the same phenomena, and only the conditions of jet and circumstellar matter differ. So far, molecular emission (e.g. MHO) has drawn attention in the NIR, alongside atomic/ionic lines in the optical, yet less attention has been brought to the [Fe II] lines in the NIR. The [Fe II] 1.644 μm line, the brightest iron line in the $H-K$ band, is reported to unveil a shocked region that is denser and/or more ionized than regions where optical lines are generated (Nisini et al. 2002). In this aspect, previous studies using frequently used molecular tracers, namely SiO, CO, and HCO⁺ in the sub-mm to mm, only revealed secondary outflows, tracing masses of low-density, distant (up to a few pc) outflows. Whereas the [Fe II] 1.644 μm line from the jet is found to extend a few au to parsec-scales in the form of a dense irradiated jet (Reiter et al. 2015).

Most previous [Fe II] outflow studies are confined to certain types of objects or regions: specific star-forming regions (Orion; Takami et al. 2002, Carina; Reiter, Smith & Bally 2016, Shinn et al. 2013) or a certain mass range of YSOs (Caratti o Garatti et al. 2006, 2015). Recently, outflow studies toward external galaxies, namely the Large Magellanic Cloud (LMC) and the Small Magellanic Cloud (SMC), became feasible (Reiter et al. 2019). These studies showed that the [Fe II] emission tends to be observed at the tip of the bipolar outflow and is rather collimated, compared to H₂ and H α which predominantly show the morphology of a ‘wake’ enclosing the [Fe II] emission (Reiter et al. 2015).

We have detected 100 YSO-IFOs (Table 3). Our result provides a large and comprehensive sample for the study of [Fe II] emission associated with YSOs. Fig. 5 shows the example of identified YSO-IFOs, displaying UKIDSS *KHJ*-band RGB images to show how the YSO-IFOs reveal unique structures in comparison to hot dust continuum structures. YSO-IFOs show diverse morphologies, diverse compared to traditionally observed/expected [Fe II] features that are located at the tip of bipolar outflows and/or are highly collimated toward the driving sources (Caratti o Garatti et al. 2006; Reiter et al. 2016).

We classified YSO-IFOs into four morphological categories; bipolar, cometary, knot-like, and amorphous. A representative case of each category would be IFO 13–14, 125, 122, and 4 in Fig. B1, respectively. Bipolar YSO-IFOs are a textbook case of star formation, consistent with the accretion-jet theory with the aid of a magnetic field (Konigl 1982; Shang et al. 2020; Frank 1999). They typically show two lobes located on opposite sides of a central source, but some show two wakes, tips, and collimated bow-shock shapes, distributed laterally from the apparent YSO jet axis. The prototypical bipolar YSO-IFOs are IFO 13 and 14. The [Fe II] 1.644 μm emission with bipolar morphology usually represents either the ‘cap’ of bow shock where an outflow collides with the ambient medium or dense, collimated jets. Cometary YSO-IFOs resemble a comet with a bright head around the driving source and a narrow faint tail-like structure. The prototypical cometary YSO-IFOs are IFO 125 and 131, both having well-defined conical structures. They are located at quite different distances, that is, 4.7 and 1.1 kpc, and the extent of the associated conical structures has very different linear scales, that is, $\sim 45\,000$ au (10 arcsec) and 5000 au (5 arcsec). For the wide-angle tails of cometary morphology,

it is possible that either (1) the jet is bending and/or precessing (Paron, Fariña & Ortega 2016), (2) a cavity structure is revealed (Hsieh, Lai & Belloche 2017), and (3) a multiple systems presence is implied (Fuente et al. 1998). Knot-like YSO-IFOs appear as knots, sometimes located symmetrically from a driving source along a certain axis. The representative knot-like YSO-IFOs, 122 and 123, are showing well-isolated compact features. These knot-like features might imply that the ejection of accreted material in the system is accompanied by sporadic bursts of accretion (Caratti o Garatti et al. 2015). Amorphous YSO-IFOs represent the remaining YSO-IFOs that are diffuse and do not have a definitive structure. The nature of the amorphous YSO is uncertain. The number of YSO-IFOs classified as bipolar, cometary, knot-like, and amorphous is 16, 18, 19, and 47, respectively.

The morphologies of YSO-IFOs are closely related to the nature of YSOs and their mass-loss histories (Caratti o Garatti et al. 2015; Paron et al. 2016). For example, the collimated and continuous jet morphology indicates a continuous ejection of accreting material from the accretion disc system (Reiter et al. 2016, 2017). The overabundance of amorphous morphologies might suggest highly varying environments or multiple systems are affecting the outflow structure. But the morphology of YSO-IFOs might depend on environments as well as foreground extinction, so detailed studies are needed of the individual objects to confirm their nature. Thirteen YSO-IFOs are associated with HH objects (Table 4). Fig. 6 shows a comparison of their [Fe II] and H α images.

YSO-IFOs constitute half the number of our catalogued sources, making YSO the most common IFO in the inner Galaxy. The number density of YSO-IFOs is 0.55 deg⁻². For comparison, the H₂ number density probed by UWISH2, which covered an almost identical area with a comparable surface brightness limit, is 2.15 deg⁻² (Froeblich et al. 2015). The flux density of YSO-IFOs ranges (2 – 820) $\times 10^{-18}$ W m⁻² with a mean of 4.3 $\times 10^{-17}$ W m⁻². This range can be compared with the results of other surveys. Caratti o Garatti et al. (2006) targeted H₂-emitting low-intermediate luminosity Class 0/I YSOs and reported that among 23, 74 per cent were also detected in [Fe II]. For the newly observed nine [Fe II] line jets in the reference, the flux range is (2.8 – 27.0) $\times 10^{-18}$ W m⁻². Caratti o Garatti et al. (2015) observed 18 intermediate to massive YSOs having H₂ and EGO counterparts, and the flux range is (2.5 – 61.9) $\times 10^{-18}$ W m⁻². Note that these fluxes are obtained from spectroscopic studies using a slit of width 1 arcsec. The majority of YSO-IFOs have flux densities comparable to those of previous studies. But a few sources are exceptionally bright. The number of YSO-IFOs brighter than outflows observed in Caratti o Garatti et al. (2015) is 10 per cent of the YSO-IFOs. Since these bright YSO-IFOs do not share certain morphologies and 40 per cent of them have RMS counterparts, they might be preferentially massive YSO outflows, which have simply not yet been identified due to the limited sky coverage of past [Fe II] observations. One possible speculation is that [Fe II] brightness does not strictly scale with driving source brightness or other outflow tracers, based on the target of previous studies, which tend to be bright *IRAS* sources accompanying outflows discovered in other tracers. This illustrates the importance of an unbiased study to correct our understanding of the characteristics of [Fe II] emitters.

The YSO-IFOs and jet-group MHOs of the UWISH2 survey can be compared one-to-one since the UWIFE survey area was fully covered by UWISH2. The spatial distribution of YSO-IFOs in Fig. 4 shows a highly clustered distribution, accompanied by the high-latitude sources in $l \sim 15^\circ\text{--}30^\circ$ and the absence of YSO-IFOs in the Galactic mid-plane at $l \sim 40^\circ\text{--}50^\circ$. This characteristic distribution is also shared in jet-group MHOs (see fig. 8 in

Table 3. IFOs associated with YSO or YSO candidates.

IFO no.	YSO/YSOc name	Morphology	d [kpc]	Reference YSO counterpart	Dist
IFO 003	YSO AGAL G007.333 – 00.016	k	2.96	ro8/u18	u18
IFO 004	YSO candidate ALLWISE J180219.38 – 213351.9	a	–	ro8	–
IFO 010	Class I YSO [RBG2009] G009.86 – 0.04 4	k	$2.36^{+0.78}_{-0.88}$	ro8/r09	r09
IFO 013	Northern lobe of YSO candidate SSTGLMC G009.7612 – 00.9575	b	–	ro8	–
IFO 014	Southern lobe of YSO candidate SSTGLMC G009.7612 – 00.9575	b	–	ro8	–
IFO 016	YSO AGAL G013.779 + 00.609	a	2.90	ro8/u18	u18
IFO 018	YSO candidate 2MASS J18140816 – 1850560	a	–	ro8	–
IFO 019	YSO candidate 2MASS J18140816 – 1850560	a	–	ro8	–
IFO 020	W 33, IRS 3 having an O6.5 star	a	$2.40^{+0.17}_{-0.15}$	b98	i13
IFO 021	W 33, IRS 1 having an O6.5 + an O7.5 or O8 star	a	$2.40^{+0.17}_{-0.15}$	b98	i13
IFO 022	W 33, [MDF2011b] c11 which encircles O6-7 star no. 23	a	$2.40^{+0.17}_{-0.15}$	m15	i13
IFO 023	1. IRAS 18114 – 1825: Class I, 2. J181421.71 – 182459.0: Class I/IIc	c	2.41	ro8/yl2/m16	yl2
IFO 029	YSO AGAL G014.414 – 00.069	c	3.1	u18	u18
IFO 030	IRAS 18144 – 1723, Class I/II binary	k	4.33	c13/v18	v18
IFO 031	1. YSO IRAS 18151 – 1208 2, 2. 2MASS J18175094 – 1208028, Class I/II YSO	a	3.00	m16	m13
IFO 032	1. 2MASS J18175094 – 1208028: Class I/II, 2. ALLWISE J181749.45 – 120751.1	a	3.00	m16	m13
IFO 033	YSO IRAS 18151 – 1208	a	3.00	v10	m13
IFO 034	[PW2010] 236, Class 0/I	a	2.10	ro8/c13/p10	p10
IFO 035	[PW2010] 236, Class 0/I	a	2.10	ro8/c13/p10	p10
IFO 036	In the middle of multiple YSOs in M 16	c	2.14	c13	b99
IFO 037	In the middle of multiple YSOs in M 16	k	2.14	c13	b99
IFO 039	Tip of column 3 of M 16, either T-Tauri star [TSH2002] S-1 or S-2	k	2.14	t02	b99
IFO 040	Edge of column 2 of M 16	k	2.14	t02	b99
IFO 045	Near the edge of M 16 Pillar V, RMS massive YSO G017.0332 + 00.7476A	a	2.14	ro8/c13	b99
IFO 046	Proximity of Class 0/I YSO [PW2010] 378	a	2.10	ro8/p10	p10
IFO 047	Feature connected to Class 0/I YSO [PW2010] 411	c	2.10	c13/p10	p10
IFO 048	In the vicinity of YSOs in M 16	a	2.14	g07	b99
IFO 049	Spatially connected to massive YSO G017.0666 + 0.6826	c	2.14	c13	b99
IFO 051	Compact feature in the crowded region of YSOs in M 17	k	1.60 ± 0.30	si	n01
IFO 052	Ditto	k	1.60 ± 0.30	si	n01
IFO 053	Ditto	k	1.60 ± 0.30	si	n01
IFO 056	Multiple YSO candidates in the northern region of M17, EB (extended bubble)	c	1.98	ro8/p09	c16
IFO 057	Ditto	a	1.98	ro8/p09	c16
IFO 059	Southern jet of IFO 060	b	1.85 ± 0.2	ro8/c13	x19
IFO 060	IRAS 18177–1405 aligned with IFO 059, 061, in M 16	b	1.85 ± 0.2	ro8/c13	x19
IFO 061	Northern jet of IFO 060	b	1.85 ± 0.2	ro8/c13	x19
IFO 063	Located at the edge of IRDC HEC G016.93 + 00.24	a	2.40	si	r10
IFO 072	Class I/II YSO IRAS 18229 – 1308	c	3.40	ro8/c13/m16	u22
IFO 074	Massive YSO IRAS 18241 – 1134. [Fe II] 1.64 μ m detected	c	12.60	c13	c13
IFO 077	Class I YSO candidate J182859.53 – 115009.6	a	–	k21	–
IFO 091	Biconical structure coincident with FIR clumps, new PN in Froebrich et al. (2015)	b	3.42	e17	t15
IFO 092	Located in IRDC 24.764 – 0.12. Proto-stellar clumps in the vicinity	k	3.57	si	t15
IFO 094	Coincident with UKIDSS source UGPS J183720.81 – 064158.4. Multiple nearby YSOs	c	–	si	–
IFO 099	North-western jet, aligned with IFO 100 and 101	b	3.50	ro8/k21	t15
IFO 100	Class I YSO candidate, previously reported as AGB candidate	b	3.50	ro8/k21	t15
IFO 101	South-eastern jet, aligned with IFO 99 and 100	b	3.50	ro8/k21	t15
IFO 104	In the middle of ALLWISE J183951.16 – 043113.8 Class III or more evolved YSO	a	–	ro8/m16/j18	–
	and semiregular variable ASASSN-V J183948.07 – 043015.9		2.20		j18
IFO 108	Proximity of pre-main-sequence star candidate GaiaDR2 4 258 232 818 679 065 216	a	2.01	v20	b18
IFO 111	Massive protostellar object [VEN2013] G029.8623 – 0.0437, [Fe II] detection reported	c	6.21	c13/a20	116
IFO 115	Spatially coincident with H ₂ , which is connected to MSX6C G031.8380 – 00.1284,	a	–	ro8/e03	–
	YSO candidate SSTGLMC G031.8361 – 00.1408 in the vicinity				
IFO 122	Shares a similar compact structure with IFO 123, aligned east to west	k	4.80	ro8/k21	u18
IFO 123	Coincident with flat-spectrum YSO candidate SSTGLMC G035.2913 + 00.8076	k	4.80	ro8/k21	u18

Table 3 – *continued*

IFO no.	YSO/YSOc name	Morphology	d [kpc]	Reference YSO counterpart	Dist
IFO 124	Matches to proto-stellar clump 34.93 + 0.338 1	a	2.90	t15	t15
IFO 125	Cometary structure coincident with massive YSO IRAS 18527 + 0301	c	4.70	m96	u18
IFO 126	Ultrawide binary Gaia2 4 280 756 726 686 953 984 is the closest, Class III or more evolved YSO ALLWISE J185522.49 + 030130.3 in 30' distance	a	–	t20/m16	–
IFO 131	Class I/II massive YSO IRAS 18555 + 0056, [Fe II] reported by p16	c	1.10	c13/p16	113
IFO 134	Close to YSO candidate SSTGLMC G035.2868 – 00.9528, proto-stellar clump is coincident	a	2.48	ro8	t15
IFO 136	Flat spectrum YSO SSTGLMC G039.2199+00.8638. G039.2060 + 00.8818 in west	a	–	ro8/k21	–
IFO 141	Compact component matches to massive YSO IRAS 19045 + 0518	c	3.60	c13	c13
IFO 144	Aligned with YSO AGAL G039.708 – 01.237 and IFO 145	b	0.60	u18	u18
IFO 145	Elongated and pointing toward YSO AGAL G039.708 – 01.237 and IFO 144	b	0.60	u18	u18
IFO 146	H α PN candidate, yet aligned with IFO 144, 145, H $_2$ knots	a	0.60	s14	u18
IFO 151	Coincident with Class I and flat-SED YSOs, [TBP2010] L673 10 and 13	a	0.60	t10	u18
IFO 152	Close to HH 1186, 42 arcsec away from [TBP2010] L673 YSO 15 and IFO 151	k	0.60	t10	u18
IFO 154	Class III/photosphere YSO SStOERC G049.1662 + 0.2159 at W 51 (or foreground, see k09)	k	5.40	s17	s17
IFO 163	Surrounding YSOs, for example, Class III/photosphere SStOERC G049.5851 – 0.3814	a	5.40	s17	113
IFO 165	Connected to Class I YSO SStOERC G049.9010 – 00.5922, spectral index of flat (FS) according to k21	c	5.40	s17	113
IFO 174	Emerges from Class I YSO [RML2017] MC2 M105	c	3.09	r17	t15
IFO 175	A southern compact jet of massive YSO IRAS 19236 + 1456	k	3.39	c13	t15
IFO 176	Southern diffuse emission from massive YSO IRAS 19236 + 1456	a	3.39	c13	t15
IFO 177	Diffuse structure in contact with YSO AGAL G052.488 – 00.172	c	1.60	u18	u18
IFO 178	Class I YSO 1 ($\sim 10 M_{\odot}$) or 2 ($\sim 5 M_{\odot}$) in k18	a	1.60	ro8/c13/k18	u18
IFO 179	In the proximity of Class I YSO SSTGLMC G053.1570 + 00.0735	a	1.60	ro8/k15	u18
IFO 180	Close to Class I YSO SSTGLMC G053.1612 + 00.0668 and multiple YSOs	a	1.60	ro8/k15	u18
IFO 181	Surrounding Class I YSO SSTGLMC G053.1266 + 00.0499	k	1.60	ro8/k15	u18
IFO 183	Coincident with MSXDC G053.25+00.04 MM6 and ISOGAL-P J192931.1 + 175954 (Class I)	a	1.60	k15	u18
IFO 184	Flat-spectrum YSO 2MASS J19293167 + 1800581	k	1.60	ro8/k15	u18
IFO 185	SSTGLMC G053.2389+00.0552 (Class I), 2MASS J19293167 + 1800581 (FS)	a	1.60	ro8/k15	u18
IFO 186	2MASS J19300219 + 1755001 (FS)	c	–	k15	–
IFO 189	Multiple compact structures surrounding ES-NW of Massive YSO MSX6C G056.3694 – 00.6333	a	6.40	c13	c13
IFO 190	Diffuse structure on South of Class II YSO SSTGLMC G058.4098 + 00.3279	a	2.80	ro8/k21	v13
IFO 192	The head of the cometary structure matches EGO G058.09 – 0.34, one of the low-mass EGOs	k	0.74	ro8/cy13	cy13
IFO 193	Two biconical structures, tails toward SE and W, Class I YSO SSTGLMC G058.4801 – 00.2205 at the centre	b	6.15	ro8/k21	m21
IFO 194	The head of cometary structure corresponds to YSO candidate SSTGLMC G059.0069 – 00.2481	c	–	ro8	–
IFO 198	Aligned with star-forming region IRAS 19410 + 2336 and IFO 199	a	2.20	c13	113
IFO 199	Amorphous IFO points toward star-forming region IRAS 19410 + 2336	a	2.20	c13	113
IFO 203	Compact IFO at the east of biconical outflow S87, emerging from $\sim 20 M_{\odot}$ pre-main-sequence object	a	2.20	b89	113

*Notes.**Column 3: morphology categories: b – bipolar, c – cometary, k – knot-like, and a – amorphous. Column 4: distance of counterpart in kpc. Column 5: references of counterpart classification and distance.

*References: a20 – Areal et al. (2020), b18 – Bailer-Jones et al. (2018), b89 – Barsony (1989), b98 – Beck, Kelly & Lacy (1998), b99 – Belikov et al. (1999), c13 – Cooper et al. (2013), c16 – Csengeri et al. (2016), cy13 – Cyganowski et al. (2013), e03 – Egan et al. (2003), e17 – Elia et al. (2017), g07 – Guarcello et al. (2007), i13 – Immer et al. (2013), j18 – Jayasinghe et al. (2018), k09 – Kang et al. (2009), k15 – Kim et al. (2015), k18 – Kim et al. (2018), k21 – Kuhn et al. (2021), l13 – Lumsden et al. (2013), l16 – Li et al. (2016), m13 – Sánchez-Monge et al. (2013), m15 – Messineo et al. (2015), m16 – Marton et al. (2016), m21 – Mège et al. (2021), m96 – Molinari et al. (1996), n01 – Nielbock et al. (2001), p09 – Povich et al. (2009), p10 – Povich & Whitney (2010), p16 – Paron et al. (2016), r09 – Ragan et al. (2009), r10 – Rygl et al. (2010), r17 – Retes-Romero et al. (2017), ro8 – Robitaille et al. (2008), s14 – Sabin et al. (2014), si – SIMBAD, s17 – Saral et al. (2017), t02 – Thompson, Smith & Hester (2002), t10 – Tsitili et al. (2010), t15 – Traficante et al. (2015), t20 – Tian et al. (2020), u18 – Urquhart et al. (2018), u22 – Urquhart et al. (2022), v13 – Veneziani et al. (2013), v18 – Varricatt et al. (2018), v20 – Vioque et al. (2020), x19 – Xu et al. (2019), yu12 – Yuan et al. (2012)




Figure 5. IFOs with YSO counterpart candidates in continuum-subtracted [Fe II] images as in Fig. 1. Only six representative IFOs are shown. The crosses denote adjacent YSOs in the field of view, while the contours are H_2 2.12 μm emission contours adopted from UWISH2. The right frames are three-colour *KHJ* UKIDSS images of the same field of view. This figure is available in its entirety in Appendix B1.

Froeblich et al. 2015). As seen in Fig. 5, about 85 per cent of YSO-IFOs accompany jet/PDR-group MHOs in the vicinity. For example, in the M 16 (Eagle nebula), 6 YSO-IFOs were identified, and a few hundreds of jet/PDR-group MHOs are also present. A detailed comparison of YSO-IFOs with jet/PDR-group MHOs discovered in the subsequent UWISH2 studies will be helpful for the comparison of different shock tracers (Ioannidis & Froeblich 2012a, b; Froeblich & Makin 2016; Makin & Froeblich 2018; Samal et al. 2018).

We can compare our results with the results of the RMS survey where NIR spectra of YSO candidates have been obtained. In the common survey area ($10^\circ \lesssim l \lesssim 62^\circ$, $|b| \lesssim 1.5^\circ$), there are 182 RMS sources, and among the 72 sources from which spectra have been obtained, 58 have [Fe II] line emission, though some of the detections could be confused with the Br 12 line. For comparison, only 17 of 182 RMS objects have been identified as YSO-IFO in our study (for some RMS sources, 2–3 IFOs correspond to one RMS source.) Among these 17 sources, the NIR spectra have been obtained for eight sources, and [Fe II] lines were detected in six sources, that is, [Fe II] lines were reported as non-detection for two sources in the RMS survey. We note that the non-detection for the two is based on a comparison of Br 11 and Br 12/[Fe II] line strengths ($\text{Br } 11 \times 0.788 > \text{Br } 12/[\text{Fe II}]$) and it might be possible that a weak [Fe II] line is in fact present but missed by low spectral resolution, as the authors noted (Cooper et al. 2013). To assess this possibility, we checked the slit configuration (central position and PA) in Cooper et al. (2013)

and compared it with YSO-IFO morphology. For both IFO 72 and 141, the RMS slit intersects the driving source but does not include the bright part of extended YSO-IFO structures. Indeed, the authors tried to include extended structures inside the slit in imaging mode prior to spectroscopy mode, yet even narrow-band [Fe II] images of UWIFE without continuum subtraction turned out to severely hinder extended emission. Therefore, most YSO-IFOs apparently do not have RMS source counterparts, which is claimed to be a 90 per cent complete list of massive protostellar populations (Lumsden et al. 2013). This seems to suggest that most of YSO-IFOs are associated with low-mass star formation. It is also worthwhile to note that the majority of YSO-IFOs (87 per cent) are not associated with HHS, which suggests that the [Fe II] emission is tracing optically hidden star-forming regions.

4.2 Compact H II regions

CHII and UCHII regions are the earlier stages of ‘classical’ H II regions. An UCHII region is a photoionized region with a diameter $\lesssim 0.1$ pc and an electron density $n_e \gtrsim 10^4 \text{ cm}^{-3}$, embedded in a molecular cloud (Wood & Churchwell 1989). In this evolutionary stage, mass accretion of the central star is thought to be insignificant (Churchwell 2002; Zinnecker & Yorke 2007). A CHII region is an H II region in the intermediate phase between UCHII and classical H II regions, having a radius $\lesssim 0.1$ pc and $n_e \gtrsim 10^3 \text{ cm}^{-3}$. The lifetime

Table 4. IFOs associated with HH objects.

IFO no.	HH name	d [kpc]	Exciting source	Region	Comment	Morp.	Reference Type	Dist
IFO 041	HH 216	2.14	HH-N	Eagle nebula	Parsec-scale HH	b	m82	b99
IFO 042					Parsec-scale HH	b	m82	b99
IFO 043					Parsec-scale HH	b	m82	b99
IFO 044					Parsec-scale HH	b	m82	b99
IFO 118	HH 722				Wrongly identified as HH 172 (Nikogossian, Magakian & Movsessian 2007)	a	c94	d92
IFO 119	GGD 30	1.70	GGD30IR	GM 2–30	Binary. One of them is Class I YSO, $A_V \sim 25$ mag, bipolar envelope SE-NW side	a	s07	s87
IFO 120					Binary. One of them is Class I YSO, $A_V \sim 25$ mag, bipolar envelope SE-NW side	a	s07	s87
IFO 121					Binary. One of them is Class I YSO, $A_V \sim 25$ mag, bipolar envelope SE-NW side	a	s07	s87
IFO 153	HH 32	0.20 ± 0.03	AS353	Aquila Rift	T-Tauri binary, H α is coincident, [Fe II] detection reported (d03)	a	h74	r06
IFO 155	HH 250A	0.30	HH 250-IRS	Aquila Rift	Class I binary, bow-shock, H α detected. Launched 3500 yr ago, adjacent to IFO 153	a	d97	s97
IFO 195	HH 803	2.40	1548C27 IRS1		SW of parsec-scale HH (7.5 pc), H α detected. $L_{\text{bol}} = 580 L_{\odot}$	k	c04	d92
IFO 196	HH 165				NE counterpart of the HH 803, very faint	a	c81	d92
IFO 200	HH 365				Central structure of parsec-scale HH, both perpendicular jet and curvature are identified	a	a97	d92

*Notes.** Column 3: distance of counterpart in kpc. Column 7: morphology categories: b – bipolar, k – cometary, k – knot-like, and a – amorphous; Column 8. References of counterpart classification and distance: a97 – Alten et al. (1997), b99 – Belikov et al. (1999), c04 – McGroarty, Ray & Bally (2004), c81 – Craine, Boeshaar & Byard (1981), c94 – Cappellaro et al. (1994), d03 – Davis et al. (2003), d92 – Dent & Aspin (1992), d97 – Devine, Reipurth & Bally (1997), h74 – Herbig (1974), m82 – Meaburn & White (1982), r06 – Rice, Prato & McLean (2006), s07 – Smith, Lawson & Wright (2007), s87 – Scoville et al. (1987), and s97 – Sakamoto et al. (1997).

of UCHII and CHII is $\sim 2\text{--}4 \times 10^5$ yr (Davies et al. 2011, Mottram et al. 2011).

In UCHII and CHII regions, [Fe II] emission can be enhanced by the interaction of stellar wind with the ambient medium. Bloomer et al. (1998) detected enhanced shell-like [Fe II] emission along the periphery of the CHII region NGC 7538 IRS 2. The observed [Fe II] 1.644 $\mu\text{m}/\text{Br } \gamma$ ratio was 0.15, which is an order of magnitude greater than that of H II regions, and it implies that the [Fe II] line emission emanates from shocked stellar wind material. Shinn et al. (2014) searched for [Fe II] 1.644 μm emission associated with UCHII regions employing the CORNISH UCHII catalogue and the UWIFE survey data. Among the 237 UCHII regions in the survey area, five and one candidate were found to have associated [Fe II] emission features, which were suggested to be shock-excited by outflows from central YSOs. Kim, Lacy & Jaffe (2017) also reported the detection of [Fe II] emission from UCHII Monoceros R2. Hereafter, we refer to IFOs associated with CHII/UCHII regions or with H II regions in even earlier evolutionary stages as CHII-IFOs.

We have detected 22 IFOs associated with 16 UCHII/CHII regions (Table 5). Six IFOs (IFO 24, 25, 26, 97, 107, and 156) had been previously reported by Shinn et al. (2014). We have discovered IFOs associated with an UCHII precursor (IFO 137) and an UCEC (IFO 138), which are thought to be earlier progenitor or less massive populations (Molinari et al. 1998; Alexander & Kobulnicky 2012). Among the 16 UCHII/CHII regions with [Fe II] emission features, 10 are catalogued in CORNISH, which corresponds to 4 per cent of the 237 UCHII regions in the CORNISH catalogue in the survey area. The detectability might be partly due to the large extinction in UCHII/CHII regions, which is typically $A_V \sim 30\text{--}50$ or $A_K \sim 3\text{--}5$ (Hanson, Luhman & Rieke 2002). Indeed, the A_V of three UCHII regions with associated IFOs had been found to have relatively low extinction ($A_V \sim 9\text{--}20$, Shinn et al. 2014).

Fig. 7 shows the 22 CHII-IFOs. CHII-IFOs have diverse morphologies, for example, jet-like, shell-like, and amorphous morphologies. A representative IFO with jet morphology is IFO 97, which appears as a collimated beam from the centre to the boundary of the H II region. The jet appears to extend beyond the radio continuum boundary (see Fig. 7), which might reflect a possible correlation with the boundary of the ionization front (Goddi et al. 2020). The representative IFO with a shell-like morphology is IFO 132. An exemplary CHII-IFO of amorphous morphology would be the IFO 138, having a diffuse structure either outside or inside of the H II region in the radio. The properties of a CHII region have been rarely investigated in [Fe II] emission. Shinn et al. (2014) proposed that some IFOs identified in the vicinity of UCHII regions (IFOs 24–26, 97, 107, and 156) are the ‘footprint’ outflow features of UCHII regions, that is, the features produced by outflowing material ejected during an earlier, active accretion phase of massive YSOs, based on the morphological relation between the [Fe II] and 5 GHz radio features, the outflow mass-loss rate, the traveltime of the [Fe II] features, and the existence of several YSO candidates near the UCHII regions. The newly discovered CHII-IFOs in this study might serve as a chance to investigate the origin of the [Fe II] emission in the vicinity of CHII regions.

4.3 H II regions

H II regions are not expected to be bright in the [Fe II] lines, since in their photoionized regions, Fe atoms are predominantly in higher ionization states, and Fe atoms are thought to be mostly locked in dust grains (Koo et al. 2016). According to theoretical models of photoionized regions, the [Fe II] emission from an H II region




Figure 6. IFOs with HH counterpart candidates. The left panels show continuum-subtracted [Fe II] images as in Fig. 1. The right panels show $H\alpha$ in the same field of view. IPHAS images were used except for IFO 41–44 where the *Hubble Space Telescope* F657N image was used. Inset on the IFO 119–121 is a magnified [Fe II]- H image of the saturated star (west of IFO 120 and north in the inset) and bright part of IFO 119 (south in the inset). Contours are H_2 2.12 μm emission adopted from UWISH2. The arrow points to the driving source of the HH object. When the driving source is out of the image field of view, the arrow points from the driving source to the IFO.

is mainly emitted in the high-density partially ionized zones near ionization fronts, predominantly excited by electron collisions (Oliva et al. 1989; Bautista & Pradhan 1998). In the Orion nebula, for example, [Fe II] images exhibit filamentary structures and diffuse emission that might be associated with ionization fronts, together with some knotty features (Takami et al. 2002). Expanding H II regions can drive shocks, but the shock velocity is low ($\sim 10 \text{ km s}^{-1}$), so [Fe II] line emission is not expected to be enhanced (e.g. Mouri et al. 2000). The [Fe II] 1.644 μm /Pa α ratio of the Orion is 0.013, which is more than two orders of magnitude smaller than those of SNRs (Oliva et al. 1989; Mouri et al. 2000). So Galactic H II regions have not been a popular target of deep and high-resolution [Fe II] imaging (Kraus et al. 2006; Bally et al. 2022). The depletion of Fe atoms in the H II region, however, is uncertain. In the Orion nebula, it has been estimated that 90 per cent of Fe is locked onto dust grains (Baldwin et al. 1991, 1996; Osterbrock, Tran & Veilleux 1992; Rodríguez 2002). But there are studies which showed that, in many H II and star-

forming regions, Fe is not depleted as heavily as in the Orion nebula (Osterbrock et al. 1992; Peimbert 1993; Rodríguez 2002; Okada et al. 2008; Peimbert & Peimbert 2010). It has been suggested that some populations of dust grains might be easily destroyed by UV radiation from OB stars and Fe atoms are released into the gas phase (Okada et al. 2008; Peißker et al. 2020). For external galaxies, Alonso-Herrero et al. (2003) did an imaging study of the starburst galaxies M 82 and NGC 253 in [Fe II] 1.644 μm and Pa α (1.87 μm) lines, and, by comparing their intensity ratios, concluded that 6 per cent–8 per cent of [Fe II] line fluxes are due to H II regions. Mouri et al. (2000), Riffel et al. (2016), Hennig et al. (2018), and Fazeli et al. (2019) suggested that some of the [Fe II] emission from external galaxies could be due to H II regions based on their low [Fe II] 1.257 μm /Pa β ratios.

We have identified 11 IFOs associated with 4 H II regions (Table 6).

All HII-IFOs are located in the well-known star-forming complexes W 31, M 17, and W 51. Fig. 8 shows the 11 HII-IFOs. We can

Table 5. IFOs associated with UCHII/CHII or UCHII/CHII candidates.

IFO no.	UCHII, CHII name	d [kpc]	Type	Comment	Type	Reference
IFO 024	IRAS 18116 – 1646	4.50	UCHII	Outer arc, east of cometary H II region	k18	u18
IFO 025	IRAS 18116 – 1646	4.50	UCHII	Diffuse filament filling inside, coincident with the boundary of cometary H II region	k18	u18
IFO 026	IRAS 18116 – 1646	4.50	UCHII	Outer arc, west of cometary H II region	k18	u18
IFO 054	M 17 UC1	1.60 ± 0.30	HCHII	Multiple shell-like features around hypercompact H II region	s04	m01
IFO 071	IRAS 18222 – 1317	4.04	UCHII	Partial shell-like + amorphous structure around UCHII	k18	114
IFO 082	IRAS 18308 – 0503	2.90	UCHII	Amorphous, compact feature north of UCHII	b96	113
IFO 083	IRAS 18308 – 0503	2.90	UCHII	Amorphous, diffuse feature south of UCHII	b96	113
IFO 086	IRAS 18314 – 0720	9.30 ± 0.40	UCHII	Shell-like structures, west of UCHII	k18	k03
IFO 087	IRAS 18314 – 0720	9.30 ± 0.40	UCHII	Multiple shell-like structures encircling UCHII	k18	k03
IFO 089	IRAS 18317 – 0757	4.60	UCHII	Amorphous structures enveloping the south-east of UCHII	k18	113
IFO 090	IRAS 18317 – 0757	4.60	UCHII	Compact knot-like structure north-east of UCHII	k18	113
IFO 097	G025.3809 – 00.1815	3.80	UCHII	Amorphous structures in the central region of UCHII + south-western jet	k18	a09
IFO 107	IRAS 18416 – 0420	3.30	UCHII	Shell-like structure stretching from the centre of UCHII to west	k18	u18
IFO 127	IRAS 18530 + 0215	9.00	UCHII	Amorphous + knot-like feature south of UCHII	k18	113
IFO 132	G35.2S	2.40 ± 0.50	CHI	Oval structure, slightly limb-brightened, brighter at east. CHII region (radius: 0.14 pc)	f11	z13
IFO 137	[CPA2006]N75	2.80	UCHII?	Northern shell-like structure. Single O9.5V star. UCHII precursors, Mol 91C1–5, along the structure	m98	z13
IFO 138	UCEC 8	2.80	UCHII?	UCEC with a radius of 0.09 pc.	a12	113
				Eight possible ionizing stars, most of them are early-B.		
IFO 148	IRAS 19111 + 1048	4.40	UCHII	About 19 central sources of spectral type earlier than B0.5 (v06)	k18	113
IFO 149	IRAS 19111 + 1048	4.40	UCHII	Diffuse emission north of IFO 148	k18	113
IFO 156	IRAS 19191 + 1538	2.10	UCHII	Partial shell-like structure in the west, compact structure superposed in the north	k18	113
IFO 202	IRAS 19442 + 2427	2.20	UCHII	Diffuse emission south of cometary UCHII, the spectral type of O9-B0 (d20)	k18	113
IFO 204	IRAS 19446 + 2505	2.20	UCHII/HII	CORNISH UCHII G061.4763+00.0892 in east, H II region G061.4758 + 00.0913 in west	k18	113

*Notes.** Column 3: distance of counterpart in kpc. Column 6: references of counterpart classification and distance: a09 – Anderson & Bania (2009), a12 – Alexander & Kobulnicky (2012), b96 – Bronfman et al. (1996), d20 – de la Fuente et al. (2020), f11 – Froebrich & Ioannidis (2011), k03 – Kolpak et al. (2003), k18 – Kalcheva et al. (2018), 113 – Lumsden et al. (2013), 114 – Leahy, Green & Tian (2014), m98 – Molinari et al. (1998), m01 – Nielbock et al. (2001), s04 – Sewilo et al. (2004), u18 – Urquhart et al. (2018), v06 – Vig et al. (2006), and z13 – Zhu et al. (2013)




Figure 7. IFOs with CHII counterpart candidates. The left panels show continuum-subtracted [Fe II] images as in Fig. 1. The right panels show a radio continuum in the same field of view. Contours on both images are the boundaries of CHII in the radio. The contours of IFO 24–26, 54, 71, 86–87, 97, 132, and 137 are from New-GPS 20 cm, IFO 89–90, 107, 127, 138, 148–149, 156, and 202–203 are from CORNISH 5 GHz, and IFO 82–83 are from the National Radio Astronomy Observatory Very Large Array (VLA) Sky Survey. Only the radio image and contour of IFO 204 are from the old-GPS 20 cm. IFO 203 is a YSO-IFO inside the field of view. The yellow crosses in both panels are the same as in Fig. 5. The red cross shows the central position of the UWISH2 H₂ emission.

see that some IFOs appear as thin filaments elongated along the radio structure (e.g. IFO 55 and IFO 62) or as diffuse amorphous emission structures within the radio structure (e.g. IFO 11 and 12), so the association of IFOs with H II regions is very likely. The filamentary structures might correspond to ionization fronts and/or boundaries of PDRs as in the Orion nebula. On the other hand, some IFOs are faint and diffuse, and they extend beyond the radio boundary of the H II regions, for example, IFO 159 and 161, so their association is uncertain and needs to be confirmed.

4.4 Planetary nebulae

PNe represent a short-lived phase near the endpoint of low- to intermediate-mass star (1–8 M_{\odot}) evolution which is preceded by the AGB, post-AGB, and pre-PN phases. The circumstellar envelope of the AGB carbon star is considered highly Fe-depleted (Maun & Huggins 2010), though Fe becomes abundant with time (Fe abundance is negatively correlated with the C/O ratio, Delgado-Inglada & Rodríguez 2014). In turn, PNe are not expected to be strong [Fe II] emitters, also having a Fe-deficit nature with <10 per cent

Table 6. IFOs associated with H II region or H II region candidates.

IFO no.	HII/HIIC name	d [kpc]	Type	Comment	Reference Type	Reference Dist
IFO 011	G10.2 – 0.3 (W 31)	3.55 ^a	HII	Diffuse structure in west of H II region	r07	ul2
IFO 012	G10.2 – 0.3 (W 31)	3.55 ^a	HII	Diffuse structure coincident with O stars/YSOs in the central region. UCHII G10.15 – 0.34 is embedded	r07	ul2
IFO 055	M 17 S	1.60 ± 0.30	HII	The south-western large-scale structure of M 17	p09	n01
IFO 058	M 17 S (south)	1.60 ± 0.30	HII	Diffuse filaments at the south of M 17	p09	n01
IFO 062	M 17 N	1.60 ± 0.30	HII	The north-eastern large-scale structure of M 17	p09	n01
IFO 064	M 17 S (south)	1.60 ± 0.30	HII	Diffuse filaments at the south of M 17	p09	n01
IFO 066	M 17 EB ^b	1.60 ± 0.30	HII?YSO?	Amorphous structure located in the extremity of M 17, star Gaia2 4 097 840 529 185 958 272 in west	p09	n01
IFO 067	[KC97c] G015.1 – 00.9	1.98	HII	The elongated structure along the radio emission of H II region [KC97c] G015.1 – 00.9	k97	r03
IFO 158	W 51A	5.40	HII	Shell-like diffuse emission at the western boundary of compact radio	f21	l13
IFO 159	W 51B	6.00	HII	Large-scale, multiple filamentary structures in W 51B	f21	k95
IFO 161	W 51A	5.50	HII	Large-scale, miscellaneous amorphous structures in W 51A	f21	c09

Notes. ^aIFO distances marked with ‘a’ are controversial values. Column 3: distance of counterpart in kpc. Column 6: references of counterpart classification and distance: c09 – Clark et al. (2009), f21 – Fujita et al. (2021), k95 – Koo, Kim & Seward (1995), k97 – Kuchar & Clark (1997), l13 – Lumsden et al. (2013), n01 – Nielbock et al. (2001), p09 – Povich et al. (2009), r03 – Russeil (2003), r07 – Roshi et al. (2017), and ul2 – Urquhart et al. (2012).

^bThe region M 17 EB (extended bubble) is defined in detail by Povich et al. (2009).

existing in gas and the remaining probably enshrouded in dust grains (Delgado-Inglada & Rodríguez 2014). Meanwhile, in the context of environmental factors, PNe could be an [Fe II] emitter, since it has a partially ionized zone where Fe⁺ is apt to exist, and at a certain point of its evolution, a low-velocity shock is expected to occur. In short, suitable ionization conditions and energy to excite Fe (Greenhouse et al. 1991) can be established in PNe, and its iron-depleted nature is a key factor to determine the existence of [Fe II] emission.

Besides the theoretical expectation, previous studies reported the detection of [Fe II] emission towards stellar objects in a variety of evolutionary stages: post-AGB (IRAS 16594 – 4656; Van de Steene & van Hoof 2003), pre-PN (M 1–92; Davis et al. 2005), and PN (Hubble 12; Welch et al. 1999, M 2–9; Smith et al. 2005, NGC 2440; Hora, Latter & Deutsch 1999). Some authors suggested a circumstellar origin (e.g. Smith et al. 2005; Clark et al. 2014), especially Baan, Imai & Orosz (2021) reported the detection of [Fe II] emission revealing the interaction of an accretion inflow, which is composed of material ejected in earlier post-AGB and pre-PN circumstellar material, and stellar outflow.

Table 7 shows PN-IFOs. They are IFOs spatially coincident with PNe, PN candidates, and sources in earlier evolutionary stages such as post-AGBs. Seventeen PN-IFOs are associated with 14 PNe; 5 PNe, 8 PN candidates, and one post-AGB candidate. For comparison, in a previous study, Lee et al. (2014) reported the detection of [Fe II] emission in six PNe among 29 known PNe. In the survey area, there are 296 HASH ‘true’ (131), likely (40), and possible (125) PNe, so that the detection rate is 4.7 per cent. If we limit the sample to the ‘true’ PN, the detection rate slightly drops to 3.8 per cent (i.e. 5 out of 131). This very low detection rate of PNe in [Fe II] emission (4.7 per cent and 3.8 per cent) contrasts with the results in H₂, where detection rates are 30 and 21 per cent, respectively (for 10° < l < 66°, $|b|$ < 1.5°, Gledhill et al. 2018). It is interesting that even with an order of magnitude larger sample of PNe in this study, our result is somewhat consistent with the former [Fe II] and H₂ detection rates of 7 per cent and 39 per cent derived from 41 PNe (Hora et al. 1999). The slightly higher detection rate of Hora et al. (1999) could be because their samples are either moderately sized or optically bright.

The number density of PN-IFOs is 0.07 deg^{−2} within 180 deg² whereas it is 1.25 deg^{−2} within 209 deg² in UWISH2 (Froebrich et al. 2015). However, unlike the previous argument (Kastner et al. 1996), not all [Fe II]-emitting PNe are seen in H₂ emission; we found 3 of our 14 PNe in Table 7 were absent from the list of PNe with H₂ emission. Also, the median flux of PN-IFOs is greater than that of the H₂-emitting PNe, that is, 6.46 × 10^{−17} versus 4.53 × 10^{−17} W m^{−2}. Therefore, our result shows that the H₂-emitting PNe are not necessarily brighter than the non-H₂-emitting PNe in [Fe II] emission.

Fig. 9 presents [Fe II]-H images of PN-IFOs. Note that there are three bipolar PNe, each of which possesses two associated IFOs (IFO 5 and IFO 6, IFO 8 and IFO 9, and IFO 129 and IFO 130). We classified the morphologies of PN-IFOs using the basic ‘ERBIAS’ classifier following Parker et al. (2006), where ‘E’ = elliptical, ‘R’ = round, ‘B’ = bipolar, ‘I’ = irregular, ‘A’ = asymmetric, and ‘S’ = quasi-stellar. Their subclassifiers ‘amprs’ are also adopted to describe detailed morphology; the main object has a one-sided enhancement/asymmetry ‘a’, has multiple shells or external structure ‘m’, exhibits point symmetry ‘p’, has a well-defined ring structure or annulus ‘r’, or resolved internal structure ‘s’. An IFO can have several ‘amprs’ subclassifications. The results are summarized in Table 7, where their morphologies in H α and H₂ are also listed (Parker et al. 2016;




Figure 8. IFOs with H II region counterpart candidates in continuum-subtracted [Fe II] images as in Fig. 1. Contours are boundaries of H II regions in the radio continuum: IFO 11, 12, 55, and 62 with New-GPS 20 cm data; IFO 58, 64, 66, and 67 with GPS 90 cm data; 158, 159, and 161 with THOR 1420 MHz continuum + VLA GPS (VGPS) H I data. Arrows point to the boundaries of IFO structures. The format for these images is the same as that of Fig. 1.

Table 7. IFOs associated with PN or PN candidates.

IFO no.	PN/PNC name	Type	d [kpc]	Size [pc]	[Fe II]	Morphology H ₂	H α	Distance	Reference Morphology	Type
IFO 005	NGC 6537	PN	2.81 ± 0.45	0.49 ^a	Bp	Bmps	Bmps	3	1,2	–
IFO 006	NGC 6537	PN	2.81 ± 0.45	0.43 ^a	Bp	Bmps	Bmps	3	1,2	–
IFO 008	HRDS G008.362 – 0.623	PNC	–	– ^a	Bas	Bmps	–	–	1	4
IFO 009	HRDS G008.362 – 0.623	PNC	–	– ^a	Bas	Bmps	–	–	1	4
IFO 027	PNG G014.5 + 00.4	PNC	–	–	B	B	S	–	1,2	5
IFO 050	SSTGLMC G015.7993 – 00.0063	PN	–	–	Ia	I?	–	–	6	–
IFO 073	G020.9 + 00.8	PNC	–	–	A	Is	Aa	–	1,2	7
IFO 081	M 1–51	PN	1.59	0.18	Bp ^b	–	B	8	2	–
IFO 095	IRAS 18348 – 0616	PNC/UCHII?	6.5	0.50	Ema	Ema	–	18	6	2019
IFO 112	IRAS 18458 – 0213	PN	4.90	0.13	Bsa	Bs	–	9	1	–
IFO 114	PNG 032.1 + 00.1	PNC/HII?	4.90	0.42	Ia	–	–	9	–	10,2
IFO 129	IRAS 18551 + 0159	PNC	4.30 ± 0.50	0.45 ^a	Br	Bs	B	11	1	4
IFO 130	IRAS 18551 + 0159	PNC	4.30 ± 0.50	0.45 ^a	Br	Bs	B	11	1	4
IFO 157	PNG 050.4 + 00.7	PNC/sym?	5.30	3.08	Bp	Bs	B	9	1	12,2
IFO 164	PNG 050.8 + 00.0	PNC	18.46 ± 2.59	0.26	I	–	–	17	–	17
IFO 173	IRAS 19234 + 1627	PN	4.70 – 9.50	0.31 – 0.64	Ears	Er	Emrs	13	1,2	–
IFO 188	IRAS 19312 + 1950	pAGBc/YSO?	2.42	0.46	Ams	As	–	14	6	1516

*Notes.** For the morphological classification, we adopted an ‘ERBIAS’ classifier (Parker et al. 2006, and references therein), and when listed, we showed the morphological classification of H α counterpart in HASH (Parker et al. 2016). For the morphological classification of the H₂ counterpart and H α classification of IFO 129, 130, and 173, we made use of Gledhill et al. (2018). The ‘ERBIAS’ represents the main structure of PN by its elliptical, round, bipolar, irregular, and asymmetric or quasi-stellar morphology. The substructure is further classified into the subclassifier ‘ampr’, which explains asymmetry, multiple structures, point symmetry, ring, and internal structure. When no counterpart was detected, we marked ‘–’. Column 3: PNC: PN candidate, sym: symbiotic star candidate, and pAGBc: post-AGB candidate. Column 4: distance of counterpart in kpc. Column 5: physical scale of an IFO in the adopted distance. Column 7: references of distance, H₂/H α morphology, counterpart type in Column 3, respectively. Ref. 1 – Gledhill et al. (2018), 2 – Parker et al. (2016), 3 – Navarro, Corradi & Mampaso (2012), 4 – Froebrich et al. (2015), 5 – Miszalski et al. (2008), 6 – this study (UWISH2), 7 – Boissay et al. (2012); Miscellaneous Emission Nebulae (MEN) list, 8 – Phillips (2004), 9 – Lumsden et al. (2013), 10 – Ferrero et al. (2015), 11 – Zhu et al. (2013), 12 – Sabin et al. (2014), 13 – Yang et al. (2016), 14 – Vickers et al. (2015), 15 – Nakashima et al. (2016), 16 – Cordiner et al. (2016), 17 – Irabor et al. (2018), 18 – Cichowolski et al. (2018), 19 – Bronfman et al. (1996), and 20 – Kanarek et al. (2015).

^aPart of bipolar lobes.

^bPN with quadrupolar morphology is included in the bipolar category considering the formation mechanisms of the two could be analogous; see Machado, Stanghellini & Guerrero (1996) for details.




Figure 9. Continuum-subtracted [Fe II] images of IFOs with PN counterpart candidates. Only six representative IFOs are shown. The units on the UWIFE [Fe II]-H are DN, with the darker colour denoting a higher DN. The right frames are three-colour *KHJ* UKIDSS images of the same field of view. The corresponding source names for each IFO are shown. The cross marks the central position of the counterpart. The images of IFO 73, 95, 112, 129, 157, and 164 are smoothed with a two-pixel Gaussian. In all images, north is at the top and east is on the left side. This figure is available in its entirety in Appendix B2.

Gledhill et al. 2018). The $H\alpha$ morphologies are from the HASH survey, while the H_2 morphologies are from the UWISH2 survey. For the PN-IFOs without a counterpart in the UWISH2 PN catalogue (IFO 50, 95, and 188), we inspected the UWISH2 data and classified their morphology in the same format (see Table 7). Some PNe have different morphologies in the [Fe II], H_2 , and $H\alpha$ emission, which implies a complex surrounding environment and/or complex mass-loss history.

The physical sizes of PN-IFOs have been derived for 10 PNe that have previously estimated distances (Table 7). The sizes of 4 IFOs associated with ‘true’ PNe range from 0.13 to 0.92 pc, and three of the PN-IFOs are larger than 0.9 pc. This contrasts with the majority of PNe in $H\alpha$ being ≤ 0.2 pc (González-Santamaría et al. 2020). This seems to suggest that the [Fe II] emission preferentially traces large, bright PNe.

For example, in PNG 050.4 + 00.7, the size of the associated IFO (IFO 157) substantially exceeds the previously known size of the counterparts (2 arcmin and 19 arcsec, respectively).

The IFO has a partial ‘S’ shape elongated along the east–west direction, with IRAS 19194 + 1548 superposed at the western part. The structure becomes gradually fainter toward the west, therefore the angular size of the partial ‘S’ shape should be considered as lower limit. The implied physical scale of 3.1 pc largely surpasses the generally accepted size of PNe (one of the oldest and largest PNe,

the Helix nebula has an outermost size of 1.76 pc). The driving source is suspected to be in a symbiotic star system (Akraş et al. 2019) and the updated size is compatible with the sizes of large shells/nebulæ around symbiotic stars (McCollum et al. 2008).

4.5 Nebulae of luminous blue variables

Infrared [Fe II] 1.644 μm emission around prominent nebulousity of LBVs is thought to be ubiquitous. Smith (2002, henceforth, S02) searched for [Fe II] 1.644 μm emission in nine well-known LBVs and found the emission in 7 of them, resulting in a detection rate of 77 per cent. This high detection rate surpasses that of SNRs (i.e. 24 per cent, Lee et al. 2019), the population that is thought to provide the most adequate environment for the existence of [Fe II] 1.644 μm emission. S02 could not pinpoint the essential condition needed for strong [Fe II] emission to exist. Shock heating and radiative heating as possible excitation mechanism of [Fe II] emission were suggested by the author.

Shock-excited [Fe II] emission can arise when the LBV’s environment meets requirements such as (i) a large difference in the outflow speed between the stellar wind and pre-existing LBV nebula and (ii) a difference of velocities between the stellar wind and ejected shell created during S Doradus outbursts or giant eruption phases. This velocity difference of 50–150 km s^{-1} (S02) is ascribed to weaker gravity

Table 8. IFOs associated with LBV nebula or LBV nebula candidates.

IFO no.	LBV/LBVc name	d [kpc]	Reference	
			Type	d
IFO 065	HD 168 625	1.55	5	4
IFO 102	26.47 + 0.02	≤ 6.5	1	1
IFO 103	26.47 + 0.02	≤ 6.5	1	1
IFO 162	[KW97] 37–17 (= LS1)	$6.0, 2.5^{+2.4}_{-1.3}$	2	3,4

*Notes.**Column 3: distance of counterpart in kpc. Column 4: references of counterpart classification and distance: 1 – Clark et al. (2003): assuming 1.8 mag kpc^{-1} , 2 – Okumura et al. (2000), 3 – Clark et al. (2009): observational and theoretical constraints + W51 membership + parallax, 4 – Bailer-Jones et al. (2018): *Gaia* DR2 parallax, and 5 – Hutsemekers et al. (1994)




Figure 10. Continuum-subtracted [Fe II] images of IFOs having LBV counterparts. The right frames are three-colour *KJH* UKIDSS images of the same field of view. Note that there is a spike pattern around a bright star, coincident with IFO 65. The format for these images is the same as that of Fig. 1.

in an active phase. When LBV evolves toward a cooler temperature (to a local temperature lower than 30 000 K), Hydrogen atoms and opacity-enhancing ions start to emerge on the surface, which is known as the ‘modified’ Eddington limit (Humphreys & Davidson

1994). The elevated opacity makes the outward radiation pressure stronger and overpowers the inward gravity force. The resultant lower effective gravity helps LBVs easily induce the aforementioned mass loss. In these S Doradus outbursts and giant eruption phases, the




Figure 11. IFOs with counterpart candidates unknown in continuum-subtracted [Fe II] images as in Fig. 1. Only six representative IFOs are presented. The right frames are three-colour *KHJ* UKIDSS images of the same field of view. The format for these images is the same as that of Fig. 1. This figure is available in its entirety in Appendix B3.

weaker gravity results in an ejected shell having a lower expansion velocity than normal stellar winds. The following post-eruption wind has a velocity higher than that of the aforementioned high mass-loss phase and eventually overtakes the ejected shell. Meanwhile, photoionized [Fe II] emission was reported from two hot (30 000 K) LBVs (AG Car and R 127, S02) which was attributed to their stronger UV flux.

We detected [Fe II] emission features associated with 3 LBVs (Table 8). So the [Fe II] detection rate of LBV nebula in our study is 14 per cent. If we include the 9 LBV samples of S02, the detection rate would be 29 per cent, that is, 9 out of 30 LBV nebulae (HD 168 625 duplicated in both studies). This new detection rate with a threefold sample is lower than the previous study, making the general physical conditions of LBVs not particularly suitable for the [Fe II] 1.644 μm line to arise but comparable to those of SNRs. The discrepancy in detection rates between this study and S02 might be due to the biased sample S02 used, which includes confirmed LBVs and candidate LBVs showing nebulosity in the Galaxy and the two most famous LBVs in LMC.

The [Fe II]-H images of identified LBV-IFOs are shown in Fig. 10. Brief information about them is listed in Table 8. In the [Fe II] emission, all identified LBV-IFOs share an elliptical/circular morphology. This is similar to their morphologies at 8 μm , but the extent appears smaller. We note that for G26.47 + 0.02 (IFO 102 and 103) the south-eastern diffuse structure was noticed in the [Fe II]-H

image. But the possibility of it being an artefact prevented us from assigning it as an IFO. The morphological coincidence of this South-East structure, IFO 102 and 103 with respect to the prominent part of the 8 μm nebula (Paron et al. 2012) implies the possibility of more extended, diffuse [Fe II] emission than seemingly identified. There are some new features revealed by [Fe II] emission: (1) IFO 65 – HD 168 625 is located at the centre of optical/IR elliptical structures that are broken toward the north-east. In the [Fe II] emission, we see a complete circular structure, the centre of which is offset toward the northeast. (2) IFO 162 – [KW97] 37–17 shows multiple shells in [Fe II] emission, forming together a much brighter elliptical structure than those in 8 μm or optical. This possibly indicates that the LBV had several active erupting phases that manufactured bright [Fe II]-emitting shells one by one.

We found that all [Fe II]-detected LBVs in the UWIFE survey also accompany nebulosities at 8 μm , but not vice versa. For example, we could not detect [Fe II] emission in three LBVs with 8 μm nebulosity (HD 168607, AFGL 2298, and GAL 024.73 + 00.69). Thus, the question of whether the LBV nebula, on account of the preceding giant eruption, is a prerequisite for the [Fe II] emission remains unanswered (S02). More comprehensive LBV samples and constrained physical properties of LBVs are needed to understand the possible relationship between the existence of the [Fe II] 1.644 μm line in LBV nebula and their past eruption histories.

Table 9. IFOs associated with SNRs.

IFO no.	SNR name		d [kpc]	F_{tot} [$10^{-17} \text{ W m}^{-2}$]
	G-name	Other name(s)		
IFO 007	G8.7 – 0.1	W 30	4.5	– ^a
IFO 015	G11.2 – 0.3	–	4.4	1090.00
IFO 038	G15.9 + 0.2	–	10.0	7.34
IFO 069	G18.1 – 0.1	–	5.6	21.50
IFO 070	G18.1 – 0.1	–	5.6	1.38
IFO 075	G18.9 – 1.1	–	2.0	14.10
IFO 076	G18.9 – 1.1	–	2.0	2.55
IFO 078	G18.9 – 1.1	–	2.0	192.00
IFO 079	G18.9 – 1.1	–	2.0	1.89
IFO 080	G21.8 – 0.6	Kes 69	5.2	627.00
IFO 084	G21.5 – 0.9	–	4.6	6.59
IFO 085	G21.5 – 0.9	–	4.6	59.70
IFO 088	G23.3 – 0.3	W 41	4.2	584.00
IFO 093 ^b	G28.8 + 1.5	–	3.4	0.39
IFO 098	G27.8 + 0.6	–	2.0	167.00
IFO 105	G27.4 + 0.0	Kes 73	8.5	274.00
IFO 106	G28.6 – 0.1	–	9.6	197.00
IFO 113	G31.9 + 0.0	3C 391	7.1	3423.55
IFO 117	G32.8 – 0.1	Kes 78	4.8	114.00
IFO 128	G34.7 – 0.4	W 44	2.8	3942.00
IFO 139	G39.2 – 0.3	3C 396	8.5	618.00
IFO 140	G41.5 + 0.4	–	4.1	1421.45
IFO 143	G41.1 – 0.3	3C 397	10.0	1691.00
IFO 147	G43.3 – 0.2	W49B	10.0	4739.00
IFO 160	G49.2 – 0.7	W51C	6.0	582.37

Notes. ^aDistances are from Lee et al. (2019). See Lee et al. (2019) for original references.

^aDue to significant background errors, flux was not derived and not included in the statistics of fluxes.

^bDistance from Shan et al. (2018).

^cPartially covered in the UWIFE survey.

4.6 Supernova remnants

SNRs are the brightest objects in [Fe II] emission. In SNRs, this line is mostly emitted from cooling gas behind radiative shocks. [Fe II] lines are strong in shocked gas because Fe abundance could be enhanced by shocks owing to grain destruction (Dinerstein 1995; Nisini 2008; Koo et al. 2016, and references therein). Before the UWIFE survey, a dozen Galactic and LMC SNRs had been observed in the NIR [Fe II] lines. The SNRs that are bright in [Fe II] emission lines may be divided into two groups: (1) middle-aged SNRs interacting with dense molecular (or atomic) clouds such as W 44 (Reach, Rho & Jarrett 2005), 3C 391 (Reach et al. 2002), and (2) young SNRs interacting with the dense circumstellar medium, such as Cas A (Koo et al. 2018), G11.2–0.3 (Moon et al. 2009), RCW 103 (Burton & Spyromilio 1993), and W49B (Lee et al. 2019). Then, Lee et al. (2019, hereafter, L19) searched for [Fe II] emission at the positions of the SNRs in the catalogue of Green (2014) using the UWIFE survey data and detected [Fe II] emission features toward 19 SNRs, more than half of which were new detections. In external galaxies, [Fe II] emission is used as a tracer of SNRs (Blair et al. 2014; Bruursema et al. 2014; Long et al. 2020), although strong [Fe II] lines may originate from sources ionized by X-rays, for example, in active galactic nuclei (Mouri et al. 2000; Morel, Doyon & St-Louis 2002).

We detected 25 IFOs associated with SNRs. All these SNR-IFOs belong to the 19 SNRs in Lee et al. (2019) except one (Table 9). It is worthwhile to point out that Lee et al. (2019) searched [Fe II] emission at 79 SNRs of the Green’s catalogue that are fully covered by the UWIFE survey. Four SNRs partially observed in the survey (i.e. G7.0 – 0.1, G13.3 – 1.3, G28.8 + 1.5, and G38.7 – 1.3) were not investigated, and our unbiased search resulted in the identification of a small [Fe II]-emitting patch inside the region of G28.8 + 1.5. Meanwhile, the Green’s catalogue of Galactic SNRs

has been updated (Green 2019), adding a new SNR G53.4 + 0.0 (partially covered in the UWIFE) and rejecting four (G20.4 + 0.1, G21.5 – 0.1, G23.6 + 0.3, and G59.8 + 1.2 that were reclassified as H II regions) in the survey area. None of the new or rejected SNRs showed [Fe II] emission features. So the new detection rate for fully covered SNRs is 25 per cent (19/75).

We note that Lee et al. (2019) compensated the [Fe II] line flux for the flux subtracted from the H band by multiplying with 1.15, whereas the fluxes in Table 9 are observed fluxes. As presented in Lee et al. (2019), IFO 147 that matches W49B is the brightest SNR-IFO. The detailed results for the 19 SNRs can be found in Lee et al. (2019).

5 SUMMARY

We have presented the first comprehensive catalogue of Galactic IFOs discovered in the UKIRT Widefield Infrared Survey for [Fe II] (UWIFE). It is the first Galactic catalogue of extended [Fe II] line emission sources using an unbiased, large-scale survey. We have discovered many previously unreported [Fe II] 1.644 μm line sources. Therefore, this catalogue provides an opportunity to broaden the horizons of the study of the shocked regions of our Galaxy, especially with the synergy of the UWISH2 survey.

We have searched for extended IFOs in the inner GP ($7^\circ < l < 62^\circ$; $|b| \lesssim 1^\circ 5$). In order for the search to be efficient, we removed point-like continuum sources from the [Fe II] 1.644 μm images using H -band images taken as part of the UKIDSS GPS survey. We identified most of the IFOs by visual inspection and added several faint IFOs with an automatic source identification which uses the same source detection algorithm as in UWISH2 (Froebrich et al. 2015). In total, 204 IFOs were identified. We measured the sizes and fluxes of these 204 IFOs and presented their properties. We have searched for the counterparts of the IFOs via positional cross-matching with previously known sources and found that the majority of IFOs are associated with SNRs, YSOs, HII regions, PNe, and LBVs. We group IFOs by their counterpart types and discuss their statistical and morphological properties. The main results are summarized as follows.

(1) In the 180 deg² GP area of the first Galactic quadrant covered by the UWIFE survey ($7^\circ < l < 62^\circ$; $|b| \lesssim 1^\circ 5$), we identified 204 IFOs. The identified IFOs are classified according to their counterparts: YSO-IFOs, HII-IFOs, CHII-IFOs, PN-IFOs, LBV-IFOs, and SNR-IFOs. There are 100 YSO-IFOs, 11 HII-IFOs, 22 CHII-IFOs, 17 PN-IFOs, 4 LBV-IFOs, and 25 SNR-IFOs. We could not identify counterparts for 25 IFOs, and they are classified as ‘unknown-IFOs’ (Fig. 11, Table 10). The majority of IFOs are new discoveries that have never been revealed in previous [Fe II] line studies.

(2) The SNR-IFOs and HII-IFOs are the brightest IFOs, and they dominate the [Fe II] 1.644 μm line flux in the GP. They contribute 96 per cent of the total [Fe II] 1.644 μm line flux of the IFOs ($2.6 \times 10^{-13} \text{ W m}^{-2}$); 76 per cent by SNR-IFOs and 20 per cent by HII/CHII-IFOs. The YSO-IFOs, PN-IFOs, and LBV-IFOs are generally orders of magnitude fainter, while the unknown-IFOs are the faintest.

(3) The average number density of IFOs is $\sim 1.1 \text{ deg}^{-2}$. The number density is highly variable spatially, especially for the IFOs associated with objects in the early-evolutionary phase, for example, IFOs associated with H II regions and YSOs. In Galactic longitude, there are prominent peaks at $l \sim 16^\circ$ and 51° , while there is a ‘void’ at $l \sim 40^\circ - 50^\circ$ where the number of IFOs is very small. The spatial

Table 10. IFOs associated with unknown-type.

IFO no.	Comment	Reference
IFO 001	Amorphous structure elongated from NE to SW. The bright star 15 arcsec east to the IFO is ALLWISE J180138.06 – 224227.8, a luminosity class of giant	s19
IFO 002	Compact IFO, H II region in the vicinity, H ₂ at south is slightly more extended	–
IFO 017	Located in between late-type star [MHS2002] 196 and dark cloud SDC G12.804 + 0.055	–
IFO 028	Biconical structure in the north-east of IRAS 18134 – 1838, however, relation is uncertain	–
IFO 068	An amorphous source in the middle of a possible evolved star J182227.66 – 171556.6 and the two stars having a high proper motion at south	m19 r10
IFO 096	Spatially coincident with large-scale radio shell G25.8 + 0.2	c18
IFO 109	IFO at the border of AllWISE J184501.48 – 001716.3, H ₂ emission at north	–
IFO 110	Amorphous IFO, possibly surrounding MIR source AllWISE J184515.66 – 031606.9	–
IFO 116	Amorphous IFO, no well-known sources in the vicinity	–
IFO 133	Cometary IFO elongated north to south, AllWISE J185905.19 + 004837.9 at the north	–
IFO 135	Amorphous IFO adjoining <i>K</i> -band source UGPS J185923.30 + 010415.0 which was not detected in <i>J</i> and <i>H</i> bands	–
IFO 142	Bow-shock morphology with apex on NW side, in between possible giant star at south and evolved/Class III YSO candidate at north	s19 m16
IFO 150	Compact structure coincident with H ₂ , ALLWISE J191530.41 + 132737.2 showing MIR photometric anomaly is located in the vicinity	s17
IFO 166	Knot-like small-scale IFO	–
IFO 167	Coincident with extended H ₂ , but with the shifted peak position	–
IFO 168	A compact, faint knot in between IFO 167 and 169	–
IFO 169	Knot-like diffuse IFO	–
IFO 170	Knot-like IFO in the middle of IFO 166 – 169 and IFO 172	–
IFO 171	Knot-like IFO in the middle of IFO 166 – 169 and IFO 172	–
IFO 172	Compact IFO, small-scale H ₂ in the vicinity	–
IFO 182	Amorphous IFO with the bright northern component coincident with MIR source GLIMPSE G052.9805 – 00.0394	–
IFO 187	Knot-like IFO. Possible evolved star WISE J193120.72 + 192006.1 and dwarf star <i>Gaia</i> DR2 1 825 442 262 813 750 912 in south	m19 s19
IFO 191	Diffuse elongated IFO from east to west. MIPS GAL source MG059.1989+00.5106 at north, MIR source ALLWISE J194013.97 + 232657.5 superposed to the IFO, red giant star 2MASS J19401620 + 2326577 at east	–
IFO 197	Slightly elongated compact IFO, the centre of NGC 6823 is located at east	–
IFO 201	Cometary IFO with its south-eastern head part coincident with ALLWISE J194611.21 + 221554.3, which is a luminosity class of dwarf candidate	s19

*Notes.**Column 3: references of counterpart candidates: c18 – Cichowski et al. (2018), m16 – Marton et al. (2016), m19 – Marton et al. (2019), r10 – Roeser, Demleitner & Schilbach (2010), s17 – Solarz et al. (2017), and s19 – Stassun et al. (2019).

*Counterpart candidates around IFO 166–172: 1: YSO candidate SSTGLMC G050.3746 – 00.4149, 2: AGB candidate SSTGLMC G050.3756 – 00.4214, 3: MHO 2624, 4: EGO G050.36 – 0.42, 5: YSO candidate SSTGLMC G050.3666 – 00.3944, 6: YSO candidate SSTGLMC G050.3647 – 00.3979, 7: YSO candidate SSTGLMC G050.3587 – 00.4123, 8: YSO candidate SSTGLMC G050.3675 – 00.4089, 9: YSO candidate SSTGLMC G050.3691 – 00.4096, 10: YSO candidate SSTGLMC G050.3704 – 00.4095, 11: YSO candidate SSTGLMC G050.3741 – 00.4083, 12: YSO AGAL G050.376 – 00.421, 13: YSO candidate SSTGLMC G050.3762 – 00.4205, and 14: YSO candidate SSTGLMC G050.3575 – 00.4182.

distribution in Galactic latitude is centred at $b = -0^\circ 12$ with a standard deviation of $0^\circ 65$.

(4) The results on the individual types of IFOs are summarized below.

(i) YSO-IFOs

We detected 100 YSO-IFOs, which constitutes half of the IFOs in our catalogue. Only 17 of those are associated with the RMS sources, which represent massive YSOs. The YSO-IFOs might be preferentially tracing low-mass YSOs. On the other hand, the majority (87 per cent) of YSO-IFOs are not associated with HH objects, suggesting that the YSO-IFOs are revealing previously hidden, optically obscured outflows in star-forming regions. YSO-IFOs have diverse morphologies, and we have classified them into four categories; bipolar, cometary, knot-like, and amorphous.

(ii) HII-IFOs and CHII-IFOs

We have identified 11 IFOs associated with 4 H II regions (Table 6). Almost all HII-IFOs are located in the well-known star-forming complexes, W 31, M 17, and W 51. Some HII-IFOs appear as either thin filaments or diffuse amorphous emission

structures within the radio structure, so their association with the H II regions is very likely. But some are faint and diffuse and extend beyond the radio boundary of the H II regions, so their association is uncertain and needs to be confirmed. We also detected 22 IFOs associated with 16 CHII, including 6 previously reported (Table 5). Among the 16 CHII regions, 10 are catalogued in CORNISH, which corresponds to 4 per cent of the 237 CHII regions in the CORNISH catalogue in the survey area. CHII-IFOs have diverse morphologies: jet-like, shell-like, and amorphous.

(iii) PN-IFOs

We detected 17 PN-IFOs. They are associated with 14 PNe (i.e. 5 PNe, 8 PN candidates, and one post-AGB candidate; Table 7), which correspond to about 4.7 per cent of the PNe in the survey area. We have classified the morphologies of PN-IFOs following Parker et al. (2006) and compared them with those in H α and H₂. Some PNe have [Fe II] morphologies different from the H α and H₂ morphologies, which implies that the [Fe II] line reveals new substructures, possibly probing additional mass-loss histories. The physical sizes of some PN-IFOs are larger than 0.9 pc.

(iv) LBV-IFOs

We detected 4 LBV-IFOs. They are associated with 3 LBVs out of 22 LBVs and their candidates in the survey area (Table 8), so the detection rate of [Fe II] emission associated with LBVs in this study is 14 per cent. All LBV-IFOs share an elliptical or circular morphology. Some show multiple shells. We found that all [Fe II]-detected LBVs in the UWIFE survey also accompany nebulosity at 8 μm , but not vice versa.

(v) SNR-IFOs

We detected 25 SNR-IFOs. They are associated with 20 SNRs, which corresponds to 25 per cent of the 75 known SNRs in the survey area. The SNR-IFOs occupy 76 per cent of the total [Fe II] flux of IFOs, and the four brightest IFOs are SNR-IFOs. On the other hand, the lowest surface brightness IFOs are also SNR-IFOs, showing the patchy [Fe II] emission in SNRs. All SNRs with [Fe II] emission features except one (G28.8 + 1.5) have been previously reported by Lee et al. (2019). The detailed results on the [Fe II] emission on the 19 SNRs can be found in Lee et al. (2019).

ACKNOWLEDGEMENTS

The authors are grateful to the anonymous referee for the comments and suggestions that improved the clarity of this paper. We also acknowledge the Cambridge Astronomical Survey Unit and the WFCAM Science Archive for the reduction and ingest of the survey data. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This research has made use of the VizieR catalogue access tool, CDS, Strasbourg, France (DOI: 10.26093/cds/vizier). The original description of the VizieR service was published in 2000, *A&AS* 143, 23. The UWIFE survey was supported by the K-GMT Science Program funded through the Korea GMT Project operated by the Korea Astronomy and Space Science Institute (KASI). B-CK acknowledges support from the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT and Future Planning (2020R1A2B5B01001994 and RS-2023-00277370). This research of H-JK was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (RS-2023-00246733). YK acknowledges support from Kim In-Ha Scholarship.

DATA AVAILABILITY

The data underlying this article are available to download from <http://gems0.kasi.re.kr/uwif/>.

REFERENCES

Akras S., Guzman-Ramirez L., Leal-Ferreira M. L., Ramos-Larios G., 2019, *ApJS*, 240, 21
 Alexander M. J., Kobulnicky H. A., 2012, *ApJ*, 755, 30
 Alonso-Herrero A., Rieke G. H., Rieke M. J., Kelly D. M., 2003, *AJ*, 125, 1210
 Alten V. P., Bally J., Devine D., Miller G. J., 1997, in Reipurth B., Bertout C.eds, Proc. IAU Symp. 182, Herbig-Haro Flows and the Birth of Stars, Kluwer, Dordrecht, p. 51
 Anderson L. D., Bania T. M., 2009, *ApJ*, 690, 706
 Areal M. B., Paron S., Fariña C., Ortega M. E., Celis Peña M., Rubio M., 2020, *A&A*, 641, A104
 Baan W. A., Imai H., Orosz G., 2021, *Res. Astron. Astrophys.*, 21, 275

Bailer-Jones C. A. L., Rybizki J., Fouesneau M., Mantelet G., Andrae R., 2018, *AJ*, 156, 58
 Baldwin J. A. et al., 1996, *ApJ*, 468, L115
 Baldwin J. A., Ferland G. J., Martin P. G., Corbin M. R., Cota S. A., Peterson B. M., Slettebak A., 1991, *ApJ*, 374, 580
 Bally J., Chia Z., Ginsburg A., Reipurth B., Tanaka K. E. I., Zinnecker H., Faulhaber J., 2022, *ApJ*, 924, 50
 Barbosa C. L., Blum R. D., Conti P. S., Damini A., Figuerêdo E., 2008, *ApJ*, 678, L55
 Barsony M., 1989, *ApJ*, 345, 268
 Bautista M. A., Pradhan A. K., 1998, *ApJ*, 492, 650
 Beck S. C., Kelly D. M., Lacy J. H., 1998, *AJ*, 115, 2504
 Belikov A. N., Kharchenko N. V., Piskunov A. E., Schilbach E., 1999, *A&AS*, 134, 525
 Beuther H. et al., 2016, *A&A*, 595, A32
 Blair W. P. et al., 2014, *ApJ*, 788, 55
 Bloomer J. D. et al., 1998, *ApJ*, 506, 727
 Blum R. D., Conti P. S., Damini A., 2000, *AJ*, 119, 1860
 Blum R. D., Damini A., Conti P. S., 2001, *AJ*, 121, 3149
 Boissay R., Parker Q. A., Frew D. J., Bojicic I., 2012, IAU Symp. 283, Planetary Nebulae: An Eye to the Future. Kluwer, Dordrecht, p. 316
 Bronfman L., Nyman L. A., May J., 1996, *A&AS*, 115, 81
 Bruursema J., Meixner M., Long K. S., Otsuka M., 2014, *AJ*, 148, 41
 Burton M., Spyromilio J., 1993, *Publ. Astron. Soc. Aust.*, 10, 327
 Cappellaro E., Benetti S., Sabbadin F., Salvadore L., Turatto M., Zanin C., 1994, *MNRAS*, 267, 871
 Caratti o Garatti A., Giannini T., Nisini B., Lorenzetti D., 2006, *A&A*, 449, 1077
 Caratti o Garatti A., Stecklum B., Linz H., Garcia Lopez R., Sanna A., 2015, *A&A*, 573, A82
 Carey S. J. et al., 2009, *PASP*, 121, 76
 Casali M. et al., 2007, *A&A*, 467, 777
 Churchwell E. et al., 2009, *PASP*, 121, 213
 Churchwell E., 2002, *ARA&A*, 40, 27
 Cichowski S., Duronea N. U., Suad L. A., Reynoso E. M., Dorda R., 2018, *MNRAS*, 474, 647
 Clark D. M., López J. A., Edwards M. L., Winge C., 2014, *AJ*, 148, 98
 Clark J. S., Davies B., Najarro F., MacKenty J., Crowther P. A., Messineo M., Thompson M. A., 2009, *A&A*, 504, 429
 Clark J. S., Egan M. P., Crowther P. A., Mizuno D. R., Larionov V. M., Arkharov A., 2003, *A&A*, 412, 185
 Cooper H. D. B. et al., 2013, *MNRAS*, 430, 1125
 Cordiner M. A. et al., 2016, *ApJ*, 828, 51
 Craine E. R., Boeshaar G. O., Byard P. L., 1981, *AJ*, 86, 751
 Csengeri T. et al., 2016, *A&A*, 586, A149
 Cyganowski C. J. et al., 2008, *AJ*, 136, 2391
 Cyganowski C. J., Koda J., Rosolowsky E., Towers S., Donovan Meyer J., Egusa F., Momose R., Robitaille T. P., 2013, *ApJ*, 764, 61
 Davies B., Hoare M. G., Lumsden S. L., Hosokawa T., Oudmaier R. D., Urquhart J. S., Mottram J. C., Stead J., 2011, *MNRAS*, 416, 972
 Davis C. J., Gell R., Khanzadyan T., Smith M. D., Jenness T., 2010, *A&A*, 511, A24
 Davis C. J., Smith M. D., Gledhill T. M., Varricatt W. P., 2005, *MNRAS*, 360, 104
 Davis C. J., Whelan E., Ray T. P., Chrysostomou A., 2003, *A&A*, 397, 693
 de la Fuente E., Porras A., Trinidad M. A., Kurtz S. E., Kemp S. N., Tafaya D., Franco J., Rodríguez-Rico C., 2020, *MNRAS*, 492, 895
 Delgado-Inglada G., Rodríguez M., 2014, *ApJ*, 784, 173
 Dent W. R. F., Aspin C., 1992, *MNRAS*, 259, 401
 Devine D., Reipurth B., Bally J., 1997, in Reipurth B., Bertout C.eds, Proc. IAU Symp. 182, Low Mass Star Formation - from Infall to Outflow. Kluwer, Dordrecht, p. 91
 Dinerstein H., 1995, in Williams R., Livio M.eds, The Analysis of Emission Lines: A Meeting in Honor of the 70th Birthdays of D. E. Osterbrock and M. J. Seaton. Cambridge University Press, Cambridge, p. 134
 Drew J. E. et al., 2005, *MNRAS*, 362, 753
 Egan M. P., Price S. D., Kraemer K. E., 2003, American Astronomical Society Meeting Abstracts, 57.08

- Elia D. et al., 2017, *MNRAS*, 471, 100
- Ellerbroek L. E., Podio L., Kaper L., Sana H., Huppenkothen D., de Koter A., Monaco L., 2013, *A&A*, 551, A5
- Fazeli N., Busch G., Valencia-S. M., Eckart A., Zajaček M., Combes F., García-Burillo S., 2019, *A&A*, 622, A128
- Ferrero L., Le Du P., Mulato L., Outters N., Zoll S., 2015, *L'Astronomie*, 129, 1.42
- Frank A., 1999, *New Astron. Rev.*, 43, 31
- Froebrich D. et al., 2011, *MNRAS*, 413, 480
- Froebrich D. et al., 2015, *MNRAS*, 454, 2586
- Froebrich D., Ioannidis G., 2011, *MNRAS*, 418, 1375
- Froebrich D., Makin S. V., 2016, *MNRAS*, 462, 1444
- Fuente A., Martín-Pintado J., Rodríguez-Franco A., Moriarty-Schieven G. D., 1998, *A&A*, 339, 575
- Fujita S. et al., 2021, *PASJ*, 73, S172
- Gledhill T. M., Froebrich D., Campbell-White J., Jones A. M., 2018, *MNRAS*, 479, 3759
- Goddi C., Ginsburg A., Maud L. T., Zhang Q., Zapata L. A., 2020, *ApJ*, 905, 25
- González-Santamaría I., Manteiga M., Manchado A., Ulla A., Dafonte C., 2020, *Galaxies*, 8, 29
- Green D. A., 2014, *Bull. Astron. Soc. India*, 42, 47
- Green D. A., 2019, *J. Astrophys. Astron.*, 40, 36
- Greenhouse M. A., Woodward C. E., Thronson Harley A. J., Rudy R. J., Rossano G. S., Erwin P., Puetter R. C., 1991, *ApJ*, 383, 164
- Guarcello M. G., Prisinzano L., Micela G., Damiani F., Peres G., Sciortino S., 2007, *A&A*, 462, 245
- Hanson M. M., Howarth I. D., Conti P. S., 1997, *ApJ*, 489, 698
- Hanson M. M., Luhman K. L., Rieke G. H., 2002, *ApJS*, 138, 35
- Helfand D. J., Becker R. H., White R. L., Fallon A., Tuttle S., 2006, *AJ*, 131, 2525
- Hennig M. G., Riffel R. A., Dors O. L., Riffel R., Storchi-Bergmann T., Colina L., 2018, *MNRAS*, 477, 1086
- Herbig G. H., 1974, *Lick Obs. Bull.*, 658, 1
- Hoare M. G. et al., 2012, *PASP*, 124, 939
- Hollenbach D. J., Chernoff D. F., McKee C. F., 1989, in Böhm-Vitense E. ed., *ESA SP-290: Infrared Spectroscopy in Astronomy*. ESA, Noordwijk, p. 245
- Hora J. L., Latter W. B., Deutsch L. K., 1999, *ApJS*, 124, 195
- Hsieh T.-H., Lai S.-P., Belloche A., 2017, *AJ*, 153, 173
- Humphreys R. M., Davidson K., 1994, *PASP*, 106, 1025
- Hutsemekers D., van Drom E., Gosset E., Melnick J., 1994, *A&A*, 290, 906
- Immer K., Reid M. J., Menten K. M., Brunthaler A., Dame T. M., 2013, *A&A*, 553, A117
- Ioannidis G., Froebrich D., 2012a, *MNRAS*, 421, 3257
- Ioannidis G., Froebrich D., 2012b, *MNRAS*, 425, 1380
- Irabor T. et al., 2018, *MNRAS*, 480, 2423
- Jayasinghe T. et al., 2018, *MNRAS*, 477, 3145
- Kalcheva I. E., Hoare M. G., Urquhart J. S., Kurtz S., Lumsden S. L., Purcell C. R., Zijlstra A. A., 2018, *A&A*, 615, A103
- Kanarek G., Shara M., Faherty J., Zurek D., Moffat A., 2015, *MNRAS*, 452, 2858
- Kang M., Bieging J. H., Povich M. S., Lee Y., 2009, *ApJ*, 706, 83
- Kastner J. H., Weintraub D. A., Gatley I., Merrill K. M., Probst R. G., 1996, *ApJ*, 462, 777
- Kim H.-J., Koo B.-C., Davis C. J., 2015, *ApJ*, 802, 59
- Kim H.-J., Koo B.-C., Pyo T.-S., Davis C. J., 2018, *ApJ*, 863, 74
- Kim H., Lacy J. H., Jaffe D. T., 2017, *American Astronomical Society Meeting Abstracts #230*. p. 215.05
- Kolpak M. A., Jackson J. M., Bania T. M., Clemens D. P., Dickey J. M., 2003, *ApJ*, 582, 756
- Konigl A., 1982, *ApJ*, 261, 115
- Koo B.-C., 2014, in Ray A., McCray R. A. eds, *Proc. IAU Symp. 296, Supernova Environmental Impacts*. Kluwer, Dordrecht, p. 214
- Koo B.-C., Kim H.-J., Lee Y.-H., Raymond J. C., Lee J.-J., Yoon S.-C., Moon D.-S., 2018, *ApJ*, 866, 139
- Koo B.-C., Kim K.-T., Seward F. D., 1995, *ApJ*, 447, 211
- Koo B.-C., Lee Y.-H., 2015, *Publ. Korean Astron. Soc.*, 30, 145
- Koo B.-C., Moon D.-S., Lee H.-G., Lee J.-J., Matthews K., 2007, *ApJ*, 657, 308
- Koo B.-C., Raymond J. C., Kim H.-J., 2016, *J. Korean Astron. Soc.*, 49, 109
- Kraus S. et al., 2006, *A&A*, 455, 521
- Kuchar T. A., Clark F. O., 1997, *ApJ*, 488, 224
- Kuhn M. A., de Souza R. S., Krone-Martins A., Castro-Ginard A., Ishida E. O., Povich M. S., Hillenbrand L. A., *COIN Collaboration*, 2021, *ApJS*, 254, 33
- Leahy D., Green K., Tian W., 2014, *MNRAS*, 438, 1813
- Lee H.-G. et al., 2013, *ApJ*, 770, 143
- Lee H.-G., Moon D.-S., Koo B.-C., Lee J.-J., Matthews K., 2009, *ApJ*, 691, 1042
- Lee J.-J. et al., 2014, *MNRAS*, 443, 2650
- Lee Y.-H., Koo B.-C., Lee J.-J., Burton M. G., Ryder S., 2019, *AJ*, 157, 123
- Li F. C., Xu Y., Wu Y. W., Yang J., Lu D. R., Menten K. M., Henkel C., 2016, *AJ*, 152, 92
- Long K. S., Blair W. P., Winkler P. F., Lacey C. K., 2020, *ApJ*, 899, 14
- Lucas P. W. et al., 2008, *MNRAS*, 391, 136
- Lumsden S. L., Hoare M. G., Urquhart J. S., Oudmaijer R. D., Davies B., Mottram J. C., Cooper H. D. B., Moore T. J. T., 2013, *ApJS*, 208, 11
- Makin S. V., Froebrich D., 2018, *ApJS*, 234, 8
- Manchado A., Stanghellini L., Guerrero M. A., 1996, *ApJ*, 466, L95
- Marton G. et al., 2019, *MNRAS*, 487, 2522
- Marton G., Tóth L. V., Paladini R., Kun M., Zahorecz S., McGehee P., Kiss C., 2016, *MNRAS*, 458, 3479
- Mauron N., Huggins P. J., 2010, *A&A*, 513, A31
- McCullum B., Bruhweiler F. C., Wahlgren G. M., Eriksson M., Verner E., 2008, *ApJ*, 682, 1087
- McGroarty F., Ray T. P., Bally J., 2004, *A&A*, 415, 189
- McKee C. F., Chernoff D. F., Hollenbach D. J., 1984, in Kessler M. F., Phillips J. Peds, *Astrophysics and Space Science Library*, Vol. 108, *Galactic and Extragalactic Infrared Spectroscopy*. D. Reidel Publishing Co., Dordrecht, p. 103
- Meaburn J., White N. J., 1982, *MNRAS*, 199, 121
- Mège P. et al., 2021, *A&A*, 646, A74
- Messineo M. et al., 2015, *ApJ*, 805, 110
- Miszalski B., Parker Q. A., Acker A., Birkby J. L., Frew D. J., Kovacevic A., 2008, *MNRAS*, 384, 525
- Molinari S. et al., 2010, *PASP*, 122, 314
- Molinari S., Brand J., Cesaroni R., Palla F., 1996, *A&A*, 308, 573
- Molinari S., Brand J., Cesaroni R., Palla F., Palumbo G. G. C., 1998, *A&A*, 336, 339
- Moon D.-S., Koo B.-C., Lee H.-G., Matthews K., Lee J.-J., Pyo T.-S., Seok J. Y., Hayashi M., 2009, *ApJ*, 703, L81
- Morel T., Doyon R., St-Louis N., 2002, *MNRAS*, 329, 398
- Mottram J. C. et al., 2011, *ApJ*, 730, L33
- Mouri H., Kawara K., Taniguchi Y., 2000, *ApJ*, 528, 186
- Nakashima J.-i., Ladeyschikov D. A., Sobolev A. M., Zhang Y., Hsia C.-H., Yung B. H. K., 2016, *ApJ*, 825, 16
- Navarro S. G., Corradi R. L. M., Mampaso A., 2012, in *Proc. IAU Symp. 283, Planetary Nebulae: An Eye to the Future*. Kluwer, Dordrecht, p. 460
- Nazé Y., Rauw G., Hutsemekers D., 2012, *A&A*, 538, 47
- Nielbock M., Chini R., Jütte M., Manthey E., 2001, *A&A*, 377, 273
- Nikogossian E. G., Magakian T. Y., Movsessian T. A., 2007, *Astron. Rep.*, 51, 735
- Nisini B., 2008, in Bacciotti F., Testi L., Whelan E. eds, *Lecture Notes in Physics*, Vol. 742, *Jets from Young Stars II*. Springer-Verlag, Berlin Heidelberg, p. 79
- Nisini B., Caratti o Garatti A., Giannini T., Lorenzetti D., 2002, *A&A*, 393, 1035
- Oh H. et al., 2016, *ApJ*, 817, 148
- Okada Y., Onaka T., Miyata T., Okamoto Y. K., Sakon I., Shibai H., Takahashi H., 2008, *ApJ*, 682, 416
- Okumura S.-i., Mori A., Nishihara E., Watanabe E., Yamashita T., 2000, *ApJ*, 543, 799
- Oliva E., Moorwood A. F. M., Danziger I. J., 1989, *A&A*, 214, 307
- Osterbrock D. E., 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*. University Science Books,

- Osterbrock D. E., Tran H. D., Veilleux S., 1992, *ApJ*, 389, 305
- Parker Q. A., Bojičić I. S., Frew D. J., 2016, *J. Phys. Conf. Ser.*, 728, 032008
- Parker Q. A. et al., 2006, *MNRAS*, 373, 79
- Paron S., Combi J. A., Petriella A., Giacani E., 2012, *A&A*, 543, A23
- Paron S., Fariña C., Ortega M. E., 2016, *A&A*, 593, A132
- Peimbert A., Peimbert M., 2010, *ApJ*, 724, 791
- Peimbert M., 1993, *Rev. Mex. Astron. Astrofis.*, 27, 9
- Peiřker F., Eckart A., Sabha N. B., Zajaček M., Bhat H., 2020, *ApJ*, 897, 28
- Pesenti N., Dougados C., Cabrit S., O'Brien D., Garcia P., Ferreira J., 2003, *A&A*, 410, 155
- Phillips J. P., 2004, *MNRAS*, 353, 589
- Povich M. S. et al., 2009, *ApJ*, 696, 1278
- Povich M. S., Whitney B. A., 2010, *ApJ*, 714, 285
- Pudritz R. E., Norman C. A., 1983, *ApJ*, 274, 677
- Pyo T.-S. et al., 2006, *ApJ*, 649, 836
- Pyo T.-S., Hayashi M., Kobayashi N., Terada H., Tokunaga A. T., 2009, *ApJ*, 694, 654
- Ragan S. E., Bergin E. A., Gutermuth R. A., 2009, *ApJ*, 698, 324
- Reach W. T., Rho J., Jarrett T. H., 2005, *ApJ*, 618, 297
- Reach W. T., Rho J., Jarrett T. H., Lagage P.-O., 2002, *ApJ*, 564, 302
- Reipurth B., 2000, *VizieR Online Data Catalog*, p. V/104
- Reiter M., Kiminki M. M., Smith N., Bally J., 2017, *MNRAS*, 470, 4671
- Reiter M., Nayak O., Meixner M., Jones O., 2019, *MNRAS*, 483, 5211
- Reiter M., Smith N., Bally J., 2016, *MNRAS*, 463, 4344
- Reiter M., Smith N., Kiminki M. M., Bally J., 2015, *MNRAS*, 450, 564
- Retes-Romero R., Mayya Y. D., Luna A., Carrasco L., 2017, *ApJ*, 839, 113
- Rice E. L., Prato L., McLean I. S., 2006, *ApJ*, 647, 432
- Riffel R. A. et al., 2016, *MNRAS*, 461, 4192
- Robitaille T. P. et al., 2008, *AJ*, 136, 2413
- Rodríguez M., 2002, *A&A*, 389, 556
- Roeser S., Demleitner M., Schilbach E., 2010, *AJ*, 139, 2440
- Roshi D. A., Churchwell E., Anderson L. D., 2017, *ApJ*, 838, 144
- Russell D., 2003, *A&A*, 397, 133
- Rygl K. L. J., Wyrowski F., Schuller F., Menten K. M., 2010, *A&A*, 515, A42
- Sabin L. et al., 2014, *MNRAS*, 443, 3388
- Sakamoto S., Hasegawa T., Handa T., Hayashi M., Oka T., 1997, *ApJ*, 486, 276
- Samal M. R., Chen W. P., Takami M., Jose J., Froebrich D., 2018, *MNRAS*, 477, 4577
- Sánchez-Monge Á., López-Sepulcre A., Cesaroni R., Walmsley C. M., Codella C., Beltrán M. T., Pestalozzi M., Molinari S., 2013, *A&A*, 557, 94
- Saral G. et al., 2017, *ApJ*, 839, 108
- Sasaki M., Mäkelä M. M., Klochkov D., Santangelo A., Suleimanov V., 2018, *MNRAS*, 479, 3033
- Scoville N. Z., Yun M. S., Clemens D. P., Sanders D. B., Waller W. H., 1987, *ApJS*, 63, 821
- Sewilo M., Churchwell E., Kurtz S., Goss W. M., Hofner P., 2004, *ApJ*, 605, 285
- Shan S. S., Zhu H., Tian W. W., Zhang M. F., Zhang H. Y., Wu D., Yang A. Y., 2018, *ApJS*, 238, 35
- Shang H., Krasnopolsky R., Liu C.-F., Wang L.-Y., 2020, *ApJ*, 905, 116
- Shinn J.-H. et al., 2013, *ApJ*, 777, 45
- Shinn J.-H. et al., 2014, *ApJS*, 214, 11
- Shu F., Najita J., Ostriker E., Wilkin F., Ruden S., Lizano S., 1994, *ApJ*, 429, 781
- Simon R., Jackson J. M., Rathborne J. M., Chambers E. T., 2006, *ApJ*, 639, 227
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Smith N., 2002, *MNRAS*, 336, L22
- Smith N., Balick B., Gehrz R. D., 2005, *AJ*, 130, 853
- Smith R. G., Lawson W. A., Wright C. M., 2007, *MNRAS*, 375, 257
- Solarz A., Bilicki M., Gromadzki M., Pollo A., Durkalec A., Wypych M., 2017, *A&A*, 606, 39
- Stassun K. G. et al., 2019, *AJ*, 158, 138
- Takami M. et al., 2002, *ApJ*, 566, 910
- Takami M. et al., 2006, *ApJ*, 641, 357
- Thompson R. I., Smith B. A., Hester J. J., 2002, *ApJ*, 570, 749
- Tian H.-J., El-Badry K., Rix H.-W., Gould A., 2020, *ApJS*, 246, 4
- Traficante A., Fuller G. A., Peretto N., Pineda J. E., Molinari S., 2015, *MNRAS*, 451, 3089
- Tsitali A. E., Bourke T. L., Peterson D. E., Myers P. C., Dunham M. M., Evans Neal J. I., Huard T. L., 2010, *ApJ*, 725, 2461
- Urquhart J. S. et al., 2022, *MNRAS*, 510, 3389
- Urquhart J. S. et al., 2012, *MNRAS*, 420, 1656
- Urquhart J. S. et al., 2018, *MNRAS*, 473, 1059
- Van de Steene G. C., van Hoof P. A. M., 2003, *A&A*, 406, 773
- Varricatt W. P., Wouterloot J. G. A., Ramsay S. K., Davis C. J., 2018, *MNRAS*, 480, 4231
- Veneziani M. et al., 2013, *A&A*, 549, 130
- Vickers S. B., Frew D. J., Parker Q. A., Bojičić I. S., 2015, *MNRAS*, 447, 1673
- Vig S., Ghosh S. K., Kulkarni V. K., Ojha D. K., Verma R. P., 2006, *ApJ*, 637, 400
- Vioque M., Oudmaijer R. D., Schreiner M., Mendigutía I., Baines D., Mowlavi N., Pérez-Martínez R., 2020, *A&A*, 638, A21
- Weis K., Bomans D. J., 2020, *Galaxies*, 8, 20
- Welch C. A., Frank A., Pipher J. L., Forrest W. J., Woodward C. E., 1999, *ApJ*, 522, L69
- White R. L., Becker R. H., Helfand D. J., 2005, *AJ*, 130, 586
- Witham A. R., Knigge C., Drew J. E., Greimel R., Steeghs D., Gänsicke B. T., Groot P. J., Mampaso A., 2008, *MNRAS*, 384, 1277
- Wood D. O. S., Churchwell E., 1989, *ApJS*, 69, 831
- Xu J.-L. et al., 2019, *A&A*, 627, A27
- Xue R., Wu Y., 2008, *ApJ*, 680, 446
- Yang A. Y., Tian W. W., Zhu H., Leahy D. A., Wu D., 2016, *ApJS*, 223, 6
- Yuan J.-H., Li J. Z., Huang Y. F., Hsia C.-H., Miao J., 2012, *A&A*, 540, 95
- Zhu H., Tian W. W., Torres D. F., Pedalotti G., Su H. Q., 2013, *ApJ*, 775, 95
- Zinnecker H., Yorke H. W., 2007, *ARA&A*, 45, 481
- Zucker C., Battersby C., Goodman A., 2018, *ApJ*, 864, 153

APPENDIX A: EXAMPLES OF ARTEFACTS




Figure A1. Example of artefacts in the [Fe II]-H image. (1) The residual of star subtraction shown as cross-stripes. (2) A variable and saturated star with a diffraction pattern. (3) Arc-shape ghosts near a very bright star. The diffraction pattern of a bright star is superposed. (4) Electronic crosstalk near a bright star. (5) Diffraction pattern of a bright star. (6) High proper-motion star. (Black: bright and white: dark.)

APPENDIX B: THE UWIFE [Fe II]-H AND GPS 3-COLOUR IMAGES OF IFOs




Figure B1. IFOs with YSO counterpart candidates in continuum-subtracted [Fe II] and three-colour *KHJ* UKIDSS images in the same field of view. The format for these images is the same as that of Fig. 1. Crosses denote adjacent YSOs in the field of view. Contours are H_2 2.12 μm emission adopted from UWISH2.




Figure B1. *Continued.*




Figure B1. *Continued.*




Figure B1. *Continued.*




Figure B1. *Continued.*




Figure B1. *Continued.*

Downloaded from <https://academic.oup.com/mnras/article/528/3/4657/7591325> by guest on 19 February 2024




Figure B1. *Continued.*




Figure B2. Continuum-subtracted [Fe II] images of IFOs with PN counterpart candidates. The units on the UWIFE [Fe II]-H are DN, with the darker colour denoting a higher DN. The corresponding source names for each IFO are shown. The cross marks the central position of the counterpart. The images of IFO 73, 095, 112, 129, 157, and 164 are smoothed with a two-pixel Gaussian. In all images, north is at the top and east is on the left side.




Figure B2. *Continued.*





Figure B3. IFOs with counterpart candidates unknown in continuum-subtracted [Fe II] images as in Fig. 1. IFO 95 is a PN-IFO inside the field of view. The format for these images is the same as that of Fig. 1.

Downloaded from https://academic.oup.com/mnras/article/528/3/4657/7591325 by guest on 19 February 2024

**Figure B3.** *Continued.*

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.