



Kent Academic Repository

Gavilev, Olena V., Katerynych, Oleg O., Ionov, Igor A., Dekhtiarova, Olena O., Griffin, Darren K. and Romanov, Michael N (2024) *Overview of triticale use for poultry feeding*. . Multidisciplinary Digital Publishing Institute (MDPI), Basel, Switzerland encyclopedia article, Encyclopedia (online), Entry 55265.

Downloaded from

<https://kar.kent.ac.uk/105057/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://encyclopedia.pub/entry/55265>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

This entry is adapted from <https://doi.org/10.3390/encyclopedia4010027>

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Triticale: A General Overview of Its Use in Poultry Production

Subjects: Agriculture, Dairy & Animal Science

Contributor: Olena V. Gaviley, Oleg O. Katerynych, Igor A. Ionov, Olena O. Dekhtiarova, Darren K. Griffin, Michael N. Romanov

Triticale, a hybrid of wheat and rye, is one of the most promising grain crops. In terms of productivity, the level of metabolizable energy, and the composition of essential amino acids, triticale surpasses rye and is not inferior to wheat. It is resistant to the most dangerous diseases and pests. In terms of nutritional value, triticale can compete with wheat, corn, sorghum, and barley. The presence, however, of antinutrients in triticale such as non-starch polysaccharides, alkylresorcinols, and trypsin inhibitors significantly reduces the biological value of this crop. In the global practice of compound feed production, there are many methods and technologies for processing grain raw materials to increase their nutritional value. Enzymatic treatment and extrusion technologies are worthy of special attention. The high content of triticale in the compound feed of poultry breeder flocks should be used effectively, taking into account the characteristics of triticale varieties and climatic conditions. An optimal triticale level in feed (15% for layer and broiler chicks) may improve body weight gain and reduce feed costs when raising replacement young stock. Layer breeder flocks fed a 20% triticale-based diet may have increased egg production, high viability, and flock uniformity. Producing triticale–soy and triticale–sunflower extrudates and supplementing the diet of poultry flocks with essential amino acids represent promising avenues for maximizing the benefits of triticale. Innovative methods of achieving this goal should be further developed and put into practice, particularly given the expansion of triticale's cultivation areas.

Keywords: triticale ; poultry farming ; feeds ; chickens ; amino acids ; antinutrients ; enzymatic treatment ; extrudates

Poultry producers consistently face the challenges of increasing flock productivity and reducing production costs. Additional issues to be kept in mind include disease control (with a minimum use of antibiotics) for both animals and food crops, animal welfare, and environmental concerns. Balanced feeding plays one of the most crucial roles in poultry production. It needs to meet all the needs of the bird in accordance with age and productivity, providing a diet consisting of the required content of basic nutrients, energy, and biologically active substances including, proteins, amino acids, and carbohydrates ^{[1][2][3][4][5][6][7][8][9][10][11][12][13][14][15]}. Moreover, in order to expand the fodder base in the worldwide poultry industry, the sustained expansion of the harvesting area of cereal crops that are more resistant to climate change is essential ^{[16][17][18]}.

Cereals are the main component of poultry feed. Conventionally, corn and wheat are most widely used for this purpose. Barley is utilized to a lesser extent, with rye and oats in smaller quantities. In recent years, the share of traditional cereals in compound feed for poultry has lowered in developed countries from 69–70% to 40–50%. This is due to both secondary ingredients of industrial processing and non-traditional feeds ^{[16][19]}. Alongside these conventional components, new cereal crops have appeared in poultry diets ^{[20][21][22][23]}, such as sorghum and triticale (*× Triticosecale* Wittmack) (**Figure 1**). The latter is a promising crop and, with its rational use, can be a valuable compound feed ingredient and successfully partially replace the cereal crops traditionally used ^{[17][23][24][25][26][27][28][29][30][31]}. This becomes especially important in connection with the growing share of cereals that will be used for the production of biofuels, i.e., ethanol, biodiesel, and solid fuel pellets (e.g., ^{[32][33][34][35]}). The purpose of this review is to consider the nutritional, economic, and environmental benefits (as well as the drawbacks) of the rye–wheat hybrid triticale in poultry production.




Figure 1. Phenotypes of wheat (*Triticum aestivum*; left), rye (*Secale cereale*; middle), and their hybrid, triticale (right): (a–c) cultivation fields; (d–f) single ears; and (g) seeds. Image sources: (a) [https://commons.wikimedia.org/wiki/File:Parcelle_De_bl%C3%A9_REGAIN_\(Yvelynes\)_Cl_J_Weber_\(23787338000\).jpg](https://commons.wikimedia.org/wiki/File:Parcelle_De_bl%C3%A9_REGAIN_(Yvelynes)_Cl_J_Weber_(23787338000).jpg), CC-BY-2.0 (accessed on 10 February 2024); (b) <https://commons.wikimedia.org/wiki/File:Rozh.jpg>, Ivan Shishkin (1878), CC-PD-Mark (accessed on 10 February 2024); (c) <https://commons.wikimedia.org/wiki/File:Triticalefeld.jpg>, CC-BY-SA-3.0 (accessed on 10 February 2024); (d) https://commons.wikimedia.org/wiki/File:Frumento_Tenero_Rieti.jpg, CC-BY-3.0 (accessed on 10 February 2024); (e) [https://commons.wikimedia.org/wiki/File:Secale_cereale_\(Roggen\)-2008b.jpg](https://commons.wikimedia.org/wiki/File:Secale_cereale_(Roggen)-2008b.jpg), CC-BY-SA-3.0,2.5,2.0,1.0 (accessed on 10 February 2024); (f) https://commons.wikimedia.org/wiki/File:LPCC-623-Espiga_de_triticale.jpg, CC-BY-SA-3.0 (accessed on 10 February 2024); and (g) https://commons.wikimedia.org/wiki/File:Wheat,_rye,_triticale_montage.jpg, PD USDA (accessed on 10 February 2024).

References

- Wu, G.; Bazer, F.W.; Dai, Z.; Li, D.; Wang, J.; Wu, Z. Amino acid nutrition in animals: Protein synthesis and beyond. *Annu. Rev. Anim. Biosci.* 2014, 2, 387–417.
- He, W.; Li, P.; Wu, G. Amino acid nutrition and metabolism in chickens. *Adv. Exp. Med. Biol.* 2021, 1285, 109–131.
- Grela, E.R.; Kowalczyk-Vasilev, E.; Świątkiewicz, M.; Skiba, G. Barley, triticale, or rye? The type of grain can affect the growth performance and meat quality of sustainable raised pigs. *Animals* 2023, 13, 1331.
- Ibatullin, I.I.; Kryvenok, M.Y.; Panasenko, Y.O.; Ilchuk, I.I. Digestibility of feed nutrients and methionine balance in the body of chickens of the parent flock at different levels of methionine in the diets. *Sučasne Ptahivnictvo Mod. Poult. Farming* 2013, 5, 2–5. (In Ukrainian)
- Otchenashko, V.V. Productivity of meat quails fed compound feeds with different levels of crude protein. *Naučnye Tr. SWorld Sci. Pap. SWorld* 2013, 4, 79–87. (In Ukrainian)
- Ionov, I.A. Vitamin E as a means of improving the quality of poultry products and the antioxidant status of the body. *Visnik agrarnoї nauki Bull. Agric. Sci.* 2010, 4, 37–39. Available online: https://agrovisnyk.com/oldpdf/visnyk_04_2010.pdf#page=37 (accessed on 20 November 2023). (In Ukrainian).
- Lemesheva, M.M. *Poultry Feeding. Slobozhanshchyna*: Sumy, Ukraine, 2003. (In Ukrainian)
- Fagundes, N.S.; Milfort, M.C.; Williams, S.M.; Da Costa, M.J.; Fuller, A.L.; Menten, J.F.; Rekaya, R.; Aggrey, S.E. Dietary methionine level alters growth, digestibility, and gene expression of amino acid transporters in meat-type chickens. *Poult. Sci.* 2020, 99, 67–75.
- Tarasewicz, Z.; Gardzielewska, J.; Szczerbińska, D.; Ligocki, M.; Jakubowska, M.; Majewska, D. The effect of feeding with low-protein feed mixes on the growth and slaughter value of young male Pharaoh quails. *Arch. Anim. Breed.* 2007, 50, 520–530.
- Surai, P.F.; Fisinin, V.I. Antioxidant Systems of the Body: From Vitamin E to Polyphenols and Beyond. In *Proceedings of the 35th Western Nutrition Conference*, Edmonton, AB, Canada, 24–25 September 2014; pp. 265–277. Available online: <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20163076348> (accessed on 9 February 2024).
- Lemesheva, M. A feed mixture of varying structure for turkeys. *Pticevodstvo Poult. Farming* 1991, 11, 11–13. Available online: <https://www.cabidigitallibrary.org/doi/full/10.5555/19931456675> (accessed on 9 February 2024). (In Russian).

12. Wilkinson, S.J. Big Data for Poultry—What Is Possible? In Proceedings of the 29th annual Australian Poultry Science Symposium, Sydney, Australia, 4–7 February 2018; Volume 29, pp. 152–158. Available online: <https://poultry-research.sydney.edu.au/wp-content/uploads/2019/08/APSS-2018-Proceedings-Final.pdf#page=167> (accessed on 10 February 2024).
13. Fearnside, P.M. Soybean cultivation as a threat to the environment in Brazil. *Environ. Conserv.* 2001, 28, 23–38.
14. Zerjal, T.; Härtle, S.; Gourichon, D.; Guillory, V.; Bruneau, N.; Laloë, D.; Pinard-van Der Laan, M.H.; Trapp, S.; Bed'hom, B.; Quéré, P. Assessment of trade-offs between feed efficiency, growth-related traits, and immune activity in experimental lines of layer chickens. *Genet. Sel. Evol.* 2021, 53, 44–60.
15. Pesti, G.M. Impact of dietary amino acid and crude protein levels in broiler feeds on biological performance. *J. Appl. Poult. Res.* 2009, 18, 477–486.
16. Kyrychenko, V.V.; Maklyak, K.M.; Varennyk, B.F.; Kutishcheva, N.M.; Trotsenko, V.I. The practical characteristics expressing of three-linear sunflower hybrids in different agro-climatic zones of Ukraine. *Visn. Sums'kogo Nac. Agrar. Univ. Ser. Agron. Biol. Bull. Sumy Natl. Agrar. Univ. Ser. Agron. Biol.* 2016, 9, 129–133. Available online: <https://web.archive.org/web/20220325181928/http://repo.snau.edu.ua/bitstream/123456789/5660/1/32.pdf> (accessed on 20 November 2023). (In Ukrainian).
17. Kchaou, R.; Benyoussef, S.; Jebari, S.; Harbaoui, K.; Berndtsson, R. Forage potential of cereal–legume mixtures as an adaptive climate change strategy under low input systems. *Sustainability* 2023, 15, 338.
18. Masenya, T.I.; Mlambo, V.; Mnisi, C.M. Complete replacement of maize grain with sorghum and pearl millet grains in Jumbo quail diets: Feed intake, physiological parameters, and meat quality traits. *PLoS ONE* 2021, 16, e0249371.
19. Gaviley, O.V. Efficiency of the Use of Triticale in Fodder for Breeding Chickens. Ph.D. Thesis, State Poultry Research Station of NAAS, Kharkov State Zooveterinary Academy, Kharkiv, Ukraine, 2018. Available online: <https://hdzva.edu.ua/svr/wp-content/uploads/sites/34/2018/06/dysertatsiya-gavilej.pdf> (accessed on 20 November 2023). (In Ukrainian).
20. Smith, R.L.; Jensen, L.S.; Hoveland, C.S.; Hanna, W.W. Use of pearl millet, sorghum, and triticale grain in broiler diets. *J. Prod. Agric.* 1989, 2, 78–82.
21. Zhukorskyi, O.; Prytulenko, O.; Tereshchenko, O.; Bratishko, N. The Presence of Soybean, Triticale and Sorghum in Compound Feed for Poultry is Fully Justified. *Zerno i khliv Grain Bread* 2012, 2, 14–16. Available online: https://www.researchgate.net/publication/342832208_Zukorskij_O_Pritulenko_O_Teresenko_O_Bratisko_N_Prisutnist_soi_tritikale_i_sorgo_2012_-_No_2_-_S_14-16?tp=eyJjb250ZXh0Ijp7ImZpcnN0UGFnZSI6InByb2ZpbGUlLCJwYXVlIjoic2VhcmNoliwicG9zaXRpb24iOiJwYXVlSGVhZGVyIn19 (accessed on 20 November 2023). (In Ukrainian).
22. Rózewicz, M. Production, use and efficiency of utilising grains of various cereal species as feed resources for poultry production. *Pol. J. Agron.* 2019, 38, 66–74.
23. Farahat, M.; Badawi, M.; Hussein, A.; Attia, G. Effect of replacing dietary corn by sorghum on the growth performance, shank skin pigmentation, carcass traits, caecal microflora and nutrient digestibility of broiler chickens. *Int. J. Poult. Sci.* 2020, 19, 424–431.
24. Bratishko, N.; Gaviley, Y.; Pritulenko, O.; Tereshchenko, A. Triticale in compound feed for chicken breeders. *Pticevod. Poult. Farming* 2008, 9, 30–32. Available online: <https://www.elibrary.ru/item.asp?id=11566990> (accessed on 20 November 2023). (In Russian).
25. Bratishko, N.I.; Gritsenko, R.B.; Pritulenko, O.V.; Tereshchenko, A.V. Triticale in Compound Feed for Breeding Chickens. *Sučasne Ptahivnictvo Mod. Poult. Farming* 2008, 4, 3–4. Available online: https://www.researchgate.net/publication/342764397_Bratisko_NI_Gricenko_RB_Pritulenko_OV_Teresenko_AV_Tritikale_v_kombikormah_virobnicij_zurnal_-_Kiiv_Nacionalnij_agrarnij_universitet_Institut_ptahivnictv (accessed on 20 November 2023). (In Russian).
26. Bratishko, N.I.; Tereshchenko, A.V.; Pritulenko, O.V.; Gaviley, E.V. Influence of Triticale on the Productivity of Chickens When it Is Included in Compound Feed in Combination with Corn or Wheat. In *Advances in Modern Poultry Science, Proceedings of the 16th International Conference, Sergiyev Posad, Russia, 19–21 May 2009*; Fisinin, V.I., Egorov, I.A., Vasilyeva, T.V., Eds.; WPSA—Russian Branch, RAAS, All-Russia Poultry Science and Technology Institute: Sergiyev Posad, Russia, 2009; pp. 83–84. (In Russian)
27. McGoverin, C.M.; Snyders, F.; Muller, N.; Botes, W.; Fox, G.; Manley, M. A review of triticale uses and the effect of growth environment on grain quality. *J. Sci. Food Agric.* 2011, 91, 1155–1165.
28. Hermes, J.C.; Johnson, R.C. Effects of feeding various levels of triticale var. Bogo in the diet of broiler and layer chickens. *J. Appl. Poult. Res.* 2004, 13, 667–672.
29. Djekic, V.; Pandurevic, T.; Mitrovic, S.; Radovic, V.; Milivojevic, J.; Djermanovic, V. Nutritional Value of Triticale (Trijumf) for Broiler Diets. In *Proceedings of the Third International Scientific Symposium “Agrosym Jahorina 2012”, Jahorina, Bosnia and Herzegovina, 15–17 November 2012*; pp. 15–17. Available online: <https://www.cabidigitalibrary.org/doi/pdf/10.5555/20143100079> (accessed on 9 February 2024).
30. Lim, C.; Poaty Ditengou, J.; Ryu, K.; Ku, J.; Park, M.; Whiting, I.M.; Pirgozliev, V. Effect of replacing maize with different levels of triticale on laying hen performance, egg quality, yolk fatty acid profile and blood options. *J. Anim. Feed Sci.* 2021, 30, 360–366.

31. Osek, M.; Milczarek, A.; Janocha, A.; Świnarska, R. Effect of triticale as a partial or complete wheat and maize substitute in broiler chicken diets on growth performance, slaughter value and meat quality. *Ann. Anim. Sci.* 2010, 10, 275–283. Available online: <https://www.cabidigitallibrary.org/doi/full/10.5555/20113025674> (accessed on 10 February 2024).
32. Gradziuk, P.; Jończyk, K.; Gradziuk, B.; Wojciechowska, A.; Trocewicz, A.; Wysokiński, M. An economic assessment of the impact on agriculture of the proposed changes in EU biofuel policy mechanisms. *Energies* 2021, 14, 6982.
33. Liu, X.; Unaegbunam, E.; Stuart, D.T. Co-production of isobutanol and ethanol from prairie grain starch using engineered *Saccharomyces cerevisiae*. *Fermentation* 2021, 7, 150.
34. Wilczyński, D.; Talaśka, K.; Wałęsa, K.; Wojtkowiak, D.; Bembenek, M. Experimental study on the cutting process of single triticale straws. *Materials* 2023, 16, 3943.
35. Obidziński, S.; Cwalina, P.; Kowczyk-Sadowy, M.; Krasowska, M.; Sienkiewicz, A.; Faszczewski, D.; Szyszlak-Bargłowicz, J. The use of bread bakery waste as a binder additive in the production of fuel pellets from straw. *Energies* 2023, 16, 7313.

Retrieved from <https://encyclopedia.pub/entry/history/show/124618>