



# Kent Academic Repository

Ramachandran, R, Hazarika, A, Gupta, S, Nag, S, Meka, J K, Thakur, Tejender S, Yashonath, S, Vishwakarma, G, Chou, S-L, Wu, Y-J and others (2024) *Amorphous 1-propanol interstellar ice beyond its melting point*. *Monthly Notices of the Royal Astronomical Society*, 530 (1). pp. 1027-1034. ISSN 0035-8711.

## Downloaded from

<https://kar.kent.ac.uk/105367/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1093/mnras/stae759>

## This document version

Author's Accepted Manuscript

## DOI for this version

## Licence for this version

CC BY (Attribution)

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

### Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

# Amorphous 1-propanol interstellar ice beyond its melting point

R. Ramachandran<sup>1\*</sup>,<sup>†</sup>, A. Hazarika<sup>2,†</sup>, S. Gupta,<sup>1</sup> S. Nag,<sup>2</sup> J. K. Meka,<sup>1</sup> Tejender S. Thakur,<sup>3</sup> S. Yashonath,<sup>4</sup> G. Vishwakarma,<sup>5</sup> S. -L. Chou,<sup>6</sup> Y. -J. Wu,<sup>6</sup> P. Janardhan,<sup>1</sup> B. N. Rajasekhar,<sup>7</sup> Anil Bhardwaj,<sup>1</sup> N. J. Mason,<sup>8</sup> B. Sivaraman,<sup>1‡</sup> and Prabal K. Maiti<sup>2§</sup>

<sup>1</sup>Physical Research Laboratory, Ahmedabad, India.

<sup>2</sup>Center for Condensed Matter Theory, Department of Physics, Indian Institute of Science, Bangalore, India.

<sup>3</sup>Biochemistry and Structural Biology Division, CSIR- Central Research Drug Institute, Lucknow, India.

<sup>4</sup>Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, India.

<sup>5</sup>DST Unit of Nanoscience (DST UNS) and Thematic Unit of Excellence (TUE), Department of Chemistry, Indian Institute of Technology Madras, Chennai, India.

<sup>6</sup>National Synchrotron Radiation Research Center, Hsinchu, Taiwan.

<sup>7</sup>Atomic and Molecular Physics Division, Bhabha Atomic Research Centre, Mumbai, India.

<sup>8</sup>School of Physical Sciences, University of Kent, CT2 7NH, Canterbury, UK.

Accepted XXX. Received YYY; in original form ZZZ

## ABSTRACT

The recent discovery of 1-propanol (CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>OH) in the interstellar medium (ISM) is of tremendous interest since fatty alcohols have been proposed as constituents of proto-cell membranes. Motivated by this discovery, we present the laboratory mid-infrared (MIR) and vacuum ultra-violet (VUV) absorption spectra of 1-propanol ice under astrochemical conditions, mimicking an icy mantle on cold dust in the ISM. Both MIR and VUV spectra were recorded at ultra-high vacuum (UHV) of  $\sim 10^{-9}$  mbar and at temperatures ranging from 10 K to sublimation. The morphology of the 1-propanol ice deposited at 10 K was amorphous. By warming the ice to temperatures of 140 K and above, with subsequent recording of IR spectra, we observe complete sublimation of 1-propanol molecules from the substrate around 170 K. No amorphous-to-crystalline phase change was observed upon warming to higher temperatures. Additionally, We observe the IR and VUV signatures of 1-propanol ice on the substrate well beyond its melting point (147 K). To the best of our knowledge, this is the first reported observation of a molecular ice staying well beyond its melting point under such conditions. This result shows that the morphology of icy mantles on ISM cold dust grains is more complex than previously thought. Our atomistic molecular dynamics (MD) simulations capture the experimental trends and shed light on the microscopic origin of this unusual phase behaviour of 1-propanol.

**Key words:** astrochemistry – molecular data – infrared: general – ISM: molecules – methods: statistical – software: simulations

## 1 INTRODUCTION

Since the identification of methanol and ethanol (Ball et al. 1970; Zuckerman et al. 1975), it has become well-established that molecules containing the hydroxyl (OH) group are present in the interstellar medium (ISM) (e.g. Turner & Apponi (2001); Hollis et al. (2004)). These molecules are also known to be the constituents of the icy surfaces found within the solar system (e.g. Despois (1992); Crovisier et al. (2004)). In fact, their role in the chemistry of complex molecular evolution is quite central as they hold large amounts of hydrogen within their structure. The physicochemical nature of methanol and ethanol in the ISM is quite well explored (Ehrenfreund & Charnley 2000). The formation of these simple alcohols is known to occur through the irradiation of mixtures of simpler molecules such as carbon monoxide, ammonia and water (Kasamatsu et al.

1997; Watanabe & Kouchi 2002; Hiraoka et al. 2006). Additionally, a pathway via hydrogenation of formaldehyde is also known to synthesise methanol (Fuchs et al. 2009).

With the discovery of simple alcohols, the search for larger alcohols has intensified. The detection of propanal in the ISM (Hollis et al. 2004) provided evidence of the possible presence of fatty alcohols in the ISM, as aldehydes can be considered proxy molecules for their corresponding alcohols. This was further supported by the identification of propanol in the comet CG-67P (Altwegg et al. 2017). Laboratory experiments have also demonstrated that propanol can be synthesized alongside simple alcohols by irradiating a mixture of carbon monoxide and methane with energetic electrons (Abplanalp & Kaiser 2019). A hydrogenation pathway from carbon monoxide during its freeze-out phase, in the presence of acetylene, is also known to synthesize 1-propanol (Altwegg et al. 2017).

Given that propanol can be synthesized through multiple pathways under the cold dust conditions of the ISM, its presence in the ISM was anticipated to be within detectable limits. As expected, focused observations confirmed the presence of 1-propanol in the ISM (Jiménez-Serra et al. 2022). This discovery is of tremendous interest

\* E-mail: ragav@prl.res.in

† Co-first authors

‡ E-mail: bhala@prl.res.in

§ E-mail: maiti@iisc.ac.in

since fatty alcohols have been proposed as constituents of proto-cell membranes (Ruiz-Mirazo et al. 2014). However, our understanding of the physico-chemical nature of 1-propanol in the ISM remains quite limited. Although reports exist on the infrared spectra of this molecule in both liquid and ice phases (Doroshenko et al. 2012; Hudson & Moore 2018; Qasim et al. 2019), detailed experiments on the temperature-dependent behaviour of 1-propanol ice are scarce. Therefore, we conducted an experimental investigation of this fatty alcohol under astrochemical icy conditions.

To support and interpret the experimental findings, we also employed molecular dynamics (MD) simulations for the 1-propanol system. While computer simulations have been extensively used to investigate this molecule in its liquid and gas phases (Zangi 2018; Pethes et al. 2021), research into its ice phase remains scarce. Previously, we have studied the ice phase of another astrochemical complex organic molecule (COM) methanol using molecular dynamics simulation (Nag et al. 2023) and explained the recent experimental observation of the same molecule (Luna et al. 2018). By employing computational techniques, we aim not only to reproduce and elucidate the experimental results but also to delineate the physico-chemical properties of 1-propanol. A comprehensive understanding of 1-propanol, achieved through a combination of experimental and computational approaches, is crucial in astrochemistry due to its relevance as a precursor to more COMs and its potential role as an indicator of life's fundamental components (Ruiz-Mirazo et al. 2014). This study will not only enhance our fundamental knowledge of this molecule but also shape the trajectory of future studies of similar organic molecules.

## 2 METHODOLOGY

Three different techniques were employed to investigate the 1-propanol ice under astrochemical conditions. The details of each technique are elaborated below. For all the experiments 1-propanol in liquid form, procured from Merck (purity  $\geq 99.9\%$ ), was used.

### 2.1 Vacuum Ultraviolet Spectroscopy

VUV experiments of 1-propanol ices were conducted using VUV light from beamline BL-03 of the Taiwan Light Source (TLS) at the National Synchrotron Radiation Research Centre (NSRRC), Taiwan. One of the end stations of the beamline is connected to an ultrahigh vacuum chamber containing a closed cycle helium cryostat with a VUV transparent lithium-fluoride (LiF) window attached to its cold finger using a copper substrate holder. This window is cooled down to 10 K and serves as the interstellar dust analogue. Detailed discussions of the experimental setup, liquid sample preparation, and the spectral acquisition have been presented in Lu et al. (2008), and the chamber setup is similar to one shown in Fig. 2 of Ramachandran et al. (2023).

A VUV spectrum, spanning 115 nm to 220 nm with a step size of 1 nm, of the blank LiF window at 10 K was recorded at the beginning of each experiment. This spectrum was used as  $I_0$  for the respective experiments. The sample was then deposited on the LiF window using a deposition technique similar to that explained in the previous section. Spectra recorded after deposition are used as  $I$ . From  $I_0$  and  $I$ , the VUV spectra of 1-propanol ice were calculated using the Beer-Lambert law. After deposition and obtaining a spectrum at 10 K, the ice was heated to higher temperatures until sublimation, with VUV spectra recorded at regular intervals.

### 2.2 Mid-Infrared Spectroscopy

The astrochemical icy mantle conditions were recreated using the Simulator for Astromolecules at Low Temperature (SALT) facility housed at the Physical Research Laboratory, Ahmedabad. The experimental system can achieve pressures of  $10^{-10}$  mbar at 10 K. A zinc-selenide (ZnSe) substrate is used as the ISM dust analogue, which, when placed on the substrate holder at the tip of the closed-cycle helium cryostat, reaches temperatures as low as 10 K while the system is under ultrahigh vacuum conditions. A cryogenic temperature controller (Lakeshore) regulates the sample's temperature and heating rate. A detailed description of this setup can be found in Ramachandran et al. (2023). In these experiments, a heating rate of  $5 \text{ K min}^{-1}$  was primarily employed. The deposited molecular ices were probed *in situ* using IR spectroscopy (Nicolet iS 50 FTIR Spectrometer), in the mid-IR range ( $4000 \text{ cm}^{-1}$  to  $650 \text{ cm}^{-1}$ ; resolution  $4 \text{ cm}^{-1}$  and integration time 86 seconds).

1-propanol from the sample holder was vapour deposited (through a differential pressure technique) onto the ZnSe window maintained at 10 K. Deposition was carried out for approximately 50 seconds at a pressure of around  $7 \times 10^{-9}$  mbar. A post-deposition IR spectrum was recorded at 10 K. The resultant ice was then heated at the rate of  $5 \text{ K min}^{-1}$  in 10 K increments until 140 K, with subsequent recording of IR spectra. After 140 K, the sample was heated to 145 K, 146 K, 147 K and 150 K, with IR spectra recorded for each temperature. Beyond 150 K the sample was heated in 5 K increments, and IR spectra were recorded until sublimation. Deposition was also performed at 130 K, 145 K, 150 K, 160 K, 170 K, 180 K, and 185 K. In all these cases, the ice was heated at the rate of  $5 \text{ K min}^{-1}$  until sublimation, with subsequent recording of IR spectra.

### 2.3 Computational Details

In order to obtain and better understanding of the phase behaviour of 1-propanol, a vacuum-deposited film of 1-propanol on potassium-bromide (KBr) surface was simulated, and subsequently, the density of states (DoS) spectra was calculated in simulation as a function of temperature. To model 1-propanol, the all-atom OPLS-AA (Jorgensen & Tirado-Rives 2005; Dodda et al. 2017b) force-field parameters were used. The LigParGen (Dodda et al. 2017a) topology generator for the aforementioned force field was utilized to generate the topology in GROMACS-compatible format, which was subsequently used for all MD simulations. Further, to take into account the Coulombic interactions, Electrostatic Potential (ESP) charges were used as atomic partial charges. These charges were obtained during Crystal Structure Prediction (CSP) calculation. All the force field parameters are tabulated in Table S1-S5 in the Supporting Information (SI). For modelling of KBr, the potential parameters developed by Barbosa (Fuentes-Azcatl & Barbosa 2018) were used. The same potential parameters for KBr had been used in our earlier work on methanol adsorbed to KBr substrate (Nag et al. 2023). Lorentz-Berthelot mixing rules were applied for nonbonded interactions between 1-propanol and KBr.

For benchmarking the applicability of OPLS-FF parameters, several thermodynamic and spectroscopic properties of 1-propanol in liquid and gas phases such as bulk density, specific heat and vibrational spectra under atmospheric pressure were obtained in simulation and compared with known values of these properties (NIST 2023) (Stromsoe et al. 1970). The methodology is described and the benchmarking results are tabulated in the SI. Using OPLS-FF as the forcefield (FF) of choice, a very good agreement could be seen between simulation and experiment (see Table S6 of SI).

After benchmarking the FF against 1-propanol, the molecule of interest, the simulated deposition of 1-propanol was performed using the Python package PyThinFilm (Stroet et al. 2022). PyThinFilm, or PyTF, acts as a Python-based frontend to run classical MD deposition simulations using GROMACS (Abraham et al. 2015) as the molecular dynamics engine. The deposition was carried out using the ‘vacuum deposition’ method which can be described as the insertion of the molecule at a distance (hence called insertion distance) away from the substrate, and letting the dynamics run until the molecule settles on the substrate; the entire system is kept under a tall column of vacuum during this entire step. This step in the process is carried out one by one on a per-molecule basis until the desired number of molecules have been successfully deposited on the surface. For our system, the insertion distance was set to 8 nm, while the column of the vacuum was chosen to be 20 nm long. All the details of the deposition simulations are provided in the SI.

All the MD simulations were performed using the GROMACS 2023.1 (Abraham et al. 2015) molecular dynamics package. The deposited system of 1-propanol in vacuum on the KBr substrate was simulated in the constant temperature (NVT) ensemble using the stochastic velocity rescaling thermostat (Bussi et al. 2007) with temperature coupling constant value fixed at 1 ps. A 1 fs integration timestep was employed to simulate the system with an updated neighbour list every 100 timesteps. For short-range LJ and electrostatic interaction, the cut-off distance was set to 12 Å. Long-range electrostatics were handled via the Particle Mesh Ewald (PME) (Darden et al. 1993) method using fourth-order interpolation (Essmann et al. 1995) while the Fourier spacing was kept at 0.16 nm.


At all temperatures, the equilibration runs were carried out for at least 20 ns and extended further depending on the convergence of potential energy to ensure thorough thermalization. This was followed by a 30 ps simulation in the NVT ensemble to compute the corresponding density of states using the 2PT algorithm (Lin et al. (2003, 2010)). During the 30 ps NVT simulation run to obtain DoS, the coordinates, velocities and energies were stored at every 2 fs.

### 3 RESULTS AND DISCUSSION

#### 3.1 Experimental Results (VUV & IR)

VUV spectra of 1-propanol ice, deposited at 10 K and subsequently warmed to higher temperatures, are shown in Figure 1 for selected temperatures. At 10 K, the ice exhibits a broad absorption, with no fine structures visible between 115 nm – 170 nm. As the ice is warmed to higher temperatures, the spectra reveal a peak around 142 nm at 120 K, which intensifies upon warming to 130 K and 140 K. The spectrum recorded upon warming the ice to 147 K (melting point of 1-propanol ice at atmospheric pressure, Tschamler et al. (1949)) remains similar to the spectrum at 140 K, with a slight reduction in absorbance intensity. We compare the melting point at atmospheric pressure with the ice characteristics at UHV because even though melting points at UHV are unavailable in literature for most molecules, it can be conjectured that they will be lower than the melting point at atmospheric pressure. Hence, in this paper, melting point implies melting point at atmospheric pressure. The heating was continued to temperatures as high as 170 K, revealing weak absorbance in the lower wavelength range of 115 nm – 145 nm.

A similar experiment employing infrared spectral probing was conducted on 1-propanol ice. The resulting spectra at a few selected temperatures are presented in Figure 2. The spectrum of the 1-propanol ice at 10 K exhibited all the expected peaks, such as



**Figure 1.** VUV spectra of 1-propanol ice recorded after deposition at 10 K. The deposited ice was then warmed to higher temperatures and spectra recorded at specific temperatures until sublimation.


those corresponding to OH symmetric stretching, CH<sub>3</sub> and CH<sub>2</sub> symmetric stretching and CH<sub>2</sub> asymmetric stretching, CH<sub>3</sub> rocking and CH<sub>2</sub>-O stretching vibrations. The ice was then warmed to higher temperatures, up to 175 K, with subsequent recording of spectra.

Upon close examination of the spectral signatures, we focus on the OH symmetric stretching and CH<sub>3</sub> rocking vibrations. As evident from Figure 2 (a), significant changes are observed in these two peaks as the ice is warmed from 10 K to 175 K. The peak corresponding to the OH stretching vibration at 10 K was much broader and featured a shoulder peak. As the ice was warmed, the shoulder peak gradually disappeared, and the OH stretching vibration appeared as a single peak, significantly narrower than the peak observed at 10 K. However, the width of the OH stretching vibration was observed to widen slightly as the ice was further warmed to 160 K. At 175 K, the peak intensity had reduced, but the peak position and width remained similar to the spectrum at 160 K.

The band at 1069 cm<sup>-1</sup> assigned to the CH<sub>3</sub> rocking vibration had a shoulder peak, around 1058 cm<sup>-1</sup>, at 10 K, which was observed to intensify. Notably, in the spectrum recorded at 50 K, the intensity increase was significant. The intensities of these two peaks were nearly equal in the spectrum recorded at 80 K. However, the intensity of the 1058 cm<sup>-1</sup> peak continued to increase, as observed in the spectra recorded at higher temperatures, from 120 K to 160 K. At 175 K, despite a significant reduction in absorbance, the intensity of the 1058 cm<sup>-1</sup> peak remained well above that of the 1069 cm<sup>-1</sup> peak. No significant changes were observed in the remaining peaks in the mid-IR spectra of 1-propanol ice from 10 K to 175 K.

While VUV spectra revealed the presence of 1-propanol ice well beyond the melting point of the ice, the only observable change was the broad spectral signature with a slight intensity variation at 140 K. However, in the IR spectral probing of 1-propanol ice, significant changes were observed in the OH stretching and CH<sub>3</sub> rocking vibrations, indicative of molecular reorientation occurring as the amorphous ice formed at 10 K was warmed to higher temperatures. The reduction in the breadth of the OH band is attributed to ice compaction. The intensity reversal observed between the 1071 cm<sup>-1</sup> and 1057 cm<sup>-1</sup> band is also indicative of ice compaction as the ice tended towards a more ordered structure. Notably, the signatures for crystalline 1-propanol ice were entirely absent in the IR spectra, as no sharpening or splitting of the bands was observed.

The deposition was also performed at 130 K, 145 K, 150 K, 160 K,



**Figure 2.** (a) MIR spectra of 1-propanol ice recorded after deposition at 10 K. The ice was then warmed to higher temperatures with subsequent recording of spectra at specific temperatures until sublimation. A spectrum around the melting point of 1-propanol ( $\sim 147$  K) was recorded. Beyond the melting point a significant amount of 1-propanol was observed to be present until sublimation around 175 K. (b) MIR spectra of 1-propanol ice deposited at 10 K and higher temperatures ranging from 130 K to 185 K respectively. These correspond to temperatures a few K above and below the melting point of 1-propanol. Molecules of 1-propanol are found to adhere and form layers on the dust analogues even beyond its melting point.

170 K, 180 K and 185 K, with the corresponding IR spectra shown in Figure 2 (b). In each case, the ice was warmed until sublimation, and IR signatures of the ice were observed up to 185 K. However, at 180 K and 185 K, the ice could only remain on the substrate for approximately one minute. The deposited ice was amorphous and maintained its amorphous state throughout the experiments.

It is noteworthy that we were able to record the VUV and IR spectrum of 1-propanol well beyond its melting point (147 K), and the ice was observed to remain amorphous throughout the warming process, from 10 K to 175 K, until sublimation. This marks the first observation of a molecule staying on a cold substrate well beyond its melting point in a UHV chamber. It is also worth mentioning that 2-propanol, a positional isomer of 1-propanol, does not show such a behaviour. It does crystallize (120 K), and sublimates (150-155 K) before melting point (184 K) (Ayling et al. 2017). In Table S8 of

the SI, we compare the melting point and sublimation temperatures under UHV conditions for molecular ices formed on cold substrates, analogous to astrochemical icy conditions. The comparison is made between samples that exist as liquids at the standard temperature and pressure (STP) condition and ices formed by extracting vapors from their liquid counterparts. To date, the molecular ices are known to either sublime or exhibit reversible phase change as they approach the melting point (Pavithraa et al. 2017). However, from the spectral signatures, it is clearly evident that 1-propanol is present well beyond its known melting point, even under UHV conditions. We suspect the low vapour pressure of 1-propanol at such low temperatures could be the reason for the presence of this molecule well beyond the melting point. This conclusion is supported by a series of experiments where 1-propanol ice was warmed from 10 K to higher temperatures, such as 150 K, 160 K, and 170 K. Rate of loss of 1-propanol molecules from the substrate under isothermal conditions was observed to follow the following order: 170 K > 160 K > 150 K.


## 3.2 Computational Results

### 3.2.1 Predicted crystal structure of 1-propanol

Under astrochemical icy conditions, most of the molecular ices, especially lighter molecules, are known to exhibit the amorphous-to-crystalline phase transition (eg. Yuan et al. (2016), Pavithraa et al. (2017), and many others) which naturally raises the question of its possibility for 1-propanol under similar conditions. As the crystal structure of 1-propanol is not yet experimentally known, the technique of crystal structure prediction to obtain the possible crystal structure of this molecule is key to investigating this phenomenon. In the context of comparing and understanding the possibility of phase transition between crystalline and amorphous phases, knowing the predicted crystal structure provides valuable insights. It serves as a reference point for understanding the differences between the two phases, from a structural and molecular point of view. Through comparisons between the crystal structure and the amorphous state of the deposited layer, one can pinpoint differences in molecular arrangement, bonding, or other structural aspects, shedding light on the factors responsible for the phase behaviour of the molecule under astrochemical conditions.

To obtain the initial crystal structure of 1-propanol, theoretical crystal structure prediction (CSP) calculations were run. The top 100 ranked structures obtained from the forcefield-based rigid CSP computations for 1-propanol were initially saved as candidates. The crystal structure search was performed using the ESPD-derived atomic charges considering  $Z' = 1$  in the top 22 space groups covering 95% of total occurrences in the Cambridge structural database. Input geometry and ESPD charges used in the CSP computation were obtained from gas phase DFT computations performed using meta-GGA M06-2X functional with 6-311++G(d,p) basis. From the 100 structures, the final choice was selected to be the best guess, as the unit cell was isostructural with *n*-pentanol (Ramírez-Cardona et al. 2005), another primary alcohol. Section S4 of SI contains the comprehensive procedure for predicting the crystal structure of 1-propanol using CSP.

The chosen crystal structure is a monoclinic unit cell as shown in Figure 3. The unit cell lengths are ( $a = 4.4354$  Å,  $b = 4.4702$  Å,  $c = 19.7348$  Å), and the angles ( $\alpha = \gamma = 90^\circ$ ,  $\beta = 95.1940^\circ$ ). To confirm the low-temperature stability of this crystal structure, a steepest descent energy minimization followed by an NPT simulation at 10 K, and atmospheric pressure was carried out for 10 ns with a timestep of 1 fs. During the simulation, the Berendsen thermostat



**Figure 3.** Predicted crystal structure for 1-propanol. Carbon, Oxygen and Hydrogen atoms are shown in cyan, red and white colour respectively.


and barostat with a temperature coupling were used with a 1.0 ps time constant. As shown in Figure S8 (shown in the SI), the cell lengths level off quickly after a few hundred picoseconds at 10 K, which provides assurance that the crystal structure is stable enough to hold the configuration at this temperature. To obtain temperature-dependent behaviour, the post-equilibration configuration at 10 K was slowly heated to the required temperature, and then equilibrated for a few nanoseconds, following which it was heated to the next required temperature. The complete step-by-step temperature ramp is outlined in Figure S6 (in the SI).

### 3.2.2 Simulated IR Spectra of 1-propanol and Comparison


The simulated density of states of 1-propanol crystal structure in bulk phase was obtained for different temperatures through the course of annealing starting from 10 K till 120 K. The spectra at the interval temperatures were computed for the range of 500 and 4000  $\text{cm}^{-1}$  (please see Figure S9 in the SI). The spectrum of the simulated solid at 10 K and 120 K was compared against the experimental IR spectrum at the same temperature, in Figure 4. There is a good agreement between the calculations and experiments in the peak positions, except the peak corresponding to the OH stretching vibrations, which appears as a broad peak centred around 3400  $\text{cm}^{-1}$  in the experiment, but in simulation, it appears as a shorter, sharper peak at  $\sim 3700 \text{ cm}^{-1}$ . The family of peaks in the 1100 - 850  $\text{cm}^{-1}$  range that shows a uniform shift of  $\sim 70 \text{ cm}^{-1}$  between simulated and experimental case correspond to the same vibrations:  $\text{CH}_3$  rocking,  $\text{CH}_3\text{-CH}_2$  stretching and  $\text{CH}_2\text{-O}$  stretching in both cases.

To interpret the experimentally observed behaviour of 1-propanol in relation to temperature from a molecular perspective, MD simulations were performed on a vacuum-deposited system of 400 1-propanol molecules on a KBr surface. To heat the system, a linear temperature ramp with a heating rate of 0.05 K/ps was employed. Following the deposition simulations and subsequent heating, the calculation of the DoS was carried out at various temperatures. The range of spectra for various temperatures is shown for the full range frequency range in Figure 5. To identify the various trends in the spectra, it is convenient to zoom into the (1500 - 700  $\text{cm}^{-1}$ ) frequency region. In this region, many trends can be observed in the form of the appearance or disappearance of peaks, and the change in the relative heights of the peaks. As highlighted in Figure 6, there are several peaks that show differences with respect to temperature in the 1500 - 700  $\text{cm}^{-1}$  range.

In Figure 6, a number of spectral features have been highlighted to




**Figure 4.** Comparison between the DoS of simulated bulk 1-propanol solid and experimentally obtained IR spectra of deposited 1-propanol at 10 K and 120 K.




**Figure 5.** Simulated DoS spectra of the system of 400 molecules of 1-propanol deposited on the KBr substrate throughout a range of temperatures.

show the temperature variation of these respective peaks. Amongst these, the pair of peaks labelled 'iv' is worth paying attention to, as they are the peaks corresponding to the  $\text{CH}_3$  rocking,  $\text{CH}_2\text{-CH}_2$  stretching and  $\text{CH}_2\text{-O}$  stretching vibrations. They are located at  $\nu_1 = 996 \text{ cm}^{-1}$  and  $\nu_2 = 961 \text{ cm}^{-1}$ . At 10 K, a lower wavelength peak is present and has a similar intensity to the higher wavelength peak. On heating,  $\nu_1$  begins to diminish in relative intensity while  $\nu_2$  gains intensity. At  $T > 120 \text{ K}$ ,  $\nu_1$  has disappeared completely and  $\nu_2$  has a higher intensity than at 10 K. A concise description of the peaks and their temperature-dependent behaviours are listed as follows:




**Figure 6.** 1600 - 600  $\text{cm}^{-1}$  range of the simulated IR spectra of the system of 400 molecules of 1-propanol deposited on the KBr substrate. Specific regions of interest have been marked and labelled.



**Figure 7.** Experimental infrared spectra zoomed in to 1600 - 600  $\text{cm}^{-1}$  range. The highlighted pair of peaks at 1069  $\text{cm}^{-1}$  and 1059  $\text{cm}^{-1}$  shows a trend of change in relative height that corresponds to a similar observation in the spectra of the simulated system.

- (i) Peak at 745  $\text{cm}^{-1}$  is absent below 100 K but appears at 100 K and becomes prominent with increasing temperature.
- (ii) Peak at 820  $\text{cm}^{-1}$  shows gradual disappearance with increasing temperature and is completely absent at  $T > 120$  K.
- (iii) Peak at 900  $\text{cm}^{-1}$  diminishes with increasing temperature and peak at 910  $\text{cm}^{-1}$  becomes more prominent as temperature increases. Both peaks are visible at 100 K and 120 K.
- (iv) Peak at 1000  $\text{cm}^{-1}$  disappears with increasing temperature. Absent at 146 K and above.
- (v) Peak at 1250  $\text{cm}^{-1}$  disappears with increasing temperature. Absent at 146 K and above.
- (vi) Peak at 1290  $\text{cm}^{-1}$  absent below 100 K, and gains intensity with increasing temperature.

From the experimentally obtained IR spectra, shown in Figure 2 and Figure 7, the highlighted pair of peaks at 1069  $\text{cm}^{-1}$  and 1059  $\text{cm}^{-1}$  show a temperature-dependent behaviour which is similar to the pair of peaks labelled as ‘iv’ in Figure 6. The lower wavelength peak shows decreasing intensity with heating in both scenarios, and the higher wavelength peak shows increasing intensity. The temperature




**Figure 8.** Comparison of the experimental IR spectra of 1-propanol (blue), with the density of states obtained from MD simulation of the deposited 1-propanol system (orange), as well as that of the deposited 1-propanol system after constraining the bond vibration of C-C and C-O (green) to remove their contributions. The experimental IR as well as the simulated DoS were obtained at a temperature of 10 K.

where the overtaking happens is about 120 K in both cases. The effect is more pronounced in the case of simulation wherein the left-side peak is completely absent at lower temperatures; whereas in the experiment, said peak is visible but of lower intensity. With the similar behaviour observed between both cases and considering the  $\sim 70$   $\text{cm}^{-1}$  shift in the family of peaks as shown in Figure 8, it can be stated that the two observations corroborate with each other.


By converting C-C and C-O bonds to constraints in the simulation topology, it is possible to single out peaks are contributions of the respective vibrations. In Figure 8, it is seen that the peaks corresponding to  $\text{CH}_2$ - $\text{CH}_2$  stretching, and  $\text{CH}_2$ -O stretching have disappeared from the constrained simulation, with the only remaining contribution from  $\text{CH}_3$  rocking. This further supports the claim that the family of peaks highlighted are the same vibrations, but with a uniform shift of  $\sim 70$   $\text{cm}^{-1}$  between the simulated and experimental observations.

At low temperatures, the thermal freedom available to the molecules in the deposited layer is low, and it increases as the temperature increases. This thermal freedom allows for the molecules to reorient themselves within the amorphous layer to achieve better close packing. This phenomenon is termed as ‘ice compaction’. Spectrally, this would manifest as the narrowing of peaks. In Figure 9, the temperature-dependent trend in the width of the peak corresponding to the OH stretching vibration is observed. At 10 K, the peak starts off wide, and on heating, it becomes narrower until it reaches a minimum at around 120 K. On further heating, the peak begins to widen again until 175 K. The trend in this spectral feature is common across both the experimental and simulated spectra. The narrowing of the peaks is an indication of ice compaction, comparable to the observations of the breadth of the OH stretching band in the experimental data.

At the molecular scale, the driving factor behind the stability of a crystalline 1-propanol solid is the hydrogen-bonding network. Due to the presence of the hydroxyl (-OH) group in 1-propanol, there is a possibility of one hydrogen bond per molecule. The number of hy-




**Figure 9.** The O-H stretching bands in the experimental and simulated spectra fitted to a Gaussian profile, and their respective full-width at half-maximum (FWHM) plotted against temperature. The error bars plotted are the standard error, defined as the square root of the corresponding diagonal element of the covariance matrix obtained during the fitting process.




**Figure 10.** Number of hydrogen bonds per molecule in the simulated system for the range of temperatures. The calculation was done across a slice of 1 nm thickness parallel to the substrate. The average was done over 1000 configurations in the simulated trajectory. The bulk crystal data corresponds to the predicted CSP structure. The hydrogen bond distance cutoff for calculation was 0.35nm while the H-D-A angle cutoff was 30°.

drogen bonds in the system was calculated for a range of temperatures and the average number of hydrogen bonds per molecule was plotted against temperature as shown in Figure 10. In the predicted crystal structure obtained via CSP, the average number of hydrogen bonds per molecule remains fixed at 1, as the hydrogen bonding network is a driving factor in the stability of the crystal structure. However, for the deposited system, which is amorphous in nature, the average number of hydrogen bonds per molecule is notably less than 1 for all temperatures. This highlights the tendency of the ice to not form a crystal as the hydrogen bonding network is not sufficient to facilitate



a) Crystalline



b) Amorphous

**Figure 11.** For the system of 1-propanol, the hydrogen bonding network is shown (a) in the crystal phase, and (b) in the amorphous phase. The dashed lines in red denote a hydrogen bond between a Hydrogen atom (coloured white) and an Oxygen atom (coloured red). Carbon atoms are coloured in cyan.

the crystal structure, similar to what is observed experimentally. In Figure 11, we show a visual comparison between the hydrogen-bonding network that is possible in the crystal arrangement versus the amorphous system. The possibility of every -OH group participating in a hydrogen bond is always satisfied in the crystal because of the orientational order; however, as we can predict and also observe from the computational results, this isn't necessarily the case for the amorphous arrangement.

#### 4 CONCLUSION

With the discovery of 1-propanol in the ISM, it is necessary to characterise the physico-chemical behaviour of the molecule at such conditions. Analysing the observational spectra strongly relies on laboratory-based data. Hence, in this paper, we presented the temperature-dependent IR spectra of 1-propanol ice, deposited at 10 K and then warmed to 170 K, revealed an amorphous morphology throughout the warming phase until sublimation. Although indications of ice compaction were observed, signatures specific to crystallization were absent. Knowing the phase of the molecule is especially

important to calculate the accurate abundance of the molecule in the region of observation. Employing the tools of classical molecular dynamics, we also conducted simulation studies of a vacuum-deposited 1-propanol ice system across the same temperature range as the experiment. From the simulations, we extracted various results including spectral and structural properties. The observed temperature-dependent trends across these results demonstrate a notable level of agreement between experiment and simulation, providing insights into the phase behaviour of the ice at a molecular scale. Most importantly, the key observation was the presence of 1-propanol on the substrate up to 170 K, despite the known melting point of 1-propanol being 147 K. It is worth noting that, to the best of our knowledge, this marks the first observation, or at least the first reporting, of a molecule staying on the substrate beyond its melting point under such conditions. Based on the experimental and simulation results, we attribute this behaviour to the low vapour pressure of the molecule at such low temperatures, which we also confirmed through a series of experiments by maintaining the deposited ice at isothermal conditions at different temperatures beyond the melting point and below the sublimation temperature. Therefore, the reported results suggest that the morphology of the interstellar icy mantle containing 1-propanol, along with other molecules (particularly water), warrants further investigation, as the sublimation temperature of 1-propanol falls well within the range of crystallisation and sublimation temperatures of several other molecules. Consequently, our future work will focus on the 1-propanol and water ice mixture.

#### ACKNOWLEDGEMENTS

RR, BNRS, AB, NJM and BS thank NSRRC for providing beamtime for the measurements. RR, SG, JK, AB and BS acknowledge the support from PRL (Dept of Space, Govt of India). AB acknowledges the J C Bose National Fellowship. BNRS, NJM and BS acknowledge the support from Sir John and Lady Mason Academic Trust. NJM acknowledges support from the Europlanet-2024-RI. Europlanet 2024 RI has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No. 871149.

We are grateful to Dr. G. R. Desiraju of the Solid State and Structural Chemistry Unit, Indian Institute of Science for his assistance in connecting with Dr. T. S. Thakur.

#### DATA AVAILABILITY

The data will be made available on request.

#### REFERENCES

- Abplanalp M. J., Kaiser R. I., 2019, *Physical Chemistry Chemical Physics*, 21, 16949
- Abraham M. J., Murtola T., Schulz R., P'all S., Smith J. C., Hess B., Lindahl E., 2015, *SoftwareX*, 1-2, 19
- Altwegg K., et al., 2017, *Monthly Notices of the Royal Astronomical Society*, 469, S130
- Ayling S. A., Burke D. J., Salter T. L., Brown W. A., 2017, *RSC advances*, 7, 51621
- Ball J. A., Gottlieb C. A., Lilley A., Radford H., 1970, *Astrophysical Journal*, 162, L203
- Bussi G., Donadio D., Parrinello M., 2007, *The Journal of Chemical Physics*, 126
- Crovisier J., Bockelée-Morvan D., Biver N., Colom P., Despois D., Lis D. C., 2004, *Astronomy & Astrophysics*, 418, L35
- Darden T., York D., Pedersen L., 1993, *The Journal of Chemical Physics*, 98, 10089
- Despois D., 1992, in *Symposium-International Astronomical Union*. pp 451–458, doi:<http://adsabs.harvard.edu/abs/1992IAUS..150..451D>
- Dodda L. S., Cabeza de Vaca I., Tirado-Rives J., Jorgensen W. L., 2017a, *Nucleic acids research*, 45, W331
- Dodda L. S., Vilseck J. Z., Tirado-Rives J., Jorgensen W. L., 2017b, *The Journal of Physical Chemistry B*, 121, 3864
- Doroshenko I., Pogorelov V., Sablinskas V., 2012, *Dataset Papers in Chemistry*, 2013
- Ehrenfreund P., Charnley S. B., 2000, *Annual Review of Astronomy and Astrophysics*, 38, 427
- Essmann U., Perera L., Berkowitz M. L., Darden T., Lee H., Pedersen L. G., 1995, *The Journal of Chemical Physics*, 103, 8577
- Fuchs G., Cuppen H., Ioppolo S., Romanzin C., Bisschop S., Andersson S., Van Dishoeck E., Linnartz H., 2009, *Astronomy & Astrophysics*, 505, 629
- Fuentes-Azcatl R., Barbosa M. C., 2018, *Physica A: Statistical Mechanics and its Applications*, 491, 480
- Hiraoka K., Mochizuki N., Wada A., 2006, in *AIP Conference Proceedings*. pp 86–99
- Hollis J. M., Jewell P. R., Lovas F. J., Remijan A., Møllendal H., 2004, *The Astrophysical Journal*, 610, L21
- Hudson R. L., Moore M. H., 2018, *The Astrophysical Journal*, 857, 89
- Jiménez-Serra I., et al., 2022, *Astronomy & Astrophysics*, 663, A181
- Jorgensen W. L., Tirado-Rives J., 2005, *Proceedings of the National Academy of Sciences*, 102, 6665
- Kasamatsu T., Kaneko T., Saito T., Kobayashi K., 1997, *Bulletin of the Chemical society of Japan*, 70, 1021
- Lin S.-T., Blanco M., Goddard W. A., 2003, *The Journal of Chemical Physics*, 119, 11792
- Lin S.-T., Maiti P. K., Goddard W. A. I., 2010, *The Journal of Physical Chemistry B*, 114, 8191
- Lu H.-C., Chen H.-K., Cheng B.-M., Ogilvie J., 2008, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 71, 1485
- Luna R., Molpeceres G., Ortigoso J., Satorre M. A., Domingo M., Maté B., 2018, *Astronomy & Astrophysics*, 617, A116
- NIST Accessed in 2023, NIST Chemistry WebBook, SRD 69, <https://webbook.nist.gov/cgi/inchi/InChI%3D1S/C3H8O/c1-2-3-4/h4H%2C2-3H2%2C1H3>
- Nag S., Majumdar J., Sivaraman B., Yashonath S., Maiti P. K., 2023, *Monthly Notices of the Royal Astronomical Society*, 522, 3656
- Pavithra S., et al., 2017, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 178, 166
- Pethes I., Pusztai L., Ohara K., Temleitner L., 2021, *Journal of Molecular Liquids*, 340, 117188
- Qasim D., et al., 2019, *Astronomy & Astrophysics*, 627, A1
- Ramachandran R., et al., 2023, *Journal of Chemical Sciences*, 135, 77
- Ramírez-Cardona M., Ventolà L., Calvet T., Cuevas-Diarte M., Rius J., Amigó J., Reventós M., 2005, *Powder diffraction*, 20, 311
- Ruiz-Mirazo K., Briones C., de la Escosura A., 2014, *Chemical reviews*, 114, 285
- Stroet M., Sanderson S., Sanzogni A. V., Nada S., Lee T., Caron B., Mark A. E., Burn P. L., 2022, *Journal of Chemical Information and Modeling*, 63, 2
- Stromsoe E., Ronne H. G., Lydersen A. L., 1970, *Journal of Chemical and Engineering Data*, 15, 286
- Tschamler H., Richter E., Wettig F., 1949, *Monatshefte für Chemie und verwandte Teile anderer Wissenschaften*, 80, 749
- Turner B. E., Apponi A. J., 2001, *The Astrophysical Journal*, 561, L207
- Watanabe N., Kouchi A., 2002, *The Astrophysical Journal*, 571, L173
- Yuan C., Smith R. S., Kay B. D., 2016, *Surface Science*, 652, 350
- Zangi R., 2018, *ACS omega*, 3, 18089
- Zuckerman B., et al., 1975, *Astrophysical Journal*, 196, L99

**APPENDIX A: SUPPORTING INFORMATION**

All the supporting informations are available online.

ORIGINAL UNEDITED MANUSCRIPT