



Kent Academic Repository

Higman, Victoria A, Płoskoń, Eliza, Thompson, Gary S. and Vuister, Geerten W (2024) *Perspective: on the importance of extensive, high-quality and reliable deposition of biomolecular NMR data in the age of artificial intelligence*. *Journal of Biomolecular NMR*, 78 (4). pp. 193-197. ISSN 0925-2738.

Downloaded from

<https://kar.kent.ac.uk/107686/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1007/s10858-024-00451-w>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Perspective: on the importance of extensive, high-quality and reliable deposition of biomolecular NMR data in the age of artificial intelligence

Victoria A. Higman¹ · Eliza Płoskoń¹ · Gary S. Thompson² · Geerten W. Vuister¹

Received: 9 May 2024 / Accepted: 25 September 2024
© The Author(s) 2024

Abstract

Artificial intelligence (AI) models are revolutionising scientific data analysis but are reliant on large training data sets. While artificial training data can be used in the context of NMR processing and data analysis methods, relating NMR parameters back to protein sequence and structure requires experimental data. In this perspective we examine what the biological NMR community needs to do, in order to store and share its data better so that we can make effective use of AI methods to further our understanding of biological molecules. We argue, first, that the community should be depositing much more of its experimental data. In particular, we should be depositing more spectra and dynamics data. Second, the NMR data deposited needs to capture the full information content required to be able to use and validate it adequately. The NMR Exchange Format (NEF) was designed several years ago to do this. The widespread adoption of NEF combined with a new proposal for dynamics data specifications come at the right time for the community to expand its deposition of data. Third, we highlight the importance of expanding and safeguarding our experimental data repository, the Biological Magnetic Resonance Data Bank (BMRB), not only in the interests of NMR spectroscopists, but biological scientists more widely. With this article we invite others in the biological NMR community to champion increased (possibly mandatory) data deposition, to get involved in designing new NEF specifications, and to advocate on behalf of the BMRB within the wider scientific community.

Keywords NMR · Dynamics · Database · NMR exchange format · BMRB

Slightly over 50 years ago, in October 1971, the Protein Data Bank (PDB) was announced as a joint effort between the Cambridge Crystallographic Data Centre in the UK and the Brookhaven National Laboratory in the US (Letter 1971). The aim was to harvest and make available to researchers the experimental atomic co-ordinates, structure factors and electron density maps of protein structures solved by X-ray crystallography. These data represented a

supplement to the publication of these structures in conventional scientific journals. In these early, pre-internet days, storage and distribution of the data was primarily done on magnetic tapes. As the initial announcement pointed out: “The success of the proposed system will depend on the response of the protein crystallographers supplying data” (Letter 1971). The response of the crystallographic scientific community proved to be very good and over the past five decades the PDB has grown and turned into the worldwide PDB (wwPDB) (Berman et al. 2003). It has added an additional distribution partner in Japan, the Biological Magnetic Resonance Data Bank (BMRB/NMRHub) for experimental NMR data, and most recently also a distribution hub in China (Xu et al. 2023). The wwPDB has also been accepting protein structures determined by other techniques such as Nuclear Magnetic Resonance (NMR) spectroscopy and Electron Microscopy (EM). These days, anyone wishing to publish an article describing a new protein structure is

✉ Victoria A. Higman
vad5@leicester.ac.uk

¹ Department of Molecular and Cell Biology, Leicester Institute of Structural and Chemical Biology, University of Leicester, Leicester LE1 7RH, UK

² Wellcome Trust Biological NMR Facility, School of Biosciences, Division of Natural Sciences, University of Kent, Canterbury, Kent CT2 7NZ, UK

required by journals to deposit the structure and associated data with the wwPDB.

Fifty years after the launch of the PDB, advances in computational power allowed DeepMind to use the by then ~180,000 deposited structures to develop AlphaFold2, an artificial intelligence model capable of predicting many protein structures with remarkable accuracy from the primary sequence alone (Jumper et al. 2021). While it is tempting to assume that the “protein structure problem has now been solved”, AlphaFold2’s successes have not signalled the end of experimental structural biology (Terwilliger et al. 2023). On the contrary, over the past five decades our understanding of protein structure, dynamics and the structure-function relationship has shifted dramatically. We now understand that not all protein sequences will fold into stable 3-dimensional structures (Wright and Dyson 1999; Arai et al. 2024) and that the dynamics of a protein, whether local or global,

is often integral to a protein’s function (Palmer 2004; Eisenmesser et al. 2005; Alderson and Kay 2021). NMR plays an exquisitely important role in characterising the dynamics of folded proteins across a variety of time scales (Palmer 2004), as well as providing a key experimental tool for studying the dynamics and residual structure of intrinsically disordered proteins (IDPs) (Milles et al. 2018; Ahmed and Forman-Kay 2022). Relaxation rates, paramagnetic relaxation enhancements (PREs), residual dipolar couplings (RDCs) and chemical shifts can all contribute to our understanding of protein dynamics and function (Palmer 2004; Milles et al. 2018; Schneider et al. 2019; Alderson and Kay 2021) including otherwise inaccessible excited state structures (Baldwin and Kay 2009).

In the same way that AlphaFold2 can predict the static structure of a protein from the primary sequence alone, might it one day also be possible to predict the dynamical behaviour of a protein from the primary sequence alone? It does not look like this will become a reality any time soon, most importantly because the data likely required to underpin such work are currently not accessible. When publishing NMR resonance assignments and/or structures, the deposition with the wwPDB and BMRB of chemical shift values, restraints used in the NMR structure calculation and atomic co-ordinates is mandatory. However, when publishing studies based on other types of NMR data, such as titrations, relaxation rates or PREs, deposition of the underlying data is unfortunately not mandatory. Sometimes data are appended as Supplementary Information, and more recently some researchers have begun using repositories such as Zenodo (<https://zenodo.org/>), an OpenScience data storage facility run by CERN, or BMRbig (<https://bmrbig.bmr.io/>), hosted by the BMRB, to deposit all of the data associated with an NMR project, including the unprocessed spectra, processing parameters, software project files etc. However, even if such data can be used for individual studies, the problem with an undocumented and non-formalised way of sharing data is that these are not easily mined using automated protocols and may not even contain the required information, thus strongly hampering future large-scale, AlphaFold-style studies. Overall, the amount of deposited NMR data such as spectra, peak lists or dynamics data in the BMRB are very low (cf. Table 1). These limitations are already hampering wide-scale AI-based analysis of NMR data. ARTINA, a novel program to assign the resonances and determine the structure of globular proteins directly from solution NMR spectra contains an AI-based method for peak picking (Klukowski et al. 2018, 2022). The algorithm was trained on a mere 100 datasets of globular proteins studied by solution NMR, since these were all that was available to the authors. By comparison, the number of non-redundant structures (<90% sequence similarity) solved by solution NMR is

Table 1 Number of deposited entries in the BMRB containing a variety of different measurable NMR parameters

Parameter ^a	Number of Entries	NEF specification ^b
Assigned Chemical Shifts	15,669	✓
Spectra	247	
Peak Lists	957 ^c	✓
Heteronuclear NOE	352	(✓)
R1 ^d	356	(✓)
R2 ^d	342	(✓)
R1rho ^d	19	(✓)
¹ H Exchange Rates	26	
¹ H Exchange Protection Factors	4	
Coupling Constants	384	
RDCs	152	✓
Chemical Shift Anisotropy	7	
Binding Data (K _d)	73	
Structures ^e	14,004	-
Restraints ^e	11,462	✓

^a It is currently not possible to retrieve PREs or pseudo contact shifts (PCSs). Titration series can be deposited as multiple chemical shifts lists measured on different samples or samples with different conditions. However, they are not identified as titration series. Only 60 entries contain at least 5 chemical shift lists, but many of these are not titration series; thus, the number of deposited titration series is liable to be relatively small.

^b Parameters for which NEF specifications exist are indicated by a tick, those for which a proposal for a NEF specification has been made have a tick in brackets. Proposals for Coupling Constant, PRE and PCS NEF specifications have not been tabled yet.

^c It should be noted that a number of these entries do not contain NMR-STAR formatted peak tables but an NMR-STAR header with the peak list data in a comment in a different format (NMRPipe, NMRView, XEASY, Sparky etc.).

^d In the BMRB these are deposited as relaxation times, but for the new NEF specification we suggest using rates.

^e Structures and Restraints are deposited with the wwPDB rather than the BMRB.

nearly 9,000. Furthermore, in a wwPDB-wide study of possible multi-state NMR protein structures, comparisons with dynamics data have been hampered by the low numbers of dynamics datasets deposited (Roland Riek, personal communication). Although simulated training data has been successfully used for several NMR-based AI-models (Beckwith et al. 2021; Shukla et al. 2023), this approach is only suitable for NMR data processing and analysis, not in relating NMR data and parameters sensitive to protein dynamics and function back to protein structure and sequence (this would require currently unfeasible quantum mechanical simulations of 10,000s of atoms across a ms-s timescale).

A further problem for the deposition of NMR data has been the deposition format. When it was shown that the quality of NMR structures could be improved through the use of more modern structure calculation protocols (Nabuurs et al. 2004; Nederveen et al. 2005), a large-scale recalculation of structures was not found to be practicable without a large-scale remediation effort of deposited restraints involving much human intervention (Doreleijers et al. 2009). The critical assessment of automated structure determination of proteins by NMR (CASD-NMR) (Rosato et al. 2009, 2012, 2015) required extensive exchange of NMR data and validation of results. Again, this often required manual intervention, e.g. where header data was ambiguous or files were missing or damaged and for certain software packages stereo-specificity of restraints could not be taken into account (Ragan et al. 2015). Deposited peak lists also present a problem, since these are not curated by the BMRB and thus present in the BMRB in a variety of different formats and do not all conform to the BMRB's NMR-STAR format. In 2015, following the second CASD-NMR round, nearly all key NMR software developers came together to develop what is now called the NMR Exchange Format (NEF; <https://github.com/NMRExchangeFormat/NEF>) in order to exchange NMR data seamlessly between software packages and deposit it with the BMRB (Gutmanas et al. 2015). NEF has been designed to be accurate in its documentation of data, e.g. stereo- and non-stereospecificity of assignments and restraints, is easily extendible, uses the STAR format (Hall 1991) already used by the wwPDB and BMRB, and is humanly readable. Crucially, we are tabling a proposal for documenting NMR dynamics data for consideration by the participants of the NEF effort (see <https://github.com/NMRExchangeFormat/NEF/tree/master/specification>) with a proposal for PRE and other data types to follow. The number of programs that are able to read and/or write NEF files now includes all the major structure calculation programs (Amber (Case et al. 2005, 2023), ARIA (Rieping et al. 2007), CS-Rosetta (Nerli and Sgourakis 2019), CYANA (Güntert and Buchner 2015), YASARA (YASARA Biosciences GmbH), XPLOR-NIH (Schwieters

et al. 2003, 2006)), and several spectrum display software packages, e.g. CcpNmr Analysis (Skinner et al. 2016) and Sparky (Lee et al. 2015) as well as wwPDB/BMRB deposition. In addition, the program NEF-Pipelines (Thompson 2024) is enabling NEF-based access to numerous other programs that are not NEF compatible, as well as general manipulation of NEF files. The advent of NEF and its use by an increasingly large number of software packages should enable the easy deposition of a wide range of NMR data which correctly preserves the information content. To misquote the original PDB announcement: whether we do so, "will depend on the response of the NMR spectroscopists supplying data". We are of the opinion that journal editors will likely have to mandate, rather than merely encourage deposition of all relevant data.

A separate format issue occurs in relation to binary (time and frequency domain) spectral data. These data are currently deposited in many different formats, depending on the spectrometer manufacturer and processing or spectral analysis software used by individual scientists. Although code libraries to read and write a variety of formats do exist (Helmus and Jaroniec 2013), a universal, simple, modern and well-documented format which is used for deposition would support both large-scale data mining and reproducibility and could become part of future efforts of the NEF Consortium.

Finally, a crucial hurdle is safeguarding the actual physical repository infrastructure, whose storage capacity will have to increase by several orders of magnitude if spectral data are to be deposited. This increase in capacity is particularly important considering that many modern spectra are recorded with non-uniform sampling which typically leads to much larger processed spectrum files. Curation of the data is essential, but expensive. The BMRB, as the wwPDB partner responsible for NMR data, is a US-funded project whose existence has too often appeared to be more precarious than it should be, given its importance to the global NMR community. It is imperative that not only the NMR community, but also the wider biological and scientific community acknowledges the importance of large repositories of scientific data in general and, therefore, also supports funding for the BMRB as securely as the wwPDB and other scientific repositories, possibly also through international efforts to provide greater robustness.

It should go without saying that the deposited data needs to be accurate, requiring good training and best practice to be in place across the community. The analysis of dynamics data, in particular, is not straight forward and robust validation methods need to be developed alongside the analysis methods themselves. We are currently developing tools to enable this within the CcpNmr AnalysisAssign program (Skinner et al. 2016). A further advantage to depositing

NMR datasets and measured data is that it opens up the possibility of reanalysing the deposited data with these and other improved methods as they become available.

While experimental structure determination may continue to be our gold-standard for the study of protein structures, programs such as AlphaFold2 represent a highly useful tool for biologists studying proteins which have so far eluded experimental structure determination. Given the importance of protein dynamics and IDP residual structure for protein function it would be highly desirable to have similar tools available for their prediction in the absence of detailed experimental data. If AlphaFold2 has taught us anything, it is the value of large, well-organised data repositories such as the wwPDB. Modern artificial intelligence systems are capable of detecting patterns in large datasets which humans are not, even with the help of the best traditional algorithms. If we want to be able to make the most of this new technology, we need to ensure that all of our experimental data are deposited in a high-quality, reliable way, starting as soon as possible (Editorial 2023; Arrowsmith 2024). We therefore invite the community to participate in the discussion about the expansion of the NEF specifications for data exchange and deposition, as well as ways to increase or mandate data deposition and safeguard our repositories for the future. This will be good not only for NMR spectroscopy as an experimental technique to study proteins and other biological molecules, but for biology and science, in general.

Acknowledgements We acknowledge funding of the Collaborative Computational Project for NMR (CCPN) project through the United Kingdom Research and Innovation (UKRI) grants MR/L000555/1, MR/P00038X/1 and MR/V000950/1. We thank Dr Kumaran Basakaran, Prof. Flemming Hansen, Dr Charles Schwieters and Dr Brian Smith for helpful discussions about the design of the NEF specification for dynamics data. For the purpose of open access, the authors have applied a Creative Commons Attribution license (CC BY) to any Author Accepted Manuscript version arising from this submission.

Author contributions V.A.H. conceived the article and gathered the data. V.A.H. and G.W.V. wrote the manuscript. V.A.H., E.A., G.S.T. and G.W.V. discussed the work, contributed ideas and revised the manuscript.

Data availability No datasets were generated or analysed during the current study.

Declarations

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless

indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ahmed R, Forman-Kay JD (2022) NMR insights into dynamic, multivalent interactions of intrinsically disordered regions: from discrete complexes to condensates. *Essays Biochem* 66:863–873. <https://doi.org/10.1042/EBC20220056>
- Alderson TR, Kay LE (2021) NMR spectroscopy captures the essential role of dynamics in regulating biomolecular function. *Cell* 184:577–595. <https://doi.org/10.1016/j.cell.2020.12.034>
- Arai M, Suetaka S, Ooka K (2024) Dynamics and interactions of intrinsically disordered proteins. *Curr Opin Struct Biol* 84:102734. <https://doi.org/10.1016/j.sbi.2023.102734>
- Arrowsmith CH (2024) Structure-guided drug discovery: back to the future. *Nat Struct Mol Biol* 31:395–396. <https://doi.org/10.1038/s41594-024-01244-3>
- Baldwin AJ, Kay LE (2009) NMR spectroscopy brings invisible protein states into focus. *Nat Chem Biol* 5:808–814. <https://doi.org/10.1038/nchembio.238>
- Beckwith MA, Erazo-Colon T, Johnson BA (2021) RING NMR dynamics: software for analysis of multiple NMR relaxation experiments. *J Biomol NMR* 75:9–23. <https://doi.org/10.1007/s10858-020-00350-w>
- Berman HM, Henrick K, Nakamura H (2003) Announcing the worldwide Protein Data Bank. *Nat Struct Biol* 10:980. <https://doi.org/10.1038/nsb1203-980>
- Case DA, Cheatham TE, Darden T, Gohlke H, Luo R, Merz KM, Onufriev A, Simmerling C, Wang B, Woods RJ (2005) The Amber biomolecular simulation programs. *J Comput Chem* 26:1668–1688. <https://doi.org/10.1002/jcc.20290>
- Case DA, Aktulga HM, Belfon K, Cerutti DS, Cisneros GA, Cruz-zeiro VWD, Forouzesh N, Giese TJ, Götz AW, Gohlke H, Izadi S, Kasavajhala K, Kaymak MC, King E, Kurtzman T, Lee TS, Li P, Liu J, Luchko T, Luo R, Manathunga M, Machado MR, Nguyen HN, O'Hearn KA, Onufriev AV, Pan F, Pantano S, Qi R, Rahnamoun A, Rishchik A, Schott-Verdugo S, Shajan A, Swalls J, Wang J, Wei H, Wu X, Wu Y, Zhang S, Zhao S, Zhu Q, Cheatham TE, Roe DR, Roitberg A, Simmerling C, York DM, Nagan MC, Merz KM (2023) AmberTools. *J Chem Inf Model* 63:6183–6191. <https://doi.org/10.1021/acs.jcim.3c01153>
- Doreleijers JF, Vranken WF, Schulte C, Lin J, Wedell JR, Penkett CJ, Vuister GW, Friend G, Markley JL, Ulrich EL (2009) The NMR restraints grid at BMRB for 5,266 protein and nucleic acid PDB entries. *J Biomol NMR* 45:389–396. <https://doi.org/10.1007/s10858-009-9378-z>
- Editorial (2023) For chemists, the AI revolution has yet to happen. *Nature* 617:438
- Eisenmesser EZ, Millet O, Labeikovsky W, Korzhnev DM, Wolf-Watz M, Bosco DA, Skalicky JJ, Kay LE, Kern D (2005) Intrinsic dynamics of an enzyme underlies catalysis. *Nature* 438:117–121. <https://doi.org/10.1038/nature04105>
- Güntert P, Buchner L (2015) Combined automated NOE assignment and structure calculation with CYANA. *J Biomol NMR* 62:453–471. <https://doi.org/10.1007/s10858-015-9924-9>
- Gutmanas A, Adams PD, Bardiaux B, Fogh RH, Güntert P, Hendrickx PMS, Herrmann T, Kleywegt GJ, Kobayashi N, Lange OF, Markley JL, Montelione GT, Nilges M, Ragan TJ, Schwieters CD, Tejero R, Ulrich EL, Velankar S, Vranken WF, Wedell JR,

- Westbrook J, Wishar DSt, Vuister GW (2015) NMR Exchange Format: a unified and open standard for representation of NMR restraint data. *Nat Struct Mol Biol* 22:433–434. <https://doi.org/10.1038/nsmb.3041>
- Hall SR (1991) The STAR file: a New Format for Electronic Data transfer and archiving. *J Chem Inf Comput Sci* 31:326–333. <https://doi.org/10.1021/ci00002a020>
- Helmus JJ, Jaroniec CP (2013) NmrGlue: an open source Python package for the analysis of multidimensional NMR data. *J Biomol NMR* 55:355–367. <https://doi.org/10.1007/s10858-013-9718-x>
- Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Židek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* 596:583–589. <https://doi.org/10.1038/s41586-021-03819-2>
- Klukowski P, Augoff M, Zieba M, Drwal M, Gonczarek A, Walczak MJ (2018) NMRNet: a deep learning approach to automated peak picking of protein NMR spectra. *Bioinformatics* 34:2590–2597. <https://doi.org/10.1093/bioinformatics/bty134>
- Klukowski P, Riek R, Güntert P (2022) Rapid protein assignments and structures from raw NMR spectra with the deep learning technique ARTINA. *Nat Commun* 13. <https://doi.org/10.1038/s41467-022-33879-5>
- Lee W, Tonelli M, Markley JL (2015) NMRFAM-SPARKY: enhanced software for biomolecular NMR spectroscopy. *Bioinformatics* 31:1325–1327. <https://doi.org/10.1093/bioinformatics/btu830>
- Letter (1971) Crystallography: Protein Data Bank. *Nat New Biol* 233:223. <https://doi.org/10.1038/newbio233223b0>
- Milles S, Salvi N, Blackledge M, Jensen MR (2018) Characterization of intrinsically disordered proteins and their dynamic complexes: from in vitro to cell-like environments. *Prog Nucl Magn Reson Spectrosc* 109:79–100. <https://doi.org/10.1016/j.pnmrs.2018.07.001>
- Nabuurs SB, Nederveen AJ, Vranken W, Doreleijers JF, Bonvin AMJJ, Vuister GW, Vriend G, Spronk CAEM (2004) DRESS: a database of Refined solution NMR structures. *Proteins Struct Funct Genet* 55:483–486. <https://doi.org/10.1002/prot.20118>
- Nederveen AJ, Doreleijers JF, Vranken W, Miller Z, Spronk CAEM, Nabuurs SB, Güntert P, Livny M, Markley MJ, Nilges M, Ulrich EL, Kaptein R, Bonvin AMJJ (2005) RECOORD: a recalculated coordinate database of 500+ proteins from the PDB using restraints from the BioMagResBank. *Proteins Struct Funct Genet* 59:662–672. <https://doi.org/10.1002/prot.20408>
- Nerli S, Sgourakis NG (2019) Cs-Rosetta. *Methods Enzymol* 614:321–362. <https://doi.org/10.1016/bs.mie.2018.07.005>
- Palmer AG (2004) NMR characterization of the dynamics of biomacromolecules. *Chem Rev* 104:3623–3640. <https://doi.org/10.1021/cr030413t>
- Ragan TJ, Fogh RH, Tejero R, Vranken W, Montelione GT, Rosato A, Vuister GW (2015) Analysis of the structural quality of the CASD-NMR 2013 entries. *J Biomol NMR* 62:527–540. <https://doi.org/10.1007/s10858-015-9949-0>
- Rieping W, Bardiaux B, Bernard A, Bernard A, Malliavin TE, Nilges M (2007) ARIA2: automated NOE assignment and data integration in NMR structure calculation. *Bioinformatics* 23:381–382. <https://doi.org/10.1093/bioinformatics/btl589>
- Rosato A, Bagaria A, Baker D, Bardiaux B, Cavalli A, Doreleijers JF, Giachetti A, Guerry P, Güntert P, Herrmann T, Huang YJ, Jonker HRA, Mao B, Malliavin TE, Montelione GT, Nilges M, Raman S, van der Schot G, Vranken WF, Vuister GW, Bonvin AMJJ (2009) CASD-NMR: critical assessment of automated structure determination by NMR. *Nat Methods* 6:625–626. <https://doi.org/10.1038/nmeth0909-625>
- Rosato A, Aramini JM, Arrowsmith C, Bagaria A, Baker D, Cavalli A, Doreleijers JF, Eletsky A, Giachetti A, Guerry P, Gutmanas A, Güntert P, He Y, Herrmann T, Huang YJ, Jaravine V, Jonker HRA, Kennedy MA, Lange OF, Liu G, Malliavin TE, Mani R, Mao B, Montelione GT, Nilges M, Rossi P, van der Schot G, Schwalbe H, Szyperki TA, Vendruscolo M, Vernon R, Vranken WF, de Vries S, Vuister GW, Wu B, Yang Y, Bonvin AMJJ (2012) Blind testing of routine, fully automated determination of protein structures from nmr data. *Structure* 20:227–236. <https://doi.org/10.1016/j.str.2012.01.002>
- Rosato A, Vranken W, Fogh RH, Ragan TJ, Tejero R, Pederson K, Lee H, Prestegard JH, Yee A, Wu B, Lemak A, Houlston S, Arrowsmith CH, Kennedy M, Acton TB, Xiao R, Liu G, Montelione GT, Vuister GW (2015) The second round of critical Assessment of Automated structure determination of proteins by NMR: CASD-NMR-2013. *J Biomol NMR* 62:413–424. <https://doi.org/10.1007/s10858-015-9953-4>
- Schneider R, Blackledge M, Jensen MR (2019) Elucidating binding mechanisms and dynamics of intrinsically disordered protein complexes using NMR spectroscopy. *Curr Opin Struct Biol* 54:10–18. <https://doi.org/10.1016/j.sbi.2018.09.007>
- Schwieters CD, Kuszewski JJ, Tjandra N, Clore GM (2003) The Xplor-NIH NMR molecular structure determination package. *J Magn Reson* 160:65–73. [https://doi.org/10.1016/S1090-7807\(02\)00014-9](https://doi.org/10.1016/S1090-7807(02)00014-9)
- Schwieters CD, Kuszewski JJ, Marius Clore G (2006) Using Xplor-NIH for NMR molecular structure determination. *Prog Nucl Magn Reson Spectrosc* 48:47–62. <https://doi.org/10.1016/j.pnmrs.2005.10.001>
- Shukla VK, Heller GT, Hansen DF (2023) Biomolecular NMR spectroscopy in the era of artificial intelligence. *Structure* 31:1360–1374. <https://doi.org/10.1016/j.str.2023.09.011>
- Skinner SP, Fogh RH, Boucher W, Ragan TJ, Mureddu LG, Vuister GW (2016) CcpNmr AnalysisAssign: a flexible platform for integrated NMR analysis. *J Biomol NMR* 66:111–124. <https://doi.org/10.1007/s10858-016-0060-y>
- Terwilliger TC, Leibschnr DL, Croll T, Williams CJ, McCoy AJ, Poon BK, Afonine PV, Oeffner RD, Richardson JS, Read RJ, Adams PD (2023) AlphaFold predictions are valuable hypotheses, and accelerate but do not replace experimental structure determination. *Nat Methods* 2022.11.21.517405. <https://doi.org/10.1038/s41592-023-02087-4>
- Thompson G (2024) NEF-Pipelines. <https://doi.org/10.5281/zenodo.11127004>
- Wright PE, Dyson HJ (1999) Intrinsically unstructured proteins: reassessing the protein structure-function paradigm. *J Mol Biol* 293:321–331. <https://doi.org/10.1006/jmbi.1999.3110>
- Xu W, Velankar S, Patwardhan A, Hoch JC, Burley SK, Kurisu G (2023) Acta Crystallogr Sect D Struct Biol 79:792–795. <https://doi.org/10.1107/S2059798323006381>. Announcing the launch of Protein Data Bank China as an Associate Member of the World-wide Protein Data Bank Partnership

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.