



Kent Academic Repository

Romanov, Michael N, Ameen, Questan A., Shaker, Ahmed S., Al-Obaidi, Rana M. and Griffin, Darren K. (2024) *Conservation genetics and breeding using molecular genetic markers in Japanese quail (Coturnix japonica)*. *Frontiers in Bioscience (Scholar Edition)*, 16 (4). ISSN 1945-0516.

Downloaded from

<https://kar.kent.ac.uk/108243/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.31083/j.fbs1604023>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Original Research

Conservation Genetics and Breeding using Molecular Genetic Markers in Japanese Quail (*Coturnix japonica*)

Michael N. Romanov^{1,2,3,*}, Questan Ali Ameen⁴, Ahmed Sami Shaker⁵,
Rana Mohammed Al-Obaidi⁶, Darren K. Griffin^{2,3}

¹L. K. Ernst Federal Research Centre for Animal Husbandry, Dubrovitsy, 142132 Podolsk, Moscow Oblast, Russia

²School of Biosciences, University of Kent, CT2 7NJ Canterbury, Kent, UK

³Animal Genomics and Bioresource Research Unit (AGB Research Unit), Faculty of Science, Kasetsart University, 10900 Chatuchak, Bangkok, Thailand

⁴Department of Animal Science, College of Agricultural Engineering Sciences, University of Sulaimani, 46001 Sulaymaniyah, Iraq

⁵Medical Laboratory Technology, Al-Qalam University College, 820102 Kirkuk, Iraq

⁶Department of Basic Sciences, College of Dentistry, University of Sulaimani, 46001 Sulaymaniyah, Iraq

*Correspondence: m.romanov@kent.ac.uk (Michael N. Romanov)

Academic Editor: Federica Finetti

Submitted: 23 July 2024 Revised: 24 September 2024 Accepted: 9 October 2024 Published: 20 December 2024

Abstract

Background: The Japanese quail (*Coturnix japonica*) is a small migratory bird whose main habitats are located in East Asia, Russia, China, Japan, Korea, and India. The Japanese quail was first introduced into the Iraqi research sector in the early 1980s. This investigation aimed to identify the genetic divergence between the available genetic lines of the Japanese quail in Iraq as a first step to conducting further conservation and breeding, benefiting from studying the genetic diversity related to productivity, adaptation, and immune susceptibility. **Methods:** In this study, we harnessed the random amplified polymorphic DNA (RAPD)-polymerase chain reaction (PCR) protocol to investigate the genetic structure, diversity, and differentiation of seven distinct genetic lines of these birds with white, brown, wild grey, and yellow plumage guided by 20 molecular genetic markers. **Results:** Our findings showed a relatively high polymorphism level of these 20 markers, which was adequate for describing genetic variation within and between the quail lines under study. Overall, the pairs of the White male and Brown male, Brown female and Grey male, and Grey female and Grey male lines were the most genetically distant. Conversely, the White male and Grey male lines were the most similar. **Conclusions:** The genetic differences established between these lines can enable us to suggest recommendations for further conservation genetics and breeding of this species. In particular, we propose that choosing animals (lines) with the greatest genetic distances, i.e., the least genetic similarities, can help preserve the highest genetic variability within the population. This proposed approach mirrors many contemporary conservation strategies, and information derived directly from this study can potentially be used to improve breeder selection regimes for additional conservation initiatives *ex situ*.

Keywords: Japanese quail; *Coturnix japonica*; conservation genetics; breeding; lines; molecular genetic markers; RAPD-PCR; genetic diversity

1. Introduction

The Japanese quail (*Coturnix japonica* Temminck & Schlegel 1848) is a small migratory galliform bird, wild populations of which are primarily found in East Asia and Russia but also can be located in China, Japan, Korea, and India [1–3]. While resident populations may spend the winter in Japan, most migrate south to places such as Vietnam, Cambodia, Laos, and southern China [4]. Furthermore, the Japanese quail habitat covers much of Africa, including Tanzania, Malawi, Kenya, Namibia, Madagascar, and the Nile River Valley region stretching from Kenya to Egypt [3]. The breeding locations are mostly concentrated in East and Central Asia [1,2], specifically in Manchuria, south-eastern Siberia, northern Japan, and the Korean Peninsula. Moreover, the Japanese quail has also been noted to breed in Turkey and other parts of Europe [3]. The bird is called by different names, such as “quail” in English-speaking na-

tions, “marai” in Iraq, or “ferry” in Syria, depending on the geographic region. Japanese quail birds are characterized by their small body size, not exceeding 20 cm³ [5]. The body weight of mature *ad libitum*-fed males ranged from 190 to 250 g and that of females from 225 to 300 g [6]. These quails are adaptable to different environmental conditions and have high variability in shape and phenotypic traits such as plumage color and performance (egg production and meat yield) [7–9]. Both sexes are tailless, with wild females distinguished by light and tan feathers with black dots in the throat and upper chest areas, whereas males are grey with a brown neck area [10]. The current conservation status of Japanese quail in the wild is classified as “Near Threatened” [11].

Coturnix japonica was first domesticated in Japan in the 12th century [12] primarily because of the males’ ability to emit a beautiful call [13]. Intensive production of this species did not begin in Japan until the 1920s, when the





Fig. 1. Japanese quail genetic lines of two studied feather color varieties. Brown: (A) male (paternal) and (B) female (maternal) lines. Grey: (C) male (paternal) and (D) female (maternal) lines. The Grey male and female lines have the wild-type plumage color and pattern, while the Brown male and female lines represents the *extended brown* mutant allele at the *E (extension)* locus controlled by the *MC1R* gene [26]. Image source: authors' photographs.

first egg breeds were developed through selection. From Japan, they were successfully introduced into America, Europe, and the Middle East in the 1930s, 1940s, and 1950s,

when specific lines were bred for egg and meat production [12,14,15]. Subsequently, the modern commercial Japanese quail is characterized by several economically im-

portant traits, including rapid growth [16,17]. Additionally, commercial Japanese quail exhibits a high egg-laying rate [12,18]; these birds can lay up to 350 eggs annually. Given that these eggs weigh 10–12 g, these remarkable animals produce around 20 times their body weight yearly [19]. Japanese quails have lower feed and space requirements than other domestic birds [20], and all these traits combine to promote a prolific dual-purpose domestic bird in terms of both egg and meat production [12,21,22]. Being a photoperiodic avian species, *Coturnix japonica* is also an excellent model for studies on reproductive physiology, particularly for studying the mechanisms through which egg formation and laying occur [23–25].

In the early 1980s, Japanese quails were introduced into the research sector of Iraq, with four major genetic varieties predominating and classified based on their feather color, i.e., white, brown (Fig. 1A,B), wild grey (Fig. 1C,D, Ref. [26]), and yellow. These quails are subject to conservation and multilateral investigations. Numerous studies have addressed the productive traits of these lines to characterize the qualitative and quantitative characteristics associated with eggs [27–31], growth, and carcasses [32,33]. In addition, the gene pool lines of these quails have been used as a model for poultry nutritional research [34–37] and for immunological selection [37,38].

Molecular genetic markers are widely used in conservation genetics [39–44]. In particular, the random amplified polymorphic DNA (RAPD)-polymerase chain reaction (PCR) analysis method is broadly implemented for molecular characterization of species and populations, biodiversity evaluation, genotoxic, and other related studies (e.g., [45,46]). For example, using multilocus and dominant RAPD markers, Romanov and Weigend [47] assessed the genetic diversity of several German and Ukrainian chicken populations, while Dehghanzadeh *et al.* [48] evaluated genetic variability and distances among five Iranian native chicken populations. Abdulrazaq and Ameen [49] investigated genetic relationships between local guinea fowl, Japanese quail, and chicken populations using the RAPD-PCR technique. This method was also instrumental in defining genetic variation among other local quails [50], phenotypic quail groups of different feather colors [51], selected Japanese quail lines [52], and those improved for meat and egg production traits [53].

This study aimed to explore the genetic structure, diversity, and differentiation of seven Iraqi quail genetic lines with white, brown, wild grey, and yellow feather colors using a set of 20 molecular genetic markers based on the RAPD-PCR protocol. These lines are intended to further produce highly productive interline crosses, with their genetic characterization paramount in developing effective germplasm conservation and breeding strategies.

2. Materials and Methods

2.1 Experimental Birds

This experiment was conducted at the Poultry Breeding Unit of the Teaching and Research Farm of the Animal Science Department, College of Agricultural Engineering Sciences, University of Sulaimani, Iraq. Seven genetic lines of Japanese quail, both paternal (male) and maternal (female), were used as follows: white male (WM), white female (WF), brown male (BM; Fig. 1A), brown female (BF; Fig. 1B), grey male (GM; Fig. 1C), grey female (GF; Fig. 1D), and yellow female (YF). These lines have been bred as closed populations for ten generations and are planned for use in interline crossing. The healthy birds were kept in large commercial cages, which allowed for free movement, and had *ad libitum* access to feed and water (Fig. 1). All conditions for quail comfort environment, maintenance, and feeding were observed in accordance with the generally accepted standards for quail rearing [54–56].

2.2 Samples and DNA Extraction

Blood samples from wing veins were collected from each genetic line. Five males and five females from each line were sampled, with no euthanasia or anesthesia methods used because this conventional blood sampling procedure in birds do not require any euthanasia or anesthesia. Collected blood samples were stored at -20°C . Genomic DNA extraction from these samples was conducted using a Zymo Research kit (Orange, CA, USA). Samples for each sex of each genetic line were pooled separately into a single sample. These samples were pooled because we deduced that since each line has been bred for more than 10 generations as closed populations, the individuals within one line were genetically very close. Therefore, creating one pooled sample was not predicted to affect the results and compromise the estimation and interpretation of the observed genetic diversity.

2.3 DNA Amplification by RAPD-PCR

The protocol of the RAPD-PCR procedure was as follows: 5 μL of de-ionized water was added to a PCR tube and then supplemented with 8 μL master mix, 2 μL RAPD primer, and 5 μL of extracted DNA (with the final concentration of 35 ng/ μL). Amplification was carried out in a thermal cycler using the following conditions: initial denaturation at 94°C for 9 min, followed by 35 cycles, including denaturation at 94°C for 1 min; annealing, according to the primer's annealing temperature (Table 1), for 1 min; extension at 72°C for 2 min; final extension at 72°C for 9 min. For random amplification of DNA fragments, 20 RAPD primers (OPA-01 to OPA-20; Operon Technologies, Alameda, CA, USA; Table 1) were selected based on our preliminary studies [49,50]; these primers were tested, validated, and used in subsequent genetic diversity analyses.

Table 1. Primer sequences of 20 RAPD markers used for DNA amplification in quails.

Primer ID	Primer nucleotide sequence	T_m (°C)	(G+C)%	PIC	Band number	Band percentage	Size of fragment (bp)
OPA-01	CAGGCCCTTC	41	70	0.290936	15	6	100–1000
OPA-02	TGCCGAGCTG	38	70	0.375510	6	2.4	100–1000
OPA-03	AGTCAGCCAC	41	60	0.155844	20	8	160–1300
OPA-04	AATCGGGCTG	34	60	0.418367	23	9.2	600–1200
OPA-05	AGGGGTCTTG	41	60	0.277551	19	7.6	500–3000
OPA-06	GGTCCCTGAC	41	70	0.340936	12	4.8	200–1300
OPA-07	GAAACGGGTG	30	60	0.299320	5	2	400–2500
OPA-08	GTGACGTAGG	36.9	60	0.103401	7	2.8	300–1000
OPA-09	GGGTAACGCC	41	70	0.341950	13	5.2	500–2000
OPA-10	GTGATCGCAG	36.9	60	0.341950	13	5.2	400–600
OPA-11	CAATCGCCGT	58.6	60	0.336143	30	12	780–2100
OPA-12	TCGGCGATAG	32	60	0.400000	9	3.6	450–2800
OPA-13	CAGCACCCAC	36	70	0.340936	5	2	500–1400
OPA-14	TTCGAGCCAG	34	60			failed	
OPA-15	TTCGAGCCAG	34	60			failed	
OPA-16	AGCCAGCGAA	32	60	0.308390	16	6.4	300–1700
OPA-17	GACCGCTTGT	30	60			failed	
OPA-18	AGGTGACCGT	36.9	60	0.371882	27	10.8	130–1500
OPA-19	CAAACGTCGG	32	60	0.097959	6	2.4	550–2300
OPA-20	GTTGCGATCC	32	60	0.383673	24	9.6	400–1600
Total						250	
Average \pm standard deviation				0.304985 \pm 0.097000		14.7 \pm 3.2	

T_m (°C), primer melting (annealing) temperature; (G+C)%, guanine–cytosine content percentage; RAPD, random amplified polymorphic DNA; PIC, polymorphic information content.

2.4 DNA Fragment and Genetic Diversity Analyses

Post amplification, electrophoresis was performed using a 1.5% agarose gel for PCR product visualization. The visualized electrophoresis results were imported into GenAEx 6.5 (The Australian National University, Acton, Australia) [57,58] for subsequent band calling and fragment analysis.

Using the GenAEx 6.5 software package [57,58], the following genetic variation indicators were subsequently computed or detected: polymorphic information content (PIC) [59], observed number of alleles (N_a ; calculated as presence or absence of a band), effective number of alleles (N_e) [60], Nei's gene diversity [61], or heterozygosity (H), and Shannon's information index (I) [62]. The genetic distance and identity matrices for the seven genetic lines of quails were computed according to Nei [63,64], respectively, and an Unweighted Pair-Cluster Method using Arithmetic Averages (UPGMA) dendrogram was constructed to infer the genetic differentiation between the lines.

3. Results

3.1 RAPD-PCR and Genetic Diversity Data

After twenty primers were randomly screened for DNA polymorphism validation, seventeen were used for further RAPD-PCR analysis, producing different fragment patterns with various bands. Three markers (OPA-14, OPA-

15, and OPA-17) failed to amplify and showed no band results for these genetic lines. A total of 250 RAPD bands were detected based on all RAPD pattern analyses of the seven Japanese quail genetic lines examined. Table 1 shows the number and percentage of polymorphic bands presented. The count of bands amplified using these primers varied from a minimum of 5 (OPA-07) to a maximum of 30 (OPA-11), while the average band number was 14.7 ± 3.2 , and band sizes ranged between 100 and 3000 bp.

Regarding polymorphic information content (PIC), its range across the 17 markers was from 0.097959 to 0.418367, with the average PIC value equal to 0.304985 ± 0.097000 . The lowest PIC values (0.097959–0.155844) were identified for the following three markers: OPA-03, OPA-08, and OPA-19.

As shown in Table 2, the average number of alleles (N_a) was 2.00 because every anonymous RAPD locus was considered for two states of a band, i.e., its presence or absence. However, the effective number of alleles (N_e) ranged between 1.6897 and 1.9600. The highest N_e value estimated for each of the 12 markers (OPA-1, OPA-2, OPA-3, OPA-4, OPA-6, OPA-7, OPA-9, OPA-10, OPA-11, OPA-13, OPA-16, and OPA-19) was 1.9600. The heterozygosity (H) was lowest for five markers (OPA-05, OPA-08, OPA-12, OPA-18, OPA-20) and equaled 0.4082. The Shannon information index (I) reflected a similar pattern of highest

Table 2. The number of alleles (N_a), the effective number of alleles (N_e), Nei's gene diversity (H), and Shannon information index (I) for the markers used.

Marker locus	Sample size	N_a	N_e	H	I
OPA-01	70	2.0000	1.9600	0.4898	0.6829
OPA-02	70	2.0000	1.9600	0.4898	0.6829
OPA-03	70	2.0000	1.9600	0.4898	0.6829
OPA-04	70	2.0000	1.9600	0.4898	0.6829
OPA-05	70	2.0000	1.6897	0.4082	0.5983
OPA-06	70	2.0000	1.9600	0.4898	0.6829
OPA-07	70	2.0000	1.9600	0.4898	0.6829
OPA-08	70	2.0000	1.6897	0.4082	0.5983
OPA-09	70	2.0000	1.9600	0.4898	0.6829
OPA-10	70	2.0000	1.9600	0.4898	0.6829
OPA-11	70	2.0000	1.9600	0.4898	0.6829
OPA-12	70	2.0000	1.6897	0.4082	0.5983
OPA-13	70	2.0000	1.9600	0.4898	0.6829
OPA-16	70	2.0000	1.9600	0.4898	0.6829
OPA-18	70	2.0000	1.6897	0.4082	0.5983
OPA-19	70	2.0000	1.9600	0.4898	0.6829
OPA-20	70	2.0000	1.6897	0.4082	0.5983
Mean \pm standard deviation	70 \pm 0.0000	2.0000 \pm 0.0000	1.8805 \pm 0.1270	0.4658 \pm 0.0383	0.6580 \pm 0.0398

Table 3. Nei's genetic distances among the seven Japanese quail lines using RAPD markers.

Line	WM	WF	BM	BF	GM	GF	YF
WM	0						
WF	0.435	0					
BM	1.447	1.042	0				
BF	1.224	0.636	0.531	0			
GM	0.194	0.348	0.887	1.447	0		
GF	1.224	1.224	0.531	0.636	1.447	0	
YF	0.887	0.636	1.042	0.435	1.042	0.636	0

Lines tested: WM, White male; WF, White female; BM, Brown male; BF, Brown female; GM, Grey male; GF, Grey female; YF, Yellow female.

and lowest values across the 17 genetic markers, with the minimum value being 0.5983 and the maximum value being 0.6829.

3.2 Between-line Genetic Distances, Identity, and Relationships

The genetic distance matrix computed according to Nei [63] is presented in Table 3. The genetic distance matrix shows that the longest genetic distance (1.447) was observed between the BM and WM, BF and GM, and GM and GF lines, while the shortest one (0.194) was found between the GM and WM lines.

The pairwise matrix of Nei's genetic identity [64] is presented in Table 4. Moderate to high levels of genetic identity were estimated. The highest identity estimation (0.824) was found between the WM and GM lines, while the lowest (0.235) was between the WM and BM, BF and

Table 4. Nei's genetic identity (similarity) matrix among the seven Japanese quail lines using RAPD markers.

Line	WM	WF	BM	BF	GM	GF	YF
WM	0						
WF	0.647	0					
BM	0.235	0.353	0				
BF	0.294	0.529	0.588	0			
GM	0.824	0.706	0.412	0.235	0		
GF	0.294	0.294	0.588	0.529	0.235	0	
YF	0.412	0.529	0.353	0.647	0.353	0.529	0

Lines tested: WM, White male; WF, White female; BM, Brown male; BF, Brown female; GM, Grey male; GF, Grey female; YF, Yellow female.

GM, and GM and GF lines. This between-line similarity pattern agreed with the calculated pairwise genetic distance values, given that, in principle, genetic distance is largely an inverted (by numeric value) indicator of genetic identity (similarity).

The UPGMA dendrogram in Fig. 2 was generated based on the RAPD-PCR analysis using the seven genetic lines of Japanese quail to restore the between-line genetic relationships and visualize the respective line differentiation. The first large dendrogram cluster consisted of the WM, GM, and WF lines, and the second major cluster included the BM, GF, BF, and YF lines.

4. Discussion

Conservation and breeding of different plumage color lines of quail are pivotal for further progress in quail production [65,66]. A good strategy to increase quail mar-




Fig. 2. The Unweighted Pair-Cluster Method using Arithmetic Averages (UPGMA) dendrogram derived from the genetic relationship analysis among the seven lines of Japanese quail based on the RAPD primers. Lines tested: WM, White male; GM, Grey male; WF, White female; BM, Brown male; GF, Grey female; BF, Brown female; YF, Yellow female.

keting potential and consumer interest is to provide value-added meat, eggs, or live birds representing distinct color lines in their plumage [67,68]. The color of the plumage or the related genetic variation might significantly impact the quality of the quail's meat and eggs [65,66]. In particular, Al-Kafajy *et al.* [65], when comparing three quail lines of different plumage colors, established that the brown line demonstrated significantly higher values for body weight, albumen height, shell thickness, and carcass dressing traits than other lines. Moreover, the white line had significantly higher egg numbers, shell and yolk weights, and carcass characteristics (heart, thigh, breast, and back weights). When examining egg traits in three color lines, Petek *et al.* [66] revealed that the black line had significantly greater egg weight, strength, and shell thickness, whereas the white line had significantly lower egg length, higher shape index, and lighter yolk. Similar observations were made by Shaker *et al.* [69] for genetic groups of Kurdish local chickens, suggesting that the plumage color can affect the egg weight and other egg characteristics. Shaker and co-authors concluded that the statistically significant differences in egg weight and traits such as egg length, breadth, and shape index across the genetic groups might be attributed to their genetic makeup, which requires further characterization of these groups using molecular markers [69].

The current study aimed to explore genetic diversity among seven genetic lines of Japanese quail that were originally classified according to the specific color of their plumage. For this purpose, we employed the commonly used polymorphic molecular genetic markers based on the conventional RAPD-PCR primers. Our data demonstrated a rather high polymorphism level related to these markers that was sufficient to characterize the genetic variation within and between the studied quail lines. For instance, considering the established PIC values, one should note that this is a measure implemented in genetic investiga-

tions to assess marker quality that has values ranging from 0 (monomorphic) to 1 (very informative) [70]. Furthermore, PIC is crucial for choosing markers for genetic research since it assesses a marker's informativeness and capacity to identify polymorphisms and forecasts the likelihood that the marker's heterozygous genotype will be passed on to the progeny [70–72]. PIC values below 0.25 are not advised, while those between 0.25 and 0.50 are implied to be moderately informative and rather useful [70]. In this study, most used RAPD markers (14 primers) had PIC values above 0.28, and only three had lower values (below 0.16). This suggests that this set of markers was reasonably informative and reflected an adequate overall polymorphism estimate and a more or less true status of genetic diversity among the quail lines studied. Similar conclusions regarding these polymorphic markers and the genetic diversity of the lines can be drawn from the values obtained for N_e , H , and I . Similar to many previous biodiversity conservation studies (e.g., [73]), if, for example, the Shannon diversity values were consistently lower, this would indicate a decline in local bird diversity when abundances were considered. Obviously, this was not the case in our study.

As determined by other RAPD studies (see review [47]), this molecular marker type has some possible limitations, including inconsistent repeatability of strong bands, challenges in evaluating weak bands, and lower reproducibility of results. To overcome these potential restrictions, it is recommended to use precisely regulated PCR settings and DNA template quality as we did in the present study. Nonetheless, RAPD markers are still very quick and inexpensive molecular tools that can characterize genetic variation within and between breeds at an initial level [47].

Over recent years, several studies have been conducted in the region to determine the genetic diversity and classification of these Iraqi quail lines. Al-Neemy *et al.* [51] studied the genetic diversity of three genetic lines

(white, black, and brown) using nine RAPD markers. The relationship between these genetic lines was similar to that observed in our study. Lajan and Al-Barzinji [74] examined three genetic lines (wild grey, white, and brown) and, using RFLP-PCR markers at three genetic loci, showed that the studied gene sites contained several alleles with different nucleotide compositions. However, both the previous studies and our current study could have been more concordant with regard to the revealed genetic differentiation in the analyzed quail lines. This lack of concordance may be due to how these lines were selected for phenotypic traits and what their genetic makeup determinants were, thus causing an output that differed in the genetic characterization of the Japanese quail in other studies. Ahmed and Al-Barzinji [50], in an experiment using 20 RAPD markers and the genetic lines of Japanese quail of three different colors (desert, brown, and white), also found that the number of fragments obtained was 310, with their size ranging from 250 to 2800 bp. The established average Nei's gene diversity between the lines was 0.1026, and the highest genetic distance (27.614) was identified for the BF line relative to both the WF and Desert female lines. Our study demonstrated a higher mean value of H (0.4658), whereas the greatest genetic distances were found between the following three line pairs: BM–WM, BF–GM, and GM–GF (Table 3). It should be acknowledged that it is difficult to make direct, plausible comparisons of results across studies, given that they used different sets of genetic lines and markers.

Overall, our findings showed that, when employing molecular markers using the RAPD-PCR procedure, it was feasible to successfully assess the genetic diversity within the local genetic quail lines and the genetic relationship among the lines. This information will be important for further conservation and breeding of these lines. However, to avoid potential drawbacks of using the RAPD-PCR protocol, more sophisticated genetic markers such as microsatellites and single nucleotide polymorphisms (SNPs) [75–80] will need to be harnessed in future studies. Regarding the conservation strategies inferred from these genetic studies, we believe that selecting animals with the least genetic similarity, i.e., with the greatest genetic distances, can help conserve the maximum variability within a total population. Moreover, this selection approach can even optimize the choice of breeders for further conservation programs *ex situ* [81]. According to the dendrogram analysis (Fig. 2) based on the pairwise comparison (Table 3), we established the greatest genetic distances between the line pairs BM–WM, BF–GM, and GM–GF that reflect the pronounced evolutionary relationships between these lines due to genetic divergence in the course of their breeding. Regarding the connection between genetic findings and practical breeding applications, more genetic improvement can result from advances in molecular genetics than phenotypic information alone. This is because markers connected to genes of interest could provide advantages over utilizing information at

the DNA level. The application of genetic markers to identify genes or genomic areas that regulate traits of interest—known as quantitative trait loci (QTLs)—is also made possible by developments in molecular genetics. Furthermore, marker-assisted selection (MAS) could use the identified QTLs to promote genetic advancement [82,83].

The future growth and competitiveness of commercial quail breeding mostly depend on genetic research, conservation, and utilization of lines and crosses with high productivity and adequate variability [84–87]. Therefore, it is hardly practicable to carry out efficient selection work intended to locate and identify desirable quail genotypes without conservation measures. Future research on the conserved gene pool genetic resources of quail should use cutting-edge, whole-genome techniques that employ SNP markers [88–90]. This will contribute to understanding key physiological and molecular genetic processes involved in developing and expressing selectively significant traits in quail.

5. Conclusions

A well-known member of the *Coturnix* genus, *Coturnix japonica*, is commercially exploited worldwide for its nutritional meat and eggs [91]. Apart from the original *Coturnix* line, small flocks of mutant or crossbred birds with plumage color variants are being raised for various purposes [92,93]. Here, using the RAPD-PCR markers, we examined seven genetic lines of different feather colors with respect to their genetic diversity indicators (PIC, N_a , N_e , H , and I). This study also revealed the genetic distances between the seven quail lines of distinct colors. Our findings suggest that these lines possess sufficient genetic variety to promote genetic advancements that can bring these lineages closer together. Breeders can utilize these results to improve their understanding of the genetic variability in the local quails, providing critical information for developing the mating strategies of quails when crossing these quail lines.

From the current genetic research, we believe that the best conservation strategy would involve selecting animals (lines) with the greatest genetic distances from one another, i.e., the least genetic similarity, to preserve the highest amount of variability within an entire population. Further *ex situ* conservation programs may also benefit by selecting breeders more effectively.

Collectively, the obtained results will form the basis for developing more detailed strategies for preserving and using the genetic and breeding resources of Japanese quail [94]. Further development of MAS and genomic selection in quails looks promising. In general, the information presented in this study will be in demand in quail conservation and breeding projects to improve existing and create new promising lines and crosses of quails with specified economically important traits.

Availability of Data and Materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Author Contributions

MNR: Conceptualization, Formal Analysis, Project administration, Software, Validation, Visualization, Writing – original draft, Writing – review & editing. QAA: Conceptualization, Project administration, Funding acquisition, Methodology, Visualization, Writing – original draft, Writing – review & editing. ASS: Data curation, Funding acquisition, Resources, Writing – original draft, Writing – review & editing. RMAO: Formal Analysis, Investigation, Funding acquisition, Writing – review & editing. DKG: Validation, Supervision, Writing – review & editing. All authors have participated sufficiently in the work to take public responsibility for appropriate portions of the content and agreed to be accountable for all aspects of the work in ensuring that questions related to its accuracy or integrity. All authors read and approved the final manuscript.

Ethics Approval and Consent to Participate

Ethics standards were considered in all steps of performing procedures and animal handling as approved by the Iraq Ethical Approval No. um.VET.2021.5. This general ethics approval, issued in 2021, serves as a comprehensive guideline for research conducted across various colleges in Iraq. This approval is applicable as a broad framework for maintaining ethical standards in research. Additionally, the Ethic Committee of College of Agricultural Engineering Sciences, University of Sulaimani, issued the scientific research approval No. 6 on 14/1/2023 that states that “the Committee has approved and accepted to proceed with this research.” This approval was provided as a specific commitment to the ethical principles established in Iraq and adopted within the College. This adoption aligns with the rules outlined in the general ethics approval and ensures compliance with the specific requirements and circumstances within the University of Sulaimani.

Acknowledgment

Not applicable.

Funding

The research work conducted by Michael N. Romanov was supported by the Russian Science Foundation (Grant No. 24-16-00294, https://rscf.ru/prjcard_int/24-16-00294).

Conflict of Interest

The authors declare no conflict of interest.

References

- [1] Barilani M, Deregnacourt S, Gellego S, Galli L, Mucci N, Piomobo R, *et al.* Detecting hybridization in wild (*Coturnix c. coturnix*) and domesticated (*Coturnix c. japonica*) quail populations. *Biological Conservation*. 2005; 126: 445–455.
- [2] Puigcerver M, Vinyoles D, Rodríguez-Teijeiro JD. Does restocking with Japanese quail or hybrids affect native populations of common quail *Coturnix coturnix*? *Biological Conservation*. 2007; 136: 628–635.
- [3] Pappas J. *Coturnix japonica* Japanese Quail. Animal Diversity Web, Museum of Zoology Regents of the University of Michigan. 2002. Available at: https://animaldiversity.org/accounts/Coturnix_japonica/ (Accessed: 22 July 2024).
- [4] BirdLife International. Species factsheet: Japanese Quail *Coturnix japonica*. Data Zone. 2024. Available at: <https://datazone.birdlife.org/species/factsheet/japanese-quail-coturnix-japonica> (Accessed: 22 July 2024).
- [5] Rehman A, Hussain J, Mahmud A, Javed K, Ghayas A, Ahmad S. Productive performance, egg quality, and hatching traits of Japanese quail lines selected for higher body weight and egg number. *Turkish Journal of Veterinary & Animal Sciences*. 2022; 46:322–335.
- [6] Reda GK, Ndunguru SF, Csernus B, Knop R, Lugata JK, Szabó C, *et al.* Dietary restriction reveals sex-specific expression of the mTOR pathway genes in Japanese quails. *Scientific Reports*. 2024; 14: 8314.
- [7] Volkovoy S, Bondarenko Yu. Japanese quail plumage rainbow. *Priusadebnoye Khozyaystvo [Allotment Husbandry]*. 1989; 5: 14–15. Available at: <https://yablonka.net/world/zh/686-raduga-opereniya-yaponskogo-perepela.html> (Accessed: 22 July 2024). (In Russian)
- [8] German NY, Vetokh AN, Dzhagaev AY, Ilyina ER, Kotova TO. Morphometric parameters of eggs from breeds quail for meat. *Veterinariâ i Kormlenie [Veterinary and Feeding]*. 2023; 2: 20–23.
- [9] Lan LT, Nhan NT, Hung LT, Diep TH, Xuan NH, Loc HT, *et al.* Relationship between plumage color and eggshell patterns with egg production and egg quality traits of Japanese quails. *Veterinary World*. 2021; 14: 897.
- [10] Mizutani M. The Japanese quail. In Chang HL, Huang YC (eds.) *Relationship between Indigenous Animals and Humans in Apec Region* (pp. 143–163). The Chinese Society of Animal Science: Tainan. 2003. Available at: <https://web.archive.org/web/20240812181743/https://www.angrin.tlri.gov.tw/apec2003/chapter5jpquail.pdf> (Accessed: 22 July 2024).
- [11] IUCN. Japanese Quail *Coturnix japonica*. IUCN Red List of Threatened Species. 2024. Available at: <https://doi.org/10.2305/IUCN.UK.2016-3.RLTS.T22678949A95209097.en> (Accessed: 9 September 2024).
- [12] Abou-Kassem DE, El-Sayiad GA, El-Samahy RA, Abd El-Hack ME, Taha AE, Kamal M, *et al.* Impacts of storage period and egg weight on hatching and growth performance of growing Japanese quails. *Poultry Science*. 2024; 103: 103772.
- [13] Kayang BB, Vignal A, Inoue-Murayama M, Miwa M, Monvoisin JL, Ito S, *et al.* A first-generation microsatellite linkage map of the Japanese quail. *Animal Genetics*. 2004; 35: 195–200.
- [14] Minvielle F. The future of Japanese quail for research and production. *World's Poultry Science Journal*. 2004; 60: 500–507.
- [15] Ashok A, Prabakaran R. Evaluation of growth traits in short term selection for different ages in Japanese quail. *Indian Journal of Fundamental and Applied Life Sciences*. 2012; 2: 152–157. Available at: <https://www.cibtech.org/J-LIFE-SCIEN CES/PUBLICATIONS/2012/Vol%20No%201/28-60%20%20Ashok%20and%20Prabakaran.pdf> (Accessed: 22 July 2024).

- 2024).
- [16] German NY, Volkova NA, Larionova PV, Vetokh AN, Volkova LA, Sermiyagin AA, *et al.* Genome-wide association studies of growth dynamics in quails *Coturnix coturnix*. *Sel'skokhozyaistvennaya Biologiya [Agricultural Biology]*. 2022; 57: 1136–1146.
- [17] Marks HL. Long-term selection for body weight in Japanese quail under different environments. *Poultry Science*. 1996; 75: 1198–1203.
- [18] Taha AE, El-Tahawy AS, Abd El-Hack ME, Swelum AA, Saadeldin IM. Impacts of various storage periods on egg quality, hatchability, post-hatching performance, and economic benefit analysis of two breeds of quail. *Poultry Science*. 2019; 98: 777–784.
- [19] Jatoi AS. Productive Performance of Four Close-bred Flocks of Japanese Quails with Different Body Weight and Its Effect on Subsequent Progeny Growth [PhD's thesis]. Department of Poultry Production, Faculty of Animal Production and Technology, University of Veterinary and Animal Sciences, Lahore, Pakistan. 2012.
- [20] Vali N, Edriss MA, Rahmani HR. Genetic parameters of body and some carcass traits in two quail strains. *International Journal of Poultry Science*. 2005; 4: 296–300.
- [21] Ahmad S, Mehmood S, Javed K, Mahmud A, Usman M, Rehman A, *et al.* Different selection strategies for the improvement of the growth performance and carcass traits of Japanese quails. *Brazilian Journal of Poultry Science*. 2018; 20: 497–506.
- [22] Arunrao KV, Kannan D, Amutha R, Thiruvankadan AK, Yakubu A. Production performance of four lines of Japanese quail reared under tropical climatic conditions of Tamil Nadu, India. *Frontiers in Genetics*. 2023; 14: 1128944.
- [23] Ottinger MA, Abdelnabi M, Li Q, Chen K, Thompson N, Harada N, *et al.* The Japanese quail: a model for studying reproductive aging of hypothalamic systems. *Experimental Gerontology*. 2004; 39: 1679–1693.
- [24] Srivastava R, Cornett LE, Chaturvedi CM. Impact of estrogen and photoperiod on arginine vasotocin and VT3 receptor expression in the shell gland of quail. *Frontiers in Bioscience (Scholar Edition)*. 2018; 10: 372–385.
- [25] Prituzhalova AO, Volkova NA, Kuzmina TI, Vetokh AN, Dzha-gaev AY. Monitoring of indicators of chromatin status in quails ovarian follicles granulosa cells of different directions of productivity. *Agrarnaâ Nauka [Agrarian Science]*. 2023; 368: 53–57.
- [26] Nadeau NJ, Minvielle F, Mundy NI. Association of a Glu92Lys substitution in MC1R with extended brown in Japanese quail (*Coturnix japonica*). *Animal Genetics*. 2006; 37: 287–289.
- [27] Hasan KH. Evaluation of productive performance of Japanese quail in summer of Iraq. *Diyala Agricultural Sciences Journal*. 2013; 5: 69–80.
- [28] Hassan KH, Abd-alsattar AR, Yassen HA, Abed HT, Wahab DA. Effect of variety, age and sex in some productive traits in Japanese quail in Iraq. *American Journal of Bio Science*. 2015; 3: 55–58.
- [29] El-Attrouny MM, Iraqi MM. Influence of selection for egg production on egg quality traits in Japanese quail. *South African Journal of Animal Science*. 2021; 51: 128–137.
- [30] Ahmed LS, Al-Barzinji YM. Comparative study of hatchability and fertility rate among local quails. *Iraqi Journal of Agricultural Sciences*. 2020; 51: 744–751.
- [31] Al-Tikriti SS, Al-Nassery HZ. Effect of egg weight and type of breeding on the productive performance of Japanese quail. *IOP Conference Series: Earth and Environmental Science*. 2023; 1213: 012079.
- [32] Mahmoud M, Iraqi MM, Nassar FS. Heritability and genetic correlations of carcass and meat quality traits in white and brown strains of Japanese quail. *Journal of World's Poultry Research*. 2024; 14: 297–307.
- [33] Mohammed DA, Salih JH, Hussen SH. Effect of different protein sources with different levels on growth performance and carcass dissection of Japanese quail (*Coturnix coturnix japonica*). *Science Journal of University of Zakho*. 2021; 9: 1–7.
- [34] Hassan KH, Fadhil MA. Genetic selection for body weight in Japanese quail (*Coturnix coturnix japonica*) under different nutritional environments. *Advances in Animal and Veterinary Sciences*. 2019; 7: 526–529.
- [35] Al-Obaidi FA, Al-Shadeedi SM. Comparison some native fowls (chicken, mallard ducks, quail and turkey) in components and chemical composition of the eggs in Iraq. *Al-Anbar Journal of Veterinary Sciences*. 2017; 10: 65–69.
- [36] Salih JH, Mohammed DA, Hussen SH. Impact of protein source and its levels on egg production and egg quality of Japanese quail (*Coturnix coturnix japonica*). *Science Journal of University of Zakho*. 2021; 9: 138–143.
- [37] Beski SS. Physiological and immunological responses of Japanese quails to oleobiotic. *The Iraqi Journal of Agricultural Science*. 2018; 49: 194–199.
- [38] Jabbar AA. Gastroprotective and immuno-supportive role of *Alcea kurdica* against stress induced lesion in Japanese quails. *Baghdad Science Journal*. 2022; 19: 716–724.
- [39] Frankham R. Genetic markers for conservation purposes. In *Genetic Management of Fragmented Animal and Plant Populations*. Online edn. Oxford Academic: Oxford. 2017.
- [40] Li Y, Liu C, Wang R, Luo S, Nong S, Wang J, *et al.* Applications of molecular markers in conserving endangered species. *Biodiversity Science*. 2020; 28: 367.
- [41] Romanov MN, Weigend S, Bondarenko YuV, Podstreshny AP, Kutnyuk PI, Sakhatsky NI. Studies on poultry germplasm diversity and conservation in Ukraine. In Preisinger R (ed.) *Proceedings of the Poultry Genetics Symposium* (p. 140). Working Group 3 of WPSA, Lohmann Tierzucht: Cuxhaven. 1999. Available at: <https://kar.kent.ac.uk/46412/> (Accessed: 22 July 2024).
- [42] Moiseyeva IG, Romanov MN, Kovalenko AT, Mosyakina TV, Bondarenko YV, Kutnyuk PI, *et al.* The Poltava chicken breed of Ukraine: its history, characterization and conservation. *Animal Genetic Resources Information*. 2007; 40: 71–78.
- [43] Modi WS, Romanov M, Green ED, Ryder O. Molecular cytogenetics of the California condor: evolutionary and conservation implications. *Cytogenetic and Genome Research*. 2009; 127: 26–32.
- [44] Romanov MN, Koriabine M, Nefedov M, de Jong PJ, Ryder OA. Construction of a California condor BAC library and first-generation chicken-condor comparative physical map as an endangered species conservation genomics resource. *Genomics*. 2006; 88: 711–718.
- [45] Rocco L, Valentino IV, Scapigliati G, Stingo V. RAPD-PCR analysis for molecular characterization and genotoxic studies of a new marine fish cell line derived from *Dicentrarchus labrax*. *Cytotechnology*. 2014; 66: 383–393.
- [46] El-Gendy EA, Helal MA, Goher NH, Mostageer A. Molecular characterization of genetic biodiversity in ducks, using RAPD-PCR analysis. *Arab Journal of Biotechnology*. 2005; 8: 253–264. Available at: <https://www.researchgate.net/publication/237703456> (Accessed: 22 July 2024).
- [47] Romanov MN, Weigend S. Using RAPD markers for assessment of genetic diversity in chickens. *Archiv für Geflügelkunde*. 2001; 65: 145–148. Available at: <https://www.researchgate.net/publication/255710825> (Accessed: 22 July 2024).
- [48] Dehghanzadeh H, Mirhoseini SZ, Romanov MN, Ghorbani A. Evaluation of genetic variability and distances among five Iranian native chicken populations using RAPD markers. *Pakistan Journal of Biological Sciences: PJBS*. 2009; 12: 866–871.

- [49] Abdulrazaq HS, Ameen QA. Genetic relationship between local guinea fowl, quail and chicken using RAPD–PCR technique. *Mesopotamia Journal of Agriculture*. 2023; 51: 39–49.
- [50] Ahmed LS, Al-Barzinji YMS. Genetic diversity among local quail using RAPD–DNA marker. *Revista Científica de la Facultad de Veterinaria*. 2019; 29: 178–185. Available at: <https://www.researchgate.net/publication/332182286> (Accessed: 22 July 2024).
- [51] Al-Neemy MA, Tawfeq EM, Hadi HM. Determine the genetic variation of phenotypic groups of quail (different with feathers color) using the chain polymerase reaction technique (PCR). *IOP Conference Series: Earth and Environmental Science*. 2021; 910: 012110.
- [52] Karabağ K, Balcioglu MS. Genetic diversity among selected Japanese quail (*Coturnix coturnix japonica*) lines using RAPD markers. *Journal of Applied Animal Research*. 2010; 38: 149–152.
- [53] Bozdoğan HM, Özcan BD. Determination of genetic variations in Japanese quails improved for meat and egg production traits using PCR-RAPD method. *Cukurova University Journal of Natural and Applied Sciences*. 2023; 2: 13–19. Available at: <https://dergipark.org.tr/en/download/article-file/3510725> (Accessed: 22 July 2024).
- [54] Podstreshnyi OP, Tereshchenko OV, Katerynych OO, Tkachyk TE, Podstreshna IO. Production of Quail Eggs and Meat: Methodical Recommendations. Tereshchenko OV (ed.). 2nd edn. Poultry Research Institute, NAAS of Ukraine: Birky. 2010. Available at: <https://www.researchgate.net/publication/342802513> (Accessed: 22 July 2024). (In Russian)
- [55] Podstreshnyi O, Tereshchenko O. Maintenance of adult quails. *Ahrarna Krayina [Agrarian Country]*. 2012; 6: 8–9. Available at: <https://www.researchgate.net/publication/342832587> (Accessed: 22 July 2024). (In Russian)
- [56] Podstreshnyi O, Tereshchenko O. Feeding young quails. *Ahrarna Krayina [Agrarian Country]*. 2012; 7: 6. Available at: <https://www.researchgate.net/publication/342832583> (Accessed: 22 July 2024). (In Russian)
- [57] Peakall RO, Smouse PE. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*. 2006; 6: 288–295.
- [58] Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics (Oxford, England)*. 2012; 28: 2537–2539.
- [59] Botstein D, White RL, Skolnick M, Davis RW. Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American Journal of Human Genetics*. 1980; 32: 314–331.
- [60] Kimura M, Crow JF. The number of alleles that can be maintained in a finite population. *Genetics*. 1964; 49: 725–738.
- [61] Nei M. Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences of the United States of America*. 1973; 70: 3321–3323.
- [62] Lewontin RC. The apportionment of human diversity. In: Dobzhansky T, Hecht MK, Steere WC (eds.) *Evolutionary Biology* (pp. 381–398). Springer: New York. 1972.
- [63] Nei M. Genetic distance between populations. *The American Naturalist*. 1972; 106: 283–292.
- [64] Nei M. The theory of genetic distance and evolution of human races. *Japanese Journal of Human Genetics*. 1978; 23: 341–369.
- [65] Al-Kafajy FR, Al-Shuhaib MB, Al-Jashami GS, Al-Thuwaini TM. Comparison of three lines of Japanese quails revealed a remarkable role of plumage color in the productivity performance determination. *Journal of World's Poultry Research*. 2018; 8: 111–119. Available at: [http://www.jwpr.science-line.com/attachments/article/47/J%20World%20Poult%20Res%208\(4\)%20111-119,%202018.pdf](http://www.jwpr.science-line.com/attachments/article/47/J%20World%20Poult%20Res%208(4)%20111-119,%202018.pdf) (Accessed: 22 July 2024).
- [66] Petek M, Çavuşoglu E, Odabaşı F, Yeşilbağ D. Comparison of egg quality characteristics of different quail lines. *Journal of Research in Veterinary Medicine*. 2022; 41: 27–31.
- [67] Panda B, Singh RP. Developments in processing quail meat and eggs. *World's Poultry Science Journal*. 1990; 46: 219–234.
- [68] Santhi D, Kalaikannan A. Japanese quail (*Coturnix coturnix japonica*) meat: characteristics and value addition. *World's Poultry Science Journal*. 2017; 73: 337–344.
- [69] Shaker AS, Hermiz HN, Al-Khatib TR, Mohammed RM. Egg shape characterization for four genetic groups of Kurdish local chickens. *Food and Nutrition Science*. 2016; 1: 20–25. Available at: <https://www.ias.org/iasar/filedownloads/fnsij/2016/013-0004.pdf> (Accessed: 22 July 2024).
- [70] Guo X, Elston RC. Linkage information content of polymorphic genetic markers. *Human Heredity*. 1999; 49: 112–118.
- [71] Elston RC. Polymorphism information content. In *Encyclopedia of Biostatistics*. 2nd edn. John Wiley & Sons: New York. 2014.
- [72] Serrote CML, Reiniger LRS, Silva KB, Rabaiolli SMDS, Stefanel CM. Determining the Polymorphism Information Content of a molecular marker. *Gene*. 2020; 726: 144175.
- [73] Garizábal-Carmona JA, Betancur JS, Montoya-Arango S, Franco-Espinosa L, Ruíz-Giraldo N, Mancera-Rodríguez NJ. Bird diversity across an Andean City: The limitation of species richness values and watershed scales. *Acta Biológica Colombiana*. 2023; 28: 506–516.
- [74] Lajan SA, Al-Barzinji YMS. Detection of quantitative loci correlation with growth traits in local quail using PCR-RFLP technique. *Iraqi Journal of Agricultural Sciences*. 2022; 53: 26–16.
- [75] Romanov MN, Weigend S. Genetic diversity in chicken populations based on microsatellite markers. In Dekkers JCM, Lamont SJ, Rothschild MF (eds.) *From Jay Lush to Genomics: Visions for Animal Breeding and Genetics, Conference [Proceedings]* (p. 174). Iowa State University, Department of Animal Science: Ames. 1999.
- [76] Nunome M, Nakano M, Tadano R, Kawahara-Miki R, Kono T, Takahashi S, et al. Genetic divergence in domestic Japanese quail inferred from mitochondrial DNA D-loop and microsatellite markers. *PloS One*. 2017; 12: e0169978.
- [77] Mathur S, Tomeček JM, Heniff A, Luna R, DeWoody JA. Evidence of genetic erosion in a peripheral population of a North American game bird: the Montezuma quail (*Cyrtonyx montezumae*). *Conservation Genetics*. 2019; 20: 1369–13681.
- [78] Dementeva NV, Romanov MN, Kudinov AA, Mitrofanova OV, Stanishevskaya OI, Terletsky VP, et al. Studying the structure of a gene pool population of the Russian White chicken breed by genome-wide SNP scan. *Sel'skokhozyaistvennaya Biologiya [Agricultural Biology]*. 2017; 52: 1166–1174.
- [79] Mathur S. An Evolutionary Genomics Study for Conservation of the Montezuma Quail. Doctoral dissertation, Purdue University: West Lafayette, IN, USA. 2020
- [80] Dementieva NV, Shcherbakov YS, Tyshchenko VI, Terletsky VP, Vakhrameev AB, Nikolaeva OA, et al. Comparative analysis of molecular RFLP and SNP markers in assessing and understanding the genetic diversity of various chicken breeds. *Genes*. 2022; 13: 1876.
- [81] Sponenberg DP, Beranger J, Martin AM, Couch CR. Conservation of rare and local breeds of livestock. *Revue Scientifique et Technique (International Office of Epizootics)*. 2018; 37: 259–267.
- [82] Albers GA, Rattink AP, Vereijken AL. The future of molecular genetics in poultry breeding. In *EPC 2006 – 12th European Poultry Conference (Paper 355)*. World's Poultry Science Association: Verona. 2006. Available at: <https://www.researchgate.net/publication/254200124> (Accessed: 22 July 2024).
- [83] Negash F, Tadesse Y, Getachew A. Genomic selection for poul-

- try breeding and its potential applications in developing countries. *Poultry Science Journal*. 2021; 9: 133–145.
- [84] Baumgartner J, Bondarenko YuV. Search for autosexing strains and crosses in Japanese quail. In Proceedings of the 8th International Symposium on Actual Problems of Avian Genetics (pp. 262–265). Slovak Society for Agriculture, Forestry, Food and Veterinary Sciences of Slovak Academy of Sciences; Poultry Research and Production Institute Bratislava; Czechoslovak Branch of WPSA; Smolenice. 1989. Available at: <https://www.cabidigitallibrary.org/doi/full/10.5555/19890171986> (Accessed: 22 July 2024).
- [85] Faraji-Arough H, Maghsoudi A, Rokouei M. Study of genetic and non-genetic effects on cumulative survival in a crossbred population of quail. *Tropical Animal Health and Production*. 2022; 55: 5.
- [86] Ibrahim FK, Sabri MA, Mohamed AA. Estimating genetic parameters for productive qualities in local and Ukrainian quail and their various crossbreeds. *NTU Journal of Agriculture and Veterinary Science*. 2024; 4. Available at: <https://journals.ntu.edu.iq/index.php/NTU-JAVS/article/view/814> (Accessed: 22 July 2024).
- [87] Khalil MH, Elattar EA, El-Seedy AS, Shebl MK. Detection of single-nucleotide polymorphisms in growth hormone and insulin-like growth factor-1 genes related to growth traits in purebred and crossbred quails. *Veterinary World*. 2024; 17: 1482–1489.
- [88] Wu Y, Zhang Y, Hou Z, Fan G, Pi J, Sun S, *et al*. Population genomic data reveal genes related to important traits of quail. *GigaScience*. 2018; 7: giy049.
- [89] Kayang BB, Fillon V, Inoue-Murayama M, Miwa M, Leroux S, Fève K, Monvoisin JL, Pitel F, Vignoles M, Mouilhayrat C, Beaumont C. Integrated maps in quail (*Coturnix japonica*) confirm the high degree of synteny conservation with chicken (*Gallus gallus*) despite 35 million years of divergence. *BMC Genomics*. 2006; 7: 1–8.
- [90] Yuan Z, Zhang X, Pang Y, Qi Y. Association analysis of PMEL gene expression and single nucleotide polymorphism with plumage color in quail. *Animal Biotechnology*. 2023; 34: 5001–5010.
- [91] Santos TC, Gates RS, Tinôco IFF, Zolnier S, Rocha KSO, Freitas LCSR. Productive performance and surface temperatures of Japanese quail exposed to different environment conditions at start of lay. *Poultry Science*. 2019; 98: 2830–2839.
- [92] Petek M, Ozen Y, Karakas E. Effects of recessive white plumage colour mutation on hatchability and growth of quail hatched from breeders of different ages. *British Poultry Science*. 2004; 45: 769–774.
- [93] Aydin CE, Petek ME, Çibik RE. Effect of recessive colour mutation on haematological characteristics of Japanese quail (*Coturnix coturnix japonica*). *Archiv für Geflügelkunde*. 2008; 72: 164–167. Available at: <https://avesis.uludag.edu.tr/yayin/d278d550-5501-4467-b53b-a5df3cd9ea31/effect-of-recessive-colour-mutation-on-haematological-characteristics-of-japanese-quail-coturnix-coturnix-japonica> (Accessed: 22 July 2024).
- [94] Ryabokon YuO, Pabat VO, Mykytyuk DM, Frolov VV, Katerynych OO, Bondarenko YuV, *et al*. Catalog of Poultry Breeding Resources of Ukraine. Ryabokon YuO (ed.) Poultry Research Institute: Kharkiv. 2005. Available at: <http://avianua.com/archiv/plevreestr/per.pdf> (Accessed: 22 July 2024). (In Russian)