



# Kent Academic Repository

**Menajovsky, Maria F, Mayor, Pedro, Bodmer, Richard, Pérez-Peña, Pedro, Ulloa, Gabriela M, Greenwood, Alex D, Montero, Stephanie, Lescano, Andrés G, Santolalla, Meddy L, Segalés, Joaquim and others (2025) *Monitoring of Selected Swine Viral Diseases in Peruvian Amazon Peccaries*. *EcoHealth*, 22 . pp. 69-78. ISSN 1612-9202.**

## Downloaded from

<https://kar.kent.ac.uk/108531/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1007/s10393-024-01692-9>

## This document version

Publisher pdf

## DOI for this version

## Licence for this version

CC BY (Attribution)

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

## Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

## Original Contribution

# Monitoring of Selected Swine Viral Diseases in Peruvian Amazon Peccaries

Maria F. Menajovsky,<sup>1</sup> Pedro Mayor<sup>1,2,3</sup>, Richard Bodmer,<sup>2,3,4</sup> Pedro Pérez-Peña,<sup>5</sup> Gabriela M. Ulloa,<sup>6,7,8</sup> Alex D. Greenwood,<sup>9,10</sup> Stephanie Montero,<sup>6</sup> Andrés G. Lescano,<sup>6,11</sup> Meddly L. Santolalla,<sup>6</sup> Joaquim Segalés,<sup>1,12,13</sup> Marina Sibila,<sup>12,13,14</sup> Oscar Cabezón,<sup>15</sup> and Johan Espunyes<sup>15</sup>

<sup>1</sup>Departament de Sanitat i Anatomia Animals, Facultat de Veterinària, Universitat Autònoma de Barcelona, Edifici V, Bellaterra, 08193 Barcelona, Spain

<sup>2</sup>ComFauna, Comunidad de Manejo de Fauna Silvestre en la Amazonía y en Latinoamérica, Iquitos, Peru

<sup>3</sup>Museo de Culturas Indígenas Amazónicas, Iquitos, Peru

<sup>4</sup>School of Anthropology and Conservation, Durrell Institute of Conservation and Ecology, University of Kent, Canterbury, UK

<sup>5</sup>Instituto de Investigaciones de la Amazonía Peruana (IIAP), Iquitos, Peru

<sup>6</sup>Emerge, Emerging Diseases and Climate Change Research Unit, School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Lima, Peru

<sup>7</sup>Programa de Pós-Graduação em Saúde e Produção Animal na Amazônia, Universidade Federal Rural da Amazônia (UFRA), Pará, Brazil

<sup>8</sup>Grupo Enfermedades Emergentes, Universidad Científica del Sur, Lima, Peru

<sup>9</sup>Department of Wildlife Diseases, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany

<sup>10</sup>School of Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

<sup>11</sup>Clima, Latin American Center of Excellence for Climate Change and Health, and Emerge, Emerging Diseases and Climate Change Research Unit, Universidad Peruana Cayetano Heredia, Lima, Peru

<sup>12</sup>Centre de Recerca en Sanitat Animal (CRESA), Unitat Mixta d'Investigació IRTA-UAB en Sanitat Animal, Campus de la Universitat Autònoma de Barcelona (UAB), 08193 Bellaterra, Catalonia, Spain

<sup>13</sup>WOAH Collaborating Centre for the Research and Control of Emerging and Re-Emerging Swine Diseases in Europe (IRTA-CreSA), 08193 Bellaterra, Spain

<sup>14</sup>IRTA. Programa de Sanitat Animal. Centre de Recerca en Sanitat Animal (CRESA), Campus de la Universitat Autònoma de Barcelona (UAB), 08193 Bellaterra, Spain

<sup>15</sup>Wildlife Conservation Medicine Research Group (WildCoM), Departament de Medicina i Cirurgia Animals, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

**Abstract:** Peccaries (collared peccary—CP—and white-lipped peccary—WLP) are an essential source of protein and income for rural communities in the Amazon region. Since 1980s, researchers in the Amazon have reported recurrent local disappearances of WLP populations. Although such disappearances impact the species conservation and the food security of rural societies, no studies have drawn consistent conclusions about the causes of these population collapses. However, it has recently been proposed that the overabundance of this species before its decline would be related to infectious disease outbreaks. In the current study, we aimed to determine the circulation (occurrence and exposure) of viruses relevant to swine health in CP and WLP

populations, namely classical swine fever virus (CSFV), Aujeszky's disease virus (ADV), swine vesicular disease virus (SVDV), and porcine circoviruses (PCV). The study was conducted in two areas of the northeastern Peruvian Amazon: the Yavarí-Mirín River basin (2008–2020), where WLPs experienced extreme population fluctuations, and the Pucacuro National Reserve (2012–2014), where no WLP disappearances have been reported. Since WLP is not easily found during population declines, we also sampled CP as an indicator of virus circulation in the area as they are likely to be susceptible to the same pathogens. CSFV and ADV antibodies were detected in both peccary species and both areas. Diseases caused by CSFV and ADV have the potential to act as ultimate causes of population collapse, especially in large WLP populations where overabundance could increase the rate of pathogen transmission. Our results were inconclusive in establishing whether or not these viruses drove the WLP population to collapse, but their potential role warrants deeper investigation, expanding the geographical coverage of studies on infectious diseases in peccaries.

**Keywords:** *Tayassu pecari*, *Pecari tajacu*, Epidemiology, Population dynamics, Classical swine fever, Aujeszky's disease

## INTRODUCTION

Peccaries (Cetartiodactyla: Tayassuidae) are native to Latin America and are distributed from Mexico to northern Argentina (Sowls, 1997). They play a pivotal role in ecosystem dynamics by rooting soils, dispersing seeds and seedlings, consuming plant and animal matter, and serving as prey for top predators (e.g., *Panthera onca* and *Puma concolor*) (Sobral et al., 2017; Weckel et al., 2006; Porfirio et al., 2017). In the Amazon, two species of peccaries coexist: the white-lipped peccary (WLP, *Tayassu pecari*) and the collared peccary (CP, *Pecari tajacu*).

Subsistence hunting is essential to traditional livelihoods and an important source of protein and income for rural communities—including indigenous and campesino communities—and cities in the Amazon region (Bodmer & Pezo, 2001; Mayor et al., 2022). In the early 2000s, the annual harvest in the rural Peruvian and the Brazilian Amazon was estimated to be approximately 14,000 WLP and 20,000 CP individuals (Bodmer & Pezo, 2001), and 611,527 WLP and 551,949 CP individuals (Peres, 2000), respectively. In addition, peccaries are also of enormous socioeconomic importance as they are the most traded wild meat species in urban markets, accounting for 43 tons per year (17.9% of the total biomass traded) of WLP meat and 97 tons (39.8%) of CP meat in Iquitos, the largest urban market in the Peruvian Amazon (Mayor et al., 2022).

Since 1980s, researchers have reported recurrent local disappearances of WLP populations across their Latin American range, spanning at least 50 million hectares (Richard-Hansen et al., 2014; Fragoso et al., 2016; 2022), and affecting the ecosystem dynamics, as well as the food

supply and economic activities of indigenous and rural communities (Fragoso et al., 2016; 2022). These population cycles appear to last 20–30 years, with rapid population decline occurring in the first 1–5 years, followed by the absence or low abundance for the next 7–12 years, and finally a slow growth over the next 20 years (Fragoso et al., 2022). However, not all populations recover. Currently, the WLP, now considered 'Vulnerable' by the *International Union for Conservation of Nature* (IUCN, 2023), is found in only 21% of its recognized distribution geographic range (Taber et al., 2021).

Early studies attributed these populations' oscillations to migrations, overhunting, and environmental changes (such as land-use changes), but none have reached consistent conclusions (Altrichter et al., 2012; Fragoso et al., 2022). The density-dependent overcompensation theory has recently gained prominence, suggesting that WLP population overabundance triggers density-dependent effects, such as increased pathogen transmission, resulting in disease outbreaks and population decline (Fragoso et al., 2022). Even though both WLP and CP are influenced by the same environmental conditions, have similar reproduction cycles (Mayor et al., 2017), and are similar hunting preferences for subsistence hunters (Pérez-Peña et al., 2021), such disappearance cycles have not been reported for CP. However, there are several ecological differences between the two species. WLP weigh 30–50 kg and form herds of around 400 individuals with a home range of 16–200 km<sup>2</sup> (Fragoso, 1998; Kiltie & Terborgh, 1983), whereas CP weigh 25 kg on average and form herds of 3–20 individuals with a home range of 1–7 km<sup>2</sup> (Keuroghlian et al., 2004). In general, WLP occupy densities ranging from 3.7

to 25 individuals/km<sup>2</sup>, while CP occupy densities ranging from 2.8 to 9.8 individuals/km<sup>2</sup> (Fragoso, 1998; Keuroghlian et al., 2004). These differences may influence how animals respond to pathogens and disease outbreaks (Fragoso et al., 2022).

Several swine viruses, including classical swine fever virus (CSFV), Aujeszky's disease virus (ADV), and swine vesicular disease virus (SVDV), as well as various porcine circoviruses (PCV), have a significant impact on swine health, affecting the reproductive, nervous, respiratory, and gastrointestinal systems, with high associated mortality rates that can reach 100% (Postel et al., 2018; Zuckermann, 2000; WOA, 2023). However, to our knowledge, there is a significant gap in information about the occurrence and health impact of infectious diseases on wildlife in the Amazon region. Indeed, diseased WLP and carcasses have been observed during periods of population decline in some regions, but further examination was not carried out (Fragoso et al., 2022). In the Southern Peruvian Amazon, studies have reported the occurrence of antibodies against ADV in free-living WLP populations (Romero Solorio, 2010). Also, ADV and porcine circovirus 2 (PCV-2) have been reported in CP and WLP populations in the Bolivian and Brazilian Amazon but ignoring their impact on their population dynamics (Karesh et al., 1998; De Castro et al., 2014). PCV-2 has also been documented among domestic pigs in the Southern Brazilian Amazon (Dutra et al., 2013). Despite these few studies, pathogens pertinent to swine health in the Amazon region remain largely underexplored (Menajovsky et al., 2023).

WLP populations can experience explosive population growth (Fang et al., 2008) and due to the risk and rate of infectious disease transmission is higher in dense populations (Tarwater & Martin, 2001), we hypothesized that highly virulent infectious diseases may be involved in the severe population declines of WLP in the Amazon. However, the logistic difficulties of studying wildlife diseases in such remote areas are a major challenge, as evidenced by a lack of information about infectious diseases and their role in the decline of wild species populations, such as WLP (Menajovsky et al., 2023).

The present study aimed to improve the knowledge about the circulation of selected infectious diseases in areas of WLP fluctuations by assessing the seroprevalence and presence of major swine pathogens in CP and WLP populations in the Peruvian Amazon.

## MATERIALS AND METHODS

### Study Area

The study was carried out in two areas of the Peruvian Amazon: the Yavarí-Mirín River (YMR) basin and the Pucacuro National Reserve. The Yavarí-Mirín River basin (YMR; 04°19'53"S; 71°57'33"W) is a remote area on the Peru-Brazil border, composed of a diverse landscape that ranges from upland forests with nutrient-poor sandy soils to flooded forests with relatively nutrient-rich soils (Ter Steege et al., 2003). The only village in YMR is Nueva Esperanza, a Yagua indigenous community of around 300 people. The main human activities of this community are traditional small-scale agriculture, fishing, logging, and subsistence hunting (Mayor et al., 2015). There is no large-scale agriculture in the study area; therefore, the crops are subsistence-based. In the YMR basin, WLPs have shown extreme population fluctuations over the last 25 years, including a decline from a high of 15 ind./km<sup>2</sup> in 2000 to 2 ind./km<sup>2</sup> in 2004 and a complete disappearance between 2005 and 2015 (Fang et al., 2008; Bodmer et al., 2024).

The Pucacuro National Reserve (PNR), located on the border with Ecuador (2°26'53"S 75°20'29"W), is composed of high terrace forests, non-flooded habitat with dissected relief in a humid tropical forest. PNR has a game species management plan that includes CP, WLP, and *Cuniculus paca*. Due to their large population sizes and relatively high reproductive capacity, management groups hunt these species for consumption and commercialization to support local indigenous hunters' economies (SERNANP, 2014). There have been no reports of WLP or CP population declines in PNR (Perez Peña et al., 2016).

Both study areas are highly biodiverse and well-preserved, with low human impact (Pitman et al., 2003; SERNANP, 2019), and the nearest pig farms are 160 and 170 km away, respectively, from YMR and PNR (Fig. 1).

### Biological Collection

Blood samples were obtained from both study areas, totaling samples from 98 WLP and 140 CP. The samples were collected by subsistence hunters and park rangers as part of a wildlife conservation program, taking advantage of the discarded material from legal subsistence hunting. Hunters impregnated blood from the cranial or caudal cava veins on either Whatman filter paper n. 3 or FTA® cards (Scheilcher & Schuell, Germany). The YMR basin was

sampled from 2008 to 2015 and 2019 to 2020, while the PNR was sampled from 2012 to 2014. Samples were collected in all seasons, and hunters recorded the species, date, location, and sex. The samples were kept at room temperature in a sheltered environment in the study areas for 15 to 100 days before being stored at  $-20^{\circ}\text{C}$  until analyses (Aston et al., 2014; Menajovsky et al., 2024; Morales et al., 2017).

### Serological Analyses

A  $132\text{ mm}^2$  area of the filter paper with blood was cut with a sterile punch (Staples®) and diluted in  $400\text{ }\mu\text{L}$  of sterile PBS (phosphate-buffered saline). These samples were vortexed for 20 s, stored at  $4^{\circ}\text{C}$  for 24 h before being vortexed for 20 s, and frozen at  $-20^{\circ}\text{C}$  until analysis. As serum represents 40% of total blood volume (Nobuto, 1966) and

the blood concentration on filter paper is  $40\text{ }\mu\text{L}/\text{cm}^2$ , we obtained a serum dilution of 1:20. The elutions were tested for antibodies against CSFV, ADV, and SVDV using the ID Screen Classical Swine Fever E2 Competition ELISA kit, the ID Screen Aujeszky gB Competition ELISA kit, and the ID Screen Swine Vesicular Disease Competition (IDvet, Montpellier, France), respectively.

### Molecular Analyses

RNA was extracted from  $200\text{ }\mu\text{L}$  of the elutions using the commercial kit IndiMag® Pathogen Kit (Indical Bioscience GmbH, Leipzig, Germany) according to the manufacturer's instructions. RNA concentrations were determined using the Qubit Fluorometric Quantification High Sensitivity Assay (Invitrogen, California, USA). Molecular detection of porcine circoviruses 1, 2, 3, and 4, as well as CSFV, was



**Figure 1.** Map of the study area, including the hunting areas of Nueva Esperanza community in the Yavari Mirin River basin and the Pucacuro National Reserve. The map also shows all pig farms registered in the region (squares; SENASA, pers. comm.).



performed by conventional and RT-PCR, following previously described protocols (Vilček et al., 1994; Quintana et al., 2002; Oliver-Ferrando et al., 2016; Franzo et al., 2020; Saporiti et al., 2021). The 95% confidence intervals for antibodies and pathogen prevalence were calculated using the Wilson score method (Thrusfield, 2018).

### Statistical Analysis

We used Fisher's test to compare seroprevalences of CSFV and ADV between all the animals analyzed in YMR and PNR, as well as between the two analyzed periods in YMR.

### Ethics

Field and laboratory procedures were previously authorized by Peruvian and International Authorities: Dirección General de Flora y Fauna Silvestre from Peru (041–2007-DGGFS-DGEFFS, 0350–2012-DGFFS-DGEFFS, 258–2019-MINAGRI-SERFOR-DGGSPFFS), the Head of the National Reserve of Pucacuro (03–2012-SERNANPRN Pucacuro), and the Institutional Animal Use Ethics Committee from the Universidad Peruana Cayetano Heredia (029–03-19, protocol #102,142) and the *Universitat Autònoma de Barcelona* (protocol #4829). Export permits were as follows: 003258-CITES-Perú, 003260-CITES-Perú, BB-00017 20I-Spain, and BB-00018 20I-Spain.

## RESULTS

Six specimens presented antibodies against CSFV, three CP and three WLP, for an overall seroprevalence of 2.14% (CI<sub>95%</sub>: 0.70–6.11%) in CP and 3.06% (CI<sub>95%</sub>: 1.05–8.62) in WLP (Table 1). Seropositive CPs were only detected in YMR in 2009, 2013, and 2014, during the WLP population crash in the area. One seropositive WLP was detected in YMR in 2020 and two seropositive individuals in PNR in 2014. Antibodies against ADV were detected in 3.06% WLPs (3/98; CI<sub>95%</sub>: 1.05–8.62). Two seropositive individuals were hunted in YMR in 2019, during WLP population recovery, and one in PNR in 2014 (Table 1). All samples were negative for antibodies against SVDV. Molecular analyses for PCV-1, PCV-2, PCV-3, PCV-4, and CSFV were all negative.

Fisher tests did not reveal any significant difference in the seroprevalence between both areas or both periods.

## DISCUSSION

Peccaries are essential to the food security of rural societies in the Amazon region (Bodmer & Pezo, 2001; Fragozo et al., 2022). Even though the extreme WLP population declines have led to its recategorization as “Vulnerable” by the IUCN (Keuroghlian et al., 2013), the impact of infectious diseases on these declines has not been assessed

**Table 1.** Results for the Detection of Antibodies Against Classical Swine Fever Virus (CSFV) and Aujeszky's Disease Virus (ADV) in White-Lipped Peccary (WLP; *Tayassu pecari*) and Collared Peccary (CP; *Pecari tajacu*) in the Yavarí-Mirín River Basin and the Pucacuro National Reserve (Peruvian Amazon).

Area	Species	Period	CSFV	ADV
Yavarí-Mirín River basin	WLP	2008–2015	0/6 (0%) [0.00–39.03%]	0/6 (0%) [0.00–39.03%]
		2019–2020	1/49 (2.04%) [0.36–10.69%]	2/49 (4.08%) [1.13–13.71%]
		Total	1/55 (1.82%) [0.32–9.61]	2/55 (3.64%) [1.00–12.32%]
	CP	2008–2015	3/98 (3.06%) [1.05–8.62]	0/98 (0%) [0.00–3.77%]
		2019–2020	0/39 (0%) [0.00–8.97%]	0/39 (0%) [0.00–8.97%]
		Total	3/137 (2.20%) [0.75–6.24]	0/137 (0%) [0.00–2.73]
Pucacuro National Reserve	WLP	2012–2014	2/43 (4.65%) [1.28–15.46]	1/43 (2.33%) [0.41–12.06]
	CP	2012–2014	0/3 (0%) [0.00–56.15]	0/3 (0%) [0.00–56.15]

The 95% confidence intervals are presented in square brackets.  
CSFV Classical swine fever virus, ADV Aujeszky's disease virus.

appropriately. Wildlife population declines are usually multifactorial processes, even if one cause can be identified as predominantly responsible (Sodhi et al., 2009). Many species are already threatened due to habitat fragmentation, declining genetic diversity, or overexploitation (Frank et al., 2006), and their combination with infectious diseases increases even more the risk of local extinctions (Smith et al. 2006).

Throughout the Amazon, at least 28 independent WLP disappearance events have been recorded (Fragoso et al., 2022). Since there is no published data on the infectious diseases' impact in peccaries, and it is not clear which pathogens may be present in peccary populations, their assessment should be carefully addressed. In this sense, a recent study highlighted the necessity of performing longitudinal studies on infectious diseases in peccaries to address this gap of knowledge (Menajovsky et al., 2023).

The present study confirmed the exposure of Peruvian peccaries to CSFV and ADV. These viruses cause disease in domestic and feral pigs, being the latest ones considered as reservoir hosts for domestic pigs worldwide (Postel et al., 2018; Zuckermann, 2000). Classical swine fever (CSF) has not been previously reported in the Amazon region, but it is considered an endemic disease in domestic pigs from other regions in Brazil and Peru (Pereda et al., 2005; De Oliveira et al., 2020), and its circulation has also been reported in CP in non-Amazon areas in Colombia (Montenegro et al., 2018). Our results present evidence of peccaries contact with the virus in the Amazon region, as both WLP and CP were exposed to CSFV in both study areas. However, our results are inconclusive in establishing whether or not these viruses drove the WLP population to collapse. On the one hand, WLP were almost absent in the YMR basin between 2008 and 2015 ( $n = 6$ ; 0% seroprevalence), when antibodies against CSFV were detected in collared peccaries ( $n = 98$ ; 3.06% seroprevalence), tipping the balance in favor of the hypothesis of CSFV as a modulator of population dynamics of WLP. Antibodies against CSFV were only detected in WLP during the following years (2019–2020) when the population had recovered. On the other hand, antibodies against CSFV were detected in WLP from the PNR, where populations did not decline. Since CSFV is highly contagious and can cause large mortality outbreaks in domestic pigs and wild boars, a significant role of CSFV in the population declines of peccaries cannot be completely ruled out and warrants deeper investigation.

Aujeszky's disease can also cause significant outbreaks in domestic pigs with high mortality and major economic consequences (Zuckermann, 2000). This suggests that the effect of the virus on the population dynamics of free-ranging wild boar appears could be more limited, probably due to the circulation of attenuated strains, as the experimental infection with virulent strains results in severe disease and death (Müller et al. 2001). As with CSFV, the susceptibility and impact of ADV on peccaries are still unknown but potentially significant in case of virulent strains. Seropositivity for antibodies against ADV has been previously observed in WLP from the Southern Peruvian and Bolivian Amazon, but not in CP (Romero Solorio, 2010; Karesh et al., 1998). In our study, antibodies were detected only in WLP and in both study areas, with similar low seroprevalences as in the previous studies. As with CSFV, antibodies were not detected in the YMR basin during the decline in the WLP population and were only detected in the following years. Similarly, these findings are inconclusive in establishing the role of ADV in the declines of WLP populations. However, they deserve further attention as the impact of the virus on peccaries is still unknown. In addition, ADV does not only affect swine species but can also cause significant mortality in carnivorous mammals, for example, in dogs and in wild species as *Puma concolor* (Zhang et al., 2015; Cunningham et al., 2021). Thus, the presence of ADV in peccaries may represent a conservation issue for both peccaries and endangered carnivore species.

We consider it unlikely that the pathogens studied were introduced through pig farms. Feral pigs have not been reported in either YMR or PNR, and pig farms are located near Iquitos, at least 160 km away from both sampling areas (Fig. 1). The maximum home range of CP is 7km<sup>2</sup> and of WLP is 100–200km<sup>2</sup> (Taber et al., 1994; Fragoso, 1998; Kiltie & Terborgh, 1983), and the presence of large rivers in the area as ecological barriers hinder contact with the aforementioned farms. Additionally, in front of our study, area lies the Brazilian territory, which includes Brazil's largest Indigenous Reserve. These are areas where farming is prohibited (Verissimo et al., 2011). Consequently, we can reasonably infer that there is no spillover of pathogens from domestic pigs in the frontier area. However, the presence of traditional small-scale pig farming cannot be ruled out and should be further evaluated to understand the potential role of small-scale farming in the introduction of swine viruses in the free-living Amazon. In fact, it is possible that previous contacts with

backyard pigs could have introduced these viruses within the free-living peccary population, where they have been naturally transmitted between and within their populations.

Antibodies against SVDV and genome from PCV-1, PCV-2, PCV-3, PCV-4, and CSFV were not detected in any sample from the study. Swine vesicular disease virus and the four porcine circoviruses are associated with different degrees of disease in pigs. From these, only PCV-2 has previously been reported in wild peccaries and domestic pigs in the Colombian Amazon (Montenegro et al., 2018) and the Brazilian Amazon and Pantanal (De Castro et al., 2014; Dutra et al., 2013). SVDV and PCV have not been reported in peccaries nor domestic pigs in the Amazon basin (Menajovsky et al., 2023), supporting our findings and evidencing their improbable role in the population dynamics of WLP.

Some technical constraints, such as sample conservation, must be considered when interpreting our results. Filter paper is a convenient technique when working with local communities in logistically challenging remote areas because it simplifies sample collection and storage, especially since an adequate cold chain is unavailable. Local communities rely on subsistence hunting for food and could become active samplers of valuable biological material that is often discarded. In previous studies, these same samples have already been used successfully to perform serological and molecular DNA analyses (Aston et al., 2014; Morales et al., 2017); however, it is important to consider that storage at room temperature under ambient conditions could potentially impact sample stability. This can lead to a higher likelihood of false negative results compared to analyzing frozen biological samples (Bevins et al., 2016), particularly when working with RNA viruses. Therefore, our results must be considered conservative, and the reported seroprevalence may be lower than reality. Furthermore, it is worth noting that PNR was sampled more than 10 years ago, and since then, the health status of the peccaries' populations in the area might have changed. Additionally, we obtained few samples in the CP sampling in PNR and during the WLP population crash in YVR, these negative results may be underestimating circulating antibodies against CSFV, ADV or SVDV.

Determining infectious diseases' impact on WLP disappearance episodes is challenging. The presence of a pathogen in a declining population does not imply that the pathogen played a significant role in the decline (McCallum, 2012). Baseline data still needs to be included to

model the epidemiology and the impact of viruses like CSFV and ADV on peccaries (Menajovsky et al., 2023). Due to viral strains with different severity of disease (Pereda et al., 2005; De Oliveira et al., 2020), isolation and phylogenetic analysis of circulating strains in free-ranging peccaries and experimental infections are required to evaluate and understand their pathogenesis and potential effects. Expanding the geographical coverage of studies on infectious diseases in peccaries to include other areas where population declines have been reported would assist in determining whether pathogens are associated to population declines. We focused our study on viruses known to have a significant impact on swine health, but bacterial, parasitic, and other viral infections may also have repercussions on populations dynamics and should be questioned further. Because the fluctuations of WLP populations are likely to be multifactorial, it is also essential to include strategies for monitoring population dynamics that allow anticipating the population decline to develop surveillance before and during a WLP disappearance event.

## ACKNOWLEDGEMENTS

---

This work has received financial support from the ERANet-LAC [grant 17/HLH-0271], and contract N° 136-2018-FONDECYT. Menajovsky MF acknowledges a doctoral scholarship from the Catalan Agency for Management of University and Research Grants [scholarship FI-SDUR EMC/3345/2020]. Ulloa GM received a fellowship from the PPGSPAA program, of the CNPq in GD modality (140312/2020-0). Santolalla ML and Lescano A are supported by training grant D43 TW007393 awarded by the Fogarty International Center of the US National Institutes of Health. Finally, the authors would like to acknowledge local Amazonian communities for their help in collecting biological samples and Saul Daniel Vilchez from the Peruvian National Agrarian Health Service for the information on regional pig farms.

## FUNDING

---

Open Access Funding provided by Universitat Autònoma de Barcelona. Fondo Nacional de Desarrollo Científico, Tecnológico y de Innovación Tecnológica, 136-2018-FONDECYT, ERANet-LAC, 17/HLH-0271.



## DECLARATIONS

**CONFLICT OF INTEREST** The authors declare no competing interests.

**ETHICAL APPROVAL** All procedures performed in studies involving human participants were in accordance with the ethical standards of the Institutional and/or National Research Committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**HUMAN OR ANIMAL RIGHTS** All applicable institutional and/or national guidelines for the care and use of animals were followed.

## OPEN ACCESS

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## REFERENCES

- Altrichter M, Taber A, Beck H, Reyna-Hurtado R, Lizarraga L, Keuroghlian A, Sanderson EW (2012) Range-wide declines of a key Neotropical ecosystem architect, the near threatened white-lipped peccary *Tayassu pecari*. *Oryx* 46(1):87–98. <https://doi.org/10.1017/S0030605311000421>
- Aston EJ, Mayor P, Bowman DD, Mohammed HO, Liotta JL, Kwok O, Dubey JP (2014) Use of filter papers to determine seroprevalence of *Toxoplasma gondii* among hunted ungulates in remote Peruvian Amazon. *International Journal for Parasitology: Parasites and Wildlife* 3(1):15–19
- Bevins S, Pappert R, Young J, Schmit B, Kohler D, Baeten L (2016) Effect of storage time and storage conditions on antibody detection in blood samples collected on filter paper. *Journal of Wildlife Diseases* 52(3):478–483
- Bodmer RE, Puertas P, Fang T, Antúnez M, Soplin S, Caro J, Mayor P (2024). Management of Subsistence Hunting of Mammals in Amazonia: A Case Study in Loreto, Peru. In *Amazonian Mammals: Current Knowledge and Conservation Priorities* (pp. 275–297). Cham: Springer International Publishing.
- Bodmer RE, Pezo E (2001) Rural development and sustainable wildlife use in Peru. *Conservation Biology* 15(4):1163–1170. <https://doi.org/10.1046/j.1523-1739.2001.0150041163.x>
- Cunningham MW, Onorato DP, Saylor KA, Leone EH, Conley KJ, Mead DG, Waltzek TB (2021) Pseudorabies (Aujeszky's disease) is an underdiagnosed cause of death in the Florida panther (puma concolor coryi). *The Journal of Wildlife Diseases* 57(4):784–798
- De Castro AMMG, Brombila T, Bersano JG, Soares HS, de Souza Silva SO, Minervino AHH, Ogata RA, Gennari SM, Richtzenhain LJ (2014) Swine infectious agents in *Tayassu pecari* and *Pecari tajacu* tissue samples from Brazil. *Journal of Wildlife Diseases* 50(2):205–209. <https://doi.org/10.7589/2013-01-021>
- De Oliveira LG, Gatto IRH, Mechler-Dreibi ML, Almeida HMS, Sonálio K, Storino GY (2020) Achievements and challenges of classical swine fever eradication in Brazil. *Viruses* 12:1327
- Dutra V, Chitarra CS, Paula AJ De, Carolina J, Carolina A, Faria S De, JX, Filho O, Marques M, Oliveira CS De, Pescador CA, Nakazato L (2013) Detection of *Pasteurella Multocida* by qPCR Associated with Pneumonic Lung in Pigs Slaughtered in Mato Grosso Brazil. *International Journal of Sciences*, 2.
- Epstein PR (2001) Climate change and emerging infectious diseases. *Microbes and Infection* 3(9):747–754. [https://doi.org/10.1016/S1286-4579\(01\)01429-0](https://doi.org/10.1016/S1286-4579(01)01429-0)
- Fang T, Bodmer R, Puertas P, Mayor P, Perez P, Acero R, Hayman D (2008) Certificación de pieles de pecaríes (*Tayassu tajacu* y *T. pecari*) en la Amazonía peruana: Una estrategia para la conservación y manejo de fauna Silvestre en la Amazonia peruana. Wust Editions-Darwin Institute, Lima, Peru. 203 pp.
- Fragoso JMV (1998) Home range and movement patterns of white-lipped peccary (*Tayassu pecari*) herds in the northern Brazilian Amazon. *Biotropica* 30(3):458–469. <https://doi.org/10.1111/j.1744-7429.1998.tb00080.x>
- Fragoso JMV, Levi T, Oliveira LFB, Luzar JB, Overman H, Read JM, Silvius KM (2016) Line transect surveys underdetect terrestrial mammals: Implications for the sustainability of subsistence hunting. *PLoS ONE* 11(4):1–18. <https://doi.org/10.1371/journal.pone.0152659>
- Fragoso JMV, Antunes AP, Silvius KM, Constantino PAL, Zapata-Rios G, Bizri HRE, et al. (2022) Large-scale population disappearances and cycling in the white-lipped peccary, a tropical forest mammal. *PLoS ONE* 17(10):e0276297
- Frank L, Hemson G, Kushnir H, Packer C (2006) Lions, conflict and conservation in Eastern and Southern Africa. In *The eastern and southern African lion conservation workshop* (pp. 11–13). Johannesburg.
- Franzo G, Ruiz A, Grassi L, Sibila M, Drigo M, Segalés J (2020) Lack of porcine circovirus 4 genome detection in pig samples from Italy and Spain. *Pathogens* 9(6):433
- International Union for the Conservation of Nature – IUCN 2023. [www.iucnredlist.org](http://www.iucnredlist.org)
- Karesh W, Uhart M, Painter L, Wallace R, Braselton E, Thomas LE, Namara TM, Gottdenker N (1998) Health evaluation of white-lipped peccary populations. In *Bolivia. 1998 Proceedings AAZV and AAWV Joint Conference*. P, December 2014, 445–449.

- Keuroghlian A, Eaton DP, Longland WS (2004) Area use by white-lipped and collared peccaries (*Tayassu pecari* and *Tayassu tajacu*) in a tropical forest fragment. *Biological Conservation* 120(3):411–425. <https://doi.org/10.1016/j.biocon.2004.03.016>
- Keuroghlian A, Desbiez A, Reyna-Hurtado R, Altrichter M, Beck H, Taber A, Fragoso JMV (2013) *Tayassu pecari*. The IUCN Red List of Threatened Species 2013: e.T41778A44051115.
- Kiltie RA, Terborgh J (1983) Observations on the behavior of rain forest peccaries in Peru: why do white-lipped peccaries form herds? *Z. Tierpsychol.* 62:241–255
- Mayor P, Pérez-Peña P, Bowler M, Puertas PE, Kirkland M, Bodmer R (2015) Effects of selective logging on large mammal populations in a remote indigenous territory in the northern Peruvian Amazon. *Ecology and Society*. <https://doi.org/10.5751/ES-08023-200436>
- Mayor P, El Bizri H, Bodmer RE, Bowler M (2017) Assessment of mammal reproduction for hunting sustainability through community-based sampling of species in the wild. *Conservation Biology* 31(4):912–923
- Mayor P, El Bizri HR, Morcatty TQ, Moya K, Bendayán N, Solis S, et al. (2022) Wild meat trade over the last 45 years in the Peruvian Amazon. *Conservation Biology* 36(2):1–13. <https://doi.org/10.1111/cobi.13801>
- McCallum H (2012) Disease and the dynamics of extinction. *Philosophical Transactions of the Royal Society b: Biological Sciences* 367(1604):2828–2839
- Menajovsky MF, Espunyes J, Cabezón O, Mayor P (2023) Infectious diseases of interest for the conservation of peccaries in the Amazon: a systematic quantitative review. *Biological Conservation* 277:109867
- Menajovsky MF, Espunyes J, Ulloa G, Calderon M, Diestra A, Malaga E, Munoz C, Montero S, Lescano AG, Santolalla ML, Cabezón O, Mayor P (2024) *Toxoplasma gondii* in a remote subsistence hunting-based indigenous community of the Peruvian Amazon. *Trop. Med. Infect. Dis.* 9:98. <https://doi.org/10.3390/tropicalmed9050098>
- Montenegro OL, Roncancio N, Soler-Tovar D, Cortés-Duque J, Contreras-Herrera J, Sabogal S, Acevedo LD, Navas-Suárez PE (2018) Serologic survey for selected viral and bacterial swine pathogens in colombian collared peccaries (*Pecari tajacu*) and feral pigs (*Sus scrofa*). *Journal of Wildlife Diseases* 54(4):700–707. <https://doi.org/10.7589/2017-09-236>
- Morales EA, Mayor P, Bowler M, Aysanoa E, Pérez-Velez ES, Pérez J, et al. (2017) Prevalence of *Trypanosoma cruzi* and other trypanosomatids in frequently-hunted wild mammals from the Peruvian Amazon. *The American Journal of Tropical Medicine and Hygiene* 97(5):1482
- Müller TF, Teuffert J, Zellmer R, Conraths FJ (2001) Experimental infection of European wild boars and domestic pigs with pseudorabies viruses with differing virulence. *Am J Vet Res.* 62(2):252–258
- Nobuto K (1966) Toxoplasmosis in animal and laboratory diagnosis. *Japan Agricultural Research Quarterly* 1(3):11–188
- Oliver-Ferrando S, Segalés J, López-Soria S, Callén A, Merdy O, Joisel F, Sibila M (2016) Evaluation of natural porcine circovirus type 2 (PCV2) subclinical infection and seroconversion dynamics in piglets vaccinated at different ages. *Veterinary Research* 47:1–11
- Pereda AJ, Greiser-Wilke I, Schmitt B, Rincon MA, Mogollon JD, Sabogal ZY, Lora AM, Sanguinetti H, Piccone ME (2005) Phylogenetic analysis of classical swine fever virus (CSFV) field isolates from outbreaks in South and Central America. *Virus Res.* 110:111–118
- Peres CA (2000) Effects of subsistence hunting on vertebrate community structure in Amazonian forests. *Conservation Biology* 14(1):240–253
- Pérez-Peña PE, Gonzáles-Tanchiva C, Trigos-Pinedo M (2016) Evaluación del plan de manejo de animales de caza en la Reserva Nacional Pucacuro. *Folia Amazónica* 25(1):1–16
- Pérez-Peña PE, Ramos-Rodríguez MC, Angulo-Perez N, Caballero-Dulce Y, Del Aguila H, Cachique MS, Montalván R (2021) Sostenibilidad de la caza de mamíferos en tres territorios indígenas de la cuenca alta del Putumayo, Nororiente de la Amazonía peruana. *Ciencia Amazónica (Iquitos)* 9(1):83–96. <https://doi.org/10.22386/ca.v9i1.322>
- Pitman N, Vriesendorp C, Moskovits D (Eds) (2003) Perú: Yavarí. Rapid Biological Inventories, Number 11, Field Museum of Natural History, Chicago, Illinois.
- Porfirio G, Sarmento P, Foster V, Fonseca C (2017) Activity patterns of jaguars and pumas and their relationship to those of their potential prey in the Brazilian Pantanal. *Mammalia* 81(4):401–404
- Postel A, Austermann-Busch S, Petrov A, Moennig V, Becher P (2018) Epidemiology, diagnosis and control of classical swine fever: recent developments and future challenges. *Transboundary and Emerging Diseases* 65:248–261. <https://doi.org/10.1111/tbed.12676>
- Quintana J, Balasch M, Segalés J, Calsamiglia M, Rodríguez-Arriola GM, Plana-Durán J, Domingo M (2002) Experimental inoculation of porcine circoviruses type 1 (PCV1) and type 2 (PCV2) in rabbits and mice. *Veterinary Research* 33(3):229–237
- Richard-Hansen C, Surugue N, Khazraie K, Le Noc M, Grenand P (2014) Long-term fluctuations of white-lipped peccary populations in French Guiana. *Mammalia*. 78:291–301
- Romero Solorio M (2010) Avaliação sanitária da presença de doenças e caracterização dos padrões de caça de subsistência do queixada (*Tayassu pecari*) de vida livre na Amazônia Peruana. Universidade de Sao Paulo.
- Saporiti V, Valls L, Maldonado J, Perez M, Correa-Fiz F, Segalés J, Sibila M (2021) Porcine circovirus 3 detection in aborted fetuses and stillborn piglets from swine reproductive failure cases. *Viruses* 13(2):264
- SERNANP (2014) Plan de manejo de animales de caza en la Reserva Nacional Pucacuro. Realizado por cazadores kichwas de la comunidad 28 de julio, Alfonso Ugarte y Asociación de Cazadores de Intuto. Servicio Nacional de Áreas Naturales Protegidas por el Estado. Lima. 38 pp.
- SERNANP (2019). Servicio Nacional de Áreas Naturales Protegidas por el Estado: Reserva Nacional Pucacuro. Available: <http://www.gob.pe/institucion/sernanp/informes-publicaciones/1749650-reserva-nacional-pucacuro> [accessed April 29, 2024].
- Smith KF, Sax DF, Lafferty KD (2006) Evidence for the role of infectious disease in species extinction and endangerment. *Conservation Biology* 20(5):1349–1357
- Sobral M, Silvius KM, Overman H, Oliveira LFB, Raab TK, Fragoso JMV (2017) Mammal diversity influences the carbon cycle through trophic interactions in the Amazon. *Nature Ecology and Evolution* 1(11):1670–1676. <https://doi.org/10.1038/s41559-017-0334-0>
- Sodhi NS, Brook BW, Bradshaw CJ (2009) Causes and consequences of species extinctions. *The Princeton Guide to Ecology* 1(1):514–520

- Sowls LK (1997) Javelinas and other peccaries: their biology, management, and use. Texas AandM University Press.
- Taber AB, Doncaster CP, Neris NN, Colman FH (1994) Ranging behavior and activity patterns of two sympatric peccaries, *Catagonus wagneri* and *Tayassu tajacu*, in the Paraguayan Chaco. *Mammalia* 58:61–71
- Taber A, Chalukian SC, Altrichter M, Minkowski K, Lizárraga L, Sanderson E, et al., (2008) El Destino de los Arquitectos de los Bosques Neotropicales: evaluación de la distribución y el estado de conservación de los pecaríes labiados y los tapires de tierras bajas. New York: Wildlife Conservation Society, Ecuador, Tapir Specialist Group, Wildlife Trust, and Grupo Especialista de la CSE/UICN. In *Cerdos, Pecaríes Y Hipopótamos*, 1–181
- Tarwater PM, Martin CF (2001) Effects of population density on the spread of disease. *Complexity* 6(6):29–36
- Ter Steege H, Pitman N, Sabatier D, Castellanos H, Van Der Hout P, Daly DC, et al. (2003) A spatial model of tree  $\alpha$ -diversity and tree density for the Amazon. *Biodiversity and Conservation* 12(11):2255–2277. <https://doi.org/10.1023/A:1024593414624>
- Thrusfield M (2018) Presenting numerical data. *Veterinary epidemiology*, 4th ed. Blackwell Science Ltd., Oxford, UK, 251–269.
- Veríssimo, A., Rolla, A., Vedoveto, M., & Futada, S. D. M. (2011). Áreas Protegidas na Amazônia Brasileira: avanços e desafios.
- Vilček Š, Herring AJ, Herring JA, Nettleton PF, Lowings JP, Paton DJ (1994) Pestiviruses isolated from pigs, cattle and sheep can be allocated into at least three genogroups using polymerase chain reaction and restriction endonuclease analysis. *Archives of Virology* 136:309–323
- Weckel M, Giuliano W, Silver S (2006) Jaguar (*Panthera onca*) feeding ecology: distribution of predator and prey through time and space. *Journal of Zoology* 270(1):25–30
- World Organisation for Animal Health – WOAHA 2023. [www.woah.org](http://www.woah.org)
- Zhang L, Zhong C, Wang J, Lu Z, Liu L, Yang W, Lyu Y (2015) Pathogenesis of natural and experimental Pseudorabies virus infections in dogs. *Virology Journal* 12(1):1–10. <https://doi.org/10.1186/s12985-015-0274-8>
- Zuckermann FA (2000) Aujeszky's disease virus: opportunities and challenges. *Veterinary Research* 31(1):121–131. <https://doi.org/10.1051/vetres:2000111>