



Kent Academic Repository

Karpenko, Larisa Yu., Bakhta, Alesya A., Babich, Olga O., Sukhikh, Stanislav A., Vorobyov, N.I., Nikonov, I N, Borisova, Solomonida D., Griffin, Darren K. and Romanov, Michael N. (2025) *Effects of a complex feed additive on productivity and blood parameters of laying hens using stochastic fractal-based neural network model*. Poultry Science Journal, 13 (2). pp. 243-252. ISSN 2345-6604. (In press)

Downloaded from

<https://kar.kent.ac.uk/109922/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.22069/psj.2025.23008.2207>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Effects of a Complex Feed Additive on Productivity and Blood Parameters of Laying Hens Using Stochastic Fractal-based Neural Network Model

Larisa Yu. Karpenko¹, Alesya A. Bakhta¹, Olga O. Babich², Stanislav A. Sukhikh², Nikolay I. Vorobyov³, Ilya N. Nikonov¹, Solomonida D. Borisova¹, Darren K. Griffin^{4,5} & Michael N. Romanov^{4,5,6}

¹ Federal State-Funded Educational Institution of Higher Education “St. Petersburg State University of Veterinary Medicine”, St. Petersburg, 196084, Russia

² Research and Education Center “Industrial Biotechnologies”, Immanuel Kant Baltic Federal University, Kaliningrad, 236016, Russia

³ All-Russian Research Institute for Agricultural Microbiology, Pushkin, St. Petersburg, 196608, Russia

⁴ School of Natural Sciences, University of Kent, Canterbury, CT2 7NJ, UK

⁵ Animal Genomics and Bioresource Research Unit (AGB Research Unit), Faculty of Science, Kasetsart University, Chatuchak, Bangkok 10900, Thailand

⁶ L. K. Ernst Federal Research Center for Animal Husbandry, Dubrovitsy, Podolsk, Moscow Oblast, 142132, Russia

Poultry Science Journal 2025, 13(2): 245-258

Keywords

Brown algae
Blood parameter
Complex feed additive
Adsorbent shungite
Gaussian neural network

Corresponding author

Michael N. Romanov
m.romanov@kent.ac.uk

Article history

Received: November 29, 2024

Revised: April 06, 2025

Accepted: May 12, 2025

Abstract

Neural networks (NNs) benefit biomedicine and agriculture, especially when relying on the specificity and implementation of stochastic fractal-supported models. In the poultry industry, a particular challenge is the search for an ideal sorbent-based complex additive to minimize the loss of valuable feed components that can be tailored to groups of gastrointestinal microorganisms. The aim of this study was thus to develop and apply a mathematical model and Gaussian NN to analyze productivity and blood parameters of laying hens when administering a complex feed additive from the mineral shungite sorbent, plus a nutritive supplement of brown seaweed meal. We developed and built a computational NN that modelled the stochastic ManyToOne relationship of an array of hens' main blood parameters and performance traits. The results presented herein were that the artificial computational stochastic fractal-based NN (EuclidNN) first effectively analyzed the profiles of operational taxonomic units (OTUs) of the physiological/biochemical blood parameters. Also, correlation coefficients were highly positive in relation to certain zootechnical indicators, suggesting that feed additive intake may have led to changes in these performance traits. Calculations suggested that when implementing the feed additive, the values of the Cognitive Salience Index (CSI) vector $vCSI2$ declined. Hereby, this vector correlates with, and affects the egg production trait. Moreover, there was a certain relationship between the feed additive intake and feed and water consumption. Further, EuclidNN computed the respective bioconsolidation indices of hens and, simultaneously, processed several profiles of OTUs for all experimental variants. It also contributed to the calculation of bioconsolidation index values for each variant, i.e., a quantitative assessment of the physiological/biochemical blood descriptors, depending on diet. Collectively, the poultry productivity prediction based on the developed model and NN is pivotal as an initial step for future improvements of economically important traits in chickens when using novel and efficient complex feed additives.

Introduction

For agricultural, and particularly poultry data analysis, neural networks (NNs) present promising and potentially applicable tools for the analysis and synthesis of numerous biological systems (Pitelinskiy and Shimansky, 2013; Jahanmiri and Parker, 2022). One of the major problems of concern is that industrial conditions of livestock exploitation create significant functional stress for the bird's body (Kochish *et al.*, 2019, 2021). Adaptive responses to external stimuli are often stressed, weakening natural defenses, and adversely affecting the health and growth rate of poultry (Young *et al.*, 2022; Kirikovich *et al.*, 2012). One of the ways to alleviate this is through using various feed additives, e.g., mineral-containing substances to stimulate metabolic barriers in the body (Yarov, 2005; Ghahri *et al.*, 2010; Sharapova, 2011). Contemporary activity in the search for ideal sorbents (i.e., insoluble materials or mixtures thereof that are used to recover liquids through absorption and/or adsorption) aims to reduce the negative effects caused by mycotoxins from the feedstuffs. Minimal loss of useful feed components and adaptability to microorganisms living in various parts of the gastrointestinal tract are also key objectives (Ghahri *et al.*, 2010; Okolelova and Mansurov, 2013; Zasorin *et al.*, 2019; Kochish *et al.*, 2020b; Tyurina, 2022). One of the promising sorbent materials is the mineral shungite (Kochish *et al.*, 2020a; Buryakov *et al.*, 2023), which has a specific set of physical properties and structural characteristics; it is composed of non-crystalline carbon with a metastable structure, being incapable of graphitization (Ignatov and Mosin, 2013; Tukhatov, 2013). Shungite rock has good adsorption properties due to its developed and active surface and is capable of acting as a catalyst or catalyst carrier in organic synthesis reactions (Gorodnichev *et al.*, 2019; Vats *et al.*, 2016; Zasorin *et al.*, 2019). Recently, attention has been also drawn to marine brown algae as a nutrient and efficient feed additive. These algae are rich in protein, vitamins and other biologically active substances that play an important role in the body's metabolism (Sharapova, 2011; Buryakov *et al.*, 2023; for review see Kochish *et al.*, 2021).

To model complex stochastic biochemical processes occurring in a bird's body, a computational NN can be employed that includes procedures of multivariate mathematical statistics, i.e., correlation, cluster and discriminant analyses (Meireles *et al.*, 2003; Colbrook *et al.*, 2022; Taye, 2023). One of the instrumental solutions for constructing appropriate mathematical models to describe certain processes, including biologically significant ones, can be stochastic fractal-based mathematical models. These comprise hybrid mixing methods for the mathematical modeling (Moroz and Maslovskaya,

2020; Zaikina *et al.*, 2022) applicable to various (e.g., biological) processes. Previously, we developed a concept of fractal conformity-based bioconsolidation index, which varies within 0 ... 1 depending on the efficiency of coherent biological processes in the body, and tested an artificial NN modelling approach (Kochish *et al.*, 2020b, 2023; Pukhalskiy *et al.*, 2023; Vorobyov *et al.*, 2023a,b), suggesting reasonable prospects of their applications in biological studies. Computational NNs are designed to model a ManyToOne relationship that, for instance, connects a multiple set of heterogeneous bird blood parameters of different dimensions with one single dimensionless numerical indicator for the cognitive significance of the data being studied, i.e., the Cognitive Salience Index ($CSI = 0 \dots 10$; Sutrop, 2001; Mascarenhas, 2018). In one particular case, the dimensionless *CSI* index can be interpreted as an indicator of an increase/decrease in the intensity of biochemical processes in the bird's body that determine the egg performance of laying hens. Furthermore, an important stage in the implementation of an NN in poultry-related research is setting up/training the NN and checking/validating it based on a correlation analysis of the *CSI* values with egg production indicators of layers. After successful NN training and validation, it is feasible to estimate the feed additive intake effects (positive or negative) on the egg performance by increasing/decreasing the *CSI* values.

Considering the remarkable adsorption properties of shungite rock and the nutritional benefits of incorporating brown algae into poultry diets, a combination of these supplements can be most attractive for practical use (Sharapova, 2011; Buryakov *et al.*, 2023). Here, we thus applied an NN and hybrid stochastic fractal-based approach to the mathematical process of analyzing the poultry productivity and blood parameters under the influence of a complex of these two feed supplements. The goal of the present study was therefore to develop the appropriate mathematical model and NN based on this to analyze the performance and blood parameters of laying hens in response to the intake of a complex additive that combined shungite and brown algae, estimating the respective *CSI* and bioconsolidation indices.

Materials and Methods

Experimental birds, performance traits and blood parameters

This study was conducted at a large poultry farm located in Leningrad Oblast, Russia, using Hisex White commercial cross layers (Laptev *et al.*, 2017) at the starting age of 49 weeks. Birds were kept in group cages and fed the basic diet containing grain, root crops, green grass, and a vitamin-mineral feed premix. According to the existing standards, there

were 5 laying hens in each cage. The cage had a standard size, with the dimensions of the total seating area being $490 \times 580 \times 550$ mm. This allowed for a regulated density of hens in the cage, which meets the general requirements that are very strictly observed at this poultry farm. The control (C) group totalled 225,378 hens. The experimental (E) group numbered 218,125 birds fed the basic diet supplemented with the following complex feed additive: 0.7 kg shungite and 0.3 kg brown algae (as dried *Fucus vesiculosus* seaweed meal) per 1-ton fodder. The objective of this feeding experiment was to assess the effect of the combination of these two supplements, i.e. their synergistic impact, considering that the two feed ingredients, different in their essence, are expected to have non-overlapping effects from their use. In particular, the mineral shungite is an inorganic sorbent substance that is not digested in the hen's body, while algae, of an organic (plant) nature, when assimilated in the chicken's body, go through digestive processes in the same way as other plant (and animal) components of the compound feed. The chosen optimal dosages and the ratio of the two supplements were based on our and other preliminary tests (Sharapova, 2011; Buryakov et al., 2023) and taking into account an overall economy of the compound feed recipe that poultry enterprises in Russia adhered to for keeping feed costs reasonable. For the NN analysis, the six most important zootechnical indicators of egg production, feeding and watering of laying hens were also taken as follows: egg production rate (in %), egg weight (in g), proportions of cracked, leaking and dirty eggs (in %), feed conversion (in kg per 10 eggs produced), feed consumption (in g per hen), and water consumption (in mL per hen), plus their four derivative indices (ratios).

We studied the effect of the complex additive on older laying hens in their third feeding phase (i.e., at 75 weeks of age and after), which is considered critical and corresponds to a physiological state of decline in egg production rate and, accordingly, overall productivity. Using 0.5 to 1 mL of blood per sample, three 75-week-old females were sampled from the control and experimental groups (designated C1 and E1, respectively). Five hens per group were then sampled to analyze 12 blood parameters (Table 1) in the middle (86 weeks of age; designated C2 and E2, respectively) and at the end of the feed additive administration study (94 weeks of age; designated C3 and E3, respectively). Accordingly, twelve physiological/biochemical blood characteristics of the laying hens were assessed (Table 1), including the following most informative indices:

- Descriptors of erythrocytes, leukocytes and platelets using Goryaev's grid-equipped counting chamber (Tietz, 1997);

- Blood leukogram using microscopy, May-Grünwald fixative and Romanowsky azure-eosin dye to stain blood smears (Tietz, 1997);
- Serum hemoglobin level using the hemoglobin cyanide method, and the formula-assisted cell-color ratio (Tietz, 1997);
- Phagocytosis indices using a microscopic method and *Staphylococcus aureus* strain 209 culture inactivated by heat and standardized via an optical turbidity standard (Menshikov, 1997).

Data analysis principles using NN

Super-new artificial intelligence (AI) computer programs (D'Addona, 2014; Gharajeh, 2018; Dall'Alba et al., 2022) were used with the respective computational NNs to search, e.g., for feed additives that ensure minimal loss of useful feed components and modulate the gastrointestinal microflora to ensure a complete symbiosis of microorganisms with the animal's body (e.g., Kochish et al. et al., 2020b, 2023; Pukhalskiy et al., 2023). In this regard, the following methods can be considered most relevant (Pitelinskiy and Shimansky, 2013; Taye, 2023):

- Standard statistical methods based on mathematical instrumental regression analysis that optimize specific experimental parameters by applying an appropriate evaluation function.
- Methods based on the use of functional expansions in a Fourier series (Garrapa, 2018). Herein, it is taken into account that the experiment is a commutative process and can be transformed by a sum of functions that describe its polynomials with a given accuracy (e.g., trigonometric functions).
- Partial modelling and formal grammars (Pitelinskiy and Shimansky, 2013).

A sequence of s elements can be considered as an "approximation" in that identical oscillations are often observed in biological systems, differing only in magnitude and time of occurrence, i.e., scale-invariance (Grosu et al., 2023). To describe this, the concept of diversity as a heterogeneous destructive body is employed. For a complete description of diversity, the entire spectrum of dimensions is necessary since, unlike a common mathematical decomposition, it is not enough to introduce only one value, the fractal index d . These dimensions are usually infinite numbers and other time intervals and frequencies of a sample can be further analyzed. Current trends, i.e., specific signal sequences, can then be examined in more detail using, for instance, frequency analysis (Casals et al., 2005; Pitelinskiy and Shimansky, 2013). Conventional experiments are based on inertial analysis and use flexible linear statistical models. However, NNs are inherently nonlinear and do not require knowledge of the correlation between input data and output data. This makes them more promising than traditional methods (Meireles et al., 2003). Although simulations involve

the use of multilayer sensor devices (Rosenblatt, 1962; Minsky and Papert, 1969/1988), the use of closed-loop NNs is not always recommended due to

short-term memory and difficulty of use (Pitelinskiy and Shimansky, 2013; Colbrook *et al.*, 2022).

Table 1: Mean values ($\pm SEM^1$) of 12 blood parameters in hens by four experiment variants (C2, C3, E2, and E3) and correlation coefficients (R) with $vCSI1$ and $vCSI2$ vectors²

Blood parameters	Experiment variants ³				R ($SEM = \pm 0.01$)	
	C2	C3	E2	E3	$vCSI1$	$vCSI2$
Erythrocyte count (mln/ μ L)	2.88 \pm 0.19	3.10 \pm 0.47	2.60 \pm 0.25	3.19 \pm 0.36	0.83 ($P = 0.021$)	-0.78 ($P = 0.040$)
Erythrocyte sedimentation rate (mm/h)	1.20 \pm 0.40	1.20 \pm 0.40	1.20 \pm 0.40	1.60 \pm 0.49	0.67 ($P = 0.098$)	-0.62 ($P = 0.137$)
Leukocyte count (thousand/ μ L)	30.26 \pm 4.13	35.85 \pm 3.93	31.22 \pm 2.78	30.34 \pm 8.74	0.44 ($P = 0.324$)	-0.39 ($P = 0.391$)
Platelet count (thousand/ μ L)	71.84 \pm 11.34	65.71 \pm 10.84	54.62 \pm 16.31	62.67 \pm 22.58	-0.07 ($P = 0.889$)	0.12 ($P = 0.801$)
Eosinophils	2.20 \pm 1.10	1.20 \pm 0.40	1.80 \pm 0.75	0.40 \pm 0.49	-0.90 ($P = 0.006$)	0.95 ($P = 0.001$)
Pseudoeosinophils	6.60 \pm 1.02	14.60 \pm 4.92	12.00 \pm 1.67*	18.80 \pm 2.93	0.93 ($P = 0.003$)	-0.88 ($P = 0.010$)
Monocytes	0.40 \pm 0.49	1.00 \pm 0.00	0.80 \pm 0.75	0.80 \pm 0.40	0.74 ($P = 0.056$)	-0.69 ($P = 0.086$)
Basophils	1.00 \pm 0.89	0.80 \pm 0.40	1.20 \pm 0.40	0.80 \pm 0.75	-0.78 ($P = 0.041$)	0.83 ($P = 0.022$)
Lymphocytes	89.60 \pm 2.33	82.40 \pm 4.80	84.20 \pm 2.64	79.20 \pm 2.79	-0.86 ($P = 0.014$)	0.91 ($P = 0.005$)
Hemoglobin (g/L)	80.00 \pm 5.48	89.20 \pm 8.70	85.60 \pm 4.18	83.00 \pm 8.85	0.50 ($P = 0.249$)	-0.45 ($P = 0.308$)
Cell-color ratio	1.68 \pm 0.21	1.79 \pm 0.44	1.99 \pm 0.11	1.59 \pm 0.28	-0.38 ($P = 0.395$)	0.44 ($P = 0.328$)
Phagocytic index (a.u.) ⁴	10.62 \pm 0.87	9.57 \pm 1.31	9.93 \pm 1.45	11.01 \pm 0.92	0.08 ($P = 0.867$)	-0.03 ($P = 0.955$)

¹ SEM , standard error of the mean. ² $vCSI1$ and $vCSI2$, vectors containing the values of the $CSI1$ and $CSI2$ cognitive salience indices for the experiment variants. ³ Experiment variants corresponded to the second and third blood sampling points (at 86 and 94 weeks of age) for the control (C2 and C3) and experimental (E2 and E3) groups of laying hens, respectively. ⁴ a.u., arbitrary units. * Significant difference between the respective C2 and E2 values (at $P < 0.05$).

General NN and stochastic fractal-based model description

The mathematical research was carried out using NN analysis and stochastic fractal-based modelling (Garrapa, 2018; Moroz and Maslovskaya, 2020; Zaikina *et al.*, 2022). The NN and its elements we report here are subject to protecting the intellectual property of the authors (Vorobyov *et al.*, 2023b). NN data mining implies a machine learning algorithm and data analysis technique that mimics the functionality of the human brain (Khlivnenko, 2015; Goodfellow *et al.*, 2016). This approach involves a large number of interconnected nodes capable of processing and transmitting information. Because of this, NNs can be typically used to process large amounts of information and identify patterns among the analyzed datasets (D'Addona, 2014), the latter being, in our case, the physiological/biochemical blood parameters of laying hens.

When examining blood parameters as operational taxonomic units (OTUs), it was important to determine how the statistics of these datasets differ from the conventional Gaussian statistics (i.e., normal distribution) (Wentzel, 1999). It was assumed

that any deviation of the actual data statistics from Gaussian statistics would be indicative of the regulation of physiological/biochemical processes in the blood, e.g., changes in the levels of erythrocytes, leukocytes, platelets, etc., that are controlled by the bird's body. Deviations from the normal distribution were assessed by the magnitude of the highest central moments in the empirical distributions of the measured values (Rosenblatt, 1962; Popov, 2013).

The physiological/biochemical characteristics of the hen's blood can be characterized by Shannon entropy indices ($IndShen$) also considered biodiversity indices (Grishanov and Grishanova, 2010; Chernov *et al.*, 2015; Gorodnichev *et al.*, 2019) and by bioconsolidation indices ($IndBconI$) of the bird's body and its immune system status (Kochish *et al.*, 2020b, 2022, 2023). With increasing entropy (increasing $IndShen$ index), the physiological/biochemical characteristics of blood across replicates are levelled out, and their dispersion decreases. Therefore, it can be assumed that the positive correlation of the $IndShen$ index with the $IndBconI$ index means that the corresponding characteristic of hen's blood may be a signal

indicator of the bioconsolidation of the immune status.

To calculate the distribution moments for the empirical blood parameter data used as OTUs and compute the respective bioconsolidation indices of the hen's body and its immune status, an artificial Gaussian NN (GNN) was implemented (Minsky and

Papert, 1969/1988; Filntisi *et al.*, 2013; Sergeev and Tarasov, 2017; Mascarenhas, 2018; Gafarov and Galimyanov, 2018) (Figure 1). This was based on a stochastic fractal-based model (Moroz and Maslovskaya, 2020; Zaikina *et al.*, 2022) for defining the flow of physiological/biochemical processes in the bird's body.




Figure 1. Artificial Gaussian neural network for calculating bioconsolidation indices of the bird's body and its immune system status. S, matrix of physiological/biochemical blood parameters; D, matrix of statistical moments of blood parameters; and C, vector of bioconsolidation indices.

Results

Constructing NN and stochastic fractal-based model

In the course of developing the artificial GNN further called *EuclidNN* (Figure 2), we built a computational NN that modelled the stochastic ManyToOne relationship of an array of hens' main blood parameters and performance traits (used as OTUs)

with one unique dimensionless characteristic, *CSI* (Sutrop, 2001; Mascarenhas, 2018). Values of the *CSI1* and *CSI2* indices for the four experiment variants were transformed into *vCSI1* and *vCSI2* vectors (Tables 1 and 2).

The most informative poultry performance indicators and correlation coefficients with vectors $vCSII$ and $vCSI2$ are summarized in Table 2.




Figure 2. Chart flow of the EuclidNN computational neural network. *mBlood* is a matrix of 12 key physiological/biochemical blood indicators (Table 1). *mEggs* is a matrix of eight physiological performance-related indicators (Table 2). *mNorm* is a matrix of normalized values of the *mBlood* matrix. *mDist* is a matrix of Euclidean distances of four experimental variants based on blood parameter data. *mCompo* is a matrix of orthogonal components in the space of blood parameter data. *mComby* is a matrix containing the results of a nonlinear combinatorial transformation of data from matrices *mDist* and *mCompo*. *vCS11* and *vCS12* are vectors containing the values of the *CS11* and *CS12* cognitive salience indices for the experiment variants. L1, L2 and L3 are layers of artificial neurons that perform matrix transformations of the initial and intermediate data.

Table 2. Mean values of eight key zootechnical indicators (i.e., hen productivity, feeding and watering) by four experiment variants (C2, C3, E2 and E3) and correlation coefficients (*R*) with vectors *vCSI1* and *vCSI2*

Indicators	Experiment variant ¹				<i>R</i> (<i>SEM</i> = ±0.01)	
	C2	C3	E2	E3	<i>vCSI1</i>	<i>vCSI2</i>
Egg production rate, %	88.7	77.9	85.4	71.8	-0.92 (<i>P</i> = 0.003)	0.98 (<i>P</i> < 0.001)
Egg weight, g	64.3	66.5	64.8	65.5	0.85 (<i>P</i> = 0.016)	-0.80 (<i>P</i> = 0.032)
Cracked eggs, %	4.0	8.8	8.8	12.4	0.83 (<i>P</i> = 0.020)	-0.78 (<i>P</i> = 0.039)
Leaking eggs, %	0.21	0.41	0.33	0.54	0.94 (<i>P</i> = 0.002)	-0.89 (<i>P</i> = 0.008)
Dirty eggs, %	2.9	9.7	4.1	12.6	0.99 (<i>P</i> < 0.001)	-0.94 (<i>P</i> = 0.002)
Feed conversion, kg/10 eggs	1.46	1.64	1.63	1.72	0.84 (<i>P</i> = 0.019)	-0.78 (<i>P</i> = 0.037)
Feed consumption, g/hen	130	128	139	124	-0.66 (<i>P</i> = 0.109)	0.71 (<i>P</i> = 0.074)
Water consumption, mL/hen	250	248	250	228	-0.68 (<i>P</i> = 0.092)	0.73 (<i>P</i> = 0.061)
<i>vCSI1</i> vector (<i>SEM</i> = ±0.2)	2.5	6.7	3.4	7.4	1.00 (<i>P</i> < 0.001)	-0.95 (<i>P</i> = 0.001)
<i>vCSI2</i> vector (<i>SEM</i> = ±0.2)	7.5	3.3	6.6	2.6	-0.95 (<i>P</i> = 0.001)	1.00 (<i>P</i> < 0.001)

¹ Experiment variants corresponded to the second and third blood sampling points (at 86 and 94 weeks of age) for the control (C2 and C3) and experimental (E2 and E3) groups of laying hens. *SEM*, standard error of the mean.

The constructed multilayer EuclidNN included three artificial neuron layers L1, L2 and L3 that executed the initial and intermediate data matrix transformations (Figure 2) as outlined below.

L1 neuron layer

This performed normalization of the *mBlood* matrix data (Figure 2) (Everitt *et al.*, 2011; Mascarenhas, 2018) that contained digital data on the blood parameters of birds (OTUs) in the four experiment variants (Table 1). The results of normalization were stored in the *mNorm* matrix according to the following Equation 1:

$$mNorm_{jk} = \frac{mBlood_{jk} - \frac{1}{4} \cdot \sum_{k=1}^4 mBlood_{jk}}{\sqrt{\sum_{k=1}^4 \left[mBlood_{jk} - \frac{1}{4} \cdot \sum_{k=1}^4 mBlood_{jk} \right]^2}} \quad (1)$$

where *mBlood_{jk}* are *mBlood* matrix values; *j* = 1, 2, ..., 12 are ordinal numbers of 12 blood parameters (Table 1); and *k* = 1, 2, ..., 4 are ordinal numbers of the four experiment variants that conformed to C2, C3, E2 and E3, respectively (Tables 1 and 2).

The L1 neuron layer also computed the *mDist* matrix, i.e., the matrix of Euclidean distances between the four experiment variants in the 12-dimensional space of birds' blood parameters (Yeung and Ruzzo, 2001; Everitt *et al.*, 2011) using the following Equation 2:

$$mDist_{mn} = \sqrt{\frac{1}{12} \sum_{j=1}^{j=12} (mNorm_{jn} - mNorm_{jm})^2}, \quad (2)$$

where *m*, *n* = 1, 2, ..., 4 are ordinal numbers of the four experiment variants that conformed to C2, C3, E2 and E3, respectively (Tables 1 and 2); and *j* = 1, 2, ..., 12 are ordinal numbers of 12 blood parameters (Table 1).

L2 neuron layer

This computed the *mCompo* matrix, i.e., the matrix of principal components (orthogonal eigenvectors) in the 12-dimensional space of birds' blood indices using the *mDist* matrix data and a standard computational procedure (Yeung and Ruzzo, 2001; Jolliffe, 2002; Schmidhuber, 2015) as follows:

$$mCompo = EigenVectors(mDist). \quad (3)$$

L3 neuron layer. This computed the *mComby* matrix containing numerical data that were obtained by different variants of the nonlinear combination of the *mDist* and *mCompo* matrix data. The *mComby* matrix data were intended for training tuning of the EuclidNN model and for calculating the desired *vCSI1* and *vCSI2* index vectors using hens' blood parameters.

When training the EuclidNN neural network (Nikolić *et al.*, 2012; Schmidhuber 2015; Widrow *et al.*, 2013; Nikolenko *et al.*, 2018), the second orthogonal component (*mCompo2*, i.e., the second row of the *mCompo* matrix) in the 12-dimensional space of blood indices was chosen because the projection of blood indices onto the *mCompo2* component maximally correlated with performance indices (Table 2) (Gao *et al.*, 2012; Batushansky *et al.*, 2016). Using the values of the *mCompo2* component, the values of the vectors *vCSI1* and

$vCSI2$ (Tables 1 and 2) were calculated using the following Equations 4 and 5:

$$vCSI1_k = mCompo2_k \cdot 4.17 + 5, \quad (4)$$

$$vCSI2_k = -mCompo2_k \cdot 4.17 + 5, \quad (5)$$

where $k = 1, 2, \dots, 4$ are ordinal numbers of the four experiment variants that conformed to C2, C3, E2 and E3, respectively (Tables 1 and 2).

In addition, the correlation coefficients (R) of $vCSII$ and $vCSI2$ vectors with blood parameters and performance indicators were calculated as summarized in Tables 1 and 2.

To compute the statistical moments of blood indices in the layers of neurons M2, M3, M4, ..., Mn, the moments ($d_{l,j}$) forming the matrix of moments D (Figure 1) were calculated using the following Equations 6 and 7:

$$d_{l,j} = \left[\frac{1}{N-1} \cdot \sum_{k=1}^{k=N} Abs(s_{l,k} - m_l)^j \right]^{1/j} / m_l \quad (6)$$

$$m_l = \frac{1}{N} \sum_{k=1}^{k=N} s_{l,k} \quad (7)$$

where m_l ($s_{l,k}$) are physiological/biochemical blood parameters; $k = 1, 2, \dots, N$ are numbers of replicates of blood parameters in samples; $j = 2, 3, \dots, N-2$ are numbers of statistical moments of blood parameters; N is the number of blood parameter values in a sample; and l is the serial number of the four experiment variants.

Subsequently, $IndBconI$ (c_l) indices were calculated in the End neuron layer (Figure 1) using the following Equation 8:

$$c_l = \frac{1}{1 + \exp(-a \cdot Abs(p_l))} \quad (8)$$

where the respective additional indices were computed as follows:

$$p_l = \sum_{j=1}^{j=N-2} \left(\frac{d_{j,l} \cdot g_{j,l}}{IndShen_l} \right), \quad g_j = \frac{1}{r_D} \cdot \left(\sum_{l=1}^{l=M} d_{l,j} - r_C \right),$$

$$r_C = \frac{1}{N} \sum_{j=2}^{j=N-2} \left(\sum_{l=1}^{l=M} d_{l,j} \right), \quad r_D = \sqrt{\sum_{j=2}^{j=N-2} \left(\sum_{l=1}^{l=M} d_{l,j} - r_C \right)^2},$$

$$IndShen_l = \frac{1}{\log(N)} \sum_{k=1}^{k=N} \left[\frac{s_{l,k}}{m_l} \cdot \log \left(\frac{s_{l,k}}{m_l} \right) \right],$$

$$m_l = \frac{1}{N} \sum_{k=1}^{k=N} s_{l,k};$$

$j = 2, 3, \dots, N-2$ are numbers of statistical moments of physiological/biochemical blood parameters; $k = 1, 2, \dots, N$ are numbers of blood parameter values in a sample; l is serial number of the four experiment variants; and a is a constant.

EuclidNN computational algorithm implementation to analyze experimental data

The input data for calculations using EuclidNN (Figures 1 and 2) (Rosenblatt, 1962; Meireles *et al.*, 2003; Anastasiadis, 2005; Filntisi *et al.*, 2013; Mascarenhas, 2018; Colbrook *et al.*, 2022; Taye, 2023) were generated in the Excel environment and are presented in Tables 1 and 2).

At the first stage, the appropriate immune bioconsolidation indices were calculated from the most informative physiological/biochemical blood parameter data using the GNN artificial intelligence process and are summarized in Table 3.

Then, the correlation coefficients between the $IndBconI$ indices and performance parameters of laying hens were computed as shown in Table 4. As can be seen from Table 4, there were high correlation coefficients (positive or negative) between certain performance indicators and blood parameter bioconsolidation indices. For example, the egg mass yield indicators, i.e., the product of mean egg weight and egg production, positively correlated with the $IndBconI$ values for hemoglobin ($R = 0.998, 0.838$ and 0.916) and platelet counts ($R = 0.878, 0.777$ and 0.993).

Finally, the values of the vectors $vCSII$ and $vCSI2$ were computed using EuclidNN (Tables 1 and 5). In particular, when administering the shungite–seaweed feed additive, the $vCSII$ vector values raised as follows: $vCSII(C2) = 2.5 < vCSII(E2) = 3.4$; and $vCSII(C3) = 6.7 < vCSII(E3) = 7.4$ (Table 5). The correlation coefficients (Table 5) for the $vCSII$ vector were highly positive in relation to certain zootechnical indicators ($R = 0.83$ to 0.99), suggesting that the feed additive intake may have led to changes in these performance indicators.

Table 3: Bioconsolidation indices ($IndBconI$) for five main blood parameters according to the four experiment variants¹

Blood indicators ²	C2	C3	E2	E3
Erythrocyte count	0.65	0.77	0.52	0.53
Leukocyte count	0.37	0.30	0.41	0.48
Platelet count	0.27	0.47	0.35	0.47
Hemoglobin	0.53	0.62	0.56	0.71
Cell-color ratio	0.81	0.92	0.50	0.67

¹ Experiment variants corresponded to the second and third blood sampling points (at 86 and 94 weeks of age) for the control (C2 and C3) and experimental (E2 and E3) groups of laying hens. $SEM = \pm 0.02$, standard error of the mean.

Table 4. Correlation coefficients (*R*) between blood parameter bioconsolidation indices (*IndBconI*) and performance indicators

Performance indicators	<i>IndBconI</i>				
	Erythrocyte count	Leukocyte count	Platelet count	Hemoglobin	Cell-color ratio
Egg production rate, %	0.77 (<i>P</i> = 0.026)	−0.39 (<i>P</i> = 0.337)	0.47 (<i>P</i> = 0.235)	0.40 (<i>P</i> = 0.323)	0.93 (<i>P</i> < 0.001)
Mean egg weight, g	0.87 (<i>P</i> = 0.005)	−0.54 (<i>P</i> = 0.167)	0.35 (<i>P</i> = 0.391)	0.23 (<i>P</i> = 0.588)	0.98 (<i>P</i> < 0.001)
Mean egg mass yield ¹	0.05 (<i>P</i> = 0.898)	0.38 (<i>P</i> = 0.355)	0.77 (<i>P</i> = 0.025)	0.93 (<i>P</i> < 0.001)	0.32 (<i>P</i> = 0.435)
Mean egg mass yield to feed consumption ratio	0.72 (<i>P</i> = 0.045)	−0.36 (<i>P</i> = 0.375)	0.73 (<i>P</i> = 0.037)	0.58 (<i>P</i> = 0.134)	0.81 (<i>P</i> = 0.014)
Mean egg mass yield to water consumption ratio	−0.61 (<i>P</i> = 0.111)	0.38 (<i>P</i> = 0.360)	−0.77 (<i>P</i> = 0.026)	−0.59 (<i>P</i> = 0.125)	−0.68 (<i>P</i> = 0.065)
Total nonmarket eggs, %	−0.18 (<i>P</i> = 0.673)	0.17 (<i>P</i> = 0.694)	−0.86 (<i>P</i> = 0.006)	−0.58 (<i>P</i> = 0.129)	−0.06 (<i>P</i> = 0.892)
Feed conversion, kg per 10 eggs	−0.79 (<i>P</i> = 0.019)	0.75 (<i>P</i> = 0.033)	−0.57 (<i>P</i> = 0.137)	−0.19 (<i>P</i> = 0.660)	−0.67 (<i>P</i> = 0.069)
Feed conversion, kg per 1 kg egg mass yield	−0.68 (<i>P</i> = 0.065)	0.48 (<i>P</i> = 0.228)	−0.73 (<i>P</i> = 0.038)	−0.50 (<i>P</i> = 0.206)	−0.71 (<i>P</i> = 0.050)
Feed consumption, g per hen	−0.79 (<i>P</i> = 0.020)	0.55 (<i>P</i> = 0.159)	0.30 (<i>P</i> = 0.476)	−0.20 (<i>P</i> = 0.637)	−0.92 (<i>P</i> = 0.001)
Water consumption, mL per hen	−0.47 (<i>P</i> = 0.238)	0.18 (<i>P</i> = 0.675)	−0.76 (<i>P</i> = 0.028)	−0.70 (<i>P</i> = 0.053)	−0.63 (<i>P</i> = 0.094)

¹ The product of mean egg weight by egg production characterizes the egg mass yield as a whole. *SEM* = −0.08 ... 0, standard error of the mean.

Table 5. Mean values of vectors *vCSI1* and *vCSI2* by four experiment variants (C2, C3, E2 and E3) and correlation coefficients (*R*) with eight key zootechnical indicators (i.e., hen productivity, feeding and watering)

Vectors	Experiment variant ¹				<i>R</i> (<i>SEM</i> = ±0.01)	
	C2	C3	E2	E3	<i>vCSI1</i>	<i>vCSI2</i>
<i>vCSI1</i> (<i>SEM</i> = ±0.2)	2.5	6.7	3.4	7.4	1	−1
<i>vCSI2</i> (<i>SEM</i> = ±0.2)	7.5	3.3	6.6	2.6	−1	1

Vectors	<i>R</i> values (<i>SEM</i> = ±0.01) for indicators							
	Egg production	Egg weight	Cracked eggs	Leaking eggs	Dirty eggs	Feed conversion	Feed consumption	Water consumption
<i>vCSI1</i>	−	0.84	0.83	0.94	0.99	0.83	−	−
<i>vCSI2</i>	0.98	−	−	−	−	−	0.69	0.74

¹ Experiment variants corresponded to the second and third blood sampling points (at 86 and 94 weeks of age) for the control (C2 and C3) and experimental (E2 and E3) groups of laying hens. *SEM*, standard error of the mean.

Calculations using EuclidNN showed that when the feed additive was implemented, the values of the *vCSI2* vector declined (Table 5): *vCSI2*(C2) = 7.5 > *vCSI2*(E2) = 6.6; and *vCSI2*(C3) = 3.3 > *vCSI2*(E3) = 2.6. Hereby, this vector correlates with, and affects the egg production trait. Also, there was a certain relationship between the feed additive intake and feed and water consumption because the correlation coefficients (*R*) of the *vCSI2* vector in relation to these two indicators were positive, though relatively not very high: 0.69 and 0.74, respectively (Table 5).

Discussion

Humankind directly interacts with, and has profound effects on, the multifaceted information environment. Consequently, the relevance of using NNs in these circumstances is increasing. NNs have been already

integrated into the lives of the contemporary population, thereby assisting us in solving a large number of problems, especially in the biomedicine area. Modern industrial poultry farming operates with a myriad of objective physiological and genetic data (big data) used to formulate an optimal diet and feeding regime for birds, as well as for the justified application of protective and modulating feed additives. Big data analysis (Gharajeh, 2018; Dall'Alba *et al.*, 2022) allows us to explore the influence of external factors and feed on the health condition and egg production of layers. Based on this analysis outcome, it is possible to develop an optimal strategy for feeding birds in an industrial environment and achieve the highest performance in commercial poultry production. Using the NN approach in this study, we evaluated the

increase/decrease in the intensity of physiological/biochemical processes in the body of an agricultural species, specifically in relation to the performance of laying hens by virtue of increasing/decreasing *CSI* index values.

The developed computational EuclidNN, through internal calculations, was instrumental in defining the nature (positive or negative) of the impact that the shungite rock–brown algae feed additive has on the physiological/biochemical processes occurring in the bird's body and, correspondingly, on the performance indicators of laying hens by utilizing the basic blood indices. The latter integrally reflects the hematopoiesis processes resulting in the formation of blood cells such as erythrocytes, leukocytes and platelets (Tietz, 1997). Their production and growth depend on nutritional and vitamin status (Menshikov, 1997; Levchenko *et al.*, 2020; Matveev and Torshkov, 2020; Milevski, 2024; IAKLRF, 2023). The conversion of feed substrates into nutrients, as well as other mediated physiological/biochemical processes in the hens' body, may have occurred more efficiently when administering the feed additive. For the same reason, the water consumption for keeping birds could also reduce; however, a relatively low correlation of the *CSI2* index with feed and water consumption ($R = 0.69$ and 0.74 , respectively) was probably a consequence of additional accidental losses while keeping the flock of egg layers.

Considering the blood parameters of birds (Tables 1 and 2), we found that they correlated to varying degrees with the *vCSI1* and *vCSI2* vectors. We conclude that the high positive correlation of blood parameters (erythrocyte count and sedimentation rate, pseudoeosinophils, monocytes, and phagocytic index) with the *vCSI1* vector implied that, when using the feed additive, the values of these indicators increased. If other blood parameters (eosinophils, basophils and lymphocytes) correlated with the *vCSI2* vector, this meant that these parameters decreased in response to the feed additive intake. At the same time, it was found that the number of platelets in the blood did not correlate with the *vCSI2* vector (Table 1), suggesting that this blood indicator was not involved in the formation of *vCSI2* vector values and, therefore, may not be measured when diagnosing the health status of birds. In terms of the zootechnical characteristics (Tables 4 and 5), one can note that the feed additive administration most likely caused such a change in physiological/biochemical processes in the bird's body that increased the egg weight while reducing the shell strength. The latter could raise the likelihood of accidental eggshell damage (i.e., cracked, leaking and dirty eggs) during transportation. Consequently, the hens' body responded to the feed additive intake by restructuring internal physiological/biochemical processes,

accompanied by an increase/decrease in the birds' blood composition and the performance indicators of laying hens. On the other hand, the NN-based mathematical model was sufficiently sensitive to reveal implicit issues that might affect egg performance and lead to an increased percentage of nonmarket eggs (Table 5).

The use of NN, stochastic fractal-based and similar models has been also assessed in other studies when implementing feed additives for raising egg layers and other farm animals (Nematinia and Abdanan Mehdizadeh, 2018; de Almeida *et al.*, 2020; Ojo *et al.*, 2022; Yang *et al.*, 2023; Buryakov *et al.*, 2023; Siriani *et al.*, 2023). In particular, Buryakov *et al.* (2023) determined the bioconsolidation indices of microorganisms in the intestines of laying hens and estimated the effect of a similar complex feed additive (shungite and *Fucus vesiculosus* seaweed meal) on the self-organization of the microbial-organismal biosystem in the intestines of birds. Calculation of bioconsolidation indices in this and other studies (Kochish *et al.*, 2020b; Buryakov *et al.*, 2023) showed that shungite, in combination with brown algae, can be a promising feed additive for a beneficial effect on the body of layers.

To reduce health risks and financial concerns, decisions regarding poultry production and health status should be made based on objective criteria (de Almeida *et al.*, 2020). Consistent with our study results, de Almeida *et al.* (2020) demonstrated that the use of artificial NNs is a valuable tool to reduce the subjectivity of analysis for predicting and managing poultry flocks and egg production.

The defective eggs we focused upon in our experiments reduce the value of laying hen egg production. Mathematical modelling, as our and other studies (Yang *et al.*, 2023) have shown, can be used to solve problems of improving egg quality and productivity. To this end, Yang *et al.* (2023) developed a convolutional NN-based model to control egg category and weight. To predict egg productivity and freshness, Nematinia and Abdanan Mehdizadeh (2018) used an artificial NN that was trained with the Levenberg-Marquardt algorithm. By implementing research findings like these, the poultry industry can reduce costs and improve productivity.

Ojo *et al.* (2022) stated that the advent of digital technologies has led to significant improvements in various areas. Modern NNs have great potential to intelligently automate current and future poultry management operations to ensure high-quality, low-cost poultry production and manage bird welfare. Siriani *et al.* (2023) tested an NN with a stochastic fractal-based model algorithm to classify the mobility and resting phases of chickens during the rearing process. The stochastic fractal-based model we have developed also allows, by assessing the zootechnical

and physiological/biochemical parameters of chicken rearing, to predict their performance and, in this perspective, resistance to diseases.

In general, many models have emerged to date in the field of computer-aided research development that is accompanied by methods for identifying fractals (Meakin, 1999; Dvoryatkina *et al.*, 2017; Kochish *et al.*, 2023; Yurkovych *et al.*, 2023). Analysis and synthesis of biological systems (Chakraverty *et al.*, 2023) is a critical element in the task of rising the analysis efficiency of experiments, especially when the sequence of successive elements is considered to be disrupted. Describing complex biological systems requires the use of multidimensional concepts and additional analysis of other time intervals and frequencies of the sample being studied (Pitelinskiy and Shimansky, 2013). Traditional experiments are based on inertial analysis and use flexible linear statistical models (Pyrhönen *et al.*, 2024). However, NNs are inherently nonlinear (Ge & Wang, 2002) and do not require knowledge of the correlation between input data and output data (Hsieh, 2000), making them more promising than traditional approaches (Pitelinskiy and Shimansky, 2013). In the current study, multilayer sensitizers (perceptrons; Przybyła-Kasperek and Marfo, 2024) were used to simulate poultry performance and analyze the tolerance of exposure to the complex feed additive based on shungite and algae. However, one should take into account the possible limitations of using closed-loop NNs (Zhu *et al.*, 2021), including short-term memory and other potential difficulties.

Collectively, EuclidNN, an NN model we developed using the mathematical stochastic fractal-based method, facilitated studying the effects of the complex feed additive on the immune state of the birds' body, on the intensity of internal physiological/biochemical processes (through their blood parameters) and, ultimately, the performance of layers. As a result, changes in *CSI* index values can be used to estimate changes in the intensity of physiological/biochemical processes in birds' bodies and laying hen productivity.

Conclusion

In the current research, a mathematical stochastic fractal-based model and GNN using this model were developed for the first time, which can be implemented to analyze poultry performance indicators under the influence of the complex feed additive from the mineral shungite and brown algae. Such mycotoxin adsorbents as shungite can have a complex effect on the body of farm animals. Its multifactorial action is associated, first of all, with the removal of negative effects caused by mycotoxins. This study exemplified an NN analysis-assisted evaluation of the effects of the shungite in

combination with brown algae on the performance and blood parameters of laying hens in an industrial poultry farm.

We suggest that high bioconsolidation values correspond to increased efficiency of physiological/biochemical processes in the bird's body in response to the feed additive intake. In future, methods for assessing fractal characteristics will be increasingly used for analyzing random processes, developing mathematical models and conducting simulations in biomedicine and in a wide variety of science and technology fields (Pitelinskiy and Tyurkin, 2007). The productivity prediction approach using the obtained mathematical model and GNN will be useful in the future improvements of the productive qualities and resilience in poultry.

Traditional poultry studies do not address the one-way ManyToOne relationship of blood parameters with the quantitative and qualitative egg production characteristics of layers, since the qualitative traits are dimensionless and, in a quantitative representation, cannot participate in correlation analysis with the measured characteristics of egg performance. To overcome this obstacle, we created the computational EuclidNN that allowed customizing its calculation procedures to determine the *CSI* values. A distinctive feature of the *CSI* index is that, on the one hand, it characterizes the entire complex of blood parameters of laying hens and, on the other, represents a qualitative or quantitative egg production indicator. Thus, EuclidNN makes it possible to evaluate the complex feed additive influence not only on the quantitative indicators but also on the qualitative egg performance indicators of chickens.

Funding: The study was supported by the Russian Science Foundation (Grant No. 23-16-00181, <https://rscf.ru/project/23-16-00181/>).

Author Contributions: Conceptualization, N.I.V. and L.Y.K.; methodology, N.I.V. and A.A.B.; validation, N.I.V. and S.D.B.; data curation, O.O.B. and I.N.N.; writing—original draft preparation, N.I.V., O.O.B. and I.N.N.; writing—review and editing, S.A.S., N.I.V., M.N.R. and D.K.G.; visualization, N.I.V.; supervision, M.N.R. and D.K.G.; project administration, I.N.N.; funding acquisition, L.Y.K. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board of the Federal State-Funded Educational Institution of Higher Education “St. Petersburg State University of Veterinary Medicine” (Protocol No. 3, dated 27.06.2022).

Informed Consent Statement: Not applicable.

Data Availability Statement: Data is contained within the article.

References

- Anastasiadis AD. 2005. Neural networks training and applications using biological data. PhD Thesis. School of Computer Science and Information Systems, University of London, London, UK.
- Batushansky A, Toubiana D & Fait A. 2016. Correlation-based network generation, visualization, and analysis as a powerful tool in biological studies: A case study in cancer cell metabolism. *BioMed Research International*, 2016: 8313272. DOI: 10.1155/2016/8313272
- Buryakov NP, Zaikina AS, Trukhachev VI, Buryakova MA, Kosolapova VG, Nikonov IN, Medvedev IK, Fathala MM & Aleshin DE. 2023. Influence of dietary addition of mineral shungite and *Fucus vesiculosus* on production performance, egg quality, nutrients digestibility, and immunity status of laying hens. *Animals*, 13: 3176. DOI: 10.3390/ani13203176
- Casals J, Jerez M & Sotoca S. 2005. Modeling and Forecasting Time Series Sampled at Different Frequencies. Office for Official Publications of the European Communities. Luxembourg. ISBN 92-79-01327-0.
- Chakraverty S, Jena RM & Jena SK. 2023. Time-fractional Order Biological Systems with Uncertain Parameters. *Synthesis Lectures on Mathematics and Statistics*. Springer Nature Switzerland AG. Cham, Switzerland. ISBN 978-3-031-01295-2. DOI: 10.1007/978-3-031-02423-8
- Chernov TI, Tkachakhova AK & Kutovaya OV. 2015. Assessment of diversity indices for the characterization of the soil prokaryotic community by metagenomic analysis. *Eurasian Soil Science*, 48: 410–415. DOI: 10.1134/S1064229315040031
- Colbrook M, Antun J, Vegard H & Anders C. 2022. The difficulty of computing stable and accurate neural networks: On the barriers of deep learning and Smale's 18th problem. *Proceedings of the National Academy of Sciences of the United States of America*, 119: e2107151119. DOI: 10.1073/pnas.2107151119
- D'Addona DM. 2014. Neural network. In: Laperrière L & Reinhart G (Eds.). *CIRP Encyclopedia of Production Engineering*. Springer. Berlin, Heidelberg, Germany. DOI: 10.1007/978-3-642-20617-7_6563
- Dall'Alba G, Casa PL, Abreu FPD, Notari DL & de Avila e Silva S. 2022. A survey of biological data in a big data perspective. *Big Data*, 10: 279–297. DOI: 10.1089/big.2020.0383

Acknowledgments: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

- de Almeida LGB, de Oliveira ÉB, Furian TQ, Borges KA, Tonini da Rocha D, Salle CTP & Moraes HL de S. 2020. Artificial neural networks on eggs production data management. *Acta Scientiae Veterinariae*, 48. DOI: 10.22456/1679-9216.101462
- Dvoryatkina S, Smirnov E & Lopukhin A. 2017. New opportunities of computer assessment of knowledge based on fractal modeling. *US-China Foreign Language*, 15: 452–459. DOI: 10.17265/1539-8080/2017.07.005
- Everitt BS, Landau S, Leese M & Stahl D. 2011. *Cluster Analysis*. 5th ed. John Wiley & Sons. Chichester, UK.
- Filntisi A, Papangelopoulos N, Bencurova E, Kasampalidis I, Matsopoulos G, Vlachakis D & Kossida S. 2013. State-of-the-art neural networks applications in biology. *International Journal of Biological Sciences*, 2: 63–85. DOI: 10.4018/ijsbtt.2013100105
- Gafarov FM & Galimyanov AF. 2018. [Artificial Neural Networks and Applications]. Kazan University Publishing House. Kazan, Russia.
- Gao X, Cassidy A, Schwarzschild MA, Rimm EB & Ascherio A. 2012. Habitual intake of dietary flavonoids and risk of Parkinson disease. *Neurology*, 78: 1138–1145. DOI: 10.1212/WNL.0b013e31824f7fc4
- Garrapa R. 2018. Numerical solution of fractional differential equations: A survey and a software tutorial. *Mathematics*, 6: 16. DOI: 10.3390/math6020016
- Ge SS & Wang C. 2002. Direct adaptive NN control of a class of nonlinear systems. *IEEE Transactions on Neural Networks*, 13: 214–221. DOI: 10.1109/72.977306
- Ghahri H, Habibian R & Fam A. 2010. Evaluation of the efficacy of esterified glucomannan, sodium bentonite, and humic acid to ameliorate the toxic effects of aflatoxin in broilers. *Turkish Journal of Veterinary & Animal Sciences*, 34: 385–391. DOI: 10.3906/vet-0903-19
- Gharajeh MS. 2018. Chapter eight – Biological big data analytics. *Advanced Computing*, 109: 321–355. DOI: 10.1016/bs.adcom.2017.08.002
- Goodfellow I, Bengio Y & Courville A. 2016. *Deep Learning*. MIT Press. Cambridge, MA, USA.
- Gorodnichev RM, Pestryakova LA, Ushnitskaya LA, Levina SN & Davydova PV. 2019. [Methods of Environmental Research. Fundamentals of Statistical Data Processing: Educational and Methodological Manual]. NEFU Publishing

- House. Yakutsk, Russia. ISBN 978-5-7513-2737-8.
- Grishanov GV & Grishanova YuN. 2010. [Methods for Studying and Assessing Biological Diversity]. I. Kant Russian University. Kaliningrad, Russia.
- Grosu GF, Hopp AV, Moca VV, Bârzan H, Ciuparu A, Ercsey-Ravasz M, Winkel M, Linde H & Mureşan RC. 2023. The fractal brain: Scale-invariance in structure and dynamics. *Cerebral Cortex*, 33: 4574–4605. DOI: 10.1093/cercor/bhac363
- Hsieh WW. 2000. Nonlinear canonical correlation analysis by neural networks. *Neural Networks*, 13: 1095–1105. DOI: 10.1016/S0893-6080(00)00067-8
- IAKI.RF. 2024. [Vitamin B2 (Riboflavin)]. LLC "Institute of Allergology and Clinical Immunology", IAKI.RF. <https://xn--80apbh.xn--plai/analyzes/vitamins/vitamin-v2-riboflavin/>.
- Ignatov I & Mosin O. 2014. [Composition and structural properties of fullerene analogous mineral shungite. Mathematical model of interaction of shungite with water molecules]. *Naukovedenie*, 2(21): 12TVN214.
- Jahanimiri F & Parker DC. 2022. An overview of fractal geometry applied to urban planning. *Land*, 11: 475. DOI: 10.3390/land11040475
- Jolliffe IT. 2002. *Principal Component Analysis*. 2nd ed. Springer. New York, NY, USA. ISBN 978-0-387-95442-4. DOI: 10.1007/b98835
- Khlivnenko LV. 2015. [Practice of Neural Network Modeling]. Voronezh State Technical University. Voronezh, Russia. ISBN 978-5-7731-0429-2.
- Kirikovich SA, Kirikovich YuK & Kurepin AA. 2012. [The influence of exogenous factors on the productivity, safety and natural resistance of animals]. [Collection of Scientific Papers of the Stavropol Research Institute of Animal Husbandry and Forage Production], 2: 264–272.
- Kochish II, Smolensky VI, Laptev GY, Romanov MN, Nikonov IN, Panin AN, Surai PF, Ilina LA, Myasnikova OV, Selina MV, Korenyuga MV. 2019. Practical Recommendations for the Use of Feed Additives to Improve the Productivity and Stress Resistance of Egg Poultry. *Sel'skokhozyaistvennye tekhnologii*. Moscow, Russia.
- Kochish II, Romanov MN, Nikonov IN. 2020a. Ways to improve the productivity of chickens (using a feed additive based on shungite as an example). In: [Materials of the XIX International Scientific and Practical Conference "Modern Trends in Agricultural Production in the World Economy"]. FSBEI HE "Kuzbass State Agricultural Academy". Kemerovo, Russia. Pages 37–44.
- Kochish II, Pozyabin SV, Vorobyov NI & Nikonov IN. 2020b. Fractal bioconsolidation of microorganisms in the intestines of laying hens due to the use of a feed additive from the mineral shungite. In: Materials of the 2nd International Scientific and Practical Conference on Molecular Genetic Technologies for Analysis of Gene Expression Related to Animal Productivity and Disease Resistance. Moscow, Russia, 25 December 2020. *Sel'skokhozyaistvennye tekhnologii*. Moscow, Russia. Pages 59–75. DOI: 10.18720/SPBPU/2/k20-5 (In Russian with English summary)
- Kochish II, Romanov MN, Myasnikova OV, Nikonov IN, Selina MV, Korenyuga MV, Zimin EE, Sharafetdinov GR & Martynov VV. 2021. Methodical Recommendations on the Use of Antimicrobial Feed Additive for the Prevention of Stress in Industrial Crosses of Laying Hens. *Sel'skokhozyaistvennye tekhnologii*. Moscow, Russia. ISBN 978-5-6047654-0-1. DOI: 10.18720/SPBPU/2/z21-33 (In Russian)
- Kochish I, Vorobyov N, Nikonov I & Selina M. 2022. Neural network fractal model to evaluate the effectiveness of antimicrobial feed additives in egg poultry farming. *International Transaction Journal of Engineering, Management, & Applied Sciences & Technologies*, 13: 13A12S. DOI: 10.14456/ITJEMAST.2022.250
- Kochish II, Brazhnik EA, Vorobyov NI, Nikonov IN, Korenyuga MV, Myasnikova OV, Griffin DK, Surai PF & Romanov MN. 2023. Features of fractal conformity and bioconsolidation in the early myogenesis gene expression and their relationship to the genetic diversity of chicken breeds. *Animals*, 13: 521. DOI: 10.3390/ani13030521
- Laptev GYu, Ilyina LA, Nikonov IN, Kochish II, Romanov MN, Smolensky VI, Panin AN, Yildirim EA, Novikova NI, Filippova VA & Dubrovin AV. 2017. [Determination of intestinal microbiocenoses of chickens of the Hisex breed by the T-RFLP method in ontogenesis]. *Acta Naturae*, 9(SI 1): 33. (In Russian)
- Levchenko PV, Zhuchok AYU, Gugushvili NN & Inyukina TA. 2020. The change in hematological indices of blood of laying hens when using photomodulation in the early post-embryonic period. *Collection of Scientific Papers of KRCAHVM*, 9: 365–369. DOI: 10.34617/17qr-r266 (In Russian with English summary)
- Mascarenhas WF. 2018. Fast and accurate normalization of vectors and quaternions. *Computational and Applied Mathematics*, 37: 4649–4660. DOI: 10.1007/s40314-018-0594-6
- Matveev OA & Torshkov AA. 2020. Morphological indicators of blood of chicken-broilers in post-public ontogenesis. *Uchenye zapiski Kazanskoy gosudarstvennoy akademii veterinarnoy mediciny im. N.É. Baumana Scientific Notes, Kazan*

- Bauman State Academy of Veterinary Medicine, 241: 138–142. DOI: 10.31588/2413-4201-1883-241-1-138-142 (In Russian with English summary)
- Meakin P. 1999. A historical introduction to computer models for fractal aggregates. *Journal of Sol-Gel Science and Technology*, 15:97–117. DOI: 10.1023/A:1008731904082
- Meireles MRG, Almeida PEM & Simoes MG. 2003. A comprehensive review for industrial applicability of artificial neural networks. *IEEE Transactions on Industrial Electronics*, 50: 585–601. DOI: 10.1109/TIE.2003.812470
- Menshikov VV. 1997. [Clinical Diagnosis – Laboratory Basics]. Labinform Publishing House. Moscow, Russia. ISBN 5-89429-002-3. (In Russian)
- Milevski I. 2024. Effect of Erythropoietin on Erythrocytogenesis. Vitamin B12 and Folic Acid in Erythropoiesis. MedUniver. (In Russian)
- Minsky M & Papert S. 1969/1988. *Perceptrons: An Introduction to Computational Geometry*. MIT Press. Cambridge, MA, USA. ISBN 978-0-262-53477-2.
- Moroz LI & Maslovskaya AG. 2020. Hybrid stochastic fractal-based approach to modeling the switching kinetics of ferroelectrics in the injection mode. *Mathematical Models and Computer Simulations*, 12: 348–356. DOI: 10.1134/S207004822003014X
- Nematnia E & Abdanan Mehdizadeh S. 2018. Assessment of egg freshness by prediction of Haugh unit and albumen pH using an artificial neural network. *Journal of Food Measurement and Characterization*, 12: 1449–1459. DOI: 10.1007/s11694-018-9760-1
- Nikolenko S, Kadurin A & Arkhangelskaya E. 2018. *Deep Learning. Dive into the World of Neural Networks*. Peter. St. Petersburg, Russia. (In Russian)
- Nikolić D, Muresan RC, Feng W & Singer W. 2012. Scaled correlation analysis: A better way to compute a cross-correlogram. *European Journal of Neuroscience*, 35: 1–21. DOI: 10.1111/j.1460-9568.2011.07987.x
- Ojo RO, Ajayi AO, Owolabi HA, Oyedele LO & Akanbi LA. 2022. Internet of things and machine learning techniques in poultry health and welfare management: A systematic literature review. *Computers and Electronics in Agriculture*, 200: 107266. DOI: 10.1016/j.compag.2022.107266
- Okolelova TM & Mansurov RF. 2013. Efficiency of adsorbents in compound feeds contaminated with mycotoxins. *Pitisevodstvo, Poultry Farming*, 11: 17–18. (In Russian)
- Pitelinskiy KV & Shimanskiy SA. 2013. Application of neural networks in socioeconomic system studies. *Vestnik Moskovskogo instituta lingvistiki [Vestnik of Moscow State Linguistic University]*, 1: 88–91. (In Russian with English summary)
- Pitelinskiy KV & Tyurkin AA. 2007. Applying of neuron networks at an estimation of the real estate. In *NTI-2007: Information Society, Intelligent Information Processing, Information Technologies, Proceedings of the International Conference Dedicated to the 55th Anniversary of VINITI*. Moscow, Russia, 24–26 October 2007. VINITI of RAS. Moscow, Russia. Pages 251–253. (In Russian with English summary)
- Popov VA. 2013. *Probability Theory. Part 2. Random Variables*. Kazan University. Kazan, Russia. (In Russian)
- Przybyła-Kasperek M & Marfo KF. 2024. A multi-layer perceptron neural network for varied conditional attributes in tabular dispersed data. *PloS One*, 19: e03110. DOI: 10.1371/journal.pone.0311041
- Pukhalskiy YaV, Vorobyev NI, Loskutov SI & Laktionov YuV. 2023. Genotypic screening for the resistance of leguminous crops to the effects of heavy metals, based on neuron profiling of their amino acid exudation. *Agrarnyy vestnik Urala [Agrarian Bulletin of the Urals]*, 05(234): 83–96. DOI: 10.32417/1997-4868-2023-234-05-83-96. (In Russian with English summary)
- Pyrhönen L, Willems T, Mikkola A & Naets F. 2024. Inertial parameter identification for closed-loop mechanisms: adaptation of linear regression for coordinate partitioning. *Journal of Computational and Nonlinear Dynamics*, 19: 051001. DOI: 10.1115/1.4064794
- Rosenblatt F. 1962. *Principles of Neurodynamics: Perceptrons and the Theory of Brain Mechanisms*. Cornell Aeronautical Laboratory, Inc., Cornell University. Buffalo, NY, USA.
- Schmidhuber J. 2015. Deep learning in neural networks: An overview. *Neural Networks*, 61: 85–117. DOI: 10.1016/j.neunet.2014.09.003
- Sergeev AP & Tarasov DA. 2017. [Introduction to Neural Network Modeling]. Ural University Publishing House. Ekaterinburg, Russia. (In Russian)
- Sharapova VV. 2011. [The use of additives from fucus algae and shungite in feeding laying hens]. Author's Abstract. Cand. Agric. Sci. Dissertation. GNU VNITIP of the Russian Agricultural Academy. Sergiev Posad, Russia. (In Russian)
- Siriani ALR, Miranda IBdC, Mehdizadeh SA & Pereira DF. 2023. Chicken tracking and individual bird activity monitoring using the BoT-SORT algorithm. *AgriEngineering*, 5: 1677–1693. DOI: 10.3390/agriengineering5040104
- Sutrop U. 2001. List task and a cognitive salience index. *Field Methods*, 13: 263–276. DOI: 10.1177/1525822X0101300303

- Taye MM. 2023. Theoretical understanding of convolutional neural network: Concepts, architectures, applications, future directions. *Computation*, 11: 52. DOI: 10.3390/computation11030052
- Tietz NW. 1997. *Encyclopedia of Clinical Laboratory Tests*. Menshikov VV, Ed. Labinform Publishing House. Moscow, Russia. (In Russian)
- Tukhbatov IA. 2013. Meat productivity formation of the broiler chickens due to the sorbent feed additive. *Kormoproizvodstvo [Fodder Journal]*, 8: 40–42. (In Russian with English summary)
- Tyurina LE. 2022. Scientific and practical justification for the use of mineral substances from the sources of the Krasnoyarsk territory in feeding farm animals and poultry, Dissertation. Krasnoyarsk State Agrarian University. Krasnoyarsk, Russia. (In Russian)
- Vats T, Dutt S, Kumar R & Siril PF. 2016. Facile synthesis of pristine graphene-palladium nanocomposites with extraordinary catalytic activities using swollen liquid crystals. *Scientific Reports*, 6: 33053. DOI: 10.1038/srep33053
- Vorobyov NI, Bogolyubova NV, Platonov AV, Nikonov IN, Selina MV, Guselnikova AA & Sidnev NY. 2023a. Effect of feed supplements on blood biochemical parameters and intensity of metabolic processes in cows: the neural network modeling method. *International Journal of Chemical and Biochemical Sciences*, 24: 165–170.
- Vorobyov NI, Selina MV, Guselnikova AA, Nikonov IN & Sidnev NYu. 2023b. Program for neural network analysis of biochemical parameters of animal blood. Certificate of state registration of a computer program: RU 2023662779. Russia.
- Wentzel ES. 1999. [Probability Theory]. 6th ed. Vysshaya Shkola. Moscow, Russia. ISBN 5-06-003650-2. (In Russian)
- Widrow B, Greenblatt A, Kim Y & Park D. 2013. The No-Prop algorithm: A new learning algorithm for multilayer neural networks. *Neural Networks*, 37: 182–188. DOI: 10.1016/j.neunet.2012.09.020
- Yang X, Bist RB, Subedi S & Chai L. 2023. A computer vision-based automatic system for egg grading and defect detection. *Animals*, 13: 2354. DOI: 10.3390/ani13142354
- Yarovani NI. 2005. Use of Khotynets natural zeolites of the Oryol Region to normalize free radical oxidation in pigs. In: *Materials of the 3rd International Symposium on Modern Problems of Veterinary Dietology and Nutritional Science*. St. Petersburg, Russia. Pages 170–171. (In Russian)
- Yeung KY & Ruzzo WL. 2001. Principal component analysis for clustering gene expression data. *Bioinformatics*, 17: 763–774. DOI: 10.1093/bioinformatics/17.9.763
- Young G, Nippgen F, Titterbrandt S & Cooney MJ. 2010. Lipid extraction from biomass using co-solvent mixtures of ionic liquids and polar covalent molecules. *Separation and Purification Technology*, 72: 118–121. DOI: 10.1016/j.seppur.2010.01.009
- Yurkovych N, Mar'yan M, Opachko M & Seben V. 2023. Methods of the fractal approach in science education: innovative technology and concepts of computer modeling. *Physical and Mathematical Education*, 38: 73–78. DOI: 10.31110/2413-1571-2023-038-3-010
- Zaikina AS, Buryakov NP, Vorobyov NI & Nikonov IN. 2022. Neural network analysis of the compliance of the microbial-organismic biosystem of poultry intestines with a fractal-stochastic model. *Permskij agrarnyj vestnik, Perm Agrarian Journal*, 4(40): 98–106. DOI: 10.47737/2307-2873_2022_40_98. (In Russian with English summary)
- Zasorin AV, Zubkov DG, Selmenskij GE, Nikonov IN & Alimpiev SV. 2019. Method for Reducing the Negative Impact of Mycotoxins in Poultry (Options). National Center for Biotechnology Information. PubChem Patent Summary for RU-2680009-C1.
- Zhu B, Shin U & Shoaran M. 2021. Closed-loop neural prostheses with on-chip intelligence: A review and a low-latency machine learning model for brain state detection. *IEEE Transactions on Biomedical Circuits and Systems*, 15: 877–897. DOI: 10.1109/TBCAS.2021.3112756