

Enguito, Marie Rosellynn C., Garnero, Analía Del Valle, Gunski, Ricardo José, de Souza, Marcelo Santos, O'Connor, Rebecca E., Srikulnath, Kornsorn, Singchat, Worapong, de Oliveira, Edivaldo Herculano Correa, Romanov, Michael N, Griffin, Darren K. and and others (2026) Comparative chromosomal analysis of the Z chromosome in South American bird species shows a high rate of intrachromosomal rearrangements. Genes, 17 (1). ISSN 2073-4425.

Downloaded from

<https://kar.kent.ac.uk/112803/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.3390/genes17010112>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Article

Comparative Chromosomal Analysis of the Z Chromosome in South American Bird Species Shows a High Rate of Intrachromosomal Rearrangements

Marie Rosellynn C. Enguito ^{1,2} , Analía Del Valle Garnero ³ , Ricardo José Gunski ³ , Marcelo Santos de Souza ⁴ , Rebecca E. O'Connor ¹ , Kornsorn Srikulnath ⁵ , Worapong Singchat ⁵ , Edivaldo Herculano Correa de Oliveira ^{6,7} , Michael N. Romanov ^{1,5,8} , Darren Karl Griffin ^{1,5,*} , and Rafael Kretschmer ⁹ 

¹ School of Natural Sciences, University of Kent, Canterbury CT2 7NJ, UK; marierosellynnndce@gmail.com (M.R.C.E.); rebeccyoc@gmail.com (R.E.O.); m.romanov@kent.ac.uk (M.N.R.)

² Misamis University, H. T. Feliciano St., Ozamiz City 7200, Philippines

³ Laboratório de Diversidade Genética Animal, Universidade Federal do Pampa, São Gabriel 97300-000, Brazil; analiagarnero@unipampa.edu.br (A.D.V.G.); ricardogunski@unipampa.edu.br (R.J.G.)

⁴ Programa de Pós-Graduação em Zoologia, Universidade Federal do Amazonas, Manaus 69077-000, Brazil; marcelodesouzabio@gmail.com

⁵ Animal Genomics and Bioresource Research Unit (AGB Research Unit), Faculty of Science, Kasetsart University, Chatuchak, Bangkok 10900, Thailand; kornsorn.s@ku.ac.th (K.S.); worapong.singc@ku.ac.th (W.S.)

⁶ Instituto de Ciências Exatas e Naturais, Universidade Federal do Pará, Belém 66075-110, Brazil; ehco@ufpa.br

⁷ Laboratório de Cultura de Tecidos e Citogenética, SEAMB, Instituto Evandro Chagas, Ananindeua 67030-000, Brazil

⁸ L. K. Ernst Federal Research Center for Animal Husbandry, Dubrovitsy, Podolsk Urban Okrug, Moscow Oblast 142132, Russia

⁹ Laboratório de Citogenética e Evolução, Departamento de Genética, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Porto Alegre 91509-900, Brazil; rafael.kretschmer@ufrgs.br

* Correspondence: d.k.griffin@kent.ac.uk

Abstract

Background: Intrachromosomal rearrangements in birds play a subtle but important role in shaping genomic evolution, phenotypic diversity and speciation. However, the avian sex chromosome system (homogametic ZZ males; heterogametic ZW females) remains relatively understudied, and evolutionary rearrangements of the Z chromosome have not been mapped in most species. To address this, we employed universally hybridizing avian Z chromosome probes to metaphases of 11 avian species from South America.

Methods: Chromosome preparations were obtained from fibroblast cell cultures of 11 birds representing nine different orders; four bacterial artificial chromosome (BAC) probes were used in our interspecies fluorescence in situ hybridization (FISH) experiments. We identified chromosomal rearrangements in the species investigated, tracing the evolution of the Z chromosome in these species through comparison with reptiles from Southeast Asia (three snake species used as an outgroup), along with two reference species: chicken (Galliformes) and zebra finch (Passeriformes). **Results:** We observed high rates of intrachromosomal rearrangements in the avian Z chromosome, with most species showing different patterns from chicken and zebra finch. *Nanopterum brasiliannum* (Suliformes) and *Jacana jacana* (Charadriiformes) showed the same BAC order as chicken, but centromere repositioning was evident. Apart from Piciformes, all other species exhibited a conserved Z chromosome size. The corresponding Z chromosome sequences were homologous to regions of the long arms of Chromosome 2 and W in snakes but not on the Z chromosomes. **Conclusions:** Comparative analysis of the Z chromosome across avian orders provides important insights into the dynamics of avian sex chromosomes and the evolution of sex chromosome systems in general.



Academic Editor: Christophe Escudé

Received: 18 November 2025

Revised: 12 December 2025

Accepted: 1 January 2026

Published: 20 January 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\) license](#).

Keywords: avian Z chromosome; interspecies fluorescence in situ hybridization (FISH); bacterial artificial chromosome (BAC) probes; South American bird species; intrachromosomal rearrangements; centromere

1. Introduction

Chromosomal organization in birds plays a fundamental role in shaping their genomic evolution, phenotypic diversity and speciation processes [1]. In addition to typically around 36–40 autosomal pairs (~10 macrochromosomes, ~30 microchromosomes), a ZW sex chromosome system predominates [2].

Recent reviews by Griffin et al. [3,4] showed that, although avian genome organization is largely conserved and dates back to the age of the dinosaurs, some orders exhibit highly derived karyotypes. For example, Falconiformes (falcons, hobbies and relatives) have a reduced diploid number of approximately 50 chromosomes [5], whereas Piciformes (woodpeckers, toucans and relatives) display strikingly high diploid numbers exceeding 100 chromosomes [3,4,6,7]. Recent large-scale studies demonstrate that phylogenomic signals differ between micro- and macrochromosomes and that assigning loci to specific chromosomes improves the resolution of complex evolutionary histories [8,9]. For instance, there are contrasting roles for microchromosomes and macrochromosomes [10], the former being gene-rich, GC-rich, tending to higher recombination rates, and evolving faster than macrochromosomes [11]. Patterns of lineage-specific chromosomal rearrangement thus not only illustrate exceptions to long-term karyotype stability but also set the stage for further structural innovations that have profound effects on avian evolution [12]. This emphasizes the fact that a detailed understanding of chromosomal organization is essential for interpreting genome evolution and for reconstructing avian evolutionary trees accurately [13].

Beyond their value for phylogenetic reconstruction, patterns of chromosomal organization actively shape how genetic variation is generated and maintained across avian genomes. Recent cytogenomic and comparative genomic studies demonstrate that chromosomal rearrangement, variation in recombination rate, and sex chromosome dynamics jointly drive lineage-specific evolutionary trajectories in birds [14–16]. For instance, inversions and other intrachromosomal rearrangements can suppress recombination in affected regions, allowing co-adapted allele combinations to persist [17]. Indeed, inversion polymorphisms often coincide with loci implicated in local adaptation and reproductive isolation. This indicates that chromosomal structure can actively promote divergence and, in some cases, speciation by preserving suites of adaptive alleles from recombination-mediated breakup [18].

Moreover, the propensity for structural chromosomal changes is modulated by genomic features that influence chromosome stability. Transposable element activity, repeat density and the presence or absence of DNA-repair genes can bias lineages toward higher or lower rates of rearrangements. For instance, parrot lineages with altered repair or gene-loss signatures show elevated fusion and fission activity, helping explain why some bird clades remain highly karyotypically stable, while others are remarkably labile [18]. Collectively, the partitioning of genomes into micro- and macrochromosomes, lineage-specific rearrangement patterns and, relevant to this study, the behavior of the Z and W sex chromosomes act as a structural scaffold for the mapping of aspects of phenotypic trait evolution. These include determining where and how variation arises, how it is shuffled by recombination, how it responds to natural selection, adaptation rates and the emergence of sex-linked phenotypes.

As reviewed by Griffin et al. [3,4], the avian ZW sex-determination system was probably in place before the divergence of Palaeognathae and Neognathae, but the size differentiation between Z and W chromosomes emerged after this divergence. In the ZW sex chromosomes of most Neognathae, the W chromosome is dynamic, both in size and morphology, because of the accumulation of repetitive sequences [19–21]. The Z chromosome, however, unlike the W chromosome, which has undergone extensive degeneration, shows relatively conserved gene content, with structural rearrangements such as inversions comprising most of its variation [2–4]. Typically, the Z chromosome is usually around the fifth-largest chromosome and conserved in size. However, there are exceptions; in Piciformes, it is unusually large, probably because of the accumulation of repetitive DNA sequences [6,22]. Indeed, it exhibits notable morphological variability even among closely related species. For instance, in Columbidae (pigeons, doves, etc.), *Columbina picui* (Picui ground dove) possesses a telocentric Z chromosome, whereas *Leptotila verreauxi* (white-tipped dove) has a metacentric Z. This highlights the role of intrachromosomal changes in its evolution [23]. Similarly, fluorescence in situ hybridization (FISH) mapping using a bacterial artificial chromosome (BAC) clone (for the *ALDOB* gene) across four Galloanseres species (*Gallus gallus*, *Coturnix coturnix*, *Chrysolophus pictus*, and *Cairina moschata*—chicken, common quail, golden pheasant, and muscovy duck) revealed species-specific signal patterns [24], with BAC-FISH analysis also being instrumental for refining centromere positions [25].

Recent reviews suggest that avian sex chromosomes undergo considerable inter- and intrachromosomal change and are thus not as stable as previously thought [1–4,10]. For instance, a multiple-sex chromosome system ($\delta^{\circ}Z1Z1Z2Z2/\varphi^{\circ}Z1Z2W$) has been described by Gunski et al. [26] in the Adélie penguin (*Pygoscelis adeliae*, Sphenisciformes) and in *Sula* species [27]. Independent autosome–sex chromosome fusions have also been reported in Sylvioidea species [28–34] and neo-sex chromosomes derived from such fusion events have been identified in parrots, cuckoos and Pacific Island birds [18,35–37]. Neo-sex chromosomes can diverge rapidly in gene content and expression; they can influence sexual dimorphism and affect patterns of selection because of sex-specific inheritance and suppressed recombination [29,32].

Given that evolutionary rates and rearrangement frequencies vary significantly by chromosome [38], genomic analyses that ignore the chromosomal context risk introducing bias. The Z chromosome is particularly critical in this regard. That is, while its gene content is conserved, its structural rearrangements can act as powerful drivers of reproductive isolation and phenotypic evolution. Therefore, characterizing these Z-specific structural patterns is essential for accurate phylogenomic interpretations [12]. This study addresses this need by investigating the role of Z chromosome rearrangements in the diversification of the class Aves, providing new insights into the evolutionary consequences of sex chromosome variability in South American birds.

Indeed, despite several studies, the extent and diversity of chromosomal rearrangements in avian Z chromosomes remain poorly explored. This is partly due to limited molecular cytogenetic tools compared to the autosomes [4,39]. A critical gap in our understanding of avian karyotype evolution is therefore apparent. South American birds particularly represent distinct evolutionary radiations, with diverse ecological pressures. Their Z chromosome organization nonetheless remains poorly characterized [22,40]. This study thus applied four universally hybridizing FISH probes derived from chicken and zebra finch genomes to the Z chromosomes of 11 South American bird species representing nine distinct orders. Comparative analysis with data from squamate reptiles (snakes) plus chicken and zebra finch provides further insight into the evolutionary origins and diversification of the avian Z chromosome, tracing its structural evolution across different lineages.

2. Materials and Methods

2.1. Sample Collection

Metaphase chromosome preparations were obtained from 11 bird species (one per species) from nine different orders (Table 1), along with, for comparison, two reference avian species (chicken and zebra finch) as well as three outgroup reptiles (snakes) (Table 1). Specimens were sourced ethically from research collaborations, ensuring minimal impact on wild populations. Tissue samples (blood or fibroblast cultures) were collected following approved institutional animal care and use protocols.

Table 1. Species used in this study, Latin name, common name, species code, shape of chromosome, diploid chromosome number and most appropriate reference with an accurate chromosome number. Arranged according to evolutionary tree, e.g., Passeriformes are most distant from Squamata.

Order	Scientific Name	3-Letter Species Code	Common Name	Z Morphology	2n	Reference
New South American avian species						
Passeriformes	<i>Myiodynastes maculatus</i>	MMA	Streaked Flycatcher	Metacentric	80	[41]
Psittaciformes	<i>Myiopsitta monachus</i>	MMO	Monk Parakeet	Submetacentric	48	[42]
Piciformes	<i>Veniliornis spilogaster</i>	VSP	White-spotted Woodpecker	Metacentric	88	[43]
Piciformes	<i>Pteroglossus inscriptus</i>	PIN	Lettered Araçari	Acrocentric	112	[6]
Trogoniformes	<i>Trogon surrucura</i>	TSU	Surucua Trogon	Metacentric	82	[7,44]
Suliformes	<i>Nannopterum brasilianum</i>	NBR	Neotropical Cormorant	Submetacentric	74	[7]
Charadriiformes	<i>Jacana jacana</i>	JJA	Wattled Jacana	Submetacentric	82	[35,45]
Gruiformes	<i>Gallinula melanops</i>	GME	Spot-flanked Gallinule	Submetacentric	80	[20]
Caprimulgiformes	<i>Hydropsalis torquata</i>	HTO	Scissor-tailed nightjar	Acrocentric	74	[7]
Columbiformes	<i>Leptotila verreauxi</i>	LVE	White-tipped Dove	Metacentric	78	[23]
Columbiformes	<i>Columbina picui</i>	CPI	Picui Ground Dove	Telocentric	76	[23]
Reference species						
Passeriformes	<i>Taeniopygia guttata</i>	TGU	Timor Zebra Finch	Metacentric	80	-
Galliformes	<i>Gallus gallus</i>	GGA	Red junglefowl	Metacentric	78	-
Squamata	<i>Lacerta agilis</i>	LAG	Sand lizard	Acrocentric	38	[46]
Squamata	<i>Daboia russelii</i>	DRU	Western Russel's Viper	Metacentric	36	[47]
Squamata	<i>Notechis scutatus</i>	NSC	Mainland Tiger Snake	Metacentric	36	[47]
Squamata	<i>Naja kaouthia</i>	NKA	Siamese cobra	Metacentric	38	[48]

2.2. Chromosome Preparation

Primary cell cultures were established from tissue samples using standard protocols. Mitotic cells were arrested at metaphase by colcemid (0.05 µg/mL) exposure for 1 h, followed by hypotonic treatment in 0.075 M KCl for 20 min to cause swelling in the cytoplasm and the nucleus, making the nuclear membrane more fragile for the release of chromosomes, generating clear metaphase spreads. Cells were fixed in a 3:1 methanol/acetic acid solution (Carnoy's fixative) to preserve cellular and chromosomal structure, and metaphase spreads were prepared on clean glass slides following previously published protocols [19]. Chromosomes were air-dried and stored at -20°C until use. The chromosome preparations used here were the same as in previous studies shown in the tables presented above.

2.3. Fluorescence In Situ Hybridization (FISH)

For interspecies chromosome mapping, we selected four BAC probes derived from the Z chromosome of chicken, *Gallus gallus* (CH261-129A16 and CH261-133M4), and zebra finch, *Taeniopygia guttata* (TGMcba-200J22 and TGMcba-27019), which showed higher hybridization efficiency in cross-species FISH experiments [39]. These probes were used to investigate intrachromosomal rearrangements. Chromosomes were counterstained with DAPI (blue), and the BAC probes were labeled with Texas Red (red) or FITC (green).

At least 10 metaphases per experiment were analyzed to confirm the FISH results. The BAC clone isolation, amplification, labeling and FISH followed O'Connor et al. [49]. This approach has been applied routinely in our respective laboratories for many years, and we used it without adaptation.

2.4. Imaging and Data Analysis

Chromosomes were counterstained before analysis with 4,6-diamidino-2-phenylindole (DAPI). FISH signals and DAPI-stained chromosomes were photographed using an Olympus BX61 epifluorescence microscope equipped with a cooled CCD camera (Olympus, Tokyo, Japan), and the images were captured at 100 \times magnification using SmartCapture 3 software (Digital Scientific UK, Cambridge, UK) [49]. Comparative analysis of signal patterns among South American birds, as well as with published data from squamate reptiles, was used to infer evolutionary rearrangements and the structural dynamics of the Z chromosome.

2.5. Comparative Analysis and Evolutionary Inference

Intrachromosomal rearrangements were categorized according to type and frequency across species, using BAC probe positions on the chicken Z chromosome as a reference. Homology comparisons with previously published FISH data on reptiles provided context for the origin and evolution of the avian Z chromosome. Observed rearrangements were interpreted in the framework of lineage-specific chromosomal evolution, sex chromosome diversification, and potential adaptive significance.

3. Results

3.1. Comparative FISH Mapping Across South American Birds

There were 33 probes successfully hybridized onto the metaphases of 11 South American bird species. When probes did not show signals, this was down to insufficient number of metaphases or unknown factors (we find this to be commonplace in work of this kind). Our results nonetheless provide evidence of a high rate of intrachromosomal rearrangements in the avian Z chromosomes of South American birds, with most species showing different patterns from chicken and zebra finch. The species exhibited all possible chromosomal morphologies for the Z chromosome (metacentric, submetacentric, acrocentric, and telocentric), as shown in Figures 1 and 2.

3.2. Mapping Intrachromosomal Changes with Avian BAC Probes

Table 2 shows comparative mapping of chicken Z chromosome BAC probes onto the chromosomes of *N. kaouthia* (monocled cobra). The chromosomes in *N. kaouthia* where the BAC probe hybridized were the long arm of W chromosome (Wq) and the long arm of chromosome 2. Parts of the chicken Z chromosome correspond to different chromosomes in *N. kaouthia*. Some Z-linked BACs mapped to W chromosome (long arm), suggesting sex chromosome homology. Other Z-linked BACs mapped to autosome 2 (long arm), indicating chromosomal rearrangements (like translocations) between species.

Table 2. Comparative mapping of chicken Z chromosome BAC probes onto the chromosomes of *Naja kaouthia*.

Chromosome Number	Chicken and ZF Chromosome	BAC Name	Chromosomal Location in <i>Daboia russeli</i>	Chromosomal Location in <i>Notechis scutatus</i>	Chromosomal Location in <i>Naja kaouthia</i>
40	Z	CH261-129A16		-	Wq
41	Z	TGMCBA-200J22		W	Wq
42	Z	TGMCBA-27019	W	W	2q
43	Z	CH261-133M4		2p, W	2q




Figure 1. FISH mapping of Z chromosome BAC probes onto metaphase of selected South American bird species. The panels show hybridizations for: (a) female (*V. spilogaster*) with zebra finch BAC probe TGM CBA-200J22 (green); (b) female (*V. spilogaster*) with chicken BAC probe CH261-129A16 (green) and zebra finch BAC probe TGM CBA-27019 (red); (c) male (*N. brasiliianum*) with zebra finch BAC probe TGM CBA-200J22 (green); (d) male (*N. brasiliianum*) with chicken BAC probes CH261-129A16 (green) and CH261-133M4 (red); (e) female (*H. torquata*) with zebra finch BAC probe TGM CBA-27019 (red) and chicken BAC probe CH261-133M4 (green); (f) male (*G. melanops*) with chicken BAC probe CH261-129A16 (green) and zebra finch BAC probe TGM CBA-27019 (red); (g) female (*P. inscriptus*) with chicken BAC probe CH261-129A16 (green) and zebra finch BAC probe TGM CBA-27019 (red); (h) male (*J. jacana*) with chicken BAC probe CH261-133M4 (red); and (i) male (*M. monachus*) with chicken BAC probe CH261-129A16 (green) and zebra finch BAC probe TGM CBA-27019 (red). In all cases chromosomes are counterstained with 4',6'-diamidinophenylindole (DAPI)—blue colour. Scale bar = 10 μ m.




Figure 2. Summary of the chromosome positions of each of the BACs established by manual measurement of fractional length from the p-terminus (FLpter). Ideograms represent the Z chromosomes, arranged according to taxonomic proximity, with colored bands indicating the mapped positions of four BAC probes. The probes utilized were derived from chicken (*G. gallus*; CH261-129A16 and CH261-133M4) and zebra finch (*T. guttata*; TGMcba-200J22 and TGMcba-27019) genomes. The adapted phylogenetic tree was sourced from TimeTree databases (<http://www.timetree.org>, accessed on 7 December 2025). Species names are represented by individual code indicated below each respective ideogram: LVE (*L. verreauxi*), CPI (*C. picui*), HTO (*H. torquata*), TSU (*T. surrucura*), NBR (*N. brasiliense*), GME (*G. melanops*), JJA (*J. jacana*), VSP (*V. spilogaster*), PIN (*P. inscriptus*), MMO (*M. monachus*), MMA (*M. maculatus*).

4. Discussion

4.1. Evolutionary Insights into the Avian Z Chromosome

The overall evolutionary dynamics of Z chromosome rearrangement in the species studied are summarized in Figure 2. This gives a global overview of how the Z chromosome has changed intrachromosomally, despite being conserved interchromosomally. Of course, there were limitations inherent in our approach. They included the fact that not all probes worked on all species, the limits of resolution of all molecular cytogenetic studies (even on very extended chromosomes), the labor-intensive nature of the work, and the fact that it all relies on the vagaries of cell culture to generate analyzable metaphases.

Of the 11 species tested, only two, *N. brasiliandum* and *J. jacana*, showed the same BAC order as chicken (Figure 2). These, however, were submetacentric chromosomes, different from those of the chicken (which has a metacentric morphology), evidencing the occurrence of a centromere repositioning. The other species showed different chromosome positions, at least for one Z BAC probe. The exception was in the Piciformes, which have the Z as the largest chromosome in the karyotype, the other species having a conserved Z chromosome size (about the fifth largest). The morphology of this chromosome is variable in birds, even in closely related species. For instance, the Z chromosome in *C. picui* is telocentric, while it is metacentric in *L. verrauxi* [23]. Moreover, the chicken Z chromosome is partially conserved in cobra species but reorganized across chromosomes. Two BACs (CH261-129A16 and TGMABA-200J22) mapped to Wq. This suggests that regions of the chicken Z are homologous to the cobra W chromosome, hinting at shared ancestral sex chromosome segments. The other two BACs mapping to 2q show intrachromosomal or interchromosomal rearrangements, supporting the idea that bird and snake genomes have undergone major rearrangements since their divergence. This result helps reconstruct evolutionary changes in sex chromosomes and autosomes between birds and snakes.

Most studies now suggest that chicken has a genome organization most closely related to the ancestor [50–52]. If we consider this as the avian outgroup, *N. brasiliandum*, *M. maculatus* and *J. jacana*, having the same BAC order as chicken, provide evidence that there were no detectable intrachromosomal rearrangements but a repositioning of the centromere (Figure 2). For each of the other species, a combination of intrachromosomal rearrangements and centromere repositioning led to the configurations seen (Figure 2).

4.2. Reptilian Origins

Our results support a model in which the chicken Z chromosome shares ancestral chromosomal segments with both the cobra W chromosome and autosome 2. This reflects both deep homology and extensive post-divergence genomic reorganization. Such findings contribute to reconstructing the evolutionary history of sex chromosomes in amniotes, demonstrating that the avian Z chromosome is not evolutionarily isolated but retains signatures of ancient chromosomal structures shared with reptiles [3]. This aligns with recent comparative studies showing that amniote sex chromosomes have evolved through repeated turnovers, partial homologies and independent recruitment of autosomal segments into sex chromosome systems [53]. Integrating cytogenetic mapping with comparative genomics [54,55] therefore provides a powerful framework to trace the origin and transformation of sex chromosomes across vertebrate evolution [56–58]. Cross-referencing with published whole-genome alignments shows that the BAC signals on squamate W and chromosome 2 match conserved synteny regions previously identified as homologous to parts of the avian Z chromosome. O'Connor et al. [59] suggested that, at least in some turtles (which are phylogenetically closer to birds than snakes), the Z chromosome and chromosome 6 (designated as an autosome, as there is no evidence of a ZZ/ZW system) are homologous. Genome alignments of other turtles, however, indicate the absence of a

strict 1:1 relationship. These data imply that the avian Z chromosome, as currently defined, originated around the time of the bird–turtle divergence and, with few exceptions, has remained largely conserved since then [60]. The avian Z sequence mapped to the squamate W rather than the Z, likely due to lineage-specific rearrangements. Despite overall degeneration, the W chromosome can retain ancestral fragments, whereas the corresponding region on the squamate Z was probably lost or reorganized after divergence. The forces of evolution therefore seem to have favored the absence of interchromosomal rearrangement but not intrachromosomal rearrangement. Indeed, O’Connor et al. [59] suggested that the Z chromosome (and most others) was present intact in non-avian dinosaurs.

4.3. Intrachromosomal Rearrangement Predominates

Our findings further support the view that the avian Z chromosome evolves through a higher rate of intrachromosomal rearrangements than autosomes. This is reflected by differences in BAC order and/or centromere position in most species analyzed, relative to chicken and zebra finch. Moreover, the Z chromosomes show every major morphological class across the sample (metacentric, submetacentric, acrocentric, telocentric). This demonstrates that, while avian genomes are relatively conserved at the autosomal karyotypic level, Z chromosomes are prone to lineage-specific internal reorganization [35]. The conserved BAC order, yet different chromosome morphology, in *N. brasiliianum*, *M. maculatus* and *J. jacana* invokes a mechanism of centromere repositioning. This is increasingly recognized as an important mechanism in avian chromosome evolution, allowing changes in chromosomal morphology independent of large-scale rearrangements [39]. The remaining species, most likely, arose because of intrachromosomal rearrangements such as inversions, transpositions or localized rearrangements. Such rearrangements can accumulate over evolutionary time and have been frequently documented in the avian Z chromosome across multiple orders [61–63]. Inversions on Z chromosomes have been previously documented and shown to affect recombination and phenotypes in passerines and other groups. Centromere shifts can convert telocentric chromosomes into (sub)metacentric forms without large-scale exchange of genetic material [64]. These processes are also often associated with the accumulation or redistribution of repetitive elements and retrotransposons on sex chromosomes, which create substrates for non-allelic homologous recombination and breakpoint formation [64]. This compares to alignments of chromosome-level genome assemblies and BAC mapping in multiple Asian species that revealed inversions affecting macrochromosomes, including the Z [65]. In contrast, Neotropical chromosome-scale assemblies reveal lineage-specific rearrangements and centromere shifts in New World taxa, even though they belong to different orders. Together, these data indicate that the processes generating Z-morphological diversity are widespread geographically and not restricted to a single avian region or clade [65]. The unusually large Z chromosome in the Piciformes reflects lineage-specific expansion in chromosome size because of accumulation of repetitive DNA, which has already been reported in other groups as well [18]. Such changes may have functional implications for gene content, recombination landscapes, and even dosage compensation mechanisms [66].

4.4. Functionally and Evolutionarily, Intrachromosomal Rearrangements on the Z Chromosome Have Three Important Features

First, inversions and centromere shifts can modify recombination landscapes along the Z. This can alter linkage relationships among Z-linked genes, thus facilitating the joint inheritance of adaptive alleles or the buildup of sex-linked differentiation. This matches recent cytogenomic and population-genomic work showing that the avian Z undergoes faster structural change than many autosomes and often accumulates inversions and centromere shifts that change recombination landscapes and local gene order. Such intrachromosomal

events can explain why even closely related species differ in Z chromosome BAC order while conserving overall gene content. These rearrangements have been documented in multiple recent bird studies and are increasingly recognized as a common mechanism driving lineage-specific Z differentiation [67].

Second, changes in gene order/position can cause position effects or alter regulatory contexts for dosage-sensitive Z genes [68]. These may have downstream impacts on sexually dimorphic traits. Hybridization of avian Z BACs to reptile metaphases provides direct cytogenetic evidence that parts of the modern avian Z derive from ancestral autosomal linkage groups that are still present, in whole or in part, in reptile genomes. The localization of BAC clones on chromosome 2 in these squamate species aligns with broader comparative mapping studies, indicating that many amniote sex chromosomes stem from a restricted set of ancestral autosomes. This suggests that different lineages have repeatedly recruited and reused overlapping genomic blocks into sex chromosomes, producing partial homologies. The avian Z is thus a mosaic assembled from ancestral autosomal segments that were differentially retained, inverted, or translocated in birds but, importantly, were retained interchromosomally because, in part, of their role in sexual dimorphism [4]. Our data therefore fit a growing consensus that sex chromosomes evolve through repeated reuse and rearrangement of a limited set of ancestral genomic building blocks [66].

Third, because the Z is primarily transmitted through males, structural changes that reduce recombination or create incompatibilities may contribute to reproductive isolation and lineage divergence [63]. Recent syntheses emphasize that sex chromosome recruitment and turnover across amniotes is non-random. That is, certain linkage groups are predisposed to sex linkage, and centromere repositioning and repeat accumulation commonly accompany these transitions. Intrachromosomal rearrangements on the Z alter recombination and linkage relationships, which can facilitate the fixation of sexually beneficial alleles, influence dosage compensation trajectories and contribute to reproductive isolation.

Detecting Z-homologous segments in reptiles provides testable hypotheses regarding ancestral gene content and which Z-linked genes might have been important early in avian sex chromosome evolution. Future work that combines BAC mapping, chromosome-scale assemblies from many regions, and gene-level analyses in the pertinent reptile taxa will enable timing of specific rearrangements and assessment of their phenotypic and evolutionary impact.

5. Conclusions

This study demonstrates that intrachromosomal rearrangements of the Z chromosome are a significant factor in avian genome evolution. By linking chromosomal structure to phylogenetic divergence, our findings provide a framework for future research aimed at integrating cytogenomics with molecular and morphological data. This will help reconstruct the evolutionary tree of birds with higher resolution and greater accuracy. The comparative chromosomal analysis of the Z chromosome in South American bird species reveals a notably high rate of intrachromosomal rearrangements, indicating that the Z chromosome is more structurally dynamic than previously assumed. That is, while avian karyotypes are generally conserved, the observed rearrangements suggest that sex chromosomes, particularly the Z chromosome, are hotspots for genomic innovation and divergence. The observation that the Z chromosome exhibits all possible morphologies—metacentric, submetacentric, acrocentric and telocentric—across the studied species reinforces notions of its high structural plasticity. Comparative mapping of chicken Z chromosome BAC probes onto snake chromosomes demonstrates that the avian Z is partially conserved but extensively reorganized in the reptile–bird divergence. This study thus provides valuable cytogenetic evidence for reconstructing the evolutionary origins and transformations of

avian sex chromosomes in the context of broader amniote genome evolution. These structural changes may contribute to species-specific adaptations, influence the evolution of sex-linked traits, potentially affect reproductive isolation and ultimately contribute to the processes of speciation.

Author Contributions: Conceptualization, D.K.G. and R.K.; methodology, M.R.C.E., E.H.C.d.O., A.D.V.G., R.J.G. and M.S.d.S.; validation, M.R.C.E., D.K.G. and R.K.; formal analysis, M.R.C.E., M.N.R., D.K.G. and R.K.; investigation, M.R.C.E., E.H.C.d.O., A.D.V.G., R.J.G., M.S.d.S., R.E.O., D.K.G. and R.K.; resources, R.E.O. and D.K.G.; data curation, M.R.C.E. and R.K.; writing—original draft preparation, M.R.C.E., D.K.G. and R.K.; writing—review and editing, K.S., W.S., M.N.R., E.H.C.d.O., D.K.G., and R.K.; visualization, M.R.C.E. and R.K.; supervision, R.E.O., K.S., W.S., D.K.G. and R.K.; project administration, K.S., D.K.G. and R.K.; funding acquisition, K.S., D.K.G. and R.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Biotechnology and Biological Sciences Research Council (BB/K008226/1 and BB/K008161/1 to D.K.G.), M.R.C.E. was funded by a personal scholarship awarded by the Department of Science and Technology—Science Education Institute (DOST-SEI) Foreign Graduate Scholarship Programme, the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (Proc. 304068/2025-0 to R.K. and 307382/2019-2 to E.H.C.d.O.) grant from Kasetsart University Research and Development Institute (FF(S-KU)17.66, FF(KU)51.67, and FF(KU)61.69) awarded to W.S. and K.S.; the Visiting Research Scholar (VRC) Grant, Faculty of Science, Kasetsart University (Contract Number VRC 4/2024), awarded to D.K.G. and K.S., support from the International SciKU Branding (ISB), Faculty of Science, Kasetsart University, awarded to W.S. and K.S., the Office of the Ministry of Higher Education, Science, Research and Innovation, and the Thailand Science Research and Innovation (TSRI) through the Kasetsart University Reinventing University Program 2025 (No. RUP_Climate68_No.9) awarded to W.S., M.N.R. and D.K.G. The APC was funded as a result of Prof. Griffin's allocation as an Editor of *Genes* as an invited contribution.

Institutional Review Board Statement: The study using birds was conducted according to the guidelines of the Sistema de Autorização e Informação em Biodiversidade [IBAMA/SISBIO 61047-1 (10 October 2018), 33860-1 (8 June 2012), 68443-1 (1 April 2019) and 44173-1 (7 July 2014)] and approved by the Ethics Committee on the use of animals of Universidade Federal do Pampa [CEUA no. 026/2012 (no approval date issued), 018 (1 July 2014) and 019 (30 November 2020)], Universidade Federal do Pará [no. 170/2013 (no approval date issued)], and Universidade Federal do Rio Grande do Sul [CEUA number 30750 (10 October 2016)]. All reptile procedures were approved by the Animal Experiment Committee of Kasetsart University, Thailand [approval nos. ACKU59-SCI-034 (22 September 2016) and ACKU61-SCI-024 (28 September 2018)].

Informed Consent Statement: Not applicable.

Data Availability Statement: All raw data is available by contacting the authors.

Acknowledgments: During the preparation of this manuscript/study, the author(s) made occasional use of ChatGPT-4 for the purposes of initial drafting and language stylistic suggestions. The authors have reviewed and edited the output and take full responsibility for the content of this publication.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Abbreviations

The following abbreviations are used in this manuscript:

FISH	Fluorescence in situ hybridization
BAC	Bacterial artificial chromosome
KCl	Potassium chloride
DAPI	4,6-diamidino-2-phenylindole

References

1. Griffin, D.K.; Kretschmer, R.; Larkin, D.M.; Srikulnath, K.; Singchat, W.; Narushin, V.G.; O'Connor, R.E.; Romanov, M.N. Avian cytogenomics: Small chromosomes, long evolutionary history. *Genes* **2025**, *16*, 1001. [[CrossRef](#)]
2. O'Connor, R.E.; Kretschmer, R.; Romanov, M.N.; Griffin, D.K. A bird's-eye view of chromosomal evolution in the Class Aves. *Cells* **2024**, *13*, 310. [[CrossRef](#)]
3. Griffin, D.K.; Larkin, D.M.; O'Connor, R.E.; Romanov, M.N. Dinosaurs: Comparative cytogenomics of their reptile cousins and avian descendants. *Animals* **2023**, *13*, 106. [[CrossRef](#)] [[PubMed](#)]
4. Griffin, D.K.; Kretschmer, R.; Srikulnath, K.; Singchat, W.; O'Connor, R.E.; Romanov, M.N. Insights into avian molecular cytogenetics—With reptilian comparisons. *Mol. Cytogenet.* **2024**, *17*, 24. [[CrossRef](#)] [[PubMed](#)]
5. Nanda, I.; Karl, E.; Volobouev, V.; Griffin, D.K.; Schartl, M.; Schmid, M. Extensive gross genomic rearrangements between chicken and Old World vultures (Falconiformes: Accipitridae). *Cytogenet. Genome Res.* **2006**, *112*, 286–295. [[CrossRef](#)] [[PubMed](#)]
6. Kretschmer, R.; Furo, I.D.O.; Cioffi, M.D.B.; Gunski, R.J.; Garnero, A.D.V.; O'Brien, P.C.; Ferguson-Smith, M.A.; De Freitas, T.R.O.; De Oliveira, E.H.C. Extensive chromosomal fissions and repetitive DNA accumulation shaped the atypical karyotypes of two Ramphastidae (Aves: Piciformes) species. *Biol. J. Linn. Soc.* **2020**, *130*, 839–849. [[CrossRef](#)]
7. Kretschmer, R.; de Souza, M.S.; Furo, I.D.O.; Romanov, M.N.; Gunski, R.J.; Garnero, A.D.V.; de Freitas, T.R.O.; de Oliveira, E.H.C.; O'Connor, R.E.; Griffin, D.K. Interspecies chromosome mapping in Caprimulgiformes, Piciformes, Suliformes, and Trogoniformes (Aves): Cytogenomic insight into microchromosome organization and karyotype evolution in birds. *Cells* **2021**, *10*, 826. [[CrossRef](#)]
8. Kiazim, L.G.; O'Connor, R.E.; Larkin, D.M.; Romanov, M.N.; Narushin, V.G.; Brazhnik, E.A.; Griffin, D.K. Comparative mapping of the macrochromosomes of eight avian species provides further insight into their phylogenetic relationships and avian karyotype evolution. *Cells* **2021**, *10*, 362. [[CrossRef](#)]
9. Modi, W.S.; Romanov, M.; Green, E.D.; Ryder, O. Molecular cytogenetics of the California condor: Evolutionary and conservation implications. *Cytogenet. Genome Res.* **2009**, *127*, 26–32. [[CrossRef](#)]
10. Alfieri, J.M.; Bolwerk, K.; Hu, Z.; Blackmon, H. From micro to macro: Avian chromosome evolution is dominated by natural selection. *bioRxiv* **2024**. [[CrossRef](#)]
11. Wu, S.; Dou, T.; Yuan, S.; Yan, S.; Xu, Z.; Liu, Y.; Jian, Z.; Zhao, J.; Zhao, R.; Zi, X.; et al. Annotations of four high-quality indigenous chicken genomes identify more than one thousand missing genes in subtelomeric regions and micro-chromosomes with high G/C contents. *BMC Genom.* **2024**, *25*, 430. [[CrossRef](#)]
12. Burssed, B.; Zamariolli, M.; Bellucco, F.T.; Melaragno, M.I. Mechanisms of structural chromosomal rearrangement formation. *Mol. Cytogenet.* **2022**, *15*, 23. [[CrossRef](#)] [[PubMed](#)]
13. Stiller, J.; Feng, S.; Chowdhury, A.A.; Rivas-González, I.; Duchêne, D.A.; Fang, Q.; Deng, Y.; Kozlov, A.; Stamatakis, A.; Claramunt, S.; et al. Complexity of avian evolution revealed by family-level genomes. *Nature* **2024**, *629*, 851–860. [[CrossRef](#)] [[PubMed](#)]
14. Griffin, D.K.; Robertson, L.B.W.; Tempest, H.G.; Skinner, B.M. The evolution of the avian genome as revealed by comparative molecular cytogenetics. *Cytogenet. Genome Res.* **2007**, *117*, 64–77. [[CrossRef](#)] [[PubMed](#)]
15. Lithgow, P.E.; O'Connor, R.; Smith, D.; Fonseka, G.; Rathje, C.; Frodsham, R.; O'Brien, P.C.; Ferguson-Smith, M.A.; Skinner, B.M.; Griffin, D.K.; et al. Novel Tools for Characterising Inter- and Intra-chromosomal Rearrangements in Avian Microchromosomes. In *Proceedings of the 2014 Meeting on Avian Model Systems, Cold Spring Harbor, NY, USA, 5–8 March 2014*; Cold Spring Harbor Laboratory: Cold Spring Harbor, NY, USA, 2014; p. 56. Available online: <https://kar.kent.ac.uk/46692/> (accessed on 31 October 2025).
16. Romanov, M.N.; Dodgson, J.B.; Gonser, R.A.; Tuttle, E.M. Comparative BAC-based mapping in the white-throated sparrow, a novel behavioral genomics model, using interspecies overgo hybridization. *BMC Res. Notes* **2011**, *4*, 211. [[CrossRef](#)]
17. Berdan, E.L.; Barton, N.H.; Butlin, R.; Charlesworth, B.; Faria, R.; Fragata, I.; Gilbert, K.J.; Jay, P.; Kapun, M.; Lotterhos, K.E.; et al. How chromosomal inversions reorient the evolutionary process. *J. Evol. Biol.* **2023**, *36*, 1761–1782. [[CrossRef](#)]
18. Huang, Z.; De, O.; Furo, I.; Liu, J.; Peona, V.; Gomes, A.J.; Cen, W.; Huang, H.; Zhang, Y.; Chen, D.; et al. Recurrent chromosome reshuffling and the evolution of neo-sex chromosomes in parrots. *Nat. Commun.* **2022**, *13*, 944. [[CrossRef](#)]
19. de Oliveira Furo, I.; Kretschmer, R.; Dos Santos, M.S.; de Lima Carvalho, C.A.; Gunski, R.J.; O'Brien, P.; Ferguson-Smith, M.A.; Cioffi, M.B.; de Oliveira, E.H. Chromosomal mapping of repetitive DNAs in *Myiopsitta monachus* and *Amazona aestiva* (Psittaciformes, Psittacidae) with emphasis on the sex chromosomes. *Cytogenet. Genome Res.* **2017**, *151*, 151–160. [[CrossRef](#)]
20. Gunski, R.J.; Kretschmer, R.; Santos de Souza, M.; de Oliveira Furo, I.; Barcellos, S.A.; Costa, A.L.; Cioffi, M.B.; de Oliveira, E.H.; del Valle Garnero, A. Evolution of bird sex chromosomes narrated by repetitive sequences: Unusual W chromosome enlargement in *Gallinula melanops* (Aves: Gruiiformes: Rallidae). *Cytogenet. Genome Res.* **2019**, *158*, 152–159. [[CrossRef](#)]
21. Blagoveshchenski, I.I.; Sazanova, A.L.; Stekol'nikova, V.A.; Fomichev, K.A.; Barkova, O.I.; Romanov, M.N.; Sazanov, A.A. Investigation of pseudoautosomal and bordering regions in avian Z and W chromosomes with the use of large insert genomic BAC clones. *Genetika* **2011**, *47*, 312–319.
22. de Oliveira Furo, I.; Kretschmer, R.; O'Brien, P.C.; Ferguson-Smith, M.A.; de Oliveira, E.H.C. Chromosomal diversity and karyotype evolution in South American macaws (Psittaciformes, Psittacidae). *PLoS ONE* **2015**, *10*, e0130157. [[CrossRef](#)]

23. Kretschmer, R.; de Oliveira, T.D.; de Oliveira Furo, I.; Oliveira Silva, F.A.; Gunski, R.J.; del Valle Garnero, A.; de Bello Cioffi, M.; de Oliveira, E.H.C.; de Freitas, T.R.O. Repetitive DNAs and shrink genomes: A chromosomal analysis in nine Columbidae species (Aves, Columbiformes). *Genet. Mol. Biol.* **2018**, *41*, 98–106. [CrossRef] [PubMed]

24. Nanda, I.; Schmid, M. Conservation of avian Z chromosomes as revealed by comparative mapping of the Z-linked aldolase B gene. *Cytogenet. Cell Genet.* **2002**, *96*, 176–178. [CrossRef] [PubMed]

25. Sazanov, A.A.; Romanov, M.N.; Wardecka, B.; Sazanova, A.L.; Korczak, M.; Stekol'nikova, V.A.; Kozyreva, A.A.; Smirnov, A.F.; Jaszczak, K.; Dodgson, J.B. Chromosomal localization of 15 large insert BAC clones containing three microsatellites on chicken chromosome 4 (GGA4) which refine its centromere position. *Anim. Genet.* **2005**, *36*, 161–163. [CrossRef] [PubMed]

26. Gunski, R.J.; Cañedo, A.D.; Garnero, A.D.V.; Ledesma, M.A.; Coria, N.; Montalti, D.; Degrandi, T.M. Multiple sex chromosome system in penguins (*Pygoscelis*, *Spheniscidae*). *Comp. Cytogenet.* **2017**, *11*, 541. [CrossRef]

27. Pozzobon, L.C.; Toma, G.A.; Cioffi, M.D.B.; de Oliveira, E.H.C.; Kretschmer, R.; de Freitas, T.R.O. Karyotype evolution of suliformes and description of a ♂Z₁Z₁Z₂Z₂/♀Z₁Z₂W multiple sex chromosome system in boobies (*Sula* spp.). *Genome* **2025**, *68*, 1–11. [CrossRef]

28. Sigeman, H.; Ponnikas, S.; Hansson, B. Whole-genome analysis across 10 songbird families within *Sylvioidea* reveals a novel autosome–sex chromosome fusion. *Biol. Lett.* **2020**, *16*, 20200082. [CrossRef]

29. Sigeman, H.; Strandh, M.; Proux-Wéra, E.; Kutschera, V.E.; Ponnikas, S.; Zhang, H.; Lundberg, M.; Soler, L.; Bunkis, I.; Tarka, M.; et al. Avian neo-sex chromosomes reveal dynamics of recombination suppression and W degeneration. *Mol. Biol. Evol.* **2021**, *38*, 5275–5291. [CrossRef]

30. Sigeman, H.; Zhang, H.; Ali Abed, S.; Hansson, B. A novel neo-sex chromosome in *Sylvietta brachyura* (Macrophenidae) adds to the extraordinary avian sex chromosome diversity among *Sylvioidea* songbirds. *J. Evol. Biol.* **2022**, *35*, 1797–1805. [CrossRef]

31. Sigeman, H.; Downing, P.A.; Zhang, H.; Hansson, B. The rate of W chromosome degeneration across multiple avian neo-sex chromosomes. *Sci. Rep.* **2024**, *14*, 16548. [CrossRef]

32. Xu, L.; Ren, Y.; Wu, J.; Cui, T.; Dong, R.; Huang, C.; Feng, Z.; Zhang, T.; Yang, P.; Yuan, J.; et al. Evolution and expression patterns of the neo-sex chromosomes of the crested ibis. *Nat. Commun.* **2024**, *15*, 1670. [CrossRef] [PubMed]

33. Shogren, E.H.; Sardell, J.M.; Muirhead, C.A.; Martí, E.; Cooper, E.A.; Moyle, R.G.; Presgraves, D.C.; Uy, J.A.C. Recent secondary contact, genome-wide admixture, and asymmetric introgression of neo-sex chromosomes between two Pacific island bird species. *PLoS Genet.* **2024**, *20*, e1011360. [CrossRef] [PubMed]

34. Hansson, B.; Sigeman, H.; Ellerstrand, S. Evolutionary dynamics of enlarged sex chromosomes and novel pseudoautosomal regions in *Sylvioidea* songbirds. *Res. Sq.* **2025**. [CrossRef]

35. Kretschmer, R.; Gunski, R.J.; Garnero, A.D.V.; de Freitas, T.R.O.; Toma, G.A.; Cioffi, M.D.B.; Oliveira, E.H.C.d.; O'Connor, R.E.; Griffin, D.K. Chromosomal analysis in *Crotophaga ani* (Aves, Cuculiformes) reveals extensive genomic reorganization and an unusual Z-autosome Robertsonian translocation. *Cells* **2021**, *10*, 4. [CrossRef] [PubMed]

36. Kretschmer, R.; Santos de Souza, M.; Gunski, R.J.; del Valle Garnero, A.; de Freitas, T.R.O.; Zefa, E.; Toma, G.A.; Cioffi, M.D.B.; Herculano Corrêa de Oliveira, E.; O'Connor, R.E.; et al. Understanding the chromosomal evolution in cuckoos (Aves, Cuculiformes): A journey through unusual rearrangements. *Genome* **2024**, *67*, 168–177. [CrossRef]

37. Muirhead, C.A.; Martí, E.; Shogren, E.H.; Uy, J.A.C.; Presgraves, D.C. Genomic origins and evolution of neo-sex chromosomes in Pacific Island birds. *Proc. Natl. Acad. Sci. USA* **2025**, *122*, e2503746122. [CrossRef]

38. Duchêne, D.A.; Chowdhury, A.A.; Yang, J.; Iglesias-Carrasco, M.; Stiller, J.; Feng, S.; Bhatt, S.; Gilbert, M.T.P.; Zhang, G.; Tobias, J.A.; et al. Drivers of avian genomic change revealed by evolutionary rate decomposition. *Nature* **2025**, *641*, 1208–1216. [CrossRef]

39. Damas, J.; Kim, J.; Farré, M.; Griffin, D.K.; Larkin, D.M. Reconstruction of avian ancestral karyotypes reveals differences in the evolutionary history of macro- and microchromosomes. *Genome Biol.* **2018**, *19*, 155. [CrossRef]

40. Wu, M.Y.; Lau, C.J.; Ng, E.Y.; Baveja, P.; Gwee, C.Y.; Sadanandan, K.; Ferasyi, T.R.; Haminuddin Ramadhan, R.; Menner, J.K.; Rheindt, F.E. Genomes from historic DNA unveil massive hidden extinction and terminal endangerment in a tropical Asian songbird radiation. *Mol. Biol. Evol.* **2022**, *39*, msac189. [CrossRef]

41. de Souza, M.S.; Barcellos, S.A.; Tura, V.; Bobrowski, V.L.; Garnero, A.D.V.; Gunski, R.J.; Griffin, D.K.; Kretschmer, R. Highly conserved microchromosomal organization in Passeriformes birds revealed via BAC-FISH analysis. *Birds* **2023**, *4*, 236–244. [CrossRef]

42. Furo, I.D.O.; Kretschmer, R.; O'Brien, P.C.; Pereira, J.C.; Garnero, A.D.V.; Gunski, R.J.; O'Connor, R.E.; Griffin, D.K.; Gomes, A.J.B.; Ferguson-Smith, M.A.; et al. Chromosomal evolution in the phylogenetic context: A remarkable karyotype reorganization in neotropical parrot *Myiopsitta monachus* (Psittacidae). *Front. Genet.* **2020**, *11*, 721. [CrossRef]

43. Alves Barcellos, S.; Kretschmer, R.; Santos de Souza, M.; Tura, V.; Pozzobon, L.C.; Ochotorena de Freitas, T.R.; Griffin, D.K.; O'Connor, R.; Gunski, R.J.; del Valle Garnero, A. Understanding microchromosomal organization and evolution in four representative woodpeckers (Picidae, Piciformes) through BAC-FISH analysis. *Genome* **2024**, *67*, 223–232. [CrossRef] [PubMed]

44. Degrandi, T.M.; del Valle Garnero, A.; O'Brien, P.; Ferguson-Smith, M.A.; Kretschmer, R.; de Oliveira, E.H.; Gunski, R.J. Chromosome painting in *Trogon s. surrucura* (Aves, Trogoniformes) reveals a karyotype derived by chromosomal fissions, fusions, and inversions. *Cytogenet. Genome Res.* **2017**, *151*, 208–215. [CrossRef] [PubMed]

45. Kretschmer, R.; Souza, M.S.; Barcellos, S.A.; Degrandi, T.M.; Pereira, J.C.; O'Brien, P.C.M.; Ferguson-Smith, M.A.; Gunski, R.J.; Garnero, A.D.V.; Oliveira, E.H.C.; et al. Novel insights into chromosome evolution of Charadriiformes: Extensive genomic reshuffling in the wattled jacana (*Jacana jacana*, Charadriiformes, Jacanidae). *Genet. Mol. Biol.* **2020**, *43*, e20190236. [CrossRef] [PubMed]

46. Srikulnath, K.; Matsubara, K.; Uno, Y.; Nishida, C.; Olsson, M.; Matsuda, Y. Identification of the linkage group of the Z sex chromosomes of the sand lizard (*Lacerta agilis*, Lacertidae) and elucidation of karyotype evolution in lacertid lizards. *Chromosoma* **2014**, *123*, 563–575. [CrossRef]

47. Singchat, W.; Ahmad, S.F.; Sillapaprayoon, S.; Muangmai, N.; Duengkao, P.; Peyachoknagul, S.; O'Connor, R.E.; Griffin, D.K.; Srikulnath, K. Partial Amniote Sex Chromosomal Linkage Homologies Shared on Snake W Sex Chromosomes Support the Ancestral Super-Sex Chromosome Evolution in Amniotes. *Front. Genet.* **2020**, *11*, 948. [CrossRef]

48. Singchat, W.; O'Connor, R.E.; Tawichasri, P.; Suntronpong, A.; Sillapaprayoon, S.; Suntrarachun, S.; Muangmai, N.; Baicharoen, S.; Peyachoknagul, S.; Chanhome, L.; et al. Chromosome map of the Siamese cobra: Did partial synteny of sex chromosomes in the amniote represent “a hypothetical ancestral super-sex chromosome” or random distribution? *BMC Genom.* **2018**, *19*, 939. [CrossRef]

49. O'Connor, R.E.; Kiazim, L.; Skinner, B.; Fonseka, G.; Joseph, S.; Jennings, R.; Larkin, D.M.; Griffin, D.K. Patterns of microchromosome organization remain highly conserved throughout avian evolution. *Chromosoma* **2019**, *128*, 21–29. [CrossRef]

50. Romanov, M.N.; Farré-Belmonte, M.; Lithgow, P.E.; O'Connor, R.; Fowler, K.E.; Larkin, D.M.; Griffin, D.K. In silico Reconstruction of Chromosomal Rearrangements and an Avian Ancestral Karyotype. In *Proceedings of the International Plant and Animal Genome XXII Conference, San Diego, CA, USA, 10–14 January 2014*; Scherago International: San Diego, CA, USA, 2014. Abstract P1106. Available online: <https://kar.kent.ac.uk/37651/> (accessed on 31 October 2025).

51. Romanov, M.N.; Farré, M.; Lithgow, P.E.; O'Connor, R.; Fowler, K.E.; Skinner, B.M.; Larkin, D.M.; Griffin, D.K. Avian ancestral karyotype reconstruction and differential rates of inter- and intrachromosomal change in different lineages. *Chromosome Res.* **2015**, *23*, 414. [CrossRef]

52. Degrandi, T.M.; Barcellos, S.A.; Costa, A.L.; Garnero, A.D.; Hass, I.; Gunski, R.J. Introducing the bird chromosome database: An overview of cytogenetic studies in birds. *Cytogenet. Genome Res.* **2020**, *160*, 199–205. [CrossRef]

53. Kratochvíl, L.; Gamble, T.; Rovatsos, M. Sex chromosome evolution among amniotes: Is the origin of sex chromosomes non-random? *Philos. Trans. R. Soc. B Biol. Sci.* **2021**, *376*, 20200108. [CrossRef] [PubMed]

54. Romanov, M.N.; Dodgson, J.B. Cross-species overgo hybridization and comparative physical mapping within avian genomes. *Anim. Genet.* **2006**, *37*, 397–399. [CrossRef] [PubMed]

55. Romanov, M.N.; Daniels, L.M.; Dodgson, J.B.; Delany, M.E. Integration of the cytogenetic and physical maps of chicken chromosome 17. *Chromosome Res.* **2005**, *13*, 215–222. [CrossRef] [PubMed]

56. Augustijnen, H.; Arias-Sardá, C.; Farré, M.; Lucek, K. A genomic update on the evolutionary impact of chromosomal rearrangements. *Mol. Ecol.* **2024**, *33*, e17602. [CrossRef]

57. Bellott, D.W.; Skaletsky, H.; Pyntikova, T.; Mardis, E.R.; Graves, T.; Kremitzki, C.; Brown, L.G.; Rozen, S.; Warren, W.C.; Wilson, R.K.; et al. Convergent evolution of chicken Z and human X chromosomes by expansion and gene acquisition. *Nature* **2010**, *466*, 612–616. [CrossRef]

58. Vallender, E.J.; Lahn, B.T. Multiple independent origins of sex chromosomes in amniotes. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 18031–18032. [CrossRef]

59. O'Connor, R.E.; Romanov, M.N.; Kiazim, L.G.; Barrett, P.M.; Farré, M.; Damas, J.; Ferguson-Smith, M.; Valenzuela, N.; Larkin, D.M.; Griffin, D.K. Reconstruction of the diapsid ancestral genome permits chromosome evolution tracing in avian and non-avian dinosaurs. *Nat. Commun.* **2018**, *9*, 1883. [CrossRef]

60. Bista, B.; Valenzuela, N. Turtle insights into the evolution of the reptilian karyotype and the genomic architecture of sex determination. *Genes* **2020**, *11*, 416. [CrossRef]

61. Zhou, Q.; Zhang, J.; Bachtrog, D.; An, N.; Huang, Q.; Jarvis, E.D.; Gilbert, M.T.P.; Zhang, G. Complex evolutionary trajectories of sex chromosomes across bird taxa. *Science* **2014**, *346*, 1246338. [CrossRef]

62. Xu, L.; Auer, G.; Peona, V.; Suh, A.; Deng, Y.; Feng, S.; Zhang, G.; Blom, M.P.; Christidis, L.; Prost, S.; et al. Dynamic evolutionary history and gene content of sex chromosomes across diverse songbirds. *Nat. Ecol. Evol.* **2019**, *3*, 834–844. [CrossRef]

63. Knief, U.; Müller, I.A.; Stryjewski, K.F.; Metzler, D.; Sorenson, M.D.; Wolf, J.B. Evolution of chromosomal inversions across an avian radiation. *Mol. Biol. Evol.* **2024**, *41*, msae092. [CrossRef]

64. Viitaniemi, H.M.; Leder, E.H.; Kauzál, O.; Stopková, R.; Stopka, P.; Lifjeld, J.T.; Albrecht, T. Impact of Z chromosome inversions on gene expression in testis and liver tissues in the zebra finch. *Mol. Ecol.* **2024**, *33*, e17236.

65. Chan, Y.F.; Lu, C.W.; Kuo, H.C.; Hung, C.M. A chromosome-level genome assembly of the Asian house martin implies potential genes associated with the feathered-foot trait. *G3* **2024**, *14*, jkae077. [[CrossRef](#)]
66. Deviatiiarov, R.; Nagai, H.; Ismagulov, G.; Stupina, A.; Wada, K.; Ide, S.; Toji, N.; Zhang, H.; Sukparangsi, W.; Intarapat, S.; et al. Dosage compensation of Z sex chromosome genes in avian fibroblast cells. *Genome Biol.* **2023**, *24*, 213. [[CrossRef](#)] [[PubMed](#)]
67. López Villavicencio, M.; Ledamoisel, J.; Poloni, R.; Lopez-Roques, C.; Debat, V.; Llaurens, V. Increased evolutionary rate in the Z chromosome of sympatric and allopatric species of Morpho butterflies. *Genome Biol. Evol.* **2024**, *16*, evae227. [[CrossRef](#)]
68. Ioannidis, J.; Taylor, G.; Zhao, D.; Liu, L.; Idoko-Akoh, A.; Gong, D.; Lovell-Badge, R.; Guioli, S.; McGrew, M.J.; Clinton, M. Primary sex determination in birds depends on DMRT1 dosage, but gonadal sex does not determine adult secondary sex characteristics. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2020909118. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.