



Kent Academic Repository

Plume, René, Eden, David J., Currie, Malcolm J., Morgan, Lawrence K., Jiang, Xue-Jian, DiFrancesco, James, Imanishi, Masatoshi, Kim, Kee-Tae, Liu, Tie, Rani, Raffaele and others (2026) *MAJORS II: HCO+ & HCN Abundances in W40*. *Monthly Notices of the Royal Astronomical Society*, 543 (7). ISSN 0035-8711.

Downloaded from

<https://kar.kent.ac.uk/112922/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1093/mnras/stag186>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

For the purpose of open access, the author(s) has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in ***Title of Journal***, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

MAJORS II: HCO⁺ and HCN abundances in W40

René Plume¹,^{1*} David J. Eden², Malcolm J. Currie,³ Lawrence K. Morgan,⁴ Xue-Jian Jiang,⁵ James DiFrancesco,^{6,7} Masatoshi Imanishi^{8,9}, Kee-Tae Kim^{10,11}, Tie Liu,¹² Raffaele Rani¹³, Alessio Traficante¹⁴, Jane Cohen,¹ Neal J. Evans II,¹⁵ Luis C. Ho,^{16,17} Eun Jung Chung,¹⁸ Sihan Jiao,¹⁹ Chang Won Lee,^{10,11} Dana Alina,²⁰ Toby Moore,²¹ Jonathan M. C. Rawlings,²² Florian Kirchschrager,²³ Sudeshna Patra,²⁴ Andrew J. Rigby,²⁵ Hsien Shang,²⁶ Jihye Hwang,²⁷ Patricio Sanhueza,²⁸ Mark G. Rawlings,²⁹ Kianoosh Tahani,³⁰ Junfeng Wang,³¹ Kate Pattle,²² James S. Urquhart,³² Quang Nguyen-Luong,³³ Sarah E. Ragan,³⁴ Yang Su,³⁵ Xindi Tang,³⁶ Agata Karska^{37,24} and Michael G. Burton³⁸

Affiliations are listed at the end of the paper

Accepted 2026 January 23. Received 2026 January 22; in original form 2025 June 19

ABSTRACT

We present observations of HCN and HCO⁺ J = 3 → 2 in the central 424 arcsec × 424 arcsec region of the W40 massive star-forming region. The observations were taken as part of a pilot project for the MAJORS large program at the JCMT telescope. By incorporating prior knowledge of N(H₂) and T_K, assuming a constant density, and using the RADEX radiative transfer code we found that the HCN and HCO⁺ abundances range from X(HCN) = 0.4–7.0 × 10^{−8} and X(HCO⁺) = 0.4–7.3 × 10^{−9}. Additional modelling using the NAUTILUS chemical evolution code, which takes H₂ density variations into account, however, suggests that the HCN and HCO⁺ abundances may be fairly constant. Careful modelling of three different positions finds X(HCN) = 1.3–1.7 × 10^{−8} and X(HCO⁺) = 1.3–3.1 × 10^{−9}. Cross-comparison of the two models also provides a crude estimate of the gas density producing the HCN and HCO⁺ emission, with H₂ densities in the range 5 × 10⁴–5 × 10⁵ cm^{−3}, suggesting that the HCN and HCO⁺ emission does indeed arise from dense gas. High UV intensity (e.g. G₀ > a few thousand) has no effect on the abundances in regions where the visual extinction is large enough to effectively shield the gas from the UV field. In regions where A_V < 6, however, the abundance of both species is lowered due to destructive reactions with species that are directly affected by the radiation field.

Key words: surveys – stars: formation – ISM: abundances – ISM: clouds – ISM: molecules.

1 INTRODUCTION

Stars form in the densest parts of molecular clouds (C. J. Lada, M. Lombardi & J. F. Alves 2010) and, in high-mass star-forming regions, the densities can reach 10⁶ cm^{−3} or higher (R. Plume et al. 1997). Typical molecular tracers like CO and ¹³CO J = 1 → 0 measure the total H₂ gas mass in molecular clouds reasonably well. However, since the low J transitions of CO can be easily excited in gas with densities < 100 cm^{−3}, they primarily trace the low-density ambient molecular gas in giant molecular clouds, not specifically the gas that will contribute to forming stars. In addition, CO is known to freeze out on to dust grains in high-density low-temperature environments (e.g. H. S. Thomas & G. A. Fuller 2008; F. Fontani et al. 2012; G. Sabatini et al. 2019) and may therefore be an unreliable tracer of the gas under such conditions.

Higher critical density molecular line tracers, such as HCN, probe the behaviour of the dense gas most closely associated with star formation very well (A. Onus, M. R. Krumholz & C.

Federrath 2018). A survey of spiral galaxies and ultra luminous infrared galaxies (ULIRGS) in HCN J = 1 → 0 revealed a tight relationship between the infrared (IR) luminosity (L_{IR}), a tracer of the star formation rate (SFR), and the HCN luminosity (L_{HCN}), a tracer of total dense molecular gas content (Y. Gao & P. M. Solomon 2004a, b; I. W. Stephens et al. 2016). When these results are extended down to nearby star-forming regions, i.e. Galactic clumps, the linear relationship survives, covering a luminosity range many orders of magnitude wide and scales from parsec-sized clumps to entire ULIRGS (J. Wu et al. 2005; I. Mendigutía, C. J. Lada & R. D. Oudmaijer 2018). There are, however, some notable exceptions. For example, J. Kauffmann et al. (2017) and M. G. Santa-Maria et al. (2023) showed that HCN J = 1 → 0 traces densities of < 10⁴ cm^{−3} in cold regions of Orion A and Orion B, respectively. Furthermore, N. J. Evans et al. (2020) demonstrated that this transition can also arise in diffuse gas (n < 100 cm^{−3}) provided the column density is sufficiently high (A_V > 8).

While simulations have shown that the L_{IR}–L_{HCN} relation is a result of the free-fall time-scale of the gas and the average gas density of molecular clouds (M. R. Krumholz, A. Dekel &

* E-mail: rplume@ucalgary.ca

C. F. McKee 2012), both Galactic and extragalactic observations have pointed towards the presence of a volume-density threshold ($> 10^4 \text{ cm}^{-3}$), with the amount of gas above this density being the important factor in regulating the SFR (C. J. Lada et al. 2012; I. Evans, A. Heiderman & N. Vutisalchavakul 2014; Z.-Y. Zhang et al. 2014; A. Heiderman et al. 2010). It is therefore clear that dense gas is the important fuel for star formation. Understanding its exact role, and how the properties of molecular clouds impact the process, however, has been elusive.

The Massive, Active, JCMT-Observed Regions of Star formation (MAJORS) survey is a large program at the 15m James Clerk Maxwell Telescope (JCMT) on Maunakea, Hawaii. MAJORS was awarded 976 h of band four weather conditions (i.e. 4–6 mm of precipitable water vapour) to simultaneously map $\text{HCO}^+ \text{ J} = 3 \rightarrow 2$ ($\nu = 267.5576259 \text{ GHz}$) and $\text{HCN J} = 3 \rightarrow 2$ ($\nu = 265.8864339 \text{ GHz}$) in a large mass-selected sample of dust-continuum traced molecular clouds in the Milky Way. The sample includes clouds in the Central Molecular Zone (CMZ), the Inner Galaxy, and the Outer Galaxy. While full details of the MAJORS survey will be presented in Eden et al. (in preparation), its primary goals are to:

- (i) understand how Galactic environment influences the physics of dense gas, enabling us to determine how dense gas is produced and how it is intrinsically linked to star formation;
- (ii) produce $L_{\text{IR}}-L_{\text{gas}}$ relationships for dense gas in Galactic clumps and molecular clouds, allowing us to connect these results to studies of extragalactic systems and ULIRGs and to assess the universality of the star-formation process;
- (iii) determine the origin of variations in the HCN and HCO^+ abundances and HCN/ HCO^+ abundance ratio, and how these variations relate to the physical conditions imposed by different Galactic environments.

HCN and $\text{HCO}^+ \text{ J} = 3 \rightarrow 2$ are excellent tracers of dense gas. In the optically thin limit, the critical densities of HCO^+ and $\text{HCN J} = 3 \rightarrow 2$ are 1.4×10^6 and $1.0 \times 10^7 \text{ cm}^{-3}$, respectively (Y. L. Shirley 2015). However, since these lines are often optically thick, radiative trapping effectively reduces their critical densities to 6.8×10^3 and $7.3 \times 10^4 \text{ cm}^{-3}$, respectively (Y. L. Shirley 2015) in which case these lines may not be tracing the innermost densest regions of the clouds. In either case, however, these transitions can probe gas with densities associated with star formation activity rather than the low-density bulk envelopes of molecular clouds.

Given the high critical densities of these transitions, many previous studies have simply correlated total HCN or HCO^+ luminosity with total IR luminosity to measure the SFR in dense gas (see e.g. I. W. Stephens et al. (2016) and references therein). However this interpretation is overly simplistic. In order to be certain that a particular transition of these molecules is actually tracing gas above the 10^4 cm^{-3} threshold, one must know their abundances; given by the quantity X , where X is calculated by dividing the molecular column density by the column density of H_2 (e.g. $X(\text{HCN}) = N(\text{HCN})/N(\text{H}_2)$ and $X(\text{HCO}^+) = N(\text{HCO}^+)/N(\text{H}_2)$).

The abundances of HCN and HCO^+ (and other molecular species) are not constant, however, and can vary from location to location. Table 1 lists HCN and HCO^+ abundances (and their ratios) for a range of Galactic and extragalactic sources observed at different resolutions. These values span nearly two orders of magnitude between sources. Some of this variation may reflect differences in beam size or in the methods used to derive molecular abundances; however, Table 1 clearly

demonstrates that intrinsic variations exist. In addition, beyond observational and methodological effects, molecular abundances are fundamentally shaped by environmental factors such as gas and dust temperature, ultraviolet radiation fields, opacity, and cosmic-ray ionization rate. To fully understand the connection between dense gas, L_{IR} , and star formation in the Galaxy, it is therefore essential to measure HCN and HCO^+ abundances (and associated gas densities) across a large diverse sample of sources spanning many Galactic environments, using a consistent methodology to minimize systematic uncertainties.

1.1 W40

As part of a pilot project for MAJORS, we mapped HCO^+ and $\text{HCN J} = 3 \rightarrow 2$ in the central region of the well-known massive star-forming region W40. Recent astrometry observations from *Gaia* (F. Comerón, A. A. Djupvik & N. Schneider 2022) place W40 at a distance of $502 \pm 4 \text{ pc}$, which makes it one of the nearest high-mass star-forming regions, comparable in distance to the Orion Molecular Cloud. W40 lies on the far side of the Aquila Rift and so, in the visible, it is partially obscured by foreground extinction (up to visual extinctions of $A_V \sim 3$ magnitudes; V. Straizys, K. Černis & S. Bartašiūtė 2003). Thus, it is best observed at IR and far-infrared (FIR) wavelengths. Fig. 1 shows a $70 \mu\text{m}$ (left) and a $500 \mu\text{m}$ (middle) image of the W40 complex using the PACS (A. Poglitsch et al. 2010) and SPIRE (M. J. Griffin et al. 2010) instruments, respectively, aboard the ESA Herschel Space Observatory¹ (G. L. Pilbratt et al. 2010). The data were obtained from the Herschel Science Archive.² Fig. 1 (right) shows the H_2 column density maps derived by the Herschel Gould Belt Survey (HGBS) (P. André et al. 2010; S. Bontemps et al. 2010; V. Könyves et al. 2015; D. Arzoumanian et al. 2019) (see Section 2). The $500 \mu\text{m}$ (Fig. 1, middle) and H_2 column density (Fig. 1: right) images show that W40 is associated with an extended molecular cloud complex with an estimated mass of $\sim 10^4 M_\odot$ (S. A. Rodney & B. Reipurth 2008). The $70 \mu\text{m}$ image shows two cavities that form an hour-glass (bipolar) shape on large scales (a few pc) with an H II region at the centre powered by an OB association. The bright-rimmed clouds at the cavity walls show clearly that dense clumps and pillars are illuminated from inside by the cluster. The main cluster is located just north-west of the intersection of the two cavities seen in Fig. 1 (left); near the tip of the ‘hook’ within the grey/green box in the H_2 column density map (Fig. 1 – right). The cluster is more clearly seen in the zoomed-in image of the 2MASS *H*-band data (M. F. Skrutskie et al. 2006) in Fig. 2 (top right). The OB association is comprised of IRS/OS1a (O9.5), IRS/OS2b (B4), and IRS/OS3a (B3) and an associated stellar cluster of pre-main-sequence stars (R. Y. Shuping et al. 2012). *Chandra* X-ray observations (M. A. Kuhn et al. 2010) have revealed over 200 young stellar objects in a $17 \text{ arcmin} \times 17 \text{ arcmin}$ area centred on the cluster. Based on the Herschel fluxes at 70 and $160 \mu\text{m}$, the average strength of the UV field in W40 is $237 G_0$ (N. Schneider et al. 2020; and references therein), where G_0 is the strength of the average interstellar radiation field in Habing units ($1.6 \times 10^{-3} \text{ erg cm}^{-2} \text{ s}^{-1}$; H. J. Habing 1968). However, the same observations show that the UV field strength can reach $\sim 8200 G_0$ near the OB association.

¹Herschel was an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

²<http://archives.esac.esa.int/hsa/whsa/>

Table 1. HCN and HCO⁺ abundances (relative to N(H₂)) for various objects in the literature.

| Object(s) | X(HCN) | X(HCO ⁺) | X(HCN)/X(HCO ⁺) | Resolution | Reference |
|--------------------------------|-----------------------|-----------------------|-----------------------------|------------------------|--|
| Low-mass protostars | 1.7×10^{-9} | 6.6×10^{-9} | 2.6 | 13–18 arcsec | J. K. Jørgensen, F. L. Schöier & E. F. van Dishoeck (2004) |
| W51 | 2.0×10^{-9} | 1.5×10^{-10} | 13.3 | 1800 arcsec (40–50 pc) | Y. Watanabe et al. (2017) |
| YSO GL2591 | 2.0×10^{-8} | 10^{-8} | 2.0 | 13 arcsec | F. F. S. der Tak et al. (1999) |
| IRDCs – quiescent | – | 2.4×10^{-8} | – | 38 arcsec | P. Sanhueza et al. (2012) |
| IRDCs – active | – | 5.7×10^{-8} | – | 38 arcsec | P. Sanhueza et al. (2012) |
| ATLASGAL clumps – quiescent | – | 4.0×10^{-9} | – | 38 arcsec | S. Hoq et al. (2013) |
| ATLASGAL clumps – protostellar | – | 6.4×10^{-9} | – | 38 arcsec | S. Hoq et al. (2013) |
| ATLASGAL clumps – H II/PDR | – | 7.5×10^{-9} | – | 38 arcsec | S. Hoq et al. (2013) |
| 70 μm quiet clumps | – | 2.5×10^{-10} | – | 27 arcsec | A. Traficante et al. (2017) |
| Starburst galaxies | – | – | 2–7 | 50 pc | J. Butterworth, S. Viti & Y. Wang (2025) |
| AGNs | – | – | 1–41 | 50 pc | J. Butterworth et al. (2025) |
| ULIRGS | – | – | 2–24 | 500 pc | J. Butterworth et al. (2025) |
| LIRGS and ULIRGS | 10^{-7} – 10^{-8} | 10^{-8} | 1–10 | 0.3 arcsec (50–500 pc) | Y. Nishimura et al. (2024) |




Figure 1. Left: public PACS-70 μm observations and (middle) SPIRE-500 μm observations from the Herschel Science Archive. Right: H₂ column density maps from the HGBS (P. André et al. 2010; S. Bontemps et al. 2010; V. Könyves et al. 2015; D. Arzoumanian et al. 2019) (see Section 2). The box shows the region that we focus on in this paper (see e.g. Fig. 2). A 1 parsec scalebar is shown at the upper left corner of each image, and a circle indicating the observational beamsize appears in the lower right.

In this paper we present observations and analysis of HCN and HCO⁺ J = 3 → 2 in W40, taken as part of the MAJORS survey at the JCMT. The main goals of this paper are to determine the HCN and HCO⁺ abundances, examine the variations in abundance and physical conditions across the entire region mapped, and to estimate the gas density associated with the HCN and HCO⁺ emission. In Section 2 we present the observations. In Section 3 we present basic analyses of the data including integrated intensity maps and the observational relationship between the HCN and HCO⁺ integrated intensities. In Section 4 we calculate the HCN and HCO⁺ abundances based on an analysis of the data utilizing both the RADEX 1D radiative transfer code (F. F. S. der Tak et al. 1999) and the NAUTILUS gas-grain chemical evolution code (M. Ruaud, V. Wakelam & F. Hersant 2016). In Section 5 we present the main conclusions of the paper.

2 OBSERVATIONS

The HCO⁺ J = 3 → 2 line at $\nu = 267.5576259$ GHz and HCN J = 3 → 2 line at $\nu = 265.8864339$ GHz were observed as part of the MAJORS large program (Eden et al. in preparation – observing program M22AL002) at the James Clerk Maxwell Telescope (JCMT). Line frequencies were obtained from the Cologne Database for Molecular Spectroscopy (CDMS)³ (H. S. P. Müller

et al. 2001, 2005; C. P. Endres et al. 2016). The data were taken using the dual polarization dual-sideband $\ddot{u}\ddot{u}$ heterodyne receiver (A. R. Kerr et al. 2014). Maps were centred at α (J2000) = 18^h31^m17^s and δ (J2000) = –02°05′00″. Full width at half-maximum beam sizes are $\theta_{\text{HCO}^+ J=3 \rightarrow 2} = 18.3$ arcsec and $\theta_{\text{HCN} J=3 \rightarrow 2} = 18.4$ arcsec. Data were obtained on 2020 August 21 as part of director’s discretionary time (observing program M20AD003) to test the MAJORS observing strategy.

The atmospheric opacity at 225 GHz ranged from 0.225 to 0.315 over the course of the observations (as measured by the JCMT’s in-cabin 183 GHz Water Vapour Monitor). Data were obtained in raster scan, position switching mode with a scan speed of 16 arcsec s^{–1} and an off position at α (J2000) = 18^h31^m53^s.7 and δ (J2000) = –1°59′35″. Two scans in orthogonal directions (i.e. ‘basket-weaved’) were used to produce a smoother output map. Maps are 424 arcsec × 424 arcsec in size with a pixel/sampling size of 8 arcsec × 8 arcsec. The median system temperature during the observations was ~300 K. The spectrometer, ACSIS, was used in the 250 MHz mode (8192 channels, 30.517578 kHz spacing) resulting in a native spectral resolution of 0.034 km s^{–1}.

Data reduction was performed with ORAC-DR (T. Jenness et al. 2015), which is built on the STARLINK (M. J. Currie et al. 2014) packages KAPPA (M. J. Currie & D. S. Berry 2014), CUPID (D. S. Berry et al. 2007), and SMURF (E. Chapin et al. 2013). We used the REDUCE_SCIENCE_NARROWLINE recipe that creates a spatial cube from the raw time-series data, removes a

³<https://cdms.astro.uni-koeln.de/classic/>




Figure 2. Top left: HGBS H_2 column density overlaid with contour levels of 1, 2, 3, $4 \times 10^{22} \text{ cm}^{-2}$. These same contours are overlaid in all following plots. The green, blue, and red boxes indicate regions selected for low H_2 column density and high temperature, intermediate $N(\text{H}_2)$ and T_K , and high $N(\text{H}_2)$ and low T_K , respectively. The Green Circle, Blue Square, and Red Triangle denote positions for which we perform chemical modelling in Section 4.3. Top middle: HGBS dust temperature. Top right: 2MASS H -band image. Bottom left: integrated intensity image for $\text{HCO}^+ J = 3 \rightarrow 2$. Bottom middle: integrated intensity image for $\text{HCN } J = 3 \rightarrow 2$. Bottom right: ratio of the HCN/HCO^+ integrated intensity. Values for each quantity are given in the colour bars to the right of each plot. A 0.1 parsec scalebar is shown at the upper left corner of the first image, and a circle indicating the observational beamsize appears in the lower right of each image (except for the 2-MASS image).

baseline from the data, and trims the ends of the frequency range to remove high-noise regions. The data cubes provided by the pipeline were subsequently converted from NDF to FITS format using the STARLINK `NDF2FITS` command to allow for further processing in CASA (J. P. McMullin et al. 2007) and PYTHON. Post-processing, the data were spectrally re-binned in STARLINK by 15 channels to provide a spectral resolution of $\sim 0.5 \text{ km s}^{-1}$. With this resolution, the average 1σ noise is 0.33 K for the $\text{HCO}^+ J = 3 \rightarrow 2$ data and 0.29 K for the $\text{HCN } J = 3 \rightarrow 2$ data (T_A^*). This noise was estimated using a robust baseline-subtracted MAD method with Savitzky–Golay smoothing of 3 km s^{-1} , polyorder = 2, σ -clipping at 2.5σ , and three iterations.

The HCO^+ and HCN data in this paper are presented in the corrected antenna temperature (T_A^*) scale, but are converted to main beam brightness ($T_{\text{mb}} = T_A^*/\eta_{\text{mb}}$) for column density calculations (Section 4.1) using a main beam efficiency of $\eta_{\text{mb}} = 0.66$.⁴ While calibration errors for the ‘ \ddot{U} ’ receiver are not fully characterized, the JCMT receiver webpage⁵ states that ‘typical uncertainties in peak flux measurements tend to be in the range of 10–20 per cent’. Based on this, we adopted an average calibration uncertainty of 15 per cent.

We note that HCN has hyperfine structure (e.g. A. M. Mullins et al. 2016; R. M. Loughnane et al. 2012) but for the $J = 3 \rightarrow 2$ transition, the inner four lines are only 0.35, 0.07, and -0.54 km s^{-1} from the central strongest line and, given the width of our observed lines ($> 1 \text{ km s}^{-1}$; see Section 3), are blended, appear as a

single component, and cannot be distinguished from one another. The outer two lines are 1.82 and -2.28 km s^{-1} from the central, strongest line but, since these are each 25 times weaker than the central component (and they are not detected), we ignore the hyperfine structure of HCN in the remainder of this paper.

For subsequent analysis, we also obtained dust/kinetic temperature and H_2 column density maps from the HGBS (P. André et al. 2010; S. Bontemps et al. 2010; V. Könyves et al. 2015; D. Arzoumanian et al. 2019; hereafter HGBS). These maps were produced by V. Könyves et al. (2015) by taking the Herschel SPIRE data at 250, 350, and $500 \mu\text{m}$, smoothing to the resolution of the SPIRE $500 \mu\text{m}$ observations ($\theta_{\text{FWHM}} = 36.3 \text{ arcsec}$) and fitting the data to a modified blackbody of the form: $I_\nu = B_\nu(T_d)\kappa_\nu\mu_{\text{H}_2}m_{\text{H}}N(\text{H}_2)$, where κ_ν is the dust mass opacity coefficient (given as $\kappa_\lambda = 0.1 \times (\lambda/300 \mu\text{m})^{-2} \text{ cm}^2 \text{ g}^{-1}$), μ_{H_2} is the mean molecular weight per hydrogen molecule (2.8), and m_{H} is the mass of atomic hydrogen. In the fit, the dust temperature (T_d) and column density ($N(\text{H}_2)$) were left as free parameters. The resulting images were sampled on a 3 arcsec pixel grid. Details of the procedure to produce the dust/kinetic temperature and H_2 column density maps are provided in V. Könyves et al. (2015). These maps were subsequently regridged to the same grid and pixel size as our HCO^+ and HCN data (i.e. $8 \text{ arcsec} \times 8 \text{ arcsec}$) using the CASA (J. P. McMullin et al. 2007) `imregrid` command.

3 RESULTS

3.1 Integrated intensity comparisons

Fig. 2 shows the H_2 column density ($N(\text{H}_2)$ – top left) and dust/kinetic temperature ($T_{\text{dust}} = T_K$ – top middle) from the HGBS, as well as the location of young stars seen in the 2MASS

⁴<https://www.eoobservatory.org/jcmt/instrumentation/heterodyne/namakanui/uu-230ghz/>

⁵<https://www.eoobservatory.org/jcmt/instrumentation/heterodyne/calibration/>




Figure 3. $N(\text{H}_2)$ versus T_K (i.e. T_{dust}) for every pixel within the coloured boxes shown in Fig. 2. The green points plot the values for pixels located within the Green Region, the blue points plot the values for pixels located within the Blue Region, and the red points plot the values for pixels located within the Red Region. Error bars are 5 per cent in T_K and 12 per cent in $N(\text{H}_2)$ based on the analysis of uncertainties by A. Roy et al. (2014).

H-band image (top right) (M. F. Skrutskie et al. 2006). Note that we will continue to use the dust temperature as a proxy for the gas kinetic temperature, but will explore the effects of this assumption on our analysis in Section 4.1.2.

The figure also shows the integrated intensity maps for HCO⁺ $J = 3 \rightarrow 2$ (bottom left) and HCN $J = 3 \rightarrow 2$ (bottom middle), and the ratio of the HCN/HCO⁺ integrated maps (bottom right). To derive the HCO⁺ and HCN integrated intensities, we fit a single Gaussian to every pixel in the HCO⁺ and HCN cubes using the Trust Region Reflective Least Squares fitter (TRFLSQFITTER) in PYTHON/ASTROPY. This fitter allows upper and lower parameter bounds and, in ASTROPY 5.3 and later, provides covariance-based parameter uncertainties. The fits yield peak temperatures (T_A^* ; K), line widths (ΔV_{FWHM} ; km s⁻¹, with $\Delta V_{\text{FWHM}} = 2\sqrt{2 \ln 2} \sigma$), line centroids (V_{LSR} ; km s⁻¹), and integrated intensities ($\int T_A^* dV$; K, km s⁻¹, with $\int T_A^* dV = T_A^* \sigma \sqrt{2\pi}$). Errors in the integrated intensity were calculated from the errors in T_A^* and σ from the equation $\sqrt{(\sigma \times \sigma_{\text{err}})^2 + (T_A^* \times T_{A,\text{err}}^*)^2} \times \sqrt{2\pi}$, where $T_{A,\text{err}}^*$ is calculated by combining the 15 per cent calibration error in quadrature with the Gaussian fitting error. We retain only fits with signal-to-noise (S/N) > 3 in both the peak and the integrated intensity; pixels failing either criterion are set to NaN. Accepted fits were visually inspected to reject spurious solutions. We adopt Gaussian-fit integrals rather than MOMENT maps because this enables simultaneous S/N cuts on peak and area; tests with the MOMENT command, however, produced consistent results. Blank regions in the integrated-intensity maps mark pixels with S/N < 3. Per-pixel uncertainties depend on the local noise but are, on average, ~25 per cent of the integrated intensity.

The green, blue, and red boxes in Fig. 2 delineate regions chosen for their locations relative to the OB association and for specific conditions. Fig. 3 shows that points within the Green




Figure 4. HCN and HCO⁺ spectra and their Gaussian fits of three representative positions marked by the Green Circle, Blue Square, and Red Triangle in Fig. 2.

Region generally have higher kinetic temperatures ($\langle T_K \rangle = 30.2\text{K}$) and lower H₂ column densities ($\langle N(\text{H}_2) \rangle = 1.26 \times 10^{22} \text{ cm}^{-2}$) than the rest of W40. In contrast, points within the Red Region have some of the lowest kinetic temperatures ($\langle T_K \rangle = 21.6\text{K}$) and highest H₂ column densities ($\langle N(\text{H}_2) \rangle = 3.5 \times 10^{22} \text{ cm}^{-2}$). Points in the Blue Region have intermediate mean values of T_K and $N(\text{H}_2)$ of 27K and $2.4 \times 10^{22} \text{ cm}^{-2}$, respectively. These regions will be utilized more fully in later discussion (see Section 4). Fig. 4 shows example spectra and Gaussian fits at the positions marked by the Green Circle, Blue Square, and Red Triangle in Fig. 2 (top left). These spectra also indicate that there is a velocity gradient across the filament, which first moment maps reveal to be quite smooth.

As can be seen in Fig. 2, the molecular gas is distributed in a curved filament wrapping around the hottest gas which is co-located with the central OB association. While the HCO⁺ and HCN have similar morphologies and extent, the HCO⁺ is slightly more extended than HCN in the low integrated intensity regions. This is particularly noticeable to the left of the Green Region (where the H₂ column density is low and the dust temperature is high) and in the Red Region (where the H₂ column density is high but the dust temperature is low). The lower number of HCN pixels in the Red Region is due to weaker HCN emission lines (see e.g. Fig. 4 – bottom) that causes the integrated intensity to fall below the 3 σ noise limit.

Fig. 5 shows the relationship between the HCN $J = 3 \rightarrow 2$ and HCO⁺ $J = 3 \rightarrow 2$ integrated intensity ($I(\text{HCN})$ and $I(\text{HCO}^+)$, respectively) for the points displayed in Fig. 2. Using the `stats.linregress` function within SCIPY we found that, for the entire data set (i.e. points both inside and outside the Red, Green, and Blue Regions in Fig. 2), $I(\text{HCN})$ (K km s⁻¹) = 0.96 $I(\text{HCO}^+)$ (K km s⁻¹) with a y-intercept of -0.70, a Pearson r -value of 0.83, and a p -value of < 10⁻⁹. This suggests that there is a strong linear correlation to the data and that the trend is statistically significant. Over the entire region $\langle I(\text{HCN}) \rangle = 4.9 \text{ K}$ and $\langle I(\text{HCO}^+) \rangle = 4.8 \text{ K}$.




Figure 5. Integrated intensities (K km s^{-1}) of HCN $J = 3 \rightarrow 2$ versus $\text{HCO}^+ J = 3 \rightarrow 2$ for all positions with a signal-to-noise ratio above 3σ (i.e. those displayed in Figs 2 and 3). Points are colour coded to indicate the blue (left panel), red (middle panel), or green (right panel) regions (seen in Fig. 2) from which they were taken. The solid black lines showing the best linear fit to the data points in each of the regions. The dotted grey line shows the $I(\text{HCN}) = I(\text{HCO}^+)$ relation for comparison.

Breaking the data down into the regions displayed in Fig. 2, the coloured points show the relationship for each of the respective coloured regions, with solid black lines showing the best linear fit. In the Blue Region (left panel) $I(\text{HCN}) = 0.89 I(\text{HCO}^+)$ with a y -intercept of 0.23, a Pearson r -value = 0.75, and a p -value of $< 10^{-9}$, again suggesting strong evidence for a statistically significant linear relationship. In this region $\langle I(\text{HCN}) \rangle = 9.6 \text{ K}$ and $\langle I(\text{HCO}^+) \rangle = 10.5 \text{ K}$. In the Red Region (middle panel) $I(\text{HCN}) = 0.14 I(\text{HCO}^+)$ with a y -intercept of 1.93, a Pearson r -value = 0.28, and a p -value of 0.05, suggesting that there is only weak evidence for a statistically significant, linear relationship. In this region $\langle I(\text{HCN}) \rangle = 2.6 \text{ K}$ and $\langle I(\text{HCO}^+) \rangle = 4.8 \text{ K}$. And, in the Green Region (green points) $I(\text{HCN}) = 1.48 I(\text{HCO}^+)$ with a y -intercept of -2.26 , a Pearson r -value = 0.81, and a p -value of $< 10^{-9}$. In this region $\langle I(\text{HCN}) \rangle = 6.6 \text{ K}$ and $\langle I(\text{HCO}^+) \rangle = 5.9 \text{ K}$. Thus, the correlation is weakest in the Red Region, where the H_2 column density is highest, the dust temperature is the lowest, and the line intensities are weakest. The Green and Blue Regions, on the other hand, have similar, and strongly correlated, relationship between the HCN and HCO^+ integrated intensities.

The relationship between our $J = 3 \rightarrow 2$ integrated intensities in the Blue Region is similar to that seen in the $J = 1 \rightarrow 0$ transition in outer Galaxy clouds, where $I(\text{HCN } J = 1 \rightarrow 0) = 0.5$ to $0.9 I(\text{HCO}^+ J = 1 \rightarrow 0)$ (S. Patra et al. 2022; J. Braine et al. 2023). In the inner Galaxy, a sample of six clouds by N. J. Evans et al. (2020) find $I(\text{HCN } J = 1 \rightarrow 0) = 1.2 I(\text{HCO}^+ J = 1 \rightarrow 0)$, more similar to the relationship seen in the warm, Green Region. Given the lower critical densities of the $1 \rightarrow 0$ transitions and the larger beamsize of these observations ($\sim 1'$), it is surprising to see such a similarity in these relationships. It should be noted, however, that in the case of N. J. Evans et al. (2020) and S. Patra et al. (2022), the authors did attempt to restrict their analysis to dense gas by only selecting high column density/extinction regions with $A_V > 8$.

The fit to the full data set closely resembles that derived for the Blue Region. This is likely because the Blue Region spans the widest range of integrated intensities and contains the brightest HCN and HCO^+ emission. As a result, the global fit is strongly influenced by the large number of high-intensity points originating in this region. This illustrates the risk of examining the

HCN– HCO^+ relationship in a purely global sense, as bright regions can mask distinct relationships that may be present in other parts of the same cloud.

3.2 Molecular line ratios

Over the entire map, the HCN/ HCO^+ integrated intensity ratio ranges from 0.35 to 1.71 with a mean value of 0.83 ($\sigma = 0.3$). As can be seen in Fig. 2, while the HCN/ HCO^+ ratio traces the general morphology of the HCN integrated intensity map, it does not peak at the same location as the HCN or HCO^+ emission peaks. In fact, the HCN/ HCO^+ ratio is moderately larger (average of ≈ 1.1) to the north of the regions of brightest HCN and HCO^+ emission (i.e. in the Green Region and the northern part of the Blue Region; towards the hot gas surrounding the OB association) than it is throughout the bulk of the filament (where the average is ≈ 0.5 to 0.6). In addition, points within the Green Region generally have elevated HCN/ HCO^+ integrated intensity ratios ($\langle I(\text{HCN})/I(\text{HCO}^+) \rangle = 1.1$). In contrast, the Red Region encompasses some of the lowest HCN/ HCO^+ integrated intensity ratios ($\langle I(\text{HCN})/I(\text{HCO}^+) \rangle = 0.5$). Points in the Blue Region have an intermediate value of $\langle I(\text{HCN})/I(\text{HCO}^+) \rangle = 0.9$.

These kinds of differences in the HCN/ HCO^+ ratio are also seen in the observations of Q. Nguyen-Luong et al. (2020) who mapped the HCN and $\text{HCO}^+ J = 1 \rightarrow 0$ emission in M17 and found that the mean HCN/ HCO^+ value is enhanced in M17-HII (1.36) versus that in M17-IRDC (1.09); a difference that they attribute to increased UV heating in M17-HII. We can make more direct comparisons, however, with HCN/ $\text{HCO}^+ J = 4 \rightarrow 3$ observations due to the similar critical densities between the $J = 3 \rightarrow 2$ and $4 \rightarrow 3$ transitions. In the Perseus cloud S. L. Walker-Smith et al. (2014) find a HCN/ $\text{HCO}^+ J = 4 \rightarrow 3$ ratio of 0.1 to 0.6, which is similar to what is seen in the bulk of the filament in W40. On larger (i.e. Galactic) scales, Q.-H. Tan et al. (2018) found that the HCN/ $\text{HCO}^+ J = 4 \rightarrow 3$ ratio in six nearby star-forming galaxies varies from 0.1 to 2.7 with a mean value of 0.9 (standard deviation of 0.6), which is very comparable to our observed values across the whole of the W40 region. Similarly,

J. Butterworth et al. (2025) studied the HCN/HCO⁺ $J = 3 \rightarrow 2$ and $J = 4 \rightarrow 3$ ratios in a sample of 80 sources (active galactic nuclei, AGNs; starburst galaxies, SB; and (U)LIRGs) at various spatial resolutions. At molecular-cloud scales (25–50 pc), the HCN/HCO⁺ $J = 3 \rightarrow 2$ ratio is < 1 in SB galaxies, consistent with our red and Blue Regions. In the Green Regions, the ratio exceeds unity but does not reach typical AGN values (> 1.5). This suggests that, while similar physical/chemical processes may influence the line ratios in the Green Regions, the environmental conditions are less extreme than in AGNs.

4 DISCUSSION

4.1 HCO⁺ and HCN column densities

To determine the HCO⁺ and HCN column densities, we utilized the Gaussian fits to the spectra as described in Section 3. The peak and integrated intensities of the valid fits were converted from T_A^* to T_{mb} units using the main beam efficiency (see Section 2). Each pixel with an accepted Gaussian fit was then assigned a gas kinetic temperature (T_K) from the regridded HGBS dust temperature map (Fig. 2) assuming that $T_K = T_{dust}$. We will explore the effects of this assumption in Section 4.1.2.

We then utilized the RADEX 1D radiative transfer code (F. F. S. der Tak et al. 2007), which, given a density, kinetic temperature, line width, and molecular column density, produces the expected peak intensity (T_{mb}) and opacity for a particular molecular transition. The HGBS maps allowed us to fix the temperature in each pixel. The Gaussian fitting also provided the observed integrated intensity ($\int T_{mb} dV$) in each pixel. We then used the standard Python script RADEX-COLUMN.PY (available on the RADEX website⁶), which begins with an initial column density of 10^{12} cm^{-2} and iteratively adjusts the molecular column density (up or down) until the RADEX model integrated intensity matches the observed value within a specified tolerance. Thus, we were able to determine the HCO⁺ and HCN column densities for every pixel in our HCO⁺ and HCN maps with accepted Gaussian fits. The only free parameter in our RADEX calculation was the (unknown) H₂ volume density ($n(\text{H}_2)$), so we sampled densities logarithmically from 10^3 to 10^8 cm^{-3} in 0.5-dex increments (i.e. 10^3 , 5×10^3 , 10^4 , 5×10^4 , ...). Accordingly, we generated HCN and HCO⁺ column density maps for each of the volume densities. The optimal density is determined later through comparison with the predictions of the chemical models, which are described in Section 4.3.

4.1.1 Effects of assuming local thermodynamic equilibrium

Although local thermodynamic equilibrium (LTE) column densities are easier to calculate, LTE assumes that the population of every level is well described by the Boltzmann distribution, an assumption that is certainly not appropriate for the densities explored here. To illustrate, we have compared an optically thin LTE analysis to a ‘fiducial’ RADEX result, which uses a volume density of $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$. We have chosen this H₂ density to represent a fiducial result because it is mid-way between the density extremes of 10^3 – 10^8 cm^{-3} used in our analysis in the previous section.




Figure 6. Map of the ratio of the kinetic temperature calculated from NH₃ observations (R. K. Friesen et al. 2017; T_{NH_3}) to the dust temperature calculated from the HGBS data (V. Könyves et al. 2015; T_{dust}). The red rectangle denotes the same ‘Red Region’ as indicated in Fig. 2.

The comparison between the optically thin LTE analysis and the fiducial RADEX results shows a large difference between the two. The RADEX HCN column densities are 50–100 times larger than the LTE values and the RADEX HCO⁺ column densities are 5–20 times larger than the LTE values. These differences are due to both excitation and opacity effects. At a density of $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$, the rotational levels are sub-thermally excited and so the Boltzmann equation does not properly account for the population in the other states. In addition, the RADEX derived opacity (for $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$) across the entire cloud is significant; with $\tau_{\text{HCN}} \sim 5$ –80 and $\tau_{\text{HCO}^+} \sim 1$ –20. Thus, at a density of $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$ the optically thin, LTE assumption is seriously in error, and one must take both opacity and non-LTE excitation into account.

4.1.2 Effects of assuming $T_K = T_{dust}$

In the previous analysis we assumed that the HCN and HCO⁺ kinetic temperature was provided by the dust temperature maps (i.e. $T_K = T_{dust}$). This assumption is probably sufficient in regions where $n(\text{H}_2) > 10^5 \text{ cm}^{-3}$ (e.g. P. Goldreich & J. Kwan 1974; E. Kruegel & C. M. Walmsley 1984; P. F. Goldsmith 2001) but, in lower density regions, this assumption breaks down due to insufficient coupling between the gas and dust.

To test this assumption, and the affect it has on our column density calculations, we utilized the results of R. K. Friesen et al. (2017), who mapped the NH₃ (1,1) and (2,2) transitions in a number of sources using the Green Bank Telescope. By modelling the NH₃ data, they were able to produce maps of the gas kinetic temperature (that we call T_{NH_3}). Sensitivity in the NH₃ observations have limited this temperature map to the Red Region of our observations. These data were also regridded on to the same grid and pixel size as the HCO⁺, HCN, and dust data using the CASA (J. P. McMullin et al. 2007) `imregrid` command.

Fig. 6 shows the ratio of the gas to dust temperature (i.e. $T_{\text{NH}_3}/T_{\text{dust}}$) in the Red Region of W40. There is a distinct difference in this ratio between the northern and southern sections of the Red Region. In the northern section, the average value of $T_{\text{NH}_3}/T_{\text{dust}}$ is 0.71 ($\sigma = 0.03$), whereas in the southern section the average value is 0.89 ($\sigma = 0.05$). The reason for this

⁶<https://sronpersonalpages.nl/vdtak/radex/index.shtml>




Figure 7. Abundance maps for (top left) HCO^+ and (top right) HCN. The bottom left image shows the HCN/ HCO^+ column density (i.e. abundance) ratio. All values were calculated from the RADEX derived column densities using $n(\text{H}_2) = 10^5$ (for illustrative purposes), divided by the HGBS H_2 column density map. Colour bars to the right of each figure provide the relevant scales. A 0.1 parsec scale bar is shown in the upper left corner of the first image, and a circle indicating the observational beam size appears in the lower right.

dichotomy is not immediately clear but it may be related to the volume density. If the volume density is lower in the northern section, which could be true given the weak HCN and HCO^+ lines in this area, then we would expect weaker coupling between the gas and dust and therefore a poorer match between the gas and dust temperatures.

The difference in temperature corresponds to a difference in the derived HCN and HCO^+ column densities as well. Using our fiducial volume density (i.e. $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$) we again used RADEX to calculate the HCN and HCO^+ column densities using both the dust temperature map ($N(T_{\text{dust}})$) and the NH_3 temperature map ($N(T_{\text{NH}_3})$). A map of $N(T_{\text{NH}_3})/N(T_{\text{dust}})$ reveals a similar distinct ratio between the northern and southern sections of the Red Region. For HCN, in the northern section the average value of $N(T_{\text{NH}_3})/N(T_{\text{dust}})$ is 1.78 ($\sigma = 0.13$), whereas, in the southern section, the average value is 1.18 ($\sigma = 0.08$). For HCO^+ , in the northern section the average value of $N(T_{\text{NH}_3})/N(T_{\text{dust}})$ is 1.68 ($\sigma = 0.17$), whereas, in the southern section, the average value is 1.22 ($\sigma = 0.10$). For both molecules, the lower NH_3 temperature in the northern section results in larger column densities.

Therefore, in regions where the gas and dust temperatures are fairly well matched (i.e. within 10 per cent), the HCN and HCO^+ column densities are also fairly well matched. In this case, using the assumption that $T_{\text{K}} = T_{\text{dust}}$, we underestimate column

densities by only ~ 20 per cent. However, in regions where the gas and dust temperatures are less well matched (i.e. within 30 per cent), the HCN and HCO^+ column densities can be underestimated by up to a factor of 2. The HCN/ HCO^+ column density **ratio**, however, is not as strongly affected, since changes in the temperature affect both column densities by about the same amount and in the same direction.

4.2 HCN and HCO^+ abundances

Our final step was to produce abundance maps by dividing the HCO^+ and HCN column density maps produced by RADEX, by the regridded HGBS H_2 column density map (Fig. 2). The HCO^+ and HCN abundance maps ($X(\text{HCO}^+)$ and $X(\text{HCN})$, respectively) are shown in Fig. 7 for our fiducial case with $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$, along with the ratio of the HCN to HCO^+ abundance ($X(\text{HCN})/X(\text{HCO}^+)$). Note that these results are meant to reflect the abundance variations that one obtains by assuming a constant volume density across the entire region; an assumption that is almost certainly not accurate and will be explored in Section 4.3. Statistics for Fig. 7 are provided in Table 2.

To estimate the abundance uncertainties, we propagated each pixel's integrated-intensity uncertainty (~ 25 per cent) through RADEX. From the fitted integrated intensity we computed a

Table 2. HCN, HCO⁺ abundances (relative to N(H₂)) and HCN/HCO⁺ abundance ratios for various regions within W40. Results are presented for our fiducial case with n(H₂) = 10⁵ cm⁻³.

| | | X(HCN) (×10 ⁻⁸) | X(HCO ⁺) (×10 ⁻⁹) | X(HCN)/X(HCO ⁺) |
|--------------|------|--------------------------------|--|-----------------------------|
| Entire Map | Mean | 2.0±0.5 | 1.9±0.5 | 9.1±0.3 |
| | Min | 0.4±0.2 | 0.4±0.2 | 2.6±0.7 |
| | Max | 7.0±1.0 | 7.3±1.1 | 23.1±9.0 |
| Green Region | Mean | 3.5±0.7 | 3.0±0.7 | 12.0±0.6 |
| | Min | 1.3±0.5 | 1.2±0.5 | 5.3±1.7 |
| | Max | 6.0±0.9 | 5.6±1.1 | 19.1±6.0 |
| Blue Region | Mean | 3.4±0.6 | 4.0±0.7 | 9.5±0.4 |
| | Min | 1.0±0.3 | 1.0±0.4 | 2.6±0.7 |
| | Max | 7.0±1.0 | 7.3±1.1 | 23.1±9.0 |
| Red Region | Mean | 0.7±0.2 | 1.3±0.3 | 6.1±0.4 |
| | Min | 0.4±0.2 | 0.4±0.2 | 3.1±1.1 |
| | Max | 1.1±0.3 | 2.7±0.5 | 14.8±8.0 |

fiducial column density, and then a maximum (minimum) column density by increasing (decreasing) the intensity by its uncertainty. We also tested a 5 per cent uncertainty in the kinetic temperature. Its effect on the derived column densities was negligible and is therefore ignored. From the resulting HCN and HCO⁺ column densities and their uncertainties, we combined the 12 per cent N(H₂) uncertainty (A. Roy et al. 2014) in quadrature to obtain fiducial HCN and HCO⁺ abundances and their uncertainties. Although the propagated column density (and hence abundance) uncertainties are slightly asymmetric, at the precision reported in Table 2 they are effectively symmetric, so we quote them as simple ± values.

In Fig. 7, we can see that the HCO⁺ and HCN abundances mostly track with the integrated intensity maps (Fig. 2) with peaks in the Blue Region (around α = 18^h31^m21^s, δ = -2°06′30″) and a ‘tail’ extending into the Red Region (where N(H₂) is high and T_K is low). There are, however, a few notable differences. The Green Region and the northern part of the Blue Region (where N(H₂) is low but T_K is high) have enhanced HCO⁺ and HCN abundances. Similar enhancements are not as clearly seen in the integrated intensity maps. In addition, the abundance ratio in these two regions is also enhanced, leading to X(HCN)/X(HCO⁺) ratios in excess of 12 (and up to ~ 20), versus <~ 10 in the rest of the cloud. Note that the region with X(HCN)/X(HCO⁺) > 15 is also the region in which I(HCN)/I(HCO⁺) > 1 (Section 3).

Table 2 highlights the regional trends discussed above. In particular, the Red Region has the lowest mean, minimum, and maximum values for both abundances and the abundance ratios (although the minimum abundance ratio is slightly larger than that seen in the Blue Region). The Blue Region has a mean HCN abundance similar to that of the Green Region but also has the highest mean HCO⁺ abundance, resulting in a mean abundance ratio between that of the Red and Green Regions. While the absolute abundances vary by a factor of 15–20 across the map, the abundance ratio varies by only a factor of 9.

The column densities derived from RADEX and therefore the abundances, scale roughly with the density. If we decrease the density to n(H₂) = 5 × 10⁴ cm⁻³ the abundances of both species

scale upwards by a factor of ≈ 2 to 5, but the abundance ratios remain approximately constant (within ~ ±20 per cent). Increasing the density to n(H₂) = 5 × 10⁵ cm⁻³ results in a decrease in the absolute abundances by a factor of approximately 2 to 5, while the abundance ratios again remain largely unchanged. So, changes in the H₂ density can help explain the abundance variations across W40, as well as the relatively small changes in the abundance ratios (e.g. Table 2). A more detailed discussion of how density affects our results, as well as the method we use to estimate the appropriate H₂ volume density, is provided in the following Section.

4.3 Chemical modelling of the HCO⁺ and HCN abundances

4.3.1 Description of the modelling

To investigate the origin of the observed variations in HCN and HCO⁺ abundances and abundance ratios, and to obtain a crude estimate of the gas density in the emitting regions, we modelled their time evolution with the NAUTILUS gas–grain chemical code (M. Ruaud et al. 2016). NAUTILUS follows chemical evolution in three phases—gas, grain surface, and grain mantle—and tracks how abundances change over time as a function of physical conditions such as: gas density, dust and gas temperature, visual extinction, ultraviolet flux, and cosmic-ray ionization (and others not discussed here). The code employs the KINetic Database for Astrochemistry (KIDA) data base (V. Wakelam et al. 2012) and solves a series of coupled differential equations for a network of more than 10 000 reactions: 7627 gas-phase reactions and 3799 grain reactions, including adsorption, desorption, surface chemistry, mantle processing, and migration between the mantle and surface. These reactions connect 510 gas-phase species, 225 grain-surface species, and 225 grain-mantle species. No deuterated species or isotopologues are included in the reaction network.

As before, the HGBS maps allow us to fix the temperature and H₂ column density in each pixel. We have again assumed that T_{dust} = T_{gas} everywhere. The H₂ column density provides us with the visual extinction in each pixel. Studies of the H₂ to A_V conversion range from A_V = $\frac{N_{H_2}}{2.1 \times 10^{21}}$ (B. L. Rachford et al. 2009; H. Zhu et al. 2017) to A_V = $\frac{N_{H_2}}{1.9 \times 10^{21}}$ (R. C. Bohlin, B. D. Savage & J. F. Drake 1978; D. C. B. Whittet 1981). In this paper, we have used the more recent value of A_V = $\frac{N_{H_2}}{2.1 \times 10^{21}}$.

For the UV field, the NAUTILUS code expresses the flux as S × 10⁸ photons cm⁻² s⁻¹ in the 6–13.6 eV range (K. I. Öberg et al. 2007; Ruaud et al. 2016), where S is a scaling factor. Setting S = 1 corresponds to G₀ ~ 1 in Habing units, where G₀ = 1.6 × 10⁻³ erg cm⁻² s⁻¹ (H. J. Habing 1968). We used a scaling factor of S = 237 (i.e. G₀ = 237) to model the average radiation field in W40 from its stellar cluster (N. Schneider et al. 2020). For the initial atomic abundances, we used the default values provided by M. Ruaud et al. (2016) (their table 1) that assumes an oxygen abundance of 2.4 × 10⁻⁴ and a C/O ratio of 0.7. We also used a standard cosmic-ray (CR) ionization rate of ζ_{H₂} = 1.3 × 10⁻¹⁷ s⁻¹.

We employed a three-phase, static model within NAUTILUS. In this framework, the cloud is assumed to have fixed physical parameters that remain constant as the chemistry evolves over time. Although we acknowledge that real molecular clouds evolve dynamically, in the absence of information about the prior state and time evolution of the gas, we are constrained to adopt static conditions. The binding energies are taken from the KIDA

data base (V. Wakelam et al. 2012), with values of 2050 K for HCN and 1150 K for CO. No binding energy is provided for HCO^+ , as NAUTILUS does not permit the freeze-out of ionic species such as HCO^+ . Instead, interactions between ions and grains are treated through ion-grain recombination reactions, for example: $\text{HCO}^+ + \text{grain}^- \rightarrow \text{CO} + \text{H} + \text{grain}^0$, rather than through an adsorption-desorption balance.

The physical parameter with the most uncertainty is the H_2 volume density and so we ran NAUTILUS with densities of $n(\text{H}_2) = 10^3\text{--}10^8 \text{ cm}^{-3}$ with the same logarithmic spacing as described in Section 4.1. Densities less than 10^4 cm^{-3} and greater than 10^6 cm^{-3} failed to match the abundances calculated by our RADEX analysis, so we restricted further analysis to $n(\text{H}_2) = 10^4, 5 \times 10^4, 10^5, 5 \times 10^5, \text{ and } 10^6 \text{ cm}^{-3}$. We compared the model abundances from NAUTILUS calculated at a particular choice of H_2 density with those calculated from RADEX at the same H_2 density. If NAUTILUS was able to produce an HCN abundance, HCO^+ abundance, and HCN/HCO^+ abundance ratio that equaled those of RADEX then we considered a solution (or a ‘match’) had been found. To decide which H_2 density provided the best solution, we performed this comparison for each H_2 density and chose the density for which the match between NAUTILUS and RADEX occurs at approximately the same time (or with the smallest time difference between them). To provide an additional constraint on the models, we required that the time at which the HCN/HCO^+ abundance ratio matched the RADEX solution fall between the times when the individual HCN and HCO^+ abundances matched the observations.

4.3.2 Results of the modelling

Fig. 8 shows the modelling performed for three specific positions in W40: the Green Circle, Blue Square, and Red Triangle in Fig. 2. These positions were chosen since they have disparate physical conditions in which to investigate abundance variations. In particular, Fig. 8 shows the time evolution of HCN (red), HCO^+ (blue), and CO (black; scaled down by a factor of 1000) abundances, and the abundance ratio (orange). In all three cases, the time interval between when the HCN and HCO^+ abundances match (i.e. the span of the vertical shaded region) is relatively small, and the time at which the HCN/HCO^+ abundance ratio matches (vertical dotted line) falls within this interval, though near its outer limit. Note that while HCN crosses the observed abundance value twice, we adopt the second crossing time as the correct solution, since it also corresponds to the solution for the HCN/HCO^+ ratio. Results for the specific positions are discussed individually below. Appendix A (Figs A1, A2, and A3) presents example model results across a range of densities and illustrates how specific density solutions were evaluated, showing why some were rejected and others accepted as the best match to the observed abundances.

The Green Circle – the low $N(\text{H}_2)$ and high T_K regime The Green Circle represents a position with a low H_2 column density ($N(\text{H}_2) = 9.9 \times 10^{21} \text{ cm}^{-2}$; $A_V = 4.7$) and high temperature ($T_K = 31.9\text{K}$). The best solution for this position occurs for an H_2 density of 10^5 cm^{-3} . At this density the RADEX derived abundances are $X(\text{HCN}) = 1.72 \times 10^{-8}$ and $X(\text{HCO}^+) = 1.33 \times 10^{-9}$ which are close to average. The HCN/HCO^+ abundance ratio, however, is relatively large (12.9). In this NAUTILUS model, both observed abundances and the abundance ratio are reached at $\sim 4\text{--}6.5 \times 10^4 \text{ yr}$. It is important to emphasize that we are not

interpreting this as the actual age of the cloud. As noted by J. M. C. Rawlings, E. Keto & P. Caselli (2024), chemical ages have limited meaning when modelling a dynamically evolving cloud under fixed physical conditions. Our goal is simply to demonstrate that if the HCN and HCO^+ emission arise from the same gas under identical physical and dynamical conditions, then the precise evolutionary time is not critical. What matters is the ability to simultaneously reproduce the observed abundances of HCN and HCO^+ . The results for the best NAUTILUS model are given in the top panel of Fig. 8.

The Blue Square – the moderate $N(\text{H}_2)$ and T_K regime The Blue Square represents a position with a moderate H_2 column density ($N(\text{H}_2) = 2.9 \times 10^{22} \text{ cm}^{-2}$; $A_V = 13.8$) and temperature ($T_K = 28.4\text{K}$). The best solution for this position occurs for an H_2 density of $5 \times 10^5 \text{ cm}^{-3}$. At this density the RADEX derived abundances are $X(\text{HCN}) = 1.29 \times 10^{-8}$ and $X(\text{HCO}^+) = 1.29 \times 10^{-9}$ which are similar to those seen at the position of the Green Circle (although the Blue Square position has a higher H_2 density). The HCN/HCO^+ abundance ratio, is also relatively large (10.0). In this NAUTILUS model, both observed abundances and the abundance ratio are reached at $\sim 3.2\text{--}4.6 \times 10^4 \text{ yr}$. The best matching solution is shown in the middle panel of Fig. 8.

The Red Triangle – the high $N(\text{H}_2)$ and low T_K regime The Red Triangle represents a position with a high H_2 column density ($N(\text{H}_2) = 4.6 \times 10^{22} \text{ cm}^{-2}$; $A_V = 21.9$) and a low temperature ($T_K = 19.6\text{K}$). It is also furthest from the stellar cluster. The best solution for this position occurs for an H_2 density of $5 \times 10^4 \text{ cm}^{-3}$. At this density the RADEX derived abundances are $X(\text{HCN}) = 1.27 \times 10^{-8}$ and $X(\text{HCO}^+) = 3.13 \times 10^{-9}$. While the HCN abundance is similar to that seen at the positions of the Green Circle and the Blue Square, the HCO^+ abundance here is a factor of ~ 3 higher resulting in a fairly low HCN/HCO^+ abundance ratio (4.1). In this NAUTILUS model, both observed abundances and the abundance ratio are reached at $\sim 1.5\text{--}2.4 \times 10^5 \text{ yr}$. The results for this NAUTILUS model are given in the bottom panel of Fig. 8.

The results for all three positions are summarized in Table 3 that shows that, in W40, the HCN abundance ranges from $X(\text{HCN}) = 1.27\text{--}1.72 \times 10^{-8}$, and HCO^+ abundance varies from $X(\text{HCO}^+) = 1.29\text{--}3.13 \times 10^{-9}$ and the $X(\text{HCN})/X(\text{HCO}^+)$ ratio varies from 4.1 – 12.9. Thus, allowing the density to vary from $5 \times 10^4\text{--}5 \times 10^5 \text{ cm}^{-3}$ results in smaller abundance variations than is seen when the density is held constant (see Fig. 7 and Table 2).

Density variations of this magnitude are reasonable within molecular clouds. For example, a study of the $\text{CS } J = 1 \rightarrow 0, J = 2 \rightarrow 1, \text{ and } J = 5 \rightarrow 4$ transitions of a number of selected clumps within Orion A found that the density varied from $10^4\text{--}2.2 \times 10^5 \text{ cm}^{-3}$ (K. Tatematsu et al. 1998). Also, using a simple assumption that the depth of the H_2 molecular cloud is equivalent to its width ($\approx 0.1\text{pc}$ —see Fig. 2) the H_2 volume density in W40 would vary from $\approx 3 \times 10^4\text{--}1.5 \times 10^5 \text{ cm}^{-3}$. Under this simplifying assumption of a constant line-of-sight depth (0.1 pc), the Red Region, having the highest column density, should also exhibit the highest volume density, not the lowest as reported in Table 3. However, an alternative geometry cannot be ruled out; the Red Region may trace the edge of a limb-brightened shell wrapping around the stellar cluster rather than a simple column. In this scenario, low H_2 volume density could exist in a region of high column density.




Figure 8. NAUTILUS model results for the three positions shown in Fig. 2 (Green Circle – top panel; Blue Square – middle panel; and Red Triangle – bottom panel) showing the time evolution of HCN (red), HCO⁺ (blue), and CO (black) abundances, and the HCN/HCO⁺ abundance ratio (orange). For ease of comparison, the CO abundances have been scaled down by a factor of 1000 to plot them on the same scale as HCN and HCO⁺. The horizontal dotted lines indicate the observed HCN (red dotted) and HCO⁺ (blue dotted) abundances, and the abundance ratio (orange dotted) as determined from RADEX. The horizontal shaded regions indicate the uncertainties in the abundances. The vertical shaded region shows the time between when the RADEX and NAUTILUS HCO⁺ abundances match and when the HCN abundances match. The vertical grey dotted line indicates the time at which the HCN/HCO⁺ abundance ratio from the NAUTILUS model matches the observed RADEX value. The orange axis on the right shows the value of HCN/HCO⁺ abundance ratio.

Table 3. Best model results for the HCN and HCO⁺ abundances.

| Position | A_V (mag) | T_K (K) | $n(\text{H}_2)$ (cm^{-3}) | Model time (yr) | $N(\text{HCN})$ ($\times 10^{14}$) cm^{-2} | $X(\text{HCN})$ ($\times 10^{-8}$) | $N(\text{HCO}^+)$ ($\times 10^{13}$) cm^{-2} | $X(\text{HCO}^+)$ ($\times 10^{-9}$) | $\frac{X(\text{HCN})}{X(\text{HCO}^+)}$ |
|--------------|----------------|--------------|---|-------------------------|--|---|--|---|---|
| Green Circle | 4.7 | 31.9 | 10^5 | $4 - 6.5 \times 10^4$ | 1.70 ± 0.4 | 1.72 ± 0.5 | 1.31 ± 0.4 | 1.33 ± 0.5 | 12.9 ± 6.1 |
| Blue Square | 13.8 | 28.4 | 5×10^5 | $3.2 - 4.6 \times 10^4$ | 3.75 ± 0.3 | 1.29 ± 0.2 | 3.75 ± 0.3 | 1.29 ± 0.2 | 10.0 ± 2.2 |
| Red Triangle | 21.9 | 19.6 | 5×10^4 | $1.5 - 2.4 \times 10^5$ | 5.87 ± 1.7 | 1.27 ± 0.4 | 14.44 ± 2.4 | 3.13 ± 0.6 | 4.1 ± 1.5 |

4.3.3 Effects of density on HCN and HCO⁺ abundances and line strengths

The H₂ density affects the time-evolution of each species due to their different formation and destruction pathways. The primary formation route for HCO⁺ at this temperature is



and the main destruction pathway is



Thus, for HCO⁺, the important factors are the amount of CO available to form HCO⁺, the amount of neutral atomic carbon available to destroy it, and the times at which these two forms dominate the carbon budget. For low densities (e.g. $n(H_2) = 5 \times 10^4 \text{ cm}^{-3}$), neutral atomic C is the dominant form of carbon between 10^2 and 3×10^4 yr. However, at higher densities (e.g. $n(H_2) = 5 \times 10^5 \text{ cm}^{-3}$), neutral atomic C is the dominant form of carbon at an earlier time, i.e. between 10 and 3×10^3 yr. Therefore, at higher densities, the conversion from C to CO occurs very early in the simulation so one simultaneously loses the (destructive) neutral atomic carbon and increases the amount of (productive) CO at much earlier times than one does at low densities.

In addition, as shown in Fig. 8, the blue position – with the highest density – reaches a large CO abundance earliest in the simulation (at ~ 6500 yr, before it temporarily depletes), allowing the HCO⁺ abundance to build up quickly and match the observed value earliest in the simulations. The red position – with the lowest density – reaches its maximum CO abundance much later (4.6×10^4 yr), and therefore reproduces the observed HCO⁺ abundance latest. As expected, the green position – with intermediate density – reaches the observed HCO⁺ abundance at a time between the blue and red positions. The overall result is that the modelled HCO⁺ abundance agrees with the observed value at earlier times in regions of higher-density gas. It is also noteworthy that the green and Blue Regions do not undergo the same CO depletion seen in the red position, likely due to their elevated temperatures.

As the H₂ density increases, the chemical model matches the observed HCN abundances at earlier times as well. This is likely because the main formation pathway for HCN is:



and the main destruction pathway is:



At higher densities there is more CH₂ at early times to efficiently form HCN and, simultaneously, less C⁺ to destroy HCN (the C⁺ combines with free electrons to form neutral carbon, affecting the HCO⁺ abundance). In comparison, in their study of CN and HCN in the Serpens cloud, A. Mirocha et al. (2021) found that equations (3) and (4) are not the dominant HCN reactions for a UV field strength of $G_0 = 237$ in $A_V = 5$ gas. In their models, these are the dominant reactions for much weaker fields (i.e. $G_0 = 10^{-3} - 10^{-1}$). The difference may lie in the fact that they used a gas temperature of 50 K in their models that can change which reactions dominate.

To investigate the effect of density on HCN and HCO⁺ line strengths, we again used RADEX to model the expected HCN and HCO⁺ J = 3 → 2 integrated intensities but now using the abundances and densities from Table 3. We modelled the HCN

and HCO⁺ J = 3 → 2 emission from the Red Triangle (the peak of the H₂ map) and Blue Square (near the peak of the HCN and HCO⁺ emission). For the Red Triangle the model produces T_{mb} (HCN J = 3 → 2) = 1.9 K and T_{mb} (HCO⁺ J = 3 → 2) = 4.4 K, which is very close to the observed values of 1.8 K and 4.5 K respectively. For the Blue Square, the model produces T_{mb} (HCN J = 3 → 2) = 7.5 K and T_{mb} (HCO⁺ J = 3 → 2) = 6.9 K, which is comparable to the observed values of 6.4 and 6.8 K, respectively. Thus, the reason the HCN and HCO⁺ integrated intensity maps peak in the lower H₂ column density Blue Region, rather than in the high H₂ column density Red Region is due to excitation and abundance effects. Our RADEX results are therefore consistent with a lower density in the Red Region.

4.4 Comparison with previous work

Fig. 9 shows our best-fitting abundances (Table 3) alongside literature values (Table 1) for comparison. The HCN abundances in W40 (Table 3) are roughly an order of magnitude *higher* than those found in low-mass protostars and in W51, but comparable to the Young Stellar Object GL2591. They also fall within the lower range of values reported for LIRGs and ULIRGs. The enhanced HCN abundances in these extreme systems have been attributed by Y. Nishimura et al. (2024) to chemical processing in shocks.

The HCO⁺ abundances we derive are a factor of a few, to an order of magnitude, *lower* than those reported in most other studies, with the exception of W51 and 70 μm – quiet sources. One caveat for W51 is that Y. Watanabe et al. (2017) adopt a uniform gas temperature of 20 K across the cloud, whereas we use dust-based temperature maps that provide T_K on a pixel-by-pixel basis.

Despite variations in the absolute HCN and HCO⁺ abundances across sources, the HCN/HCO⁺ abundance ratios remain relatively consistent, differing by only a factor of a few between studies. AGN and ULIRG sources, however, show a broader range (e.g. M. Imanishi et al. 2023a, b; Y. Nishimura et al. 2024; J. Butterworth et al. 2025), which could be attributed to elevated HCN abundances in some systems – potentially driven by enhanced cosmic-ray ionization (J. Butterworth et al. 2025) or, again, by chemical processing in shocks (Y. Nishimura et al. 2024).

Thus, our modelling of the HCN and HCO⁺ abundances have revealed a few interesting results:

(i) The HCN and HCO⁺ abundances in W40 differ from those reported in both low-mass and high-mass star-forming regions, as well as in LIRGs and ULIRGs. These discrepancies may arise from variations in physical conditions (e.g. H₂ density, temperature, or the presence of shocks), which can alter the rates and efficiencies of key chemical reactions.

(ii) The HCN/HCO⁺ abundance ratios are comparatively uniform across sources and appear to be less sensitive to differences in physical conditions.

(iii) Abundance maps produced under the assumption of a constant density – such as those in Fig. 7, which suggest variations of nearly a factor of ~ 20 across W40 – may not accurately represent the true chemical structure of the region.

(iv) By incorporating a priori knowledge of the physical environment (notably $N(H_2)$ and T_K from the HGBS maps), the combined use of RADEX and NAUTILUS enables us to estimate the H₂ volume density and derive self-consistent HCN and HCO⁺




Figure 9. Comparison of HCN and HCO⁺ abundances and abundance ratios in W40 with values reported for a range of Galactic and extragalactic environments. *Top:* fractional HCN abundance $X(\text{HCN})$ relative to H₂. *Middle:* fractional HCO⁺ abundance $X(\text{HCO}^+)$. *Bottom:* abundance ratio $X(\text{HCN})/X(\text{HCO}^+)$. The coloured symbols with error bars show the three W40 positions (Green Circle, Blue Square, and Red Triangle), while the black symbols show literature values for low-mass protostars; W51; GL 2591; quiescent and active IRDCs; quiescent, protostellar, and HII/PDR ATLASGAL clumps; 70 μm-quiet clumps; and integrated measurements for starburst galaxies, AGNs, ULIRGs (J. Butterworth et al. 2025), and a combined study of LIRGs + ULIRGs (Y. Nishimura et al. 2024). See Table 1 for complete references for each of the columns. The vertical error bars denote 1σ uncertainties for the W40 points and the range of values reported for each comparison sample, where available. The abundance panels (top and middle) use logarithmic y-axes, while the ratio panel (bottom) uses a linear scale.

abundances in W40 that remain relatively uniform across the cloud.

4.5 Effects of the UV field and cosmic-ray ionization rate

The abundances of many species including HCN and HCO⁺ may be affected by the strength of the UV field since, in increased UV and/or low A_V environments, photochemical reactions may be enhanced. The NAUTILUS code calculates the rate coefficients for photochemical processes via the following equation

(V. Wakelam et al. 2012):

$$k_{photo} = Ae^{-CA_V} s^{-1}, \quad (5)$$

where A is the unattenuated photodissociation rate, which scales with the strength of the external UV field, C is a coefficient that accounts for the photodissociation/photoionization threshold of the molecular species, and A_V is the visual extinction produced by the dust. Thus, the exponential factor accounts for dust attenuation of the external UV field. Increasing the external UV field will therefore increase the parameter A , which increases the rate

coefficient. Conversely, increasing A_V will increase the attenuation of the UV field thereby decreasing the rate coefficient.

The value of A_V is fixed in each pixel from the **HGBS** maps of $N(\text{H}_2)$. The strength of the UV field is likewise fixed by the Herschel observations that show that the average strength of the UV field in W40 is $237 G_o$ (N. Schneider et al. (2020); and references therein). However, the same observations show that the UV field strength can reach $\sim 8200 G_o$ near the OB association. These higher UV field strengths might be important, particularly at the position of the Green Circle (Fig. 2), which is closest to the OB association, where the value of A_V is lowest, and where the porous and fractal properties of molecular clouds may allow for increased UV penetration.

To test the effects of a stronger UV field on our results, we reran the NAUTILUS code for each of the three selected positions using a UV scaling factor of 8200. The results for the Blue Square and Red Triangle positions are unchanged. This is unsurprising given the large value of A_V at these positions (13.3 and 21.9, respectively) that attenuates the radiation field by a factor of 1.7×10^{-6} and 3×10^{-10} , respectively, meaning that photoprocesses are unimportant for either choice of the UV field strength. A similar conclusion was reached by W. A. Baan, A. F. Loenen & M. Spaans (2010) who modelled the chemical evolution of the ISM in luminous IR galaxies and found that in high column density regions, changes in the UV radiation field only leads to small changes in the HCN/ HCO^+ line ratios, because the UV flux is largely attenuated.

At the position of the Green Circle, a radiation field of $8200 G_o$, however, is unable to produce the observed HCN and HCO^+ abundances. In particular, over the timespan of the simulation, the model produces less HCN (up to a factor of ~ 80) and HCO^+ (up to a factor of ~ 20) than is observed. This is likely because the value of A_V is only 4.7, which provides an attenuation of only 9×10^{-3} , meaning that photoprocesses are likely important. The main processes lowering the abundances of HCN and HCO^+ are not direct photodissociations as might be expected but, instead, are equation (4) and



for HCN, and



for HCO^+ . The magnitude of the UV field, coupled with the weak attenuation, means that photoprocesses produce an ample amount of C^+ , H^+ , and e^- to efficiently destroy both species through gas-phase reactions.

These results are slightly different from those found by A. Mirocha et al. (2021) using $G_o = 1 - 10^6$, $T_K = 50$ K, and $A_V = 5$, in which the dominant destruction reaction is




instead of equations (4) and (6). In their models, equations (4) and (6) are the dominant destruction reactions for much weaker fields (i.e. $G_o = 10^{-3} - 10^{-1}$). Again the difference may be due to the higher temperatures that they use in their simulations.

We also explored the value of A_V required to significantly reduce the impact of the UV field and, by $A_V = 6$, the HCN abundance is decreased by, at most, a factor of ~ 3 and HCO^+ by ~ 1.5 . Note, however, that only 54 pixels (16 per cent) have values of $A_V \leq 6$. Of these, 42 lie to the left and below of the Green Region, 12 lie within the Green Region (including the pixel corresponding to the Green Circle), and none lie in the Blue or Red

Regions. Thus, given the high value of A_V across most of W40, enhanced UV field strengths are probably irrelevant to our overall conclusions.

In order to match the observed abundances and the abundance ratio at the position of the Green Circle using a UV field of $8200 G_o$ and the observed A_V of 4.7, we have to increase the H_2 density by a factor of 10. For a density of $n(\text{H}_2) = 10^6 \text{ cm}^{-3}$ the RADEX derived abundances are $X(\text{HCN}) = 1.3 \times 10^{-9}$ and $X(\text{HCO}^+) = 2.0 \times 10^{-10}$, which are 13.2 times and 6.7 times (respectively) lower than the values in Table 3 for $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$, but, oddly, similar to the abundances found by A. Traficante et al. (2017) in $70 \mu\text{m}$ **quiet** sources, and in W51 (Y. Watanabe et al. 2017) (albeit for a different H_2 density). The fact that the HCN abundance decreases more than that of HCO^+ is probably due to the fact that, for the first 10^5 yr, the HCN destruction rate from equation (4) is an order of magnitude larger than that for HCO^+ (equation (7)). The HCN/ HCO^+ abundance ratio therefore is decreased by a factor ~ 2 from 12.9 to 6.7. In this NAUTILUS model, both observed abundances and the abundance ratio are reached at $\sim 10^4$ yr.

The HCO^+ abundance is also sensitive to the cosmic-ray (CR) ionization rate, since its primary formation route is via H_3^+ (see equation 1) which is itself strongly dependent on the cosmic-ray flux. The two steps leading up to equation (1) are:



and



In our models we adopt a ‘standard’ cosmic-ray ionization rate of $\zeta_{\text{H}_2} = 1.3 \times 10^{-17} \text{ s}^{-1}$, which is appropriate for dense molecular clouds where attenuation reduces the CR flux (P. Caselli et al. 1998; M. Padovani, D. Galli & A. E. Glassgold 2009). However, observations along several lines of sight toward massive star-forming regions (N. Indriolo et al. 2015) indicate a higher mean value of $\zeta_{\text{H}_2} = 1.78 \times 10^{-16} \text{ s}^{-1}$. To test the impact of an enhanced CR ionization rate, we ran additional models using the N. Indriolo et al. (2015) value together with our fiducial UV field strength ($237 G_o$).

While there are differences in the time evolution of each species (and their ratio) between the two models, the principal effect of increasing ζ_{H_2} to $1.78 \times 10^{-16} \text{ s}^{-1}$ is that the steady-state abundances of both HCN and HCO^+ (at times $\gtrsim 10^5$ yr) are higher. However, with the elevated ionization rate, the Red Triangle and Green Circle density solutions remain unchanged (i.e. 5×10^5 and 10^5 cm^{-3} respectively). The Blue Square, however, is best reproduced by a NAUTILUS model with a density of 10^5 cm^{-3} rather than $5 \times 10^5 \text{ cm}^{-3}$. See Appendix B (Figs B1, B2, and B3) for further details on the model results using the enhanced cosmic-ray ionization rate.

From this analysis therefore we can draw some additional conclusions:

- (i) In the presence of a strong UV field and **high** A_V the abundances of HCN and HCO^+ are *not* affected due to attenuation of the radiation field in regions of high H_2 column density.
- (ii) In the presence of a strong UV field and **low** A_V the abundance of both HCN and HCO^+ are *strongly* affected. The HCN abundance, however, is decreased by a factor of ~ 2 more than the HCO^+ abundance.

(iii) Increasing the cosmic-ray ionization rate changes the detailed time evolution of HCN and HCO⁺ and produces higher steady-state abundances, but it does not strongly affect the density solutions of our models.

5 SUMMARY AND CONCLUSIONS

As part of a pilot project for the Massive, Active, JCMT-Observed Regions of Star Formation (MAJORS) Large Program at the JCMT, we mapped the HCN and HCO⁺ $J = 3 \rightarrow 2$ transitions across the central 424 arcsec \times 424 arcsec region of the massive star-forming complex W40. Using results from the Herschel Gould Belt Survey (HGBS), we assigned a kinetic temperature (assuming $T_K = T_{\text{dust}}$) and an H₂ column density $N(\text{H}_2)$ to every pixel in our HCN and HCO⁺ maps. With T_K and the integrated intensities of HCN and HCO⁺ as inputs to the RADEX 1D radiative transfer code, we computed the corresponding HCN and HCO⁺ column densities for a range of H₂ densities (10^4 – 10^6 cm⁻³). Abundances in each pixel were then obtained by dividing these column densities by the HGBS-derived values of $N(\text{H}_2)$. Because the RADEX-derived column densities and abundances depend on the assumed H₂ density, we subsequently used the NAUTILUS gas–grain chemical evolution code to determine the H₂ density at which the NAUTILUS abundances, and the HCN/HCO⁺ abundance ratios, match those obtained from RADEX.

The main results from our study are as follows:

(i) The bulk of the HCN and HCO⁺ emission in W40 is not co-located with the region of maximum H₂ column density but, instead, arises in the Blue Region (see Fig. 2) which, from our modelling, contains the highest H₂ volume density and intermediate dust temperatures (i.e. 26–28 K). In addition, the ratio of the HCN to HCO⁺ integrated intensities, $I(\text{HCN})/I(\text{HCO}^+)$, increases with H₂ density. However, care must be taken in assigning a specific line intensity and $I(\text{HCN})/I(\text{HCO}^+)$ ratio to a specific H₂ density, since different excitation conditions and molecular abundances will produce different intensities and ratios at the same density.

(ii) By incorporating a priori knowledge of the physical conditions (notably the $N(\text{H}_2)$ and T_K data from the HGBS maps) and assuming a constant density of $n(\text{H}_2) = 10^5$ cm⁻³ we used RADEX to calculate the HCN and HCO⁺ abundances in every pixel in our W40 map. The HCN abundances range from 0.4 – 7.0×10^{-8} and the HCO⁺ abundances range from 0.4 – 7.3×10^{-9} . The corresponding HCN/HCO⁺ abundance ratio range is 2.6–23.1.

(iii) Using the NAUTILUS chemical code to find the H₂ density that produces the best match to the RADEX abundances (rather than a constant density) suggests, instead, that the HCN and HCO⁺ abundances may be fairly constant across W40. Careful modelling of three different positions (with disparate physical conditions) finds $X(\text{HCN}) = 1.3$ – 1.7×10^{-8} , $X(\text{HCO}^+) = 1.3$ – 3.1×10^{-9} . The HCN/HCO⁺ abundance ratio ranges from 4.1–12.9.

(iv) By cross-comparing the results of RADEX and NAUTILUS, we obtain a crude estimate of the gas density responsible for the HCN and HCO⁺ emission, finding H₂ densities in the range 5×10^4 – 5×10^5 cm⁻³.

(v) High UV intensity has little effect on the HCN and HCO⁺ abundance in regions where the visual extinction is large enough to effectively shield the gas from the UV field. In regions where A_V is low (i.e. < 6), however, the abundance of both species is

lowered due to destructive reactions with species which are directly affected by the radiation field (e.g. H⁺, C⁺, and e⁻). In such regions, our models indicate that the HCN abundance decreases by a factor of ~ 13 whereas the HCO⁺ abundance decreases by a factor of 6. Thus, decreased HCN and HCO⁺ abundances, coupled with decreased HCN/HCO⁺ abundance ratios, may indicate the presence of a strong UV field (and low extinction). Care must be taken with this interpretation, however, since variations in these quantities can also be a result of varying physical conditions (e.g. $n(\text{H}_2)$ and T_K) in a weak UV field.

ACKNOWLEDGEMENTS

The James Clerk Maxwell Telescope is operated by the East Asian Observatory on behalf of The National Astronomical Observatory of Japan; Academia Sinica Institute of Astronomy and Astrophysics; the Korea Astronomy and Space Science Institute; centre for Astronomical Mega-Science (as well as the National Key R&D Program of China with No. 2017YFA0402700). Additional funding support is provided by the Science and Technology Facilities Council of the United Kingdom and participating universities in the United Kingdom and Canada. The James Clerk Maxwell Telescope has historically been operated by the Joint Astronomy Centre on behalf of the Science and Technology Facilities Council of the United Kingdom, the National Research Council of Canada and the Netherlands Organisation for Scientific Research.

This research has made use of data from the Herschel Gould Belt survey (HGBS) project (<http://gouldbelt-herschel.cea.fr>). The HGBS is a Herschel Key Programme jointly carried out by SPIRE Specialist Astronomy Group 3 (SAG 3), scientists of several institutes in the PACS Consortium (CEA Saclay, INAF-IFSI Rome and INAF-Arcetri, KU Leuven, MPIA Heidelberg), and scientists of the Herschel Science centre (HSC). This publication also makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis centre/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

RP acknowledges the support of the Natural Sciences and Engineering Research Council of Canada (NSERC), through the Discovery Grant program and JC’s work was supported by the NSERC USRA program. Researchers at the University of Calgary also acknowledge and pay tribute to the traditional territories of the peoples of Treaty 7, which include the Blackfoot Confederacy (comprised of the Siksika, the Piikani, and the Kainai First Nations), the Tsuut’ina First Nation, and the Stoney Nakoda (including Chiniki, Bearspaw, and Goodstoney First Nations). The City of Calgary is also home to the Métis Nation of Alberta Region 3. The authors also wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

LCH was supported by the China Manned Space Program (CMS-CSST-2025-A09), the National Science Foundation of China (12233001), and the National Key R&D Program of China (2022YFF0503401). DA acknowledges NU FCD RP grant 201223FDD8821. PS was partially supported by a Grant-in-Aid for Scientific Research (KAKENHI Number JP22H01271 and JP23H01221) of JSPS. PS was supported by Yoshinori Ohsumi Fund (Yoshinori Ohsumi Award for Fundamental Research). KP is a Royal Society University Research Fellow, supported by

grant number URF\R1\211322. SER acknowledge support from consolidated grant (ST\W000830\1) from the UK Science and Technology Facilities Council (STFC). The work of MGR is supported by the international Gemini Observatory, a program of NSF NOIRLab, which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the U.S. National Science Foundation, on behalf of the Gemini partnership of Argentina, Brazil, Canada, Chile, the Republic of Korea, and the United States of America. CWL is supported by the Basic Science Research Program through the NRF funded by the Ministry of Education, Science and Technology (grant No. NRF-2019R1A2C1010851) and by the Korea Astronomy and Space Science Institute grant funded by the Korea government (MSIT; project No. 2024-1-841-00). MGB acknowledges the support of the Department for Communities in Northern Ireland for support of Armagh's contribution to the JCMT and to this research programme. AK acknowledges support from the Polish National Science Center grant No. 2024/54/E/ST9/00314.

DATA AVAILABILITY

The MAJORS survey and its data products are described fully in Eden et al. (in preparation). The associated download instructions are within that paper, or per request to the lead author. Observational data from the JCMT are publicly available through the Canadian Astronomy Data Centre (CADC) at <https://www.cadc-ccda.hia-ihp.nrc-cnrc.gc.ca/en/> under the proposal IDs M22AL002 and M20AD003.

REFERENCES

- André P. et al., 2010, *A&A*, 518, L102
 Arzoumanian D. et al., 2019, *A&A*, 621, A42
 Baan W. A., Loenen A. F., Spaans M., 2010, *A&A*, 516, A40
 Berry D. S., Reinhold K., Jenness T., Economou F., 2007, in Shaw R. A., Hill F., Bell D. J., eds, ASP Conf. Ser. Vol. 376, *Astronomical Data Analysis Software and Systems XVI*. Astron. Soc. Pac., San Francisco, p. 425
 Bohlin R. C., Savage B. D., Drake J. F., 1978, *ApJ*, 224, 132
 Bontemps S. et al., 2010, *A&A*, 518, L85
 Braine J., Sun Y., Shimajiri Y., van der Tak F. F. S., Fang M., André P., Chen H., Gao Y., 2023, *A&A*, 676, A27
 Butterworth J., Viti S., Wang Y., 2025, *A&A*, 697, A18
 Caselli P., Walmsley C. M., Terzieva R., Herbst E., 1998, *ApJ*, 499, 234
 Chapin E., Gibb A. G., Jenness T., Berry D. S., Scott D., Tilanus R. P. J., 2013, *Starlink User Note*, 258
 Comerón F., Djupvik A. A., Schneider N., 2022, *A&A*, 665, A76
 Currie M. J., Berry D. S., 2014, *Astrophysics Source Code Library*, record ascl:1403.022
 Currie M. J., Berry D. S., Jenness T., Gibb A. G., Bell G. S., Draper P. W., 2014, in Manset N., Forshay P., eds, ASP Conf. Ser. Vol. 485, *Astronomical Data Analysis Software and Systems XXIII*. Astron. Soc. Pac., San Francisco, p. 391
 Endres C. P., Schlemmer S., Schilke P., Stutzki J., Müller H. S. P., 2016, *J. Mol. Spect.*, 327, 95
 Evans N. J. II, Heiderman A., Vutisalchavakul N., 2014, *ApJ*, 782, 114
 Evans N. J. II, Kim K.-T., Wu J., Chao Z., Heyer M., Liu T., Nguyen-Lu'ong Q., Kauffmann J., 2020, *ApJ*, 894, 103
 Fontani F., Giannetti A., Beltrán M. T., Dodson R., Rioja M., Brand J., Caselli P., Cesaroni R., 2012, *MNRAS*, 423, 2342
 Friesen R. K. et al., 2017, *ApJ*, 843, 63
 Gao Y., Solomon P. M., 2004a, *ApJS*, 152, 63
 Gao Y., Solomon P. M., 2004b, *ApJ*, 606, 271
 Goldreich P., Kwan J., 1974, *ApJ*, 189, 441
 Goldsmith P. F., 2001, *ApJ*, 557, 736
 Griffin M. J. et al., 2010, *A&A*, 518, L3
 Habing H. J., 1968, *Bull. Astron. Inst. Netherlands*, 19, 421
 Heiderman A., Evans N. J. II, Allen L. E., Huard T., Heyer M., 2010, *ApJ*, 723, 1019
 Hoq S. et al., 2013, *ApJ*, 777, 157
 Imanishi M., Baba S., Nakanishi K., Izumi T., 2023a, *ApJ*, 950, 75
 Imanishi M., Baba S., Nakanishi K., Izumi T., 2023b, *ApJ*, 954, 148
 Indriolo N. et al., 2015, *ApJ*, 800, 40
 Jenness T., Currie M. J., Tilanus R. P. J., Cavanagh B., Berry D. S., Leech J., Rizzi L., 2015, *MNRAS*, 453, 73
 Jørgensen J. K., Schöier F. L., van Dishoeck E. F., 2004, *A&A*, 416, 603
 Kauffmann J., Goldsmith P. F., Melnick G., Tolls V., Guzman A., Menten K. M., 2017, *A&A*, 605, L5
 Kerr A. R., Pan S.-K., Claude S. M. X., Dindo P., Lichtenberger A. W., Effland J. E., Lauria E. F., 2014, *IEEE T. Terahertz Sci. Technol.*, 4, 201
 Könyves V. et al., 2015, *A&A*, 584, A91
 Kruegel E., Walmsley C. M., 1984, *A&A*, 130, 5
 Krumholz M. R., Dekel A., McKee C. F., 2012, *ApJ*, 745, 69
 Kuhn M. A., Getman K. V., Feigelson E. D., Reipurth B., Rodney S. A., Garmire G. P., 2010, *ApJ*, 725, 2485
 Lada C. J., Lombardi M., Alves J. F., 2010, *ApJ*, 724, 687
 Lada C. J., Forbrich J., Lombardi M., Alves J. F., 2012, *ApJ*, 745, 190
 Loughnane R. M., Redman M. P., Thompson M. A., Lo N., O'Dwyer B., Cunningham M. R., 2012, *MNRAS*, 420, 1367
 McMullin J. P., Waters B., Schiebel D., Young W., Golap K., 2007, in Shaw R. A., Hill F., Bell D. J., eds, ASP Conf. Ser. Vol. 376, *Astronomical Data Analysis Software and Systems XVI*. Astron. Soc. Pac., San Francisco, p. 127
 Mendigutía I., Lada C. J., Oudmaijer R. D., 2018, *A&A*, 618, A119
 Mirocha A. et al., 2021, *A&A*, 656, A146
 Müller H. S. P., Thorwirth S., Roth D. A., Winnewisser G., 2001, *A&A*, 370, L49
 Müller H. S. P., Schlöder F., Stutzki J., Winnewisser G., 2005, *J. Mol. Struct.*, 742, 215
 Mullins A. M., Loughnane R. M., Redman M. P., Wiles B., Guegan N., Barrett J., Keto E. R., 2016, *MNRAS*, 459, 2882
 Nguyen-Luong Q. et al., 2020, *ApJ*, 891, 66
 Nishimura Y. et al., 2024, *A&A*, 686, A48
 Öberg K. I., Fuchs G. W., Awad Z., Fraser H. J., Schlemmer S., van Dishoeck E. F., Linnartz H., 2007, *ApJ*, 662, L23
 Onus A., Krumholz M. R., Federrath C., 2018, *MNRAS*, 479, 1702
 Padovani M., Galli D., Glassgold A. E., 2009, *A&A*, 501, 619
 Patra S., Evans N. J. II, Kim K.-T., Heyer M., Kauffmann J., Jose J., Samal M. R., Das S. R., 2022, *AJ*, 164, 129
 Pilbratt G. L. et al., 2010, *A&A*, 518, L1
 Plume R., Jaffe D. T., Evans N. J. II, Martín-Pintado J., Gómez-González J., 1997, *ApJ*, 476, 730
 Poglitsch A. et al., 2010, *A&A*, 518, L2
 Rachford B. L. et al., 2009, *ApJS*, 180, 125
 Rawlings J. M. C., Keto E., Caselli P., 2024, *MNRAS*, 530, 3986
 Rodney S. A., Reipurth B., 2008, in Reipurth B., ed., *Handbook of Star Forming Regions*, Vol. II. The Southern Sky ASP Monograph Publications, p. 683
 Roy A. et al., 2014, *A&A*, 562, A138
 Ruaud M., Wakelam V., Hersant F., 2016, *MNRAS*, 459, 3756
 Sabatini G., Giannetti A., Bovino S., Brand J., Leurini S., Schisano E., Pillai T., Menten K. M., 2019, *MNRAS*, 490, 4489
 Sanhueza P., Jackson J. M., Foster J. B., Garay G., Silva A., Finn S. C., 2012, *ApJ*, 756, 60
 Santa-Maria M. G. et al., 2023, *A&A*, 679, A4
 Schneider N. et al., 2020, *PASP*, 132, 104301
 Shirley Y. L., 2015, *PASP*, 127, 299
 Shuping R. Y., Vacca W. D., Kassis M., Yu K. C., 2012, *AJ*, 144, 116
 Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
 Stephens I. W., Jackson J. M., Whitaker J. S., Contreras Y., Guzmán A. E., Sanhueza P., Foster J. B., Rathborne J. M., 2016, *ApJ*, 824, 29
 Straizys V., Černis K., Bartašiūtė S., 2003, *A&A*, 405, 585

- Tan Q.-H. et al., 2018, *ApJ*, 860, 165
- Tatematsu K., Umemoto T., Heyer M. H., Hirano N., Kameya O., Jaffe D. T., 1998, *ApJS*, 118, 517
- Thomas H. S., Fuller G. A., 2008, *A&A*, 479, 751
- Traficante A., Fuller G. A., Billot N., Duarte-Cabral A., Merello M., Molinari S., Peretto N., Schisano E., 2017, *MNRAS*, 470, 3882
- van der Tak F. F. S., van Dishoeck E. F., Evans N. J. II, Bakker E. J., Blake G. A., 1999, *ApJ*, 522, 991
- van der Tak F. F. S., Black J. H., Schöier F. L., Jansen D. J., van Dishoeck E. F., 2007, *A&A*, 468, 627
- Wakelam V. et al., 2012, *ApJS*, 199, 21
- Walker-Smith S. L., Richer J. S., Buckle J. V., Hatchell J., Drabek-Maunder E., 2014, *MNRAS*, 440, 3568
- Watanabe Y., Nishimura Y., Harada N., Sakai N., Shimonishi T., Aikawa Y., Kawamura A., Yamamoto S., 2017, *ApJ*, 845, 116
- Whittet D. C. B., 1981, *MNRAS*, 196, 469
- Wu J., Evans N. J. II, Gao Y., Solomon P. M., Shirley Y. L., Vanden Bout P. A., 2005, *ApJ*, 635, L173
- Zhang Z.-Y., Gao Y., Henkel C., Zhao Y., Wang J., Menten K. M., Güsten R., 2014, *ApJ*, 784, L31
- Zhu H., Tian W., Li A., Zhang M., 2017, *MNRAS*, 471, 3494

APPENDIX A: MODEL FITTING DETAILS

In this appendix, we present the model results for each of our three positions at three different densities and demonstrate how the best-matching solution is selected. Descriptions are provided in the figure captions.




Figure A1. NAUTILUS model results for the Green Circle showing the time evolution of HCN (red), HCO⁺ (blue), and CO (black) abundances, and the HCN/HCO⁺ abundance ratio (orange) for three different H₂ densities – 5 × 10⁴ (top), 10⁵ (middle), and 5 × 10⁵ (bottom) cm⁻³. This figure follows the same conventions as Fig. 8, with identical meanings for the vertical and horizontal dashed lines as well as the shaded regions. This figure shows that an H₂ density of 10⁵ cm⁻³ (middle) provides the best match to the observed abundances. At a density of 5 × 10⁴ cm⁻³ (top), the NAUTILUS model fails to reproduce the observed HCO⁺ abundance at any time. While the grey shaded regions for densities of 10⁵ cm⁻³ (middle) and 5 × 10⁵ cm⁻³ (bottom) span similar absolute time intervals (~ 2.5 × 10⁴ yr), we select the middle panel as the best match because the HCN crossing time is only 1.6 times larger than the HCO⁺ crossing time, compared with 4.1 times larger in the bottom panel.




Figure A2. Same as Fig. A1 except that results are shown for the Blue Square. This figure clearly shows that a density of $5 \times 10^5 \text{ cm}^{-3}$ (bottom) provides the best match to the abundances since, at the other two densities, the NAUTILUS model fails to reproduce the observed HCO⁺ abundances at any time.




Figure A3. Same as Fig. A1 except that results are shown for the Red Triangle. This figure clearly shows that a density of $5 \times 10^4 \text{ cm}^{-3}$ (top) provides the best match to the abundances since, at the other two densities, the NAUTILUS model fails to reproduce the observed HCN/HCO⁺ abundance ratios at any time.

APPENDIX B: ENHANCED COSMIC-RAY IONIZATION RATE MODELS

In this appendix, we present the model results for each of our three positions at three different densities using an enhanced cosmic-ray ionization rate of $\zeta_{H_2} = 1.78 \times 10^{-16} \text{ s}^{-1}$. Descriptions are provided in the figure captions.




Figure B1. NAUTILUS model results (using an enhanced cosmic-ray ionization rate of $\zeta_{H_2} = 1.78 \times 10^{-16} \text{ s}^{-1}$) for the Green Circle showing the time evolution of HCN (red), HCO⁺ (blue), and CO (black) abundances, and the HCN/HCO⁺ abundance ratio (orange) for three different H₂ densities: 5×10^4 (top), 10^5 (middle), and 5×10^5 (bottom) cm⁻³. This figure follows the same conventions as Fig. 8, with identical meanings for the vertical and horizontal dashed lines as well as the shaded regions. This figure shows that an H₂ density of 10^5 cm⁻³ (middle) provides the best match to the observed abundances. At a density of 5×10^4 cm⁻³ (top), the NAUTILUS model fails to reproduce the observed HCN abundance at any time, while the grey shaded region for a density of 5×10^5 cm⁻³ (bottom) spans a larger time interval than that for 10^5 cm⁻³ (middle). This density solution is the same as found for the ‘standard’ cosmic-ray ionization rate of $\zeta_{H_2} = 1.3 \times 10^{-17} \text{ s}^{-1}$ (see Fig. 8).




Figure B2. Same as Fig. B1 except that results are shown for the Blue Square. This figure clearly shows that a density of 10^5 cm^{-3} (middle) provides the best match to the abundances. The top panel ($5 \times 10^4 \text{ cm}^{-3}$) exhibits the largest time difference between the HCN and HCO⁺ model crossing times. The bottom panel ($5 \times 10^5 \text{ cm}^{-3}$) has the shortest time difference, but the abundance ratio falls outside this overlap window. Notably, this preferred density differs from the solution found for the ‘standard’ cosmic-ray ionization rate of $\zeta_{\text{H}_2} = 1.3 \times 10^{-17} \text{ s}^{-1}$, which favoured $5 \times 10^5 \text{ cm}^{-3}$ (see Fig. 8).




Figure B3. Same as Fig. B1 except that results are shown for the Red Triangle. This figure shows that there are two times at which the HCN/HCO⁺ abundance ratio matches the observed value. In the bottom panel, both ratio solutions fall within a broad time interval over which the HCN and HCO⁺ models cross the observed values – spanning nearly three orders of magnitude in time. In the middle panel, the difference between the HCN and HCO⁺ model crossing times exceeds two orders of magnitude. In the top panel, there are two distinct possible solutions: one between 1.2 and 8.4×10^4 yr and another between 1.5 and 3.4×10^5 yr, each spanning a factor of a few in time. Thus, as in the case of the ‘standard’ cosmic-ray ionization rate (Fig. 8), the best-fitting density is $5 \times 10^4 \text{ cm}^{-3}$ (top panel).

¹Department of Physics and Astronomy, University of Calgary, 2500 University Dr NW, Calgary, AB T2N 1N4, Canada

²Department of Physics, University of Bath, Claverton Down, Bath BA2 7AY, UK

³STFC RAL Space, Rutherford Appleton Laboratory, Harwell Campus, Didcot, Oxfordshire OX11 0QX, UK

⁴Green Bank Observatory, 155 Observatory Rd, Green Bank, WV 24944, USA

⁵Research Centre for Astronomical Computing, Zhejiang Laboratory, Hangzhou 311121, P. R. China

⁶NRC Herzberg Astronomy and Astrophysics, 5071 West Saanich Rd, Victoria, BC V9E 2E7, Canada

⁷Department of Physics and Astronomy, University of Victoria, Victoria, BC V8W 2Y2, Canada

⁸National Astronomical Observatory of Japan, Osawa, Mitaka, Tokyo 181-8588, Japan

⁹Department of Astronomy, School of Science, Graduate University for Advanced Studies (SOKENDAI), Mitaka, Tokyo 181-8588, Japan

¹⁰Korea Astronomy and Space Science Institute, 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Republic of Korea

¹¹University of Science and Technology, Korea (UST), 217 Gajeong-ro, Yuseong-gu, Daejeon 34113, Republic of Korea

¹²Shanghai Astronomical Observatory, Shanghai, CN, P. R. China

¹³Institute of Astronomy, National Tsing Hua University, No. 101, Section 2, Kuang-Fu Road, Hsinchu 30013, Taiwan

¹⁴IAPS – INAF, Via Fosso del Cavaliere, 100, I-00133 Rome, Italy

¹⁵Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400 Austin, TX 78712-1205, USA

¹⁶Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China

¹⁷Department of Astronomy, School of Physics, Peking University, Beijing 100871, China

¹⁸Korea Astronomy and Space Science Institute, 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Republic of Korea

¹⁹National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing 100012, China

²⁰Physics Department, Nazarbayev University, Astana, Kazakhstan

²¹Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, 146 Brownlow Hill, Liverpool L3 5RF, UK

²²Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK

²³Sterrenkundig Observatorium, Ghent University, Krijgslaan 281-S9, B-9000 Gent, Belgium

²⁴Institute for Advanced Studies, Nicolaus Copernicus University in Toruń, Wileńska 4, PL-87-100 Toruń, Poland

²⁵School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK

²⁶Institute of Astronomy and Astrophysics, Academia Sinica: Taipei, TW, Taiwan

²⁷Korea Astronomy and Space Science Institute (KASI), 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Republic of Korea

²⁸Department of Astronomy, School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

²⁹Gemini Observatory/NSF NOIRLab, 670 N. A'ohōkū Place, Hilo, HI 96720, USA

³⁰Department of Physics & Astronomy, Kwantlen Polytechnic University, 12666 72nd Avenue, Surrey BC V3W 2M8, Canada

³¹Department of Astronomy, Xiamen University, Xiamen 361005, Fujian, China

³²Centre for Astrophysics and Planetary Science, University of Kent, Canterbury CT2 7NH, UK

³³Université Paris-Saclay, Université Paris Cité, CEA, CNRS, AIM, 91191, Gif-sur-Yvette, France

³⁴Cardiff Hub for Astrophysics Research & Technology, School of Physics & Astronomy, Cardiff University, Queen's Buildings, Cardiff CF24 3AA, UK

³⁵Purple Mountain Observatory, Yuanhua Road 10, Nanjing 210023, China

³⁶Xinjiang Astronomical Observatory, CAS 150, Science 1-Street, Urumqi 830011, P. R. China

³⁷Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

³⁸Armagh Observatory and Planetarium, College Hill, Armagh BT61 9DB, UK

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.