



Kent Academic Repository

Caro, Tim, Rashid, Rashid Suleiman, Hamad, Ali Issa, Oates, Simon, Jackson, Isla, Entwistle, Abigail, Robinson, Janine and Mildenstein, Tammy (2026) *The Pemba flying fox Pteropus voeltzkowi: status, distribution and increasing urbanization*. *Oryx* . ISSN 1365-3008.

Downloaded from

<https://kar.kent.ac.uk/113629/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1017/s0030605325102238>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

The Pemba flying fox *Pteropus voeltzkowi*: status, distribution and increasing urbanization

TIM CARO^{1*} , RASHID SULEIMAN RASHID², ALI ISSA HAMAD²
SIMON OATES¹, ISLA JACKSON¹, ABIGAIL ENTWISTLE³
JANINE ROBINSON^{4,5} and TAMMY MILDENSTEIN⁶

Abstract Conservation of the Pemba flying fox *Pteropus voeltzkowi*, endemic to Pemba, Zanzibar, has been a success story. Low numbers last century prompted a local conservation response that likely resulted in a 400% increase in the population. But, with project funding having been gradually reduced over 15 years, it is unclear whether the population has maintained its recovery. Here we report findings from a new survey, in 2024, showing that the population size is that of the recovered population in 2008. However, many *P. voeltzkowi* have moved to urban centres, probably to avoid disturbance, primarily hunting by children. Only half the people interviewed remembered the earlier conservation campaign, but nearly all would support an environmental education campaign aimed at children in schools.

Keywords Fruit bat, monitoring, Pemba, *Pteropus voeltzkowi*, roosting locations, urbanization, Zanzibar

The supplementary material for this article is available at doi.org/10.1017/S0030605325102238

Introduction

Old World fruit bats (family Pteropodidae) are important pollinators and seed dispersers of over 163 economically useful plants (Fujita & Tuttle, 1991), notably pollinating species of the Fabaceae, Lythraceae, Malvaceae, Musaceae and Myrtaceae families, and dispersing seeds of *Ficus*, *Syzygium* and *Diospyros* species (Aziz et al., 2021). Within the Old World fruit bat family, flying foxes (genera *Acerodon*, *Desmalopex*, *Eidolon*, *Mirimiri*, *Neopteryx*, *Pteralopex*, *Pteropus*, *Styloctenium*) are the larger-sized bats, able to carry large fruits and traverse great distances, making them prominent seed dispersers (Mwinyi & Nnko, 2022). Despite their ecological importance, 51 of the 73 flying fox species are threatened because their wide-ranging

habits make them susceptible to deforestation and habitat loss (Meyer et al., 2016), hunting (Mildenstein et al., 2016) and land-use changes, including agricultural development and urbanization (Jung & Threlfall, 2016). Unfortunately, few species of flying foxes have received sustained conservation attention, so little is known about population sizes and trends.

The Pemba flying fox *Pteropus voeltzkowi* is endemic to the island of Pemba, Zanzibar, an autonomous region of Tanzania. The species is important to Pemba because it is a significant pollinator and seed disperser of wild and cultivated fruit trees (Entwistle & Corp, 1997b; Mwinyi & Nnko, 2022; Seltzer et al., 2013). The population of *P. voeltzkowi* has been surveyed twice since 1995, allowing us to monitor its trajectory over the past 3 decades. In addition to this unusual long-term conservation attention, the species appears to have shown a population recovery from very low numbers during most of the 20th century (Andersen, 1912). In 1992, an island-wide survey documented 2,400–3,600 individuals (Seehausen, 1991), and a subsequent more comprehensive survey, in 1995, counted 4,608–5,450 individuals (Entwistle & Corp, 1997a). Concerns about the small population led to an awareness and outreach programme in which initial awareness activities in 1992 were expanded in 1995 by the former Department of Commercial Crops, Fruit and Forestry, now the Department of Forestry, in conjunction with Flora & Fauna International and Lube Bat Conservancy, to establish community-based environmental clubs, convene village-level meetings and secure commitments to ban shotgun hunting and reduce roost site disturbance (Entwistle, 2002). These community-supported efforts collectively resulted in a 400% increase in the population, which reached 18,200–22,100 individuals, as assessed through two total population counts, in March and May 2008 (Robinson et al., 2010). A partial count in 2012, based on censusing the four largest roosts at the time and extrapolating, estimated 9,360–10,630 individuals (Grose & Clemmens, 2012).

Following cessation of direct conservation promotion activities with communities in 2007, it is unclear how far the shift in attitudes that favoured protection of the bats has been maintained. It is therefore unclear whether the core messages of the earlier conservation programme have been retained sufficiently to maintain a healthy population size.

*Corresponding author, tim.caro@bristol.ac.uk

¹School of Biological Sciences, University of Bristol, Bristol, UK

²Department of Forestry, Wete, Pemba, Zanzibar, Tanzania

³Flora & Fauna, Cambridge, UK

⁴Durrell Institute of Conservation and Ecology, University of Kent, Canterbury, UK

⁵Joint Nature Conservation Committee, Peterborough, UK

⁶Bat Conservation International, Austin, Texas, USA

Received 16 April 2025. Revision requested 13 August 2025.

Accepted 30 September 2025.

The objectives of this research were to provide current information on the population size and distribution of *P. voeltzkowi* colonies by visiting every roost site on the island, to examine current threats, and determine the extent to which Pembans still remember the conservation intervention programmes from 2 decades ago.

Study area

Pemba is a continental island 50 km off the coast of mainland Tanzania. Covering 988 km², it is one of two major islands comprising the Zanzibar Archipelago. It is dominated in the west by heavily vegetated ridges and valleys of primary and secondary forest and clove trees, with baobab *Adansonia digitata*, palms and coastal bush in the flatter east. There are a few small cities on the island, mainly but not exclusively the four District headquarters, but the majority of residents live in the many small villages scattered across most of the island and outlying islets. We identified six roost sites in urban areas (i.e. locations with > 10,000 people). These were the three District headquarters Wete, Chake Chake and Mkoani, plus Ole, Kengeja and Tumbe. In addition to *P. voeltzkowi*, several other vertebrate species are endemic to the island including the Pemba sunbird *Cinnyris pembae*, Pemba green-pigeon *Treron pembaensis*, Pemba white-eye *Zosterops vaughani* and Pemba scops owl *Otus pembaensis*.

Methods

Fieldwork was conducted on Pemba over consecutive days during 22 July–4 August 2024 by TC, RSR, AIH and TM. We investigated 81 reported roost sites, of which 67 were visited in person and for 14 a knowledgeable person was questioned by telephone (none of the latter contained bats). We first identified colony locations from a list of 55 roost sites visited in the last survey, in 2008 (Robinson et al., 2010). Following a snowball methodology, we added new colony locations to the list whenever these were reported by people whom we encountered at those sites. Two of the authors (RSR and AIH) are employees of the Department of Forestry; RSR knows the island intimately and was involved in previous surveys of the species. We are reasonably certain we visited nearly all occupied and unoccupied roosts. At each roost site, the village or town name, ward name, geographical coordinates, the number of bats in the roost and the characteristics of the roost were recorded; i.e. the trees in which bats were roosted, the forest type, the roost surroundings and reports or observations of any disturbance.

We were brought to the roosting areas by the Sheha (the government appointed executive officer) and local community members who were familiar with the bats' roosting locations. With their help, we identified the best vantage points for surveying the roost and conducted a daytime

roost count following a standardized direct censusing protocol for flying foxes (Mildenstein et al., 2005; Mildenstein, 2010; Kunz et al., 2009), optimized for single survey population abundance assessment on islands (Mickleburgh et al., 2009). The exact number of bats was counted using binoculars, and the colony size range was categorized as 0, 1–50, 51–100, 101–500, 501–1,000, 1,001–2,500, 2,501–5000 and > 5000 bats, to facilitate comparison with previous surveys. Roost counts are expected to have high probabilities of observing most individuals (Kunz et al., 2009), but it is possible some bats were missed, and it was more difficult to see all the bats within the diurnal roosting area at larger roosting colonies. Thus, in addition to direct roost counts, we also conducted evening dispersal counts between 18.50 and 19.40 at the two largest roosts (Chake Chake and Wete), following a standardized exit count protocol (Mildenstein et al., 2005; Mildenstein, 2010; Kunz et al., 2009). Exit counts are known to be variable, and best practices for population abundance assessment recommend multiple evening dispersal counts across consecutive evenings (Mildenstein, 2010), so our single evening dispersal counts of Chake Chake and Wete are likely to be minimum estimates.

We asked residents over 15 years of age living near each roost if they would like to take part voluntarily in a short interview, in Swahili, which they could leave at any time. Questions included the history of the occupancy of the site, whether there was roost disturbance by people, and what kind, such as tree felling, hunting, general harassment (noise and stone throwing) or fruit picking, usually by children; what foods they believed the bats to feed on; and whether they hunted and/or ate the bats. In situations where roosts were empty, interviewees were asked to approximate how many bats had roosted there in the past. Interviewees were also asked whether they remembered the previous fruit bat education campaign (that is, the initial 1992 education campaign and the large-scale engagement efforts led by the Department of Forestry during 1998–2007), and if a new campaign was needed.

Subsequently, we compared *P. voeltzkowi* population size and locations from our survey to surveys by the Department of Forestry personnel during 1992–2002, an island-wide survey in 1995 (Entwistle & Corp, 1997a) and two island-wide surveys conducted in 2004 (Robinson et al., 2010). The roost characteristics, threat assessments and interviews from our survey provide baseline information for 2024.

Results

Population status We counted a total of 25,328 *P. voeltzkowi* individuals across 81 roosting locations, 29 of which were inhabited and 52 empty (Supplementary Material 1). The modal colony size category was 101–500 bats (13 sites), eight sites had < 100 individuals, and two




FIG. 1 Number of roosts in each Pemba flying fox *Pteropus voeltzkowi* colony size category on Pemba Island, Zanzibar, in 2024.

contained 2,500–5,000 bats (Fig. 1). These two large roosts were in the cities of Chake Chake (human population size 52,000 in 2022) and Wete (36,000 in 2022) that together held 58.7% of the island's total *P. voeltzkowi* population at the time of the survey. Daytime counts at these two sites were 3,098 and 3,926 bats, respectively, whereas evening dispersal counts were 8,649 and 7,560 bats, respectively, indicating daytime counts were lower by a minimum of 64% and 48%, respectively (see Discussion).

Distribution and roost characteristics We found *P. voeltzkowi* across Pemba and on outlying islands (Fig. 2). Roost size was significantly larger in urban centres ($n = 6$, mean = 3,062) than in rural areas ($n = 55$, mean = 1,276; Mann–Whitney $U = 303$, $P < 0.001$). The roosts in Chake Chake and Wete had $> 2,501$ bats. One roost with 1,001–2,500 bats was in the centre of the small city of Mkoani; another was on agricultural land at Kidike. There were three roosts with 501–1,000 bats, just outside the town of Kengeja, and in Ngezi Forest Reserve (Supplementary Material 1). Bats were observed roosting in 32 species of tree, with 13 of these at more than one roost site (Table 1). Mango trees *Mangifera indica* were the most common roost tree species ($n = 30$) and, with tropical almond *Terminalia catappa* and ordeal *Erythrophleum suaveolens* trees, comprised 49.6% of 113 roost trees. Bats were reported to feed on 25 food species (16 reported more than once; Table 1). Mango was the most reported food species eaten ($n = 52$) and, with the fruit from the broom cluster fig *Ficus sur* and almond, comprised 66.9% of 181 reports.

Changes from previous surveys In 1995 bats were found in relatively low numbers at relatively few sites across Pemba but in 2008 and in our survey in 2024 they were found in greater numbers and at more locations (Fig. 3). The 2008 survey revealed large aggregations of bats at five sites, Mgoconi (south of Ngezi Forest Reserve), two outside the city of Mkoani, and in Kidike and Wete; all except Wete




FIG. 2 Locations and size categories of *P. voeltzkowi* roost sites recorded on Pemba in 2024. Note that some closely adjacent sites are aggregated here for the purpose of illustration.

were rural sites. In 2024, we found bats at the six large sites described earlier, two of which were the same as in 2008. The number of bats in urban and rural roosts changed between 2008 and 2024 (Table 2), with 72.5% of bats found in urban roosts in 2024, compared with 27.6% and 25.4% in the two 2008 surveys, and 21.8% in the 1995 survey. In contrast to our 2024 survey, there were no significant differences in the number of bats counted at urban and rural sites in either of the 2008 surveys (May: $n = 54$ sites, $U = 105.5$, $P = 0.288$; March: $n = 54$, $U = 105$, $P = 0.306$), or in 1995 ($n = 38$, $U = 64$, $P = 0.071$).

Disturbance In 2024, interviewees reported disturbance of bats at 42.2% of the 81 roost sites (83 reports, as two people were interviewed for each of two roosts; Table 3). Empty roosts seemed to have suffered more from disturbance than occupied roosts (50.0% vs 31.4%, respectively). Hunting had occurred at 16.9% of roosts, with more reported at now empty roosts than at occupied roosts (25.0% vs 5.7%, respectively). Disturbance appeared to have fallen from 43.9% in 1995 to 23.9% in 2000 following the onset of the conservation project and adoption of local byelaws to protect the bats (Table 3). Hunting occurred at 12.2% and 12.8% of roosts in 1995 and 2000, respectively, and at 16.9% of roost sites in 2024.

TABLE 1 Number of reports of tree species used as roosts and as a source of fruits by the Pemba flying fox *Pteropus voeltzkowi*, as indicated by interviewees on Pemba in 2024. Scientific names are provided where known.

Species	Local name	Roost	Fruits eaten
Mango <i>Mangifera indica</i>	Maembe	30	53
Almond <i>Terminalia catappa</i>	Kungu	16	34
Ordeal <i>Erythrophleum suaveolens</i>	Wavi	10	
Fig <i>Ficus</i> sp.	Kuyu	9	
Baobab <i>Adansonia digitata</i>	Buyu	7	2
African teak <i>Milicia</i> sp.	Vule	5	
Fig <i>Ficus</i> sp.	Gulele	4	
Black mangrove <i>Avicennia germinans</i>	Mangonde	3	
<i>Pachystela cinerea</i>	Chocha	2	
Fig <i>Ficus</i> sp.		2	
False mopane <i>Guibourtia coleosperma</i>	Michenza msitu	2	
Umbrella tree	Sisi	2	
Java plum <i>Syzygium cumini</i>	Zanzibaraoa/Mfuu	2	4
Fig <i>Ficus</i> sp.	Mkuyu		37
Breadfruit <i>Artocarpus altilis</i>	Masheli		15
Plum	Mbura		9
<i>Saba comorensis</i>	Mabungo		4
Laurel	Mtondoo		3
Papaya <i>Carica papaya</i>	Mapapai		2
Apple guava <i>Psidium guajava</i>	Mapera		2
Kei apple <i>Dovyalis afra</i>	Mchongoma		2
Banana <i>Musa</i> sp.	Ndizi		2
Fig <i>Ficus</i> sp.	Mlandege		2
Neem <i>Azadirachta indica</i>	Mwarubaini		2
<i>Searsia longipes</i>	Tangele		2




FIG. 3 The size and distribution of occupied *P. voeltzkowi* roosts on Pemba Island recorded in (a) 1995 (Entwistle & Corp, 1997a), (b) 2008 (Robinson et al., 2010) and (c) in our survey in 2024.

Conservation project When interviewees were asked about their recollection of the original education campaign and/or environmental club engagement, 28.4%

remembered the project, 45.7% did not, and 13.6% had only heard of it indirectly, with some not answering. When asked whether a new campaign was needed, 88.9%

TABLE 2 Total numbers (and per cent) of *P. voeltzkowi* counted in rural and urban roosts on Pemba (Fig. 2) in our survey in 2024 and in three earlier surveys, in May and March 2008 (Robinson et al., 2010) and 1995 (Entwistle & Corp, 1997a). Comparisons: 2024 vs March 2008 $\chi^2 = 8,393.5$, $df = 1$, $P < 0.00001$; 2024 vs May 2008 $\chi^2 = 1,162.1$, $df = 1$, $P < 0.00001$; 2024 vs 1995 $\chi^2 = 4,116.7$, $df = 1$, $P < 0.00001$.

	2024	May 2008	March 2008	1995
Rural	6,958 (27.5)	12,577 (72.4)	16,541 (74.6)	3,313 (78.2)
Urban	18,370 (72.5)	4,784 (27.6)	5,646 (25.4)	925 (21.8)
Total	25,328	17,361	22,187	4,238

TABLE 3 Notes recorded at roost sites, with numbers of roosts at which each issue was observed in 1995 (Entwistle & Corp, 1997a), 2000 (Entwistle, 2002) and during our survey in 2024. Blank cells indicate data were not recorded.

	Occupied sites in 1995	Occupied sites in April–May 2000	Occupied sites in 2024	Empty sites in 1995	Empty sites in April–May 2000	Empty sites in 2024
Disturbance (e.g. noise or stone throwing)	2	0	5	6	6	3
Hunting of bats	1	1	2	4	5	12
Trees felled	1	1	3	4	3	5
Site degraded as a result of human activities	0	0		1	1	
Vegetation clearance for various purposes	1	1		1	1	
Deserted	1	2	0	3	6	1
Seasonal/temporary	1	1	0	1	6	4
Picking of fruit	0	0	1	0	0	4
Uninhabited at present	1	2	1	1	0	0
Protected by community or government	2	3	7	0	0	1
Reestablished	1	1		0	1	
No information	5	9	3	3	17	5
No disturbance	1		13	0		13

answered yes. Reasons given for desiring a new campaign were to protect bats (22.2% of respondents), to protect the environment (25.9%), to improve education about bats (19.8%) to attract visitors to Pemba (8.6%) and because bats benefit the environment (11.1%).

Discussion

Population size

As a result of obstructions and limited visibility, counting bats while they are roosting can be challenging. Some previous surveys have used patch counts (Thomas & Laval, 1988), and then scaled up estimates at roosts with poor visibility. We used direct counts at small sites and evening dispersal counts at the two largest roost sites. With more time, evening dispersal counts at additional sites would have enabled us to generate a more accurate correction factor to apply to the other roosts, but evening dispersal counts over many nights (only one roost can be counted per night) would have lengthened the survey and could result in bats being double-counted as a result of inter-roost movement.

Instead, we tried an alternative method to address the issue of potential undercounting at large roosts. Specifically, we used a quadratic equation ($y = 0.003947x^2 + 0.92106x$) to account for undercounting at any roost site with > 200 bats, except Chake Chake and Wete, where we used the evening dispersal counts we made there. A quadratic function was employed because we reasoned that errors would increase disproportionately with roost size. This meant 14 of 81 roost sites had an adjusted estimate. At sites with ≤ 200 bats, we were confident of our direct daytime counts and used them. Using these plus the adjusted counts, we estimated there were 27,412 bats on Pemba. In summary, the 2024 *P. voeltzkowi* population on Pemba lies between 25,328 and 27,412 individuals, comparable to the estimate in 2008 (18,200–22,100).

Distribution

Both previously published surveys concluded that *P. voeltzkowi* roosts were most likely to be found in forest habitats (Entwistle & Corp, 1997a; Robinson et al., 2010). In the 1995 survey, nine of the 10 roosts containing > 100

bats were in primary or secondary forest (Entwistle, 2002). Our survey suggests that by 2024 there had been a significant movement of bats into towns, with 58.7% of the population in two large city roosts and with mean roost size in towns twenty times greater than in forests (Table 2).

Many of the roost sites we visited were also surveyed in 1995 and 2008. In 2008, four sites (Kidike, Mbiji 1, Mbiji 2 and Wete), accounted for 87% of the total bat population on Pemba. Of these, only Wete has retained its large population, which has also increased. At Kidike, once a significant forest roost and former tourist attraction, there has been a substantial population decline, with a count of only 1,491 bats in 2024 compared to 3,394 and 4,978 in the two 2008 counts. The two forest roosts of Mbiji 1 and Mbiji 2 were empty in 2024 but had counts of 4,678 and 7,001, respectively, in 2008. Local reports attributed this decline to disturbance. These shifts illustrate the abandonment of forest roosts and movement to urban environments. Movement to cities was also implicated in changes in a 2012 study on Pemba (Grose & Clemmens, 2012).

Other *Pteropus* species are also moving into urban areas, and species can now be found in multiple cities in Australia (Markus & Hall, 2004; Parris & Hazell, 2005; Williams et al., 2006), Egypt (Egert-Berg et al., 2021), Singapore (Chan et al., 2021) and throughout Southeast Asia (T. Mildenstein, unpubl. data). Although increased food availability can in some cases account for the movement of bats into urban areas (Williams et al., 2006), the primary food species of *P. voeltzkowi*, mango, breadfruit *Artocarpus altilis* and *Ficus* spp. (Entwistle & Corp, 1997b; Table 1), are not more abundant in cities on Pemba, and this is therefore an unlikely explanation for bat movements on the island. The most likely explanation for this phenomenon in areas where bats are hunted or disturbed is the added protection afforded in urban settings, where hunters are not tolerated by city dwellers (Mildenstein, 2010).

Ephemeral nature of roosts

Many of the roosts we visited were empty (see also Grose & Clemmens, 2012) but were reportedly occupied recently (corroborated by evidence of bat droppings in the understorey). Temporary roost occupancy appears to be a characteristic of *P. voeltzkowi* on Pemba, although the bats show strong fidelity to some roost sites, such as those in Ngezi forest and Kengeja (Pakenham, 1984; Kingdon, 1974; Bentjee, 1990). Up to eight bat censuses were conducted by Department of Forestry personnel from just before 1992 up to February 2002, with numbers available for 46 sites visited more than once. Seasonal shifts were pronounced: bats were always present at 12 of these sites, and were always absent at six, 15 were vacated over the course of the censuses, five were established, two were vacated and then re-established, one was established but then vacated, and

five flip-flopped between bats being present and absent (Entwistle, 2002).

When we arrived at an empty roost, we asked people how many bats were there previously. This was an open question but assuming any answer referred to the last time the bats were present, 27 of 46 empty roosts had reportedly housed 100–500 bats, and four sites were reported to have contained > 1,000 bats. Only rural habitats contained empty roosts; all town roosts were occupied, again suggesting bats were concentrated in towns at the time of this study in 2024.

Reasons for movements appeared twofold: disturbance and seasonal movements as a result of changing food distribution. We found that disturbance, hunting, tree felling and fruit picking were associated with 50% of empty roost sites (Table 3), and an earlier survey found a negative correlation between proximity to human settlements and colony size (Robinson et al., 2010). Absence of bats was associated with disturbance, hunting and tree cutting in both 2000 and 1995 (Table 3; Entwistle, 2002). Our information regarding changing food distribution was anecdotal, taken from people's opinions, but mango trees, an important food source, occur across the island and bats will reportedly travel up to 15 km to food sources each night, suggesting food distribution may not necessarily be a major cause of changes in roost occupancy.

Hunting of bats for food on Pemba remains a problem, with bats captured at c. 17% of roosts, mostly by boys using long sticks with sticky seed pods or catapults (Walsh, 1995). Hunting has been a consistent threat to this species (Entwistle & Juma, 2016), exacerbated by the introduction of shotguns (Entwistle & Corp, 1997a). Hunting was more prevalent than other disturbances: it was reported at 12 of 48 empty roosts, and 18.5% of 81 local people admitted to hunting bats and 28.4% said they had eaten them. These statistics, along with two reports stating that people had visited roost sites to hunt bats with shotguns in the week previous to our survey, indicate hunting is still a threat to this species.

We believe that hunting, felling of trees, and throwing stones to knock down fruit in rural areas is driving *P. voeltzkowi* into large urban roosts where these activities are suppressed by the disapproval of passersby, and by policing. For example, there was a large roost at Msitu wa Mbiji, 2.5 km outside Mkoani, in the 2008 census, but this was empty in 2024, with the bats having likely taken up residence in trees within the District Government residence in the centre of Mkoani, reportedly as a result of hunting and human disturbance at Mbiji (see also Grose & Clemmens, 2012).

Study limitations

Our study suffered from at least two limitations. Firstly, it was a single, dry season study, which limits our ability to

make unequivocal statements regarding whether rural roosts have been abandoned permanently for urban sites; bats may roost in urban areas temporarily, influenced by temporal variation in food sources. In 2025, numbers at Chake Chake were lower than they were in 2024 as a result of movement from the roost (RSR, pers. obs.). Secondly, we did not take measures of disturbance, proximity to human habitation or roost tree species into account in our analyses. It is challenging to measure disturbance quantitatively as it varies over time and in extent, requiring frequent visits to sites.

Conservation implications

The 1992 education campaign to conserve *P. voeltzkowi* was centred on the species' role in seed dispersal and forest regeneration, with the campaign attempting to encompass both formalized and community-based educational outlets (Robinson et al., 2010). Initial awareness activities in 1992 were built upon in 1995, with surveys in schools demonstrating high understanding of the ecological role of the bats, but a lack of recognition of its endemism or threatened status (Entwistle & Corp, 1997a; Trehwella et al., 2005). Interviews demonstrated the cultural importance of this species locally (Bowen-Jones & Entwistle, 2002). Subsequent work by the Department of Forestry and NGOs extended education and awareness activities (including production of a video), established community-based environmental clubs in several villages and convened village-level meetings that resulted in commitments to protect the bats, including bans on hunting with shotguns, and reducing roost site disturbance (Entwistle, 2002). The campaign showed the potential for protection of bats to be internalized within village environmental clubs, and some villages established local byelaws to protect bats and their roosts, enforced by club members (Entwistle, 2001, Rice & Hamdan, 2002); children were used as ambassadors to take the message to their parents. Evaluations of the project indicated an increase in bat population size over the period (albeit with some caution with regard to survey rigour), and decreased hunting at roost sites protected by environmental clubs (Rice & Hamdan, 2002). Further project funding during 2003–2007 supported ongoing awareness raising activities, community-led monitoring activities and development of bat ecotourism at one environmental club, Kidike, along with ongoing survey work. Further population increases were reported over this period (Fauna & Flora International, 2007), leading to a recategorization of the species on the IUCN Red List from Critically Endangered (Mickleburgh et al., 2004) to Vulnerable (Entwistle & Juma, 2016).

Yet, in 2024, less than half of interviewees had heard of these educational efforts, but nearly everyone thought a new campaign was necessary and that most people would listen and learn from it. Pembans recognize that fruit bats

play important ecological roles as pollinators and seed dispersers on the island (Mwinyi & Nnko, 2022); 37% of interviewees talked, either directly or indirectly, about the role of bats in the environment and this being a reason why a new campaign is needed.

Although the *P. voeltzkowi* population appears to have been stable since 2008, there are indications that bats are moving from rural to urban locations. Our working hypothesis is that this movement is driven by human activities, namely hunting and roost disturbance, which are less frequent in cities because of widespread disapproval of these activities. Interview responses indicated that a substantial number of local people are likely to be receptive to a new conservation project. Any campaign should continue to educate the younger generation in schools, not only because children hunt the species, but to emphasize the key roles of bats in plant pollination and seed dispersal, both of value to humans (Fujita & Tuttle, 1991), and to remind people that the species is endemic to Pemba and is therefore a threatened cultural icon. Given that bats are now an unavoidable presence in towns, there is an opportunity for the Department of Forestry to reinstate the community-level environmental clubs, which were previously successful, as an effective mechanism to discourage interference with bat roosts in urban environments but also in rural areas where they are more threatened by disturbance.

Recommendations for future surveys

Our examination of several surveys conducted by different people and organizations revealed nine challenges involved in counting *P. voeltzkowi*. (1) There is inconsistency in the time of year in which surveys are conducted. (2) There is inconsistency in choice of sites surveyed, with number and localities changing both between and within years (e.g. Kisiwa Panza and Matumbini, see Supplementary Material 1). (3) There can be confusion with site names, and where there is more than one site within a community area the roosts are not currently distinguished between each other; e.g. Mgelema (Mgelima); this could result in incorrect recording of data. (4) There is a lack of reconnaissance, training and survey experience among surveyors conducting the counts, which introduces observer error. (5) There is a lack of time allocated for replication using direct and exit counts. (6) Surveys are usually conducted just once and at one time of year, but multiple island-wide counts across seasons are required to better understand bat roosting choices and movements. (7) At sites with small numbers of *P. voeltzkowi* it is possible to count each individual, but at large roosting sites this is infeasible, and no standardization has been employed in techniques to count these populations, increasing the margin for error. (8) In some surveys, each team is allocated a section of the island to survey; this extends the length of time each team

spends in one geographical area, which could increase the risk of double-counting individuals or groups if they move between roost sites. (9) Although assessment of habitat and disturbance is made at roost sites during surveys, there is scope for improving assessment methods and collecting more detailed information. Addressing these challenges could lead to more robust surveys in the future.

Author contributions Project design: TC, TM; fieldwork: TC, RSR, AIH, TM; analyses, map production: SO, IJ; writing: TC; addition of historical text: AE, JR; revision: TM.

Acknowledgements We thank Bat Conservation International for funding, Matt Clark for advice on mapping, and two anonymous reviewers for comments.

Conflicts of interest None.

Ethical standards Observational research was conducted with permission from the Government of Zanzibar, ethical approval was obtained from the University of Bristol (UIN-24-056), and the study abided by the *Oryx* ethical standards.

Data availability Data are provided in the supplementary material.

References

- ANDERSEN, K. (1912) *Catalogue of the Chiroptera in the Collections of the British Museum. I. Megachiroptera, 2nd edition*. British Museum (Natural History), London, UK.
- AZIZ, S.A., MCCONKEY, K.R., TANALGO, K., SRITONGCHUAY, T., LOW, M.-R., YONG, J.Y. et al. (2021) The critical importance of Old World fruit bats for healthy ecosystems and economies. *Frontiers in Ecology and Evolution*, 9, 641411.
- BENTJEE, H.J. (1990) *Botanical Assessment of the Ngezi Forest, Pemba*. Forestry Development Project, Technical Paper 1. Forestry Department, Zanzibar/FINNIDA.
- BOWEN-JONES, E. & ENTWISTLE, A. (2002) Identifying appropriate flagship species: the importance of culture and local contexts. *Oryx*, 36, 189–195.
- CHAN, A.A.Q., AZIZ, S.A., CLARE, E.L. & COLEMAN, J.L. (2021) Diet, ecological role and potential ecosystem services of the fruit bat, *Cynopterus brachyotis*, in a tropical city. *Urban Ecosystems*, 24, 251–263.
- EGERT-BERG, K., HANDEL, M., GOLDSHTEIN, A., EITAN, O., BORISSOV, I. & YOVEL, Y. (2021) Fruit bats adjust their foraging strategies to urban environments to diversify their diet. *BMC Biology*, 19, 123.
- ENTWISTLE, A. (2001) Community-based protection successful for the Pemba flying fox. *Oryx*, 35, 355–356.
- ENTWISTLE, A. (2002) *Pemba flying fox conservation project: Final report to the Lubee Foundation, August 2002*. Fauna & Flora International, Cambridge, UK.
- ENTWISTLE, A. & CORP, N. (1997a) Status and distribution of the Pemba flying fox *Pteropus voeltzkowi*. *Oryx*, 31, 135–142.
- ENTWISTLE, A. & CORP, N. (1997b) The diet of *Pteropus voeltzkowi*, an endangered fruit bat endemic to Pemba Island, Tanzania. *African Journal of Ecology*, 35, 351–360.
- ENTWISTLE, A.C. & JUMA, J. (2016) *Pteropus voeltzkowi*. In *The IUCN Red List of Threatened Species 2016*. [dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T18768A22089205.en](https://doi.org/10.2305/IUCN.UK.2016-1.RLTS.T18768A22089205.en).
- FAUNA & FLORA INTERNATIONAL (2007) *CEPF Final project Completion Report*. cepf.net/sites/default/files/final_faunaflora_international_pembaisland.pdf [accessed 17 December 2025].
- FUJITA, M.S. & TUTTLE, M.D. (1991) Flying foxes (Chiroptera: Pteropodidae): threatened animals of key ecological and economic importance. *Conservation Biology*, 5, 455–463.
- GROSE, H. & CLEMMENS, C.G. (2012) *Who Else Would Plant The Trees? A Status Update on the Pemba Flying Fox*. Independent Study Project (ISP) Collection, School for International Training. digitalcollections.sit.edu/isp_collection/1419 [accessed 17 December 2025].
- JUNG, K. & THRELFALL, C.G. (2016) Urbanisation and its effects on bats—A global meta-analysis. In *Bats in the Anthropocene: Conservation of Bats in a Changing World* (eds C.C. Voigt & T. Kingston), pp. 13–34. Springer, Cham, Switzerland.
- KINGDON, J. (1974) *East African Mammals. An Atlas of Evolution in Africa. Vol. IIA. Insectivores and Bats*. University of Chicago Press, Chicago, USA.
- KUNZ, T.H., BETKE, M., HRISTOV, N.I. & VONHOF, M.J. (2009) Methods for assessing colony size, population size, and relative abundance of bats. In *Ecological and Behavioral Methods for the Study of Bats* (eds T.H. Kunz & S. Parsons), pp. 133–157. Johns Hopkins University Press, Baltimore, USA.
- MARKUS, N. & HALL, L. (2004) Foraging behaviour of the black flying-fox (*Pteropus alecto*) in the urban landscape of Brisbane, Queensland. *Wildlife Research*, 31, 345–355.
- MEYER, C.F.J., STRUEBIG, M.J. & WILLIG, M.R. (2016) Responses of tropical bats to habitat fragmentation, logging, and deforestation. In *Bats in the Anthropocene: Conservation of Bats in a Changing World* (eds C.C. Voigt & T. Kingston), pp. 63–104. Springer, Cham, Switzerland.
- MICKLEBURGH, S.P., HUTSON, A.M. & RACEY, P.A. (1992) *Old World Fruit Bats. An Action Plan for their Conservation*. IUCN, Gland, Switzerland, and Cambridge, UK.
- MICKLEBURGH, S., HUTSON, A., BERGMANS, W. & HOWELL, K. (2004) *Pteropus voeltzkowi*. In *The IUCN Red List of Threatened Species 2004*. IUCN, Gland, Switzerland, and Cambridge, UK. iucnredlist.org/species/18768/8599011 [accessed March 2006].
- MICKLEBURGH, S., WAYLEN, K. & RACEY, P. (2009) Bats as bushmeat: a global review. *Oryx*, 43, 217–234.
- MILDENSTEIN, T.L. (2010) Bat population abundance assessment and monitoring. In *Investigating the Role of Bats in Emerging Zoonoses: Balancing Ecology, Conservation, and Public Health Interest* (eds S.H. Newman, H. Field, J.H. Epstein & C. de Jong), pp. 29–46. FAO Animal Production and Health Manual, No.11. Food and Agriculture Organization of the United Nations, Rome, Italy. fao.org/3/a-i2407e.pdf [accessed 17 December 2025].
- MILDENSTEIN, T.L. & BOLAND, J. (2010) *Development of Mariana fruit bat survey protocols for Mariana Islands surveys. Final Report – Rota Island pilot surveys*. USFWS report, Region 1, Pacific Islands Office.
- MILDENSTEIN, T.L., STIER, S.C., NUEVO-DIEGO, C.E. & MILLS, L.S. (2005) Habitat selection of endangered and endemic large flying-foxes in Subic Bay, Philippines. *Biological Conservation*, 126, 93–102.
- MILDENSTEIN, T., TANSHI, I. & RACEY, P.A. (2016) Exploitation of bats for bushmeat and medicine. In *Bats in the Anthropocene: Conservation of Bats in a Changing World* (eds C.C. Voigt & T. Kingston), pp. 325–376. Springer, Cham, Switzerland.
- MWINYI, A.M. & NKO, H.J. (2022) Seasonality of Pemba flying fox (*Pteropus voeltzkowi*) in selected habitats of Pemba Island, and its roles in seed dispersal and germination. *African Journal of Ecology*, 60, 481–491.

- PAKENHAM, R.H.W. (1984) *The Mammals of Zanzibar and Pemba Islands*. Privately published by the author.
- PARRIS, K.M. & HAZELL, D.L. (2005) Biotic effects of climate change in urban environments: the case of the grey-headed flying-fox (*Pteropus poliocephalus*) in Melbourne, Australia. *Biological Conservation*, 124, 267–276.
- RICE, M. & HAMDAN, S.I. (2002) *Pemba flying fox project DCCFF and FFI Project evaluation report. August 2002*. Fauna and Flora International, Cambridge, UK.
- ROBINSON, J.E., BELL, D.J., SALEH, F.M., SULEIMAN, A.A. & BARR, I. (2010) Recovery of the vulnerable Pemba flying fox *Pteropus voeltzkowi*: population and conservation status. *Oryx*, 44, 416–423.
- SEEHAUSEN, O. (1991) The Pemba fruit bat—on the edge of extinction? *Oryx*, 25, 110–112.
- SELTZER, C.E., NDANGALASI, H.J. & CORDEIRO, N.J. (2013) Seed dispersal in the dark: shedding light on the role of fruit bats in Africa. *Biotropica*, 45, 450–456.
- TAIT, J., PEROTTO-BALDIVIESO, H.L., MCKEOWN, A. & WESTCOTT, D.A. (2014) Are flying-foxes coming to town? Urbanisation of the spectacled flying-fox (*Pteropus conspicillatus*) in Australia. *PLOS One*, 9, e109810.
- THOMAS, D.W. & LAVAL, R.K. (1988) *Survey and Census Methods in Ecological and Behavioural Methods for the Study of Bats*. Smithsonian Press, Washington, DC, USA.
- TREWHELLA, W.J., RODRIGUEZ-CLARK, K.M., CORP, N., ENTWISTLE, A., GARRETT, S.R.T., GRANER, E. et al. (2005) Environmental education as a component of multidisciplinary conservation programs: lessons from conservation initiatives for Critically Endangered fruit bats in the Western Indian Ocean. *Conservation Biology*, 19, 75–85.
- WALSH, M.T. (1995) Eating bats on Pemba Island: a local innovation or cultural borrowing. *Mvita: Newsletter of the Regional Centre Study for the Archaeology of Eastern & Southern Africa*, 6, 15–18.
- WILLIAMS, N., MCDONNELL, M., PHELAN, G., KEIM, L. & VAN DER REE, R. (2006) Range expansion due to urbanization: increased food resources attract grey-headed flying-foxes (*Pteropus poliocephalus*) to Melbourne. *Austral Ecology*, 31, 190–198.