



Kent Academic Repository

Bolton, Lauren and Petch, Joel (2025) *Non-pharmacological interventions to tackle delirium in elderly hospitalised patients? A scoping review*. *Advanced Journal of Professional Practice* . ISSN 2059-3198.

Downloaded from

<https://kar.kent.ac.uk/114826/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://journals.kent.ac.uk/index.php/ajpp/article/view/1287/2358>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Individual Research Project Presentations Day 10th June 2024, Kent and Medway Medical School.

Non- pharmacological interventions to tackle delirium in elderly hospitalised patients? A scoping review.

Student^a: Lauren Bolton | **Supervisor(s)^b:** Joel Petch

Abstract

Background: Delirium is an acute neuropsychiatric syndrome that is characterised by fluctuations in attention, awareness and cognition. Delirium has a high prevalence within hospitals and is associated with increased morbidity and mortality. This incurs substantial healthcare associated costs and often prolongs the duration of hospital admissions, placing strain on the patient and their family and impacting upon the availability of hospital resources, a widespread issue across numerous hospital trusts.

Methods: A scoping review was conducted of literature exploring the current knowledge base of non- pharmacological interventions for the prevention of delirium and their efficacy using the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) checklist. Sources were subject to thematic analysis.

Results: Four hundred and thirty-nine articles were identified through a database search on APA PsycINFO, Embase, Medline and PubMed. A total of fifteen papers were included in the final synthesis. Thematic analysis identified five major themes, including intervention type, efficacy, staff adherence and education, cost efficacy and requirements for prospective research.

Conclusions: Overall, multicomponent interventions were evidenced to be effective at preventing delirium. Adherence across protocols was good and staff education was prioritised. While the field of delirium prevention lacks an abundance of high-quality experimental evidence, recent literature has begun to address the gaps in knowledge. However, further research is vital to comprehensively assess the efficacy of these interventions and expand the knowledge base of delirium prevention. Promisingly, the rising number of prospective clinical trials suggests great potential for future research in the field.

Keywords: Delirium | Prevention | Non-pharmacological | Elderly

^a 4th Year Medical Student, Kent and Medway Medical School, Canterbury, United Kingdom

^b Senior Lecturer, Kent and Medway Medical School, Canterbury, United Kingdom

Main contact email: l.bolton25@kmms.ac.uk