



Kent Academic Repository

Alexander, Stephen P. H., Striessnig, Jörg, Gibb, Alasdair J., Mathie, Alistair A., Veale, Emma L., Kelly, Eamonn, Peach, Chloe J., Armstrong, Jane F., Faccenda, Elena, Harding, Simon D. and others (2025) *The Concise Guide to PHARMACOLOGY 2025/26: Ion channels*. *British Journal of Pharmacology*, 182 (S1). S152-S241. ISSN 0007-1188.

Downloaded from

<https://kar.kent.ac.uk/114869/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1111/bph.70231>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

The Concise Guide to PHARMACOLOGY 2025/26: Ion channels

Stephen P. H. Alexander¹ , Jörg Striessnig² , Alasdair J. Gibb³ , Alistair A. Mathie⁴ , Emma L. Veale⁵ , Eamonn Kelly⁶ , Chloe J. Peach⁷ , Jane F. Armstrong⁸ , Elena Faccenda⁸ , Simon D. Harding⁸ , Christopher Southan⁸ , Jamie A. Davies⁸ , Richard W. Aldrich⁸ , Bernard Attali⁹ , Austin M. Baggetta¹⁰ , Elvir Becirovic¹¹ , David J. Beech¹² , Martin Biel¹¹ , Roslyn M. Bill¹³ , Ana I. Caceres¹⁴, William A. Catterall¹⁵ , Han Chow Chua¹⁶ , Alex C. Conner¹⁷ , Henry Danahay¹⁸, Paul Davies¹⁹, Katrien De Clercq²⁰, Markus Delling²¹ , Francesco Di Virgilio²² , Simonetta Falzoni²², Wei Fang²³, Stefanie Fenske¹¹ , Anna Fortuny-Gomez²⁴, Samuel Fountain²⁴ , George K. Chandy²⁵ , Mohammad-Reza Ghovanloo²⁶ , Steve A. N. Goldstein²⁷ , Martin Gosling¹⁸, Christian Grimm¹¹, Stephan Grissmer²⁸ , Kotdaji Ha²¹ , Tim G. Hales²⁹ , Verena Hammelmann¹¹, Israel Hanukoglu²³ , Meiqin Hu³⁰, Ad P. Ijzerman³¹ , Sairam V. Jabba¹⁴ , Mike Jarvis³², Anders A. Jensen³³ , Sven E. Jordt¹⁴ , Leonard K. Kaczmarek³⁴ , Stephan Kellenberger³⁵ , Charles Kennedy³⁶ , Brian King³⁷ , Philip Kitchen¹³ , Qiang Liu³⁰, Joseph W. Lynch³⁸, Rian W. Manville²⁷ , Jessica Meades²⁴, Verena Mehlfeld¹¹, Annette Nicke¹¹, Stefan Offermanns³⁹ , Edward Pérez-Reyes⁴⁰ , Leigh D. Plant¹⁰ , Stephan A. Pless³³ , Lachlan D. Rash³⁸ , Dejian Ren⁴¹ , Hussein N. Rubaiy⁴² , Mootaz M. Salman⁴³ , Werner Sieghart⁴⁴ , Lucia G. Sivilotti³⁷, Trevor G. Smart³⁷ , Terrance P. Snutch⁴⁵ , Paolo Tammaro⁴³ , Jinbin Tian⁴⁶ , James S. Trimmer⁴⁷ , Charlotte Van den Eynde²⁰ , Paola Vergani⁴⁸ , Joris Vriens²⁰ , Stephen G. Waxman³⁴ , Aguan D. Wei¹⁵ , Brenda T. Winn¹⁰, Heike Wulff⁴⁷ , Haoxing Xu⁴⁹ , Fan Yang³⁰, Lixia Yue⁵⁰ , Xiaoli Zhang⁴⁹  and Michael Zhu⁴⁶ 



¹Division of Physiology, Pharmacology & Neuroscience, School of Life Sciences, University of Nottingham Medical School, Nottingham, NG7 2UH, UK, ²Pharmacology and Toxicology, Institute of Pharmacy, University of Innsbruck, A-6020, Innsbruck, Austria, ³Research Department of Neuroscience, Physiology and Pharmacology, Division of Biosciences, University College London, Gower Street, London, WC1E 6BT, UK, ⁴School of Life Sciences, University of Westminster, London, W1W 6UW, UK, ⁵Medway School of Pharmacy, Universities of Kent and Greenwich, Chatham Maritime, Kent, ME4 4TB, UK, ⁶School of Psychology and Neuroscience, University of Bristol, Bristol, BS8 1TD, UK, ⁷Division of Physiology, Pharmacology & Neuroscience, School of Life Sciences, Centre of Membrane Proteins and Receptors (COMPARE), University of Nottingham Medical School, Nottingham, NG7 2UH, UK, ⁸Institute for Neuroscience and Cardiovascular Research, University of Edinburgh, Edinburgh, EH8 9XD, UK, ⁸Stanford University, Austin, USA, ⁹Tel Aviv University, Tel Aviv, Israel, ¹⁰Northeastern University, Boston, USA, ¹¹Ludwig Maximilian University of Munich, Munich, Germany, ¹²University of Leeds, Leeds, UK, ¹³Aston University, Birmingham, UK, ¹⁴Duke University, Durham, USA, ¹⁵University of Washington, Seattle, USA, ¹⁶University of Sydney, Sydney, Australia, ¹⁷University of Birmingham, Birmingham, UK, ¹⁸University of Sussex, Brighton, UK, ¹⁹Tufts University School of Medicine, Boston, USA, ²⁰Catholic University of Leuven, Leuven, Belgium, ²¹University of California San Francisco, San Francisco, USA, ²²University of Ferrara, Ferrara, Italy, ²³Ariel University, Jerusalem, Israel, ²⁴University of East Anglia, Norwich, UK, ²⁵Nanyang Technological University, Singapore, Singapore, ²⁶Yale School of Medicine, New Haven, USA, ²⁷University of California Irvine, Irvine, USA, ²⁸Ulm University, Ulm, Germany, ²⁹University of Dundee, Dundee, UK, ³⁰Liangzhu Laboratory & Zhejiang University, Hangzhou, China, ³¹Leiden University, Leiden, Netherlands, ³²University of Illinois, Chicago, USA, ³³University of Copenhagen, Copenhagen, Denmark, ³⁴Yale University, New Haven, USA, ³⁵University of Lausanne, Lausanne, Switzerland, ³⁶Strathclyde University, Glasgow, UK, ³⁷University College London, London, UK, ³⁸University of Queensland, St. Lucia, Australia, ³⁹Max Planck Institute for Heart and Lung Research/JW Goethe University, Bad Nauheim/Frankfurt, Germany, ⁴⁰University of Virginia, Charlottesville, USA, ⁴¹University of Pennsylvania, Philadelphia, USA, ⁴²Karolinska Institute, Stockholm, Sweden, ⁴³University of Oxford, Oxford, UK, ⁴⁴Medical University of Vienna, Vienna, Austria, ⁴⁵University of British Columbia, Vancouver, Canada, ⁴⁶University of Texas at Houston, Houston, USA, ⁴⁷University of California Davis, Davis, USA, ⁴⁸Imperial College, London, UK, ⁴⁹University of Michigan, Ann Arbor, USA, ⁵⁰University of Connecticut, Farmington, USA

Abstract

The Concise Guide to Pharmacology 2025/26 marks the seventh edition in this series of biennial publications in the *British Journal of Pharmacology*. Presented in landscape format, the guide provides a comparative overview of the pharmacology of drug target families. The concise nature of the Concise Guide refers to the style of presentation, being clear, accessible, and well-structured, rather than the scope of the content, which spans approximately 500 pages. The Concise Guide summarises the key pharmacological properties of around 1900 human drug targets, and nearly 7000 interactions, involving around 4400 ligands. While the content is a substantially condensed version of the more detailed information and links available at the www.guidetopharmacology.org website, the printed guide serves as a permanent, citable, point-in-time record, that remains stable despite ongoing updates to the online database. The full contents of this publication can be found at <https://bpspubs.onlinelibrary.wiley.com/doi/10.1111/bph.70231>.

The Concise Guides provide expert-curated recommendations of 'Gold Standard' selective pharmacological tools, available either commercially or as donations, which enable the identification of individual drug targets or families of drug targets. While the Concise Guide offers a more streamlined overview, more comprehensive information, including detailed pharmacological profiles and links to multiple online databases, is available through the Guide to Pharmacology website. The 2025/26 edition of the Concise Guide is based on material current as of mid-2025, and supersedes all previous editions, including the 2023/24 Guide, and earlier Guides to Receptors and Channels. It is produced in close conjunction with the Nomenclature and Standards Committee of the International Union of Basic and Clinical Pharmacology (NC-IUPHAR), and as such provides official IUPHAR classification and nomenclature for human drug targets, where applicable.

Ion channels are one of the six major pharmacological targets into which the Guide is divided, with the others being: G protein-coupled receptors, nuclear hormone receptors, catalytic receptors, enzymes and transporters. Each section includes nomenclature guidance, concise summaries, information of the best available pharmacological tools, key references, and suggestions for further reading.

Conflict of interest

The authors state that there are no conflicts of interest to disclose.

© 2025 The Author(s). *British Journal of Pharmacology* published by John Wiley & Sons Ltd on behalf of British Pharmacological Society.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Introduction: Ion channels are pore-forming proteins that allow the flow of ions across membranes, either plasma membranes, or the membranes of intracellular organelles [545]. Many ion channels (such as most Na, K, Ca and some Cl channels) are gated by voltage but others (such as certain K and Cl channels, TRP channels, ryanodine receptors and IP₃ receptors) are relatively voltage-insensitive and are gated by second messengers and other intracellular and/or extracellular mediators. As such, there is some blurring of the boundaries between "ion channels" and "ligand-gated channels" which are compiled separately in the Guide. Resolution of ion channel structures, beginning with K channels [340] then Cl channels [358] and most recently Na channels [1088] has greatly improved understanding of the structural basis behind ion channel function. Many ion channels (*e.g.*, K, Na, Ca, HCN and TRP channels) share several structural similarities. These channels are thought to have evolved from a common ancestor and have been classified together as the "voltage-gated-like (VGL) ion channel chanome" (see [1583]). Other ion channels, however, such as Cl channels, aquaporins and connexins, have completely different structural properties to the VGL channels, having evolved quite separately.

Currently, ion channels (including ligand-gated ion channels) represent the second largest target for existing drugs after G protein-coupled receptors [1064]. However, the advent of novel, faster screening techniques for compounds acting on ion channels [351] suggests that these proteins represent promising targets for the development of additional, novel therapeutic agents for the near future.

Family structure

S154	Ligand-gated ion channels	S183	Cyclic nucleotide-regulated channels (CNG)	S219	Aquaporins
S154	5-HT ₃ receptors	S185	Potassium channels	S221	Chloride channels
S156	Acid-sensing (proton-gated) ion channels (ASICs)	S185	Calcium- and sodium-activated potassium channels (K _{Ca} , K _{Na})	S222	ClC family
S158	Epithelial sodium channel (ENaC)	S187	Inwardly rectifying potassium channels (K _{IR})	S224	CFTR
S161	GABA _A receptors	S189	Two-pore domain potassium channels (K _{2P})	S225	Calcium activated chloride channel (CaCC)
S167	Glycine receptors	S192	Voltage-gated potassium channels (K _V)	S226	Maxi chloride channel
S169	Ionotropic glutamate receptors	S196	Ryanodine receptors (RyR)	S227	Volume regulated chloride channels (VRAC)
S174	IP ₃ receptors	S198	Transient Receptor Potential channels (TRP)	S228	Connexins and Pannexins
S175	Nicotinic acetylcholine receptors (nACh)	S213	Voltage-gated calcium channels (Ca _V)	–	Mitochondrial calcium uniporter (MCU) complex
S178	P2X receptors	S215	Voltage-gated proton channel (H _V 1)	S230	Piezo channels
S180	ZAC	S216	Voltage-gated sodium channels (Na _V)	S231	Sodium leak channel, non-selective (NALCN)
S181	Voltage-gated ion channels	S219	Other ion channels	–	Store-operated ion channels
S181	CatSper and Two-Pore channels (TPC)			S232	Orai channels

Searchable database: <http://www.guidetopharmacology.org/>

Full Contents of ConciseGuide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.70231/full>

Ion channels S153

Ligand-gated ion channels

[Ion channels](#) → [Ligand-gated ion channels](#)

Overview: Ligand-gated ion channels (LGICs) are integral membrane proteins that contain a pore which allows the regulated flow of selected ions across the plasma membrane. Ion flux is passive and driven by the electrochemical gradient for the permeant ions. These channels are open, or gated, by the binding of a neurotransmitter to an orthosteric site(s) that triggers a conformational change that results in the conducting state. Modulation of gating can occur by the binding of endogenous, or exogenous, modulators to allosteric sites. LGICs mediate fast synaptic transmission, on a millisecond time scale, in the nervous system and at the somatic neuromuscular junction. Such transmission involves the release of a neurotransmitter from a pre-synaptic neurone and the subsequent activation of post-synaptically located receptors that mediate a rapid, phasic, electrical signal (the excitatory, or inhibitory, post-synaptic potential). However, in addition to their traditional

role in phasic neurotransmission, it is now established that some LGICs mediate a tonic form of neuronal regulation that results from the activation of extra-synaptic receptors by ambient levels of neurotransmitter. The expression of some LGICs by non-excitable cells is suggestive of additional functions.

By convention, the LGICs comprise the excitatory, cation-selective, nicotinic acetylcholine [195, 951], 5-HT₃ [80, 1463], ionotropic glutamate [853, 1404] and P2X receptors [624, 1342] and the inhibitory, anion-selective, GABA_A [104, 1051] and glycine receptors [873, 1573]. The nicotinic acetylcholine, 5-HT₃, GABA_A and glycine receptors (and an additional zinc-activated channel) are pentameric structures and are frequently referred to as the Cys-loop receptors due to the presence of a defining loop of residues formed by a disulphide bond in the extracellular domain of their constituent subunits [955, 1378]. However, the

prokaryotic ancestors of these receptors contain no such loop and the term pentameric ligand-gated ion channel (pLGIC) is gaining acceptance in the literature [541]. The ionotropic glutamate and P2X receptors are tetrameric and trimeric structures, respectively. Multiple genes encode the subunits of LGICs and the majority of these receptors are heteromultimers. Such combinational diversity results, within each class of LGIC, in a wide range of receptors with differing pharmacological and biophysical properties and varying patterns of expression within the nervous system and other tissues. The LGICs thus present attractive targets for new therapeutic agents with improved discrimination between receptor isoforms and a reduced propensity for off-target effects. The development of novel, faster screening techniques for compounds acting on LGICs [351] will greatly aid in the development of such agents.

5-HT₃ receptors

[Ion channels](#) → [Ligand-gated ion channels](#) → [5-HT₃ receptors](#)

Overview: The 5-HT₃ receptor (**nomenclature as agreed by the NC-IUPHAR Subcommittee on 5-Hydroxytryptamine (serotonin) receptors** [576]) is a ligand-gated ion channel of the Cys-loop family that includes the zinc-activated channels, nicotinic acetylcholine, GABA_A and strychnine-sensitive glycine receptors. The receptor exists as a pentamer of 4 transmembrane (TM) subunits that form an intrinsic cation selective channel [80], but may also form intermediary tetramers in the cell membrane during assembly [600]. Five human 5-HT₃ receptor subunits have been cloned and homo-oligomeric assemblies of 5-HT_{3A} and hetero-oligomeric assemblies of 5-HT_{3A} and 5-HT_{3B} subunits have been characterised in detail. The 5-HT_{3C} (*HTR3C*, *Q8WXA8*), 5-HT_{3D} (*HTR3D*, *Q70Z44*) and 5-HT_{3E} (*HTR3E*, *ASX5Y0*) subunits [678, 1012], like the 5-HT_{3B} subunit, do not form functional homomers, but are reported to assemble with

the 5-HT_{3A} subunit to influence its functional expression rather than pharmacological profile [559, 1014, 1462]. 5-HT_{3A}, -C, -D, and -E subunits also interact with the chaperone RIC-3 which predominantly enhances the surface expression of homomeric 5-HT_{3A} receptor [330, 1462]. The co-expression of 5-HT_{3A} and 5-HT_{3C-E} subunits has been demonstrated in human colon [672]. A recombinant hetero-oligomeric 5-HT_{3AB} receptor has been reported to contain two copies of the 5-HT_{3A} subunit and three copies of the 5-HT_{3B} subunit in the order B-B-A-B-A [85], but this is inconsistent with recent reports which show at least one A-A interface [852, 1382]. The 5-HT_{3B} subunit imparts distinctive biophysical properties upon hetero-oligomeric 5-HT_{3AB} versus homo-oligomeric 5-HT_{3A} recombinant receptors [283, 347, 506, 627, 693, 1107, 1310], influences the potency of channel blockers, but generally has only a modest effect upon the apparent

affinity of agonists, or the affinity of antagonists ([147], but see [280, 301, 347]) which may be explained by the orthosteric binding site residing at an interface formed between 5-HT_{3A} subunits [852, 1382]. However, 5-HT_{3A} and 5-HT_{3AB} receptors differ in their allosteric regulation by some general anaesthetic agents, small alcohols and indoles [579, 1204, 1301]. The potential diversity of 5-HT₃ receptors is increased by alternative splicing of the genes *HTR3A* and *HTR3E* [159, 565, 1011, 1013, 1014]. In addition, the use of tissue-specific promoters driving expression from different transcriptional start sites has been reported for the *HTR3A*, *HTR3B*, *HTR3D* and *HTR3E* genes, which could result in 5-HT₃ subunits harbouring different N-termini [627, 1011, 1410]. To date, inclusion of the 5-HT_{3A} subunit appears imperative for 5-HT₃ receptor function.

Complexes

Nomenclature	5-HT ₃ AB	5-HT ₃ A
Subunits	5-HT ₃ A, 5-HT ₃ B	5-HT ₃ A
Agonists	CSTI-300 (Partial agonist) [1179], vortioxetine (Partial agonist) [74, 762]	CSTI-300 (Partial agonist) [1179]
Selective agonists	–	(S)-zacopride (Partial agonist) [147, 1006], meta-chlorophenylbiguanide [103, 283, 771, 963, 964], 2-methyl-5-HT [103, 283, 771, 963], SR57227A [361] – Rat, 1-phenylbiguanide [103]
Antagonists	–	metoclopramide (pK _i 6–6.4) [147, 566]
Selective antagonists	–	palonosetron (pK _i 10.5) [977], alosetron (pK _i 9.5) [548], granisetron (pK _i ~8.6–8.8) [566, 963], tropisetron (pK _i 8.5–8.8) [771, 963], ondansetron (pK _i ~7.8–8.3) [147, 566, 963]
Channel blockers	picrotoxinin (pIC ₅₀ 4.2) [1377], bilobalide (pIC ₅₀ 2.5) [1377], ginkgolide B (pIC ₅₀ 2.4) [1377]	picrotoxinin (pIC ₅₀ 5) [1376], TMB-8 (pIC ₅₀ 4.9) [1335], diltiazem (pIC ₅₀ 4.7) [1376], bilobalide (pIC ₅₀ 3.3) [1376], ginkgolide B (pIC ₅₀ 3.1) [1376]
Labelled ligands	–	[³ H]ramosetron (Antagonist) (pK _d 9.8) [963], [³ H]GR65630 (Antagonist) (pK _d 8.6–9.3) [548, 771], [³ H]granisetron (Antagonist) (pK _d 8.9) [147, 566], [³ H](S)-zacopride (Antagonist) (pK _d 8.7) [1078], [³ H]LY278584 (Antagonist) (pK _d 8.5) [5]
Functional Characteristics	γ = 0.4–0.8 pS [+ 5-HT ₃ B, γ = 16 pS]; inwardly rectifying current [+ 5-HT ₃ B, rectification reduced]; n _H 2–3 [+ 5-HT ₃ B 1–2]; relative permeability to divalent cations reduced by co-expression of the 5-HT ₃ B subunit	γ = 0.4–0.8 pS [+ 5-HT ₃ B, γ = 16 pS]; inwardly rectifying current [+ 5-HT ₃ B, rectification reduced]; n _H 2–3 [+ 5-HT ₃ B 1–2]; relative permeability to divalent cations reduced by co-expression of the 5-HT ₃ B subunit

Subunits

Nomenclature	5-HT ₃ A	5-HT ₃ B	5-HT ₃ C	5-HT ₃ D	5-HT ₃ E
HGNC, UniProt	HTR3A , P46098	HTR3B , O95264	HTR3C , Q8WXA8	HTR3D , Q70Z44	HTR3E , A5X5Y0
Functional Characteristics	γ = 0.4–0.8 pS [+ 5-HT ₃ B, γ = 16 pS]; inwardly rectifying current [+ 5-HT ₃ B, rectification reduced]; n _H 2–3 [+ 5-HT ₃ B 1–2]; relative permeability to divalent cations reduced by co-expression of the 5-HT ₃ B subunit	γ = 0.4–0.8 pS [+ 5-HT ₃ B, γ = 16 pS]; inwardly rectifying current [+ 5-HT ₃ B, rectification reduced]; n _H 2–3 [+ 5-HT ₃ B 1–2]; relative permeability to divalent cations reduced by co-expression of the 5-HT ₃ B subunit	–	–	–

Comments: Quantitative data in the table refer to homooligomeric assemblies of the human 5-HT₃A subunit, or the receptor native to human tissues. Significant changes introduced by co-expression of the 5-HT₃B subunit are indicated in parenthesis. Although not a selective antagonist, **methadone** displays multimodal and subunit-dependent antagonism of 5-HT₃ receptors [301]. Similarly, **TMB-8**, **diltiazem**, **picrotoxin**, **bilobalide** and **ginkgolide B** are not selective for 5-HT₃ receptors (*e.g.* [1377]). The anti-malarial drugs **mefloquine** and **quinine** exert a modestly

more potent block of 5-HT₃A versus 5-HT₃AB receptor-mediated responses [1379]. Known better as a partial agonist of nicotinic acetylcholine α4β2 receptors, **varenicline** is also an agonist of the 5-HT₃A receptor [869]. Human [103, 963], rat [605], mouse [910], guinea-pig [771] ferret [965] and canine [631] orthologues of the 5-HT₃A receptor subunit have been cloned that exhibit intraspecies variations in receptor pharmacology. Notably, most ligands display significantly reduced affinities at the guinea-pig 5-HT₃ receptor in comparison with other species. In addition to the

agents listed in the table, native and recombinant 5-HT₃ receptors are subject to allosteric modulation by extracellular divalent cations, alcohols, several general anaesthetics and 5-hydroxy- and halide-substituted indoles (see reviews [32, 33, 1079, 1380, 1381, 1463]). The use of molecular dynamic simulation revealed cryptic binding pockets in cryo-EM structures of 5-HT₃AR, identifying a binding site for the positive allosteric modulator BrAmp, which was validated using site-directed mutagenesis with electrophysiological analysis [148, 503].

Further reading on 5-HT₃ receptors

- Andrews PL *et al.* (2014) Nausea and the quest for the perfect anti-emetic. *Eur J Pharmacol* **722**: 108-21 [PMID:24157981]
- Barnes NM *et al.* (2021) International Union of Basic and Clinical Pharmacology. CX. Classification of Receptors for 5-hydroxytryptamine; Pharmacology and Function. *Pharmacol Rev* **73**: 310-520 [PMID:33370241]
- Chang-Halabi Y *et al.* (2023) Crosstalk interactions between P2X₄ and 5-HT_{3A} receptors. *Neuropharmacology* **236**: 109574 [PMID:37156336]
- Fakhfour G *et al.* (2015) From Chemotherapy-Induced Emesis to Neuroprotection: Therapeutic Opportunities for 5-HT₃ Receptor Antagonists. *Mol Neurobiol* **52**: 1670-1679 [PMID:25377794]
- Gibbs E *et al.* (2021) Structure, Function and Physiology of 5-Hydroxytryptamine Receptors Subtype 3. *Subcell Biochem* **96**: 373-408 [PMID:33252737]
- Gupta D *et al.* (2016) 5HT₃ receptors: Target for new antidepressant drugs. *Neurosci Biobehav Rev* **64**: 311-25 [PMID:26976353]
- Lochner M *et al.* (2015) A review of fluorescent ligands for studying 5-HT₃ receptors. *Neuropharmacology* **98**: 31-40 [PMID:25892507]
- Merecz K *et al.* (2023) An overview of 5-HT₃ receptor antagonists as a treatment option for irritable bowel syndrome with diarrhea. *Expert Opin Pharmacother* **24**: 1189-1198 [PMID:37173833]
- Rojas C *et al.* (2014) Molecular mechanisms of 5-HT₃ and NK(1) receptor antagonists in prevention of emesis. *Eur J Pharmacol* **722**: 26-37 [PMID:24184669]

Acid-sensing (proton-gated) ion channels (ASICs)

Ion channels → **Ligand-gated ion channels** → **Acid-sensing (proton-gated) ion channels (ASICs)**

Overview: Acid-sensing ion channels (ASICs, **nomenclature as agreed by NC-IUPHAR** [32, 33, 692]) are members of a Na⁺ channel superfamily that includes the epithelial Na⁺ channel (ENaC), the FMRF-amide activated channel (FaNaC) of invertebrates, the degenerins (DEG) of *Caenorhabditis elegans*, channels in *Drosophila melanogaster* and the mammalian bile acid-activated ion channel BASIC [1511], previously known as BLINaC [1213] and INaC [1231]. ASIC subunits contain 2 TM domains and a large extracellular part whose shape resembles that of a hand, as shown by high-resolution structures of chicken and human ASIC1a [62, 461, 626, 1334, 1576, 1577]. They assemble as homo- or heterotrimers to form proton-gated, voltage-insensitive, Na⁺ permeable, channels that are activated by levels of acidosis occurring in both physiological and pathophysiological conditions with ASIC3 also playing a role in mechanosensation (reviewed in [225, 484, 692, 1187, 1506]). Splice variants of ASIC1 [termed ASIC1a (ASIC, ASIC α , BNaC2 α) [1455], ASIC1b (ASIC β , BNaC2 β) [207] and ASIC1b2 (ASIC β 2) [1412]; note that ASIC1a is also permeable to Ca²⁺], ASIC2 [termed ASIC2a (MDEG1, BNaC1 α , BNC1 α) [429, 1138, 1457] and ASIC2b (MDEG2, BNaC1 β) [832]] differ in the

first third of the protein. Unlike ASIC2a (listed in table), heterologous expression of ASIC2b alone does not support H⁺-gated currents. A third member, ASIC3 (DRASIC, TNaC1) [1454] is one of the most pH-sensitive isoforms (along with ASIC1a) and has the fastest activation and desensitisation kinetics, however can also carry small sustained currents. ASIC4 (SPASIC) evolved as a proton-sensitive channel but seems to have lost this function in mammals [871]. Mammalian ASIC4 does not support a proton-gated channel in heterologous expression systems but is reported to downregulate the expression of ASIC1a and ASIC3 [19, 228, 335, 483, 828]. ASICs channels are primarily expressed in central (ASIC1a, -2a, 2b and -4) and peripheral neurons including nociceptors (ASIC1-3) where they participate in neuronal sensitivity to acidosis. Humans express, in contrast to rodents, ASIC3 also in the brain [304]. ASICs have also been detected in photoreceptors and retinal cells (ASIC1-3), cochlear hair cells (ASIC1b), testis (hASIC3), pituitary gland (ASIC4), lung epithelial cells (ASIC1a and -3), urothelial cells, adipose cells (ASIC3), vascular smooth muscle cells (ASIC1-3), immune cells (ASIC1,-3 and -4) and bone (ASIC1-3) (ASIC distribution is reviewed in [312, 482, 829]). A

neurotransmitter-like function of protons has been suggested, involving postsynaptically located ASICs of the CNS in functions such as learning and fear perception [344, 747, 1633] and of the PNS in mechanoreceptor-neurite transmission [1550, 1551]. ASIC activation also contributes to cell damage in focal ischemia [1171, 1538] and autoimmune inflammation (arthritis and multiple sclerosis) [415, 1549], as well as neuron activation during seizures and pain [134, 313, 314, 327, 1506]. Heterologously expressed heteromultimers form ion channels with differences in kinetics, ion selectivity, pH- sensitivity and sensitivity to blockers that resemble some of the native proton activated currents recorded from neurons [60, 84, 375, 832]. In general, the known small molecule inhibitors of ASICs are non-selective or partially selective, whereas the venom peptide inhibitors have substantially higher selectivity and potency. Several clinically used drugs are known to inhibit ASICs, however they are generally more potent at other targets (e.g. **amiloride** at ENaCs, **ibuprofen** at COX enzymes) [1060, 1168]. The information in the tables below are for the effects of inhibitors on homomeric channels, for information of known effects on heteromeric channels see the comments below.

Nomenclature	ASIC1	ASIC2
HGNC, UniProt	ASIC1 , P78348	ASIC2 , Q16515
Endogenous activators	Extracellular H⁺ (ASIC1a) (pEC ₅₀ ~6.2–6.8), Extracellular H⁺ (ASIC1b) (pEC ₅₀ ~5.1–6.2)	Extracellular H⁺ (pEC ₅₀ ~4.1–5)
Channel blockers	Pi-hexatoxin-Hi1a (ASIC1a) (pIC ₅₀ ~9.3) [158], psalmotoxin 1 (ASIC1a) (pIC ₅₀ 9) [375], Pi-theraphotoxin-Hm3a (ASIC1a) (pIC ₅₀ ~8.5) [369], Zn²⁺ (ASIC1a) (pIC ₅₀ ~8.2) [234], JNJ-799760 (ASIC1a @pH6.0) (pIC ₅₀ 7.6) [849], compound 5b (ASIC1a @pH6.7) (pIC ₅₀ 7.6) [165], JNJ-67869386 (ASIC1a @pH6.0) (pIC ₅₀ 7.5) [849], mambalgin-1 (ASIC1a) (pIC ₅₀ ~7.3) [327], 6-iodoamiloride (ASIC1a) (pIC ₅₀ 7.1) [403], ASC06-IgG1 (Inhibition) (pIC ₅₀ ~7.1) [1147], mambalgin-1 (ASIC1b) (pIC ₅₀ ~7) [81], diminazene (ASIC1a & ASIC1b) (pIC ₅₀ ~6.5) [743, 783, 1237], NS383 (ASIC1a @pH6.5) (pIC ₅₀ 6.4) [982], compound 3b (ASIC1a @pH6.0) (pIC ₅₀ 6.4) [466], Pb²⁺ (ASIC1b) (pIC ₅₀ ~5.8), A-317567 (ASIC1a) (pIC ₅₀ ~5.7) [346] – Rat, Pb²⁺ (ASIC1a) (pIC ₅₀ ~5.4) [1482], compound 5b (ASIC1a @pH5.0) (pIC ₅₀ 5.2) [165], amiloride (ASIC1a) (pIC ₅₀ 5), benzamil (ASIC1a) (pIC ₅₀ 5) [1455], ethylisopropylamiloride (ASIC1a) (pIC ₅₀ 5) [1455], nafamostat (ASIC1a) (pIC ₅₀ ~4.9) [1411], amiloride (ASIC1b) (pIC ₅₀ 4.7) [1455], flurbiprofen (ASIC1a) (pIC ₅₀ 3.5) [1444] – Rat, ibuprofen (ASIC1a) (pIC ₅₀ ~3.5) [872, 1444]	diminazene (pIC ₅₀ ~6.1) [783], amiloride (pIC ₅₀ 4.6) [1457], A-317567 (pIC ₅₀ ~4.5) [346], nafamostat (pIC ₅₀ ~4.2) [1411], Cd²⁺ (Partial inhibition) (pIC ₅₀ ~3) [1309]
Labelled ligands	[¹²⁵ I]psalmotoxin 1 (ASIC1a) (pK _d 9.7) [1218]	–
Functional Characteristics	ASIC1a : γ =14 pS P _{Na} /P _K = 5–13, P _{Na} /P _{Ca} =2.5 rapid activation rate (5.8–13.7 ms), rapid inactivation rate (1.2–4 s) @ pH 6.0, slow recovery (5.3–13s) @ pH 7.4 ASIC1b : γ =19 pS P _{Na} /P _K =14.0, P _{Na} ≫P _{Ca} rapid activation rate (9.9 ms), rapid inactivation rate (0.9–1.7 s) @ pH 6.0, slow recovery (4.4–7.7 s) @ pH 7.4	γ =10.4–13.4 pS P _{Na} /P _K =10, P _{Na} /P _{Ca} = 20 rapid activation rate, moderate inactivation rate (3.3–5.5 s) @ pH 5
Comments	ASIC1a and ASIC1b are activated by the heteromeric Texas coral snake toxin MitTx, with pEC ₅₀ values of ~8 and ~7.6 respectively [134]. The inhibition by amiloride and diminazene decreases at more positive voltages. The inhibitory potency of compound 5b, JNJ-67869386, JNJ-799760, ibuprofen and NS383 is decreased at more acidic pH [165, 849, 982].	ASIC2 is also blocked by other diarylamidines [216]. The inhibition by amiloride and diminazene decreases at more positive voltages.

Nomenclature	ASIC3
HGNC, UniProt	ASIC3 , Q9UHC3
Endogenous activators	Extracellular H⁺ (transient component) (pEC ₅₀ ~6.2–6.7), lysophosphatidylcholine (Partial agonist) (pEC ₅₀ 5.4) [912], Extracellular H⁺ (sustained component) (pEC ₅₀ ~3.5–4.3)
Activators	GMQ (largly non-desensitizing; at pH 7.4) (pEC ₅₀ ~3) [1590], arcaine (at pH 7.4) (pEC ₅₀ ~2.9) [813], agmatine (at pH 7.4) (pEC ₅₀ ~2) [813]
Channel blockers	APETx2 (transient component only) (pIC ₅₀ 7.2) [326], diminazene (pIC ₅₀ ~6.5) [783], A-317567 (pIC ₅₀ 6) [755], NS383 (@ pH6.5; inhibition of rat but not human ASIC3) (pIC ₅₀ 5.7) [982] – Rat, nafamostat (transient component) (pIC ₅₀ ~5.6) [1411], Ugr 9-1 (transient component) (pIC ₅₀ 5) [1061], amiloride (transient component only - sustained component enhanced by 200μM amiloride at pH 4) (pIC ₅₀ 4.2–4.8) [1454], Gd³⁺ (pIC ₅₀ 4.4) [60], Zn²⁺ (pIC ₅₀ 4.2) [637], aspirin (sustained component) (pIC ₅₀ 4) [1444], diclofenac (sustained component) (pIC ₅₀ 4) [1444], salicylic acid (sustained component) (pIC ₅₀ 3.6) [1444]
Functional Characteristics	γ =13–15 pS; biphasic response consisting of rapidly inactivating transient and sustained components; very rapid activation (<5 ms) and inactivation (0.4 s); fast recovery (0.4–0.6 s) @ pH 7.4, transient component partially inactivated at pH 7.2
Comments	ASIC3 is activated by Mit-Toxin (pEC ₅₀ ~6.1) [134]. The inhibition by amiloride and diminazene decreases at more positive voltages. The inhibitory potency of NS383 is decreased at more acidic pH [982].

Comments: **Psalmotoxin 1** (PcTx1) inhibits ASIC1a by increasing the affinity to H⁺ and promoting channel desensitization [214, 375]. PcTx1 has little effect on ASIC2a, ASIC3 or ASIC1a expressed as a heteromultimer with ASIC3 or ASIC2 if the holding pH between acidic stimulations is 7.4 [375, 644]. PcTx1 and π -Hm3a potentiate ASIC1b currents [215, 369]. ASIC1-containing homo- and heteromers are inhibited by Mambalgins, toxins contained in the black mamba venom, which induce in ASIC1a an acidic shift of the pH dependence of activation [327]. π -Hi1a is selective for ASIC1a with mild potentiating activity at ASIC1b. It inhibits channel activation and is very slowly reversible [198]. **APETx2** most potently blocks homomeric ASIC3 channels, but also ASIC2b+ASIC3, ASIC1b+ASIC3, and ASIC1a+ASIC3 heteromeric channels with IC₅₀ values of 117 nM, 900 nM and 2 μ M, respectively. **APETx2** has no effect on ASIC1a or ASIC2a+ASIC3, however, it does potentiate ASIC1b and ASIC2a homomers in the low micromolar range (1–10 μ M) [326, 328, 783]. **APETx2** however also inhibits voltage-gated Na⁺ channels [125, 1097]. The antibody **ASC06-IgG1** binds to the structurally intact channel in the upper part of the extracellular domain with substantial contact on the finger domain and is highly selective for ASIC1a over other subtypes [1147]. IC₅₀ value for **A-317567** was determined using high throughput electrophysiology on human ASIC3 expressed in HEK293 cells [755]. For some of the newer small molecule inhibitors it is not known whether they inhibit ion channels in addition to ASICs [165, 849, 982]. The pEC₅₀ values for proton activation of ASIC channels are influenced

by numerous factors including extracellular di- and poly-valent ions, Zn²⁺, protein kinase C and serine proteases (reviewed in [692, 1506]). Rapid acidification is required for activation of ASIC1 and ASIC3 due to fast inactivation/desensitization. pEC₅₀ values for H⁺-activation of either transient, or sustained, currents mediated by ASIC3 vary in the literature and may reflect species and/or methodological differences [61, 290, 1454]. The transient ASIC current component is Na⁺-selective (P_{Na}/P_K of about 10) [1454, 1561] whereas the sustained current component that is observed with ASIC3 and some ASIC heteromers is non-selective between Na⁺ and K⁺ [290]. The reducing agents dithiothreitol (DTT) and **glutathione** (GSH) increase ASIC1a currents expressed in CHO cells and ASIC-like currents in sensory ganglia and central neurons [45, 233] whereas oxidation, through the formation of intersubunit disulphide bonds, reduces currents mediated by ASIC1a [1601]. ASIC1a is also irreversibly modulated by extracellular serine proteases, such as trypsin, through proteolytic cleavage [1449]. Non-steroidal anti-inflammatory drugs (NSAIDs) are direct inhibitors of ASIC currents (reviewed in [82]). Extracellular Zn²⁺ potentiates proton activation of homomeric and heteromeric channels incorporating ASIC2a, but not homomeric ASIC1a or ASIC3 channels [83]. However, removal of contaminating Zn²⁺ by chelation reveals a high affinity block of homomeric ASIC1a and heteromeric ASIC1a+ASIC2 channels by Zn²⁺ indicating complex biphasic actions of the divalent [234]. Nitric oxide potentiates submaximal currents activated by H⁺ mediated by ASIC1a, ASIC1b, ASIC2a and ASIC3 [168].

The positive modulation of homomeric, heteromeric and native ASIC channels by the peptide **FMRFamide** and related substances, such as neuropeptides FF and SF, is reviewed in detail in [469, 1433]. Inflammatory conditions and particular pro-inflammatory mediators such as **arachidonic acid** induce overexpression of ASIC-encoding genes and enhance ASIC currents [314, 907, 1289]. The sustained current component mediated by ASIC3 is potentiated by hypertonic solutions in a manner that is synergistic with the effect of **arachidonic acid** [314]. ASIC3 is partially activated by the lipids lysophosphatidylcholine (LPC) and arachidonic acid [912]. Mit-Toxin, which is contained in the venom of the Texas coral snake, activates several ASIC subtypes [134]. Selective activation of ASIC3 by GMQ, likely by binding to the central vestibule, is potentiated by mild acidosis and reduced extracellular Ca²⁺ [1590]. *Additional notes on the channels:* Until recently they were thought to be vertebrate specific channels, however are now known to have evolved over 600 million years ago and appear to be conserved throughout the superphylum of animals known as deuterostomes (which includes vertebrates, tunicates, starfish, sea urchins, sea cucumbers and acorn worms) [871]. Recently an ion-conducting-independent signaling mechanism has been revealed for ASIC1a whereby the acidosis-activated channel recruits RIPK1 to its C-terminus resulting in RIPK1 phosphorylation and activation of necroptosis. This pathway is suggested to be the primary cause of ASIC-mediated neuronal cell death in ischemic stroke [1475, 1491].

Further reading on Acid-sensing (proton-gated) ion channels (ASICs)

Gründer S *et al.* (2024) Acid-sensing ion channels and downstream signalling in cancer cells: is there a mechanistic link? *Pflugers Arch* **476**: 659–672 [PMID:38175291]
 Gründer S. (2020) *In The Oxford Handbook of Neuronal Ion Channels* Oxford University Press:
 Kellenberger S *et al.* (2015) International Union of Basic and Clinical Pharmacology. XCI. structure, function, and pharmacology of acid-sensing ion channels and the epithelial Na⁺ channel. *Pharmacol Rev* **67**: 1–35 [PMID:25287517]

Rook ML *et al.* (2021) Coupling structure with function in acid-sensing ion channels: challenges in pursuit of proton sensors. *J Physiol* **599**: 417–430 [PMID:32306405]
 Verkest C *et al.* (2022) Mechanisms of Action of the Peptide Toxins Targeting Human and Rodent Acid-Sensing Ion Channels and Relevance to Their In Vivo Analgesic Effects. *Toxins (Basel)* **14**: [PMID:36287977]

Epithelial sodium channel (ENaC)

Ion channels → Ligand-gated ion channels → Epithelial sodium channel (ENaC)

Overview

Epithelial sodium channels (ENaC) are located on the apical membrane of epithelial cells in the kidney tubules, lungs, respiratory tract, male and female reproductive tracts, sweat and salivary

glands, placenta, colon, and several other organs [171, 348, 512, 513, 1260]. In these epithelia, Na⁺ ions enter epithelial cells from the extracellular fluid *via* ENaC and are subsequently pumped out

into the interstitial fluid by the Na⁺/K⁺-ATPase on the basolateral membrane [1123]. Because sodium is a major electrolyte in the extracellular fluid (ECF), the osmotic changes caused by sodium

Searchable database: <http://www.guidetopharmacology.org/>

Full Contents of ConciseGuide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.70231/full>

Epithelial sodium channel (ENaC) S158

flux are accompanied by parallel water movement [143]. Thus, ENaC plays a central role in regulating ECF volume and blood pressure, primarily through its function in the kidney [1190]. The expression of ENaC subunits- and therefore its activity- is controlled by the renin-angiotensin-aldosterone system and other factors involved in electrolyte homeostasis [722, 1190].

Genetic studies of systemic pseudohypoaldosteronism type I revealed that ENaC activity depends on three essential subunits encoded by three separate genes encoding homologous proteins [194, 513]. Within the wider protein superfamily that includes ENaC, the first crystal structure determined was that of ASIC, which revealed a trimeric structure with a large extracellular domain anchored in the membrane by a bundle of six transmembrane helices (two per subunit) [62, 626]. The first 3D structure of human ENaC was determined using single-particle cryo-electron

microscopy at 3.7Å resolution [1031], later improved to 3.0Å [1032]. These structures confirmed that ENaC has a quaternary structure similar to ASIC. ENaC assembles as a heterotrimer, with α -, γ -, and β -subunits arranged in clockwise order when viewed from above [252]. In contrast to ASIC1, which can form a functional homotrimer, ENaC is only fully functional as a heterotrimer composed of either $\alpha\beta\gamma$ or $\delta\beta\gamma$ [692]. Recently, Houser and Bacongus co-expressed human δ , β , and γ and determined the structures of complexes using single-particle cryoelectron microscopy. The structures showed that β and γ positions are conserved among the different complexes while the α position in the $\alpha\beta\gamma$ trimer is occupied by either δ or another β [573].

In the respiratory and female reproductive tracts, large regions of the epithelium consist of multiciliated cells with a microtubule-based cytoskeleton. In these cells, ENaC is distributed along the

entire length of the cilia [368]. This localization substantially increases ENaC density on the cell surface and enables precise regulation of periciliary fluid osmolarity throughout its depth [368]. In the *vas deferens* of the male reproductive tract, the luminal surface is covered with microvilli and stereocilia supported by actin bundles [1260]. In these cells, both ENaC and the aquaporin AQP9 are localized to the projections as well as to the basal and smooth muscle layers [1260]. In contrast, CFTR- the chloride channel defective in cystic fibrosis- is confined to the apical cell surface but is absent from cilia and microvilli [368, 1260]. Collectively, ENaC function regulates epithelial fluid volume, which is essential for mucociliary clearance in the respiratory tract, gamete transport, fertilization, implantation, and cell migration [368, 513, 919].

Genes and Phylogeny

The human genome contains four homologous genes (*SCNNIA*, *SCNNIB*, *SCNNIG*, and *SCNNID*) encoding the α -, β -, γ -, and δ -ENaC subunits, respectively [172, 833, 1229, 1456]. These subunits share 23-34% sequence identity and <20% identity with ASIC subunits [513]. Genes encoding all four ENaC subunits are

present in bony vertebrates, except in ray-finned fishes, which have lost them entirely. The mouse genome has also lost *SCNNID*, the gene for δ -ENaC [457, 513]. The α -, β -, and γ -ENaC genes are present in jawless vertebrates (*e.g.*, lampreys) and cartilaginous fishes (*e.g.*, sharks) [513].

Methylation analysis of the 5'-flanking regions of *SCNNIA*, *SCNNIB*, and *SCNNIG* in human cells revealed an inverse correlation between gene expression and DNA methylation, suggesting epigenetic transcriptional control of ENaC genes [1115].

Channel biogenesis, assembly and function

ENaC subunit expression is regulated primarily by aldosterone and by numerous other extracellular and intracellular factors [719, 1069, 1190]. Most studies indicate that expression of the three subunits is not tightly coordinated [166]. However, transport of the subunits to the membrane requires all three intact subunits, and even a single missense mutation can reduce the number of assembled channels on the cell surface [359]. ENaC is constitutively active, meaning Na^+ flow does not require

an activating factor. Thus, heterologous cells expressing ENaC (*e.g.*, human cRNAs in *Xenopus* oocytes) must be maintained in amiloride-containing solutions to block channel activity. ENaC activity is then measured by replacing the bath with amiloride-free solution. The channel alternates between two states: 1) open and 2) closed. The probability of ENaC being open is referred to as open probability (P_o). ENaC regulation involves two key parameters: (1) membrane channel density and (2) open probability [680, 692].

Open probability is markedly reduced by extracellular Na^+ in a process known as sodium self-inhibition [123, 568, 1265]. A key regulatory feature is that the α - and γ -subunits contain conserved extracellular serine protease cleavage sites [513]. Proteolytic cleavage by enzymes such as furin and plasmin activates ENaC [37, 718, 1191].

Diseases associated with ENaC mutations

Mutations in *SCNNIA*, *SCNNIB*, or *SCNNIG* can cause partial or complete loss of ENaC activity [194, 510]. Such loss-of-function mutations are associated with systemic or multi-system autosomal recessive pseudohypoaldosteronism type I (OMIM abbreviation: PHA1B) [194, 368, 509, 513, 1228, 1599]. No PHA-causing mutations have been identified in *SCNNID*. Patients with PHA experience severe salt wasting in all aldosterone target organs ex-

pressing ENaC, including kidney, sweat glands, salivary glands, and respiratory tract. In infancy and early childhood, the resulting electrolyte disturbances, dehydration, and acidosis often require recurrent hospitalization. The frequency and severity of salt-wasting episodes generally improve with age [511]. PHA1B also affects female reproductive system function [132, 368]. The ENaC carboxy-terminal region contains a short consensus

sequence called the PY motif. Mutations in this motif in *SCNNIB* and *SCNNIG* are associated with Liddle syndrome, a disorder marked by early-onset hypertension [131, 1270]. The PY motif is recognized by Nedd4-2, a ubiquitin ligase. Mutations that disrupt this recognition reduce ENaC ubiquitylation, leading to channel accumulation at the membrane and increased ENaC activity [1192].

ENaC expression in tumors

Intracellular sodium concentrations are often elevated in cancer cells compared with normal cells, leading to the hypothesis that ENaC overexpression may contribute to metastasis [799]. However, RNA-seq analysis of ENaC genes and clinical data of

cervical cancer patients from The Cancer Genome Atlas (TCGA) revealed a negative correlation with histologic grades of tumor [1302]. Similarly, in breast cancer cells, overexpression or siRNA-mediated knockdown of α -ENaC showed that higher α -ENaC

levels suppress cell proliferation [1493]. In contrast, TCGA data showed that elevated *SCNNIA* expression correlates with poor prognosis in ovarian cancer [858]. Thus, the role of ENaC in tumorigenesis appears to be tissue-specific.

COVID-19

SARS-CoV-2 virions, the cause of COVID-19, are covered with glycosylated spike (S) proteins. These proteins bind to membrane-bound ACE2 as the first step of viral entry. Entry depends on S-

protein cleavage (at Arg-667/Ser-668) by a serine protease. Anand *et al.* identified a sequence motif at this cleavage site homologous to the furin cleavage site in ENaC α [38]. A comprehensive review

of COVID-19 pathophysiology suggests a role for ENaC in the early stages of infection in respiratory epithelia [441].

ENaC Inhibitors for Cystic Fibrosis

Cystic fibrosis (CF) is the most common life-limiting autosomal recessive disorder among Caucasians. CF is caused by mutations in the gene that codes for CFTR (Cystic Fibrosis Transmembrane Conductance Regulator). CFTR is a chloride and bicarbonate channel located on the apical membrane [262]. CFTR-mediated movement of Cl⁻ and HCO₃⁻ ions into the lumen also drives water flow into the lumen by osmosis. CFTR is expressed in many tissues but the most severe effect of mutated CFTR is observed in the respiratory tract in the form of airway surface liquid (ASL) depletion which leads to mucus accumulation, inflammation and bacterial infections that lead to mortality.

Normal CFTR activity inhibits ENaC by causing a reduction in

surface expression of ENaC as well as its P_o [1169]. In many epithelia, ENaC and CFTR are not co-localized on the apical membrane indicating that the two channels do not directly interact [368, 1258, 1260]. In the absence of a functional CFTR, ENaC activity increases (also named as “hyperactivated ENaC”), leading to increased Na⁺ and water absorption and consequently ASL depletion. These observations have led to the development of inhibitors targeting ENaC in the airways to ameliorate ASL dehydration [794, 903, 1261]. Most of the ENaC inhibitors in development are designed for application by inhalation and improved lung retention to avoid damaging the vital activity of ENaC in other tissues [277, 749]. Despite the development of many ENaC

inhibitors for CF [275], nearly all drug candidates were discontinued at Preclinical, Phase 1 or Phase 2 stage of clinical trials [277, 794, 903]. However, there is one candidate, EDT001, that is still under Phase 2 clinical trial [277]. The persistence of the scientists and the drug companies and the versatility of their approaches provide hope that an appropriate useful treatment will emerge to help CF patients.

Thus, insights into the physiological and pathological roles of ENaC across diverse tissues continue to guide the development of targeted inhibitors, with cystic fibrosis representing a prominent example where modulation of ENaC activity may translate fundamental biology into therapeutic benefit.

Complexes

Nomenclature	ENaC $\alpha\beta\gamma$
Subunits	ENaC α , ENaC β , ENaC γ
Activators	S3969 (pEC ₅₀ 5.9) [864]
Inhibitors	B11265162 (pIC ₅₀ 8.5) [525] – Mouse, EDT001 (pIC ₅₀ 7.2) [275]
Channel blockers	P552-02 (pIC ₅₀ 8.1), benzamil (pIC ₅₀ ~8), amiloride (pIC ₅₀ 6.7–7), triamterene (pIC ₅₀ ~5.3) [172, 691]

Subunits

Nomenclature	ENaC α	ENaC β	ENaC γ	ENaC δ
HGNC, UniProt	<i>SCNNIA</i> , P37088	<i>SCNN1B</i> , P51168	<i>SCNN1G</i> , P51170	<i>SCNN1D</i> , P51172

Further reading on Epithelial sodium channel (ENaC)

- Hanukoglu I. (2017) ASIC and ENaC type sodium channels: conformational states and the structures of the ion selectivity filters. *FEBS J* **284**: 525-545 [PMID:27580245]
- Houser A *et al.* (2025) Structural insights into subunit-dependent functional regulation in epithelial sodium channels. *Structure* **33**: 349-362.e4 [PMID:39667931]
- Kellenberger S *et al.* (2015) International Union of Basic and Clinical Pharmacology. XCI. structure, function, and pharmacology of acid-sensing ion channels and the epithelial Na⁺ channel. *Pharmacol Rev* **67**: 1-35 [PMID:25287517]
- Kleyman TR *et al.* (2020) Regulating ENaC's gate. *Am J Physiol Cell Physiol* **318**: C150-C162 [PMID:31721612]
- Rossier BC *et al.* (2015) Epithelial sodium transport and its control by aldosterone: the story of our internal environment revisited. *Physiol Rev* **95**: 297-340 [PMID:25540145]
- Sharma S *et al.* (2020) High-resolution imaging of the actin cytoskeleton and epithelial sodium channel, CFTR, and aquaporin-9 localization in the vas deferens. *Mol Reprod Dev* **87**: 305-319 [PMID:31950584]

GABA_A receptors

Ion channels → Ligand-gated ion channels → GABA_A receptors

Overview: The GABA_A receptor is a ligand-gated ion channel of the Cys-loop family that includes the nicotinic acetylcholine, 5-HT₃ and strychnine-sensitive glycine receptors. GABA_A receptor-mediated inhibition within the CNS occurs by fast synaptic transmission, sustained tonic inhibition and temporally intermediate events that have been termed 'GABA_A, slow' [179]. GABA_A receptors exist as pentamers of 4TM subunits that form an intrinsic anion selective channel. Sequences of six α , three β , three γ , one δ , three ρ , one ϵ , one π and one θ GABA_A receptor subunits have been reported in mammals [1051, 1052, 1274, 1278]. The π -subunit is restricted to reproductive tissue. Alternatively spliced versions of many subunits exist (e.g. $\alpha 4$ - and $\alpha 6$ - (both not functional) $\alpha 5$ -, $\beta 2$ -, $\beta 3$ - and $\gamma 2$), along with RNA editing of the $\alpha 3$ subunit [279]. The three ρ -subunits, ($\rho 1$ -3) function as either homo- or hetero-oligomeric assemblies [205, 1602]. Receptors formed from ρ -subunits, because of their distinctive pharmacology that includes insensitivity to bicuculline, benzodiazepines and barbiturates, have sometimes been termed GABA_C receptors [1602], **but they are classified as GABA_A receptors by NC-IUPHAR on the basis of structural and functional criteria** [79, 1051, 1052].

Many GABA_A receptor subtypes contain α -, β - and γ -subunits with the likely stoichiometry $2\alpha.2\beta.1\gamma$ [733, 1051]. It is thought that the majority of GABA_A receptors harbour a single type of α - and β -subunit variant. The $\alpha 1\beta 2\gamma 2$ hetero-oligomer constitutes the

largest population of GABA_A receptors in the CNS, followed by the $\alpha 2\beta 3\gamma 2$ and $\alpha 3\beta 3\gamma 2$ isoforms. Receptors that incorporate the $\alpha 4$ - $\alpha 5$ - or $\alpha 6$ -subunit, or the $\beta 1$ -, $\gamma 1$ -, $\gamma 3$ -, δ -, ϵ - and θ -subunits, are less numerous, but they may nonetheless serve important functions. For example, extrasynaptically located receptors that contain $\alpha 6$ - and δ -subunits in cerebellar granule cells, or an $\alpha 4$ - and δ -subunit in dentate gyrus granule cells and thalamic neurones, mediate a tonic current that is important for neuronal excitability in response to ambient concentrations of GABA [104, 388, 966, 1251, 1295]. GABA binding occurs at the β +/ α -subunit interface and the homologous γ +/ α -subunits interface creates the benzodiazepine site. A second site for benzodiazepine binding has recently been postulated to occur at the α +/ β - interface ([1161]; reviewed by [1277]). The particular α - and γ -subunit isoforms exhibit marked effects on recognition and/or efficacy at the benzodiazepine site. Thus, receptors incorporating either $\alpha 4$ - or $\alpha 6$ -subunits are not recognised by 'classical' benzodiazepines, such as flunitrazepam (but see [1578]). The trafficking, cell surface expression, internalisation and function of GABA_A receptors and their subunits are discussed in detail in several recent reviews [222, 613, 870, 1438] but one point worthy of note is that receptors incorporating the $\gamma 2$ subunit (except when associated with $\alpha 5$) cluster at the postsynaptic membrane (but may distribute dynamically between synaptic and extrasynaptic locations), whereas those incorporating the δ subunit appear to be exclusive-

ly extrasynaptic.

NC-IUPHAR [32, 33, 79, 1051] class the GABA_A receptors according to their subunit structure, pharmacology and receptor function. Currently, eleven native GABA_A receptors are classed as conclusively identified (*i.e.*, $\alpha 1\beta 2\gamma 2$, $\alpha 2\beta\gamma 2$, $\alpha 3\beta\gamma 2$, $\alpha 4\beta\gamma 2$, $\alpha 4\beta 2\delta$, $\alpha 4\beta 3\delta$, $\alpha 5\beta\gamma 2$, $\alpha 6\beta\gamma 2$, $\alpha 6\beta 2\delta$, $\alpha 6\beta 3\delta$ and ρ) with further receptor isoforms occurring with high probability, or only tentatively [1051, 1052]. It is beyond the scope of this Guide to discuss the pharmacology of individual GABA_A receptor isoforms in detail; such information can be gleaned in the reviews [54, 55, 79, 417, 647, 733, 750, 967, 1051, 1052, 1274, 1276]. Agents that discriminate between α -subunit isoforms are noted in the table and additional agents that demonstrate selectivity between receptor isoforms, for example *via* β -subunit selectivity, are indicated in the text below. The distinctive agonist and antagonist pharmacology of ρ receptors is summarised in the table and additional aspects are reviewed in [205, 648, 1008, 1602].

Several high-resolution cryo-electron microscopy structures have been described in which the full-length human $\alpha 1\beta 3\gamma 2L$ GABA_A receptor in lipid nanodiscs is bound to the channel-blocker picrotoxin, the competitive antagonist bicuculline, the agonist GABA (γ -aminobutyric acid), and the classical benzodiazepines **alprazolam** and **diazepam** [918].

Nomenclature	GABA _A receptor α 1 subunit	GABA _A receptor α 2 subunit
HGNC, UniProt	<i>GABRA1</i> , P14867	<i>GABRA2</i> , P47869
Agonists	gaboxadol [GABA site], isoguvacine [GABA site], isonipecotin acid [GABA site], muscimol [GABA site], piperidine-4-sulphonic acid [GABA site]	gaboxadol [GABA site], isoguvacine [GABA site], isonipecotin acid [GABA site], muscimol [GABA site], piperidine-4-sulphonic acid [GABA site]
Antagonists	bicuculline [GABA site], gabazine [GABA site]	bicuculline [GABA site], gabazine [GABA site]
Channel blockers	TBPS , picrotoxin	TBPS , picrotoxin
Allosteric modulators	flumazenil [benzodiazepine site] (Antagonist) (p <i>K</i> _i 9.1) [1276], α3IA [benzodiazepine site] (Inverse agonist) [1276], α5IA [benzodiazepine site] (Inverse agonist) [1276], DMCM [benzodiazepine site] (Inverse agonist) [1160]	flumazenil [benzodiazepine site] (Antagonist at α 1 receptors, but allosteric modulator at other subtypes.) (p <i>K</i> _i 9.1) [1276], α3IA [benzodiazepine site] (Inverse agonist) [1276], α5IA [benzodiazepine site] (Inverse agonist) [1276], DMCM [benzodiazepine site] (Inverse agonist) [1160]
Endogenous allosteric modulators	5α-pregnan-3α-ol-20-one (Potentiation), Zn²⁺ (Inhibition), tetrahydrodeoxycorticosterone (Potentiation)	5α-pregnan-3α-ol-20-one (Potentiation), Zn²⁺ (Inhibition), tetrahydrodeoxycorticosterone (Potentiation)
Allosteric modulators (Positive)	clonazepam [benzodiazepine site] (p <i>K</i> _i 8.9) [1139], flunitrazepam [benzodiazepine site] (p <i>K</i> _i 8.3) [497], diazepam [benzodiazepine site] (p <i>K</i> _i 7.8) [1139], alprazolam [benzodiazepine site] (pEC ₅₀ 7.4) [27]	clonazepam [benzodiazepine site] (p <i>K</i> _i 8.8) [1139], flunitrazepam [benzodiazepine site] (p <i>K</i> _i 8.3) [497], alprazolam [benzodiazepine site] (pEC ₅₀ 7.9) [27], diazepam [benzodiazepine site] (p <i>K</i> _i 7.8) [1139]
Selective allosteric modulators	zolpidem (Positive) (p <i>K</i> _i 7.4–7.7) [498, 1240, 1276], L838417 [benzodiazepine site] (Antagonist) [1276]	L838417 [benzodiazepine site] (Partial agonist) [1276], TPA023 [benzodiazepine site] (Partial agonist) [1276]
Labelled ligands	[¹¹C]flumazenil [benzodiazepine site] (Allosteric modulator, Antagonist), [¹⁸F]fluoroethylflumazenil [benzodiazepine site] (Allosteric modulator), [³⁵S]TBPS [anion channel] (Channel blocker), [³H]CGS8216 [benzodiazepine site] (Allosteric modulator, Mixed), [³H]flunitrazepam [benzodiazepine site] (Allosteric modulator), [³H]gabazine [GABA site] (Antagonist), [³H]muscimol [GABA site] (Agonist), [³H]zolpidem [benzodiazepine site] (Allosteric modulator)	[¹¹C]flumazenil [benzodiazepine site] (Allosteric modulator), [¹⁸F]fluoroethylflumazenil [benzodiazepine site] (Allosteric modulator), [³⁵S]TBPS [anion channel] (Channel blocker), [³H]CGS8216 [benzodiazepine site] (Allosteric modulator, Mixed), [³H]flunitrazepam [benzodiazepine site] (Allosteric modulator, Full agonist), [³H]gabazine [GABA site] (Antagonist), [³H]muscimol [GABA site] (Agonist)
Comments	Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]	Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]

Nomenclature	GABA _A receptor α 3 subunit	GABA _A receptor α 4 subunit
HGNC, UniProt	GABRA3, P34903	GABRA4, P48169
Agonists	gaboxadol [GABA site], isoguvacine [GABA site], isonipecotic acid [GABA site], muscimol [GABA site], piperidine-4-sulphonic acid [GABA site]	gaboxadol [GABA site], isoguvacine [GABA site], muscimol [GABA site], piperidine-4-sulphonic acid [GABA site] (low efficacy)
Selective agonists	–	isonipecotic acid [GABA site] (relatively high efficacy, partially selective)
Antagonists	bicuculline [GABA site], gabazine [GABA site]	bicuculline [GABA site], gabazine [GABA site]
Channel blockers	TBPS, picrotoxin	TBPS, picrotoxin
Allosteric modulators	flumazenil [benzodiazepine site] (Antagonist at α 1 receptors, but allosteric modulator at other subtypes.) (pK _i 9) [1160, 1276], α 5IA [benzodiazepine site] (Inverse agonist) [1276], DMCM [benzodiazepine site] (Inverse agonist) [1160]	bretazenil [benzodiazepine site] (Full agonist) [1160]
Endogenous allosteric modulators	5 α -pregnan-3 α -ol-20-one (Potentiation), Zn ²⁺ (Inhibition), tetrahydrodeoxycorticosterone (Potentiation)	5 α -pregnan-3 α -ol-20-one (Potentiation), Zn ²⁺ (Inhibition), tetrahydrodeoxycorticosterone (Potentiation)
Allosteric modulators (Positive)	clonazepam [benzodiazepine site] (pK _i 8.7) [1139], flunitrazepam [benzodiazepine site] (pK _i 7.8) [497], diazepam [benzodiazepine site] (pK _i 7.8) [1139], alprazolam [benzodiazepine site] (pEC ₅₀ 7.2) [27]	–
Selective allosteric modulators	α 3IA [benzodiazepine site] (higher affinity, partially selective) [1276], L838417 [benzodiazepine site] (Partial agonist) [1276], Ro19-4603 [benzodiazepine site] (Inverse agonist), TP003 [benzodiazepine site] (Partial agonist) [1276], TPA023 [benzodiazepine site] (Partial agonist) [1276]	Ro15-4513 [benzodiazepine site] (Full agonist)
Labelled ligands	[¹¹ C]flumazenil [benzodiazepine site] (Allosteric modulator), [¹⁸ F]fluoroethyl-flumazenil [benzodiazepine site] (Allosteric modulator), [³⁵ S]TBPS [anion channel] (Channel blocker), [³ H]CGS8216 [benzodiazepine site] (Allosteric modulator, Mixed), [³ H]flunitrazepam [benzodiazepine site] (Allosteric modulator, Full agonist), [³ H]gabazine [GABA site] (Antagonist), [³ H]muscimol [GABA site] (Agonist)	[¹¹ C]flumazenil [benzodiazepine site] (Allosteric modulator, Partial agonist), [¹⁸ F]fluoroethylflumazenil [benzodiazepine site] (Allosteric modulator), [³⁵ S]TBPS [anion channel] (Channel blocker), [³ H]CGS8216 [benzodiazepine site] (Allosteric modulator, Mixed), [³ H]Ro154513 [benzodiazepine site] (Allosteric modulator, Full agonist), [³ H]gabazine [GABA site] (Antagonist), [³ H]muscimol [GABA site] (Agonist)
Comments	Zn ²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]	Diazepam and flunitrazepam are not active at this subunit. Zn ²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]. [³ H]Ro154513 labels α 4 β 2 and α 6 β 2 receptors in the presence of a saturating concentration of a 'classical' benzodiazepine (<i>e.g.</i> diazepam).

Nomenclature	GABA_A receptor α5 subunit	GABA_A receptor α6 subunit
HGNC, UniProt	GABRA5, P31644	GABRA6, Q16445
Agonists	gaboxadol [GABA site], isoguvacine [GABA site], isonipetric acid [GABA site], muscimol [GABA site], piperidine-4-sulphonic acid [GABA site]	gaboxadol [GABA site], isoguvacine [GABA site], muscimol [GABA site], piperidine-4-sulphonic acid [GABA site] (low efficacy)
Selective agonists	–	isonipetric acid [GABA site] (relatively high efficacy, relatively selective)
Antagonists	bicuculline [GABA site], gabazine [GABA site]	bicuculline [GABA site], gabazine [GABA site]
Channel blockers	TBPS , picrotoxin	TBPS , picrotoxin
Allosteric modulators	flumazenil [benzodiazepine site] (Antagonist at α1 receptors, but allosteric modulator at other subtypes.) (pK _i 9.2) [1276], α3IA [benzodiazepine site] (Inverse agonist) [1276], DMCM [benzodiazepine site] (Inverse agonist) [1160]	flumazenil [benzodiazepine site] (Partial agonist) (pK _i 6.8) [1276], bretazenil [benzodiazepine site] (Full agonist) [1160]
Endogenous allosteric modulators	5α-pregnan-3α-ol-20-one (Potentiation), Zn²⁺ (Inhibition), tetrahydrodeoxycorticosterone (Potentiation)	5α-pregnan-3α-ol-20-one (Potentiation), Zn²⁺ (Inhibition), tetrahydrodeoxycorticosterone (Potentiation)
Allosteric modulators (Positive)	flunitrazepam [benzodiazepine site] (pK _i 8.3) [497], alprazolam [benzodiazepine site] (pEC ₅₀ 8) [27]	–
Selective allosteric modulators	α5IA [benzodiazepine site] (Inverse agonist) [1276], L655708 [benzodiazepine site] (Inverse agonist) [1160], L838417 [benzodiazepine site] (Partial agonist) [1276], MRK016 [benzodiazepine site] (Inverse agonist) [1276], RO4938581 [benzodiazepine site] (Inverse agonist) [1276], RY024 [benzodiazepine site] (Inverse agonist) [1276]	LAU159 (Full agonist) [1275, 1276], LAU463 (Full agonist) [1275, 1276], PZ-II-029 (Full agonist) [1275, 1276], Ro15-4513 [benzodiazepine site] (Full agonist), amiloride (Antagonist) [1276], furosemide (Antagonist) [1276]
Labelled ligands	[³H]RY80 [benzodiazepine site] (Selective Binding) (pK _d 9.2) [1284] – Rat, [¹¹C]flumazenil [benzodiazepine site] (Allosteric modulator), [¹⁸F]fluoroethylflumazenil [benzodiazepine site] (Allosteric modulator), [³⁵S]TBPS [anion channel] (Channel blocker), [³H]CGS8216 [benzodiazepine site] (Allosteric modulator, Mixed), [³H]L655708 [benzodiazepine site] (Allosteric modulator, Inverse agonist), [³H]flunitrazepam [benzodiazepine site] (Allosteric modulator, Full agonist), [³H]gabazine [GABA site] (Antagonist), [³H]muscimol [GABA site] (Agonist)	[¹¹C]flumazenil [benzodiazepine site] (Allosteric modulator, Partial agonist), [¹⁸F]fluoroethylflumazenil [benzodiazepine site] (Allosteric modulator), [³⁵S]TBPS [anion channel] (Channel blocker), [³H]CGS8216 [benzodiazepine site] (Allosteric modulator, Mixed), [³H]Ro154513 [benzodiazepine site] (Allosteric modulator, Full agonist), [³H]gabazine [GABA site] (Antagonist), [³H]muscimol [GABA site] (Agonist)
Comments	Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]	Diazepam and flunitrazepam are not active at channels containing this subunit. Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]. [³H]Ro154513 selectively labels α6-subunit and α4-subunit-containing receptors in the presence of a saturating concentration of a 'classical' benzodiazepine (<i>e.g.</i> diazepam). Sieghart <i>et al.</i> (2022) provides a review of the pharmacology of α6-containing GABA _A receptors.

Nomenclature	GABA_A receptor β1 subunit	GABA_A receptor β2 subunit	GABA_A receptor β3 subunit
HGNC, UniProt	GABRB1, P18505	GABRB2, P47870	GABRB3, P28472
Channel blockers	TBPS, picrotoxin	TBPS, picrotoxin	TBPS, picrotoxin
Allosteric modulators	–	–	etazolate (Binding) (pIC₅₀ 5.5) [1600]
Comments	Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]		

Nomenclature	GABA_A receptor γ1 subunit	GABA_A receptor γ2 subunit	GABA_A receptor γ3 subunit
HGNC, UniProt	GABRG1, Q8N1C3	GABRG2, P18507	GABRG3, Q99928
Channel blockers	TBPS, picrotoxin	TBPS, picrotoxin	TBPS, picrotoxin
Comments	Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively [748]. So far, the only publication investigating the pharmacology of more than three or four ligands at γ1 receptors was Khom <i>et al.</i> (2006) [700].		

Nomenclature	GABA_A receptor δ subunit	GABA_A receptor ε subunit	GABA_A receptor θ subunit
HGNC, UniProt	GABRD, O14764	GABRE, P78334	GABRQ, Q9UN88
Selective agonists	DS2 [1276] , gaboxadol [GABA site] , tracazolate [1276]	–	–
Channel blockers	TBPS, picrotoxin	TBPS, picrotoxin	TBPS, picrotoxin
Comments	Zn²⁺ is an endogenous allosteric regulator and causes potent inhibition of receptors formed from binary combinations of α and β subunits, incorporation of a δ or γ subunit causes a modest, or pronounced, reduction in inhibitory potency, respectively		

Further reading on GABA_A receptors

- Atack JR. (2008) GABA(A) receptor subtype-selective efficacy: TPA023, an alpha2/alpha3 selective non-sedating anxiolytic and alpha5IA, an alpha5 selective cognition enhancer. *CNS Neurosci Ther* **14**: 25-35 [PMID:18482097]
- Braat S *et al.* (2015) The GABAA Receptor as a Therapeutic Target for Neurodevelopmental Disorders. *Neuron* **86**: 1119-30 [PMID:26050032]
- Calvo DJ *et al.* (2016) Dynamic Regulation of the GABAA Receptor Function by Redox Mechanisms. *Mol Pharmacol* **90**: 326-33 [PMID:27439531]

- Masiulis S *et al.* (2019) GABA_A receptor signalling mechanisms revealed by structural pharmacology. *Nature* **565**: 454-459 [PMID:30602790]
- Mele M *et al.* (2016) Role of GABA_A R trafficking in the plasticity of inhibitory synapses. *J Neurochem* **139**: 997-1018 [PMID:27424566]
- Sieghart W *et al.* (2018) International Union of Basic and Clinical Pharmacology. CVI: GABA_A Receptor Subtype- and Function-selective Ligands: Key Issues in Translation to Humans. *Pharmacol Rev* **70**: 836-878 [PMID:30275042]

Glycine receptors

Ion channels → Ligand-gated ion channels → Glycine receptors

Overview: The inhibitory glycine receptor (**nomenclature as agreed by the NC-IUPHAR Subcommittee on Glycine Receptors**) is a member of the Cys-loop superfamily of transmitter-gated ion channels that includes the GABA_A, nicotinic acetylcholine and 5-HT₃ receptors and Zn²⁺-activated channels. The glycine receptor is expressed either as a homo-pentamer of α subunits, or a complex of 4α and 1β subunits [1630], that contains an intrinsic anion channel. Four differentially expressed isoforms of the α-subunit (α1-α4) and one variant of the β-subunit (β1, *GLRB*, P48167) have been identified by genomic and cDNA cloning. Further diversity originates from alternative splicing of the primary gene transcripts for α1 (α1^{INS} and α1^{del}), α2 (α2A and α2B), α3 (α3S and α3L) and β (βΔ7) subunits and by mRNA editing of the α2 and α3 subunit [364, 936, 1041]. Both α2 splicing and α3

mRNA editing can produce subunits (*i.e.*, α2B and α3P185L) with enhanced agonist sensitivity. Predominantly, the adult form of the receptor contains α1 (or α3) and β subunits whereas the immature form is mostly composed of only α2 subunits [905]. The α4 subunit is a pseudogene in humans [775]. High resolution molecular structures are available for α1 homomeric, α3 homomeric, and αβ heteromeric receptors in a variety of ligand-induced conformations [343, 343, 584, 585, 586, 1584]. As in other Cys-loop receptors, the orthosteric binding site for agonists and the competitive antagonist **strychnine** is formed at the interfaces between the subunits' extracellular domains. Inclusion of the β-subunit in the pentameric glycine receptor contributes to agonist binding, reduces single channel conductance and alters pharmacology. The β-subunit also anchors the receptor, *via* an amphipathic se-

quence within the large intracellular loop region, to gephyrin. This a cytoskeletal attachment protein that binds to a number of subsynaptic proteins involved in cytoskeletal structure and thus clusters and anchors hetero-oligomeric receptors to the synapse [708, 975]. G protein βγ subunits enhance the open state probability of native and recombinant glycine receptors by association with domains within the large intracellular loop [1571, 1572]. Intracellular chloride concentration modulates the kinetics of native and recombinant glycine receptors [1124]. Intracellular Ca²⁺ appears to increase native and recombinant glycine receptor affinity, prolonging channel open events, by a mechanism that does not involve phosphorylation [418]. Extracellular Zn²⁺ potentiates GlyR function at nanomolar concentrations [954], and causes inhibition at higher micromolar concentrations (17).

Nomenclature	glycine receptor α1 subunit	glycine receptor α2 subunit
HGNC, UniProt	<i>GLRA1</i> , P23415	<i>GLRA2</i> , P23416
Selective agonists (potency order)	glycine > β-alanine > taurine	glycine > β-alanine > taurine
Selective antagonists	ginkgolide X (pIC ₅₀ 6.1), pregnenolone sulphate (pK _i 5.7), nifedipine (pIC ₅₀ 5.5), bilobalide (pIC ₅₀ 4.7), tropisetron (pK _i 4.1), colchicine (pIC ₅₀ 3.5), PMBA, onternabez (weak inhibition), strychnine	HU-210 (pIC ₅₀ 7), WIN55212-2 (pIC ₅₀ 6.7), onternabez (pIC ₅₀ 6), ginkgolide X (pIC ₅₀ 5.6), pregnenolone sulphate (pK _i 5.3), bilobalide (pIC ₅₀ 5.1), tropisetron (pK _i 4.9), colchicine (pIC ₅₀ 4.2), 5,7-dichlorokynurenic acid (pIC ₅₀ 3.7), PMBA, strychnine
Channel blockers	ginkgolide B (pIC ₅₀ 5.1-6.2), cyanotriphenylborate (pIC ₅₀ 5.9) [1201], picrotin (pIC ₅₀ 5.3), picrotoxinin (pIC ₅₀ 5.3), picrotoxin (pIC ₅₀ 5.2)	picrotoxinin (pIC ₅₀ 6.4), picrotoxin (pIC ₅₀ 5.6), ginkgolide B (pIC ₅₀ 4.9-5.4), picrotin (pIC ₅₀ 4.9), cyanotriphenylborate (pIC ₅₀ >4.7) [1201]
Endogenous allosteric modulators	Zn ²⁺ (Potentiation) (pEC ₅₀ 7.4), Cu ²⁺ (Inhibition) (pIC ₅₀ 4.8-5.4), Zn ²⁺ (Inhibition) (pIC ₅₀ 4.8), Extracellular H ⁺ (Inhibition)	Zn ²⁺ (Potentiation) (pEC ₅₀ 6.3), Cu ²⁺ (Inhibition) (pIC ₅₀ 4.8), Zn ²⁺ (Inhibition) (pIC ₅₀ 3.4)
Selective allosteric modulators	anandamide (Potentiation) (pEC ₅₀ 7.4), HU-210 (Potentiation) (pEC ₅₀ 6.6), Δ ⁹ -tetrahydrocannabinol (Potentiation) (pEC ₅₀ ~5.5)	Δ ⁹ -tetrahydrocannabinol (Potentiation) (pEC ₅₀ ~6)
Labelled ligands	[³ H]strychnine (Antagonist)	[³ H]strychnine (Antagonist)
Functional Characteristics	γ = 86 pS (main state); (+ β = 44 pS)	γ = 111 pS (main state); (+ β = 54 pS)

Nomenclature	glycine receptor $\alpha 3$ subunit	glycine receptor $\alpha 4$ subunit (pseudogene in humans)	glycine receptor β subunit
HGNC, UniProt	<i>GLRA3</i> , O75311	<i>GLRA4</i> , Q5JXX5	<i>GLRB</i> , P48167
Selective agonists (potency order)	glycine > β -alanine > taurine	–	–
Selective antagonists	HU-210 (pIC ₅₀ 7.3), WIN55212-2 (pIC ₅₀ 7), ontenabaz (pIC ₅₀ 7), (12E,20Z,18S)-8-hydroxyvariabilin (pIC ₅₀ 5.2), nifedipine (pIC ₅₀ 4.5), strychnine	–	nifedipine (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ 5.9), pregnenolone sulphate (when co-expressed with the $\alpha 1$ subunit) (pK _i 5.6), tropisetron (when co-expressed with the $\alpha 2$ subunit) (pK _i 5.3), pregnenolone sulphate (when co-expressed with the $\alpha 2$ subunit) (pK _i 5), nifedipine (when co-expressed with the $\alpha 3$ subunit) (pIC ₅₀ 4.9), bilobalide (when co-expressed with the $\alpha 2$ subunit) (pIC ₅₀ 4.3), bilobalide (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ 3.7), ginkgolide X (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ >3.5), ginkgolide X (when co-expressed with the $\alpha 2$ subunit) (pIC ₅₀ >3.5)
Channel blockers	picROTOXIN (pIC ₅₀ 6.4), ginkgolide B (pIC ₅₀ 5.7), picROtin (pIC ₅₀ 5.2), picROtoxin (block is weaker when β subunit is co-expressed)	–	ginkgolide B (when co-expressed with the $\alpha 2$ subunit) (pIC ₅₀ 6.1–6.9), ginkgolide B (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ 5.6–6.7), ginkgolide B (when co-expressed with the $\alpha 3$ subunit) (pIC ₅₀ 6.3), cyanotriphenylborate (when co-expressed with the human $\alpha 1$ subunit) (pIC ₅₀ 5.6) [1201] – Rat, cyanotriphenylborate (when co-expressed with the human $\alpha 2$ subunit) (pIC ₅₀ 5.1) [1201] – Rat, picROTOXIN (when co-expressed with the $\alpha 3$ subunit) (pIC ₅₀ 5.1), picROtin (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ 4.6), picROtin (when co-expressed with the $\alpha 3$ subunit) (pIC ₅₀ 4.6), picROTOXIN (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ 4.6), picROTOXIN (when co-expressed with the $\alpha 2$ subunit) (pIC ₅₀ 4.5), picROTOXIN (when co-expressed with the $\alpha 1$ subunit) (pIC ₅₀ 3.7)
Endogenous allosteric modulators	Cu²⁺ (Inhibition) (pIC ₅₀ 5), Zn²⁺ (Inhibition) (pIC ₅₀ 3.8)	–	Zn²⁺ (Inhibition) (pIC ₅₀ 4.9), Zn²⁺ (Inhibition) (pIC ₅₀ 3.7)
Selective allosteric modulators	Δ^9 - tetrahydrocannabinol (Potentiation) (pEC ₅₀ ~5.3)	–	–
Labelled ligands	[³ H] strychnine (Antagonist)	–	–
Functional Characteristics	$\gamma = 105$ pS (main state); (+ $\beta = 48$ pS)	–	–
Comments	–	–	Ligand interaction data for hetero-oligomer receptors containing the β subunit are also listed under the α subunit

Comments: Data in the table refer to homo-oligomeric assemblies of the α -subunit, significant changes introduced by co-expression of the $\beta 1$ subunit are indicated in parenthesis. Not all glycine receptor ligands are listed within the table, but some that may be useful in distinguishing between glycine receptor isoforms are indicated (see detailed view pages for each subunit: $\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 4$, β). **Pregnenolone sulphate**, **tropisetron** and **colchicine**, for example, although not selective antagonists of glycine receptors, are included for this purpose. **Strychnine** is a potent and selective competitive glycine

receptor antagonist with affinities in the range 5–15 nM. **RU5135** demonstrates comparable potency, but additionally blocks GABA_A receptors. There are conflicting reports concerning the ability of cannabinoids to inhibit [860], or potentiate and at high concentrations activate [15, 310, 527, 1537, 1563] glycine receptors. Nonetheless, cannabinoid analogues may hold promise in distinguishing between glycine receptor subtypes [1563]. In addition, potentiation of glycine receptor activity by cannabinoids has been claimed to contribute to cannabis-induced analgesia relying on Ser296/307

($\alpha 1/\alpha 3$) in M3 [1537]. Several analogues of **muscimol** and **piperidine** act as agonists and antagonists of both glycine and GABA_A receptors. **Picrotoxin** acts as an allosteric inhibitor that appears to bind within the pore, and shows strong selectivity towards homomeric receptors. While its components, **picROTOXIN** and **picROtin**, have equal potencies at $\alpha 1$ receptors, their potencies at $\alpha 2$ and $\alpha 3$ receptors differ modestly and may allow some distinction between different receptor types [1564]. Binding of picrotoxin within the pore has been demonstrated in the crystal structure of the related *C. elegans*

GluCl Cys-loop receptor [538]. In addition to the compounds listed in the table, numerous agents act as allosteric regulators of glycine receptors (comprehensively reviewed in [773, 874, 1501, 1573]). Zn²⁺ acts through distinct binding sites of high- and low-affinity to allosterically enhance channel function at low (<10 μM) concentrations and inhibits responses at higher concentrations in a subunit selective manner [953]. The effect of Zn²⁺ is somewhat mimicked by Ni²⁺. Endogenous Zn²⁺ is essential for normal glycinergic neurotransmission mediated by α1 subunit-containing receptors [549]. Elevation of intracellular Ca²⁺ produces fast potentiation of glycine receptor-mediated responses. Dideoxyforskolin (4 μM) and tamoxifen (0.2–5 μM) both potentiate responses to low glycine concentra-

tions (15 μM), but act as inhibitors at higher glycine concentrations (100 μM). Additional modulatory agents that enhance glycine receptor function include inhalational, and several intravenous general anaesthetics (*e.g.* minaxolone, propofol and pentobarbitone) and certain neurosteroids. Ethanol and higher order n-alcohols also enhance glycine receptor function although whether this occurs by a direct allosteric action at the receptor [917], or through βγ subunits [1570] is debated. Recent crystal structures of the bacterial homologue, GLIC, have identified transmembrane binding pockets for both anaesthetics [1036] and alcohols [575]. Solvents inhaled as drugs of abuse (*e.g.* toluene, 1-1-1-trichloroethane) may act at sites that overlap with those recognising alcohols and volatile anaesthet-

ics to produce potentiation of glycine receptor function. The function of glycine receptors formed as homomeric complexes of α1 or α2 subunits, or hetero-oligomers of α1/β or α2/β subunits, is differentially affected by the 5-HT₃ receptor antagonist tropisetron (ICS 205-930) which may evoke potentiation (which may occur within the femtomolar range at the homomeric glycine α1 receptor), or inhibition, depending upon the subunit composition of the receptor and the concentrations of the modulator and glycine employed. Potentiation and inhibition by tropeines involves different binding modes [899]. Additional tropeines, including atropine, modulate glycine receptor activity.

Further reading on Glycine receptors

Burgos CF *et al.* (2016) Structure and Pharmacologic Modulation of Inhibitory Glycine Receptors. *Mol Pharmacol* **90**: 318-25 [PMID:27401877]
 Dutre S *et al.* (2012) Inhibitory glycine receptors: an update. *J Biol Chem* **287**: 40216-23 [PMID:23038260]
 Lynch JW. (2004) Molecular structure and function of the glycine receptor chloride channel. *Physiol Rev* **84**: 1051-95 [PMID:15383648]

Perkins DI *et al.* (2010) Molecular targets and mechanisms for ethanol action in glycine receptors. *Pharmacol Ther* **127**: 53-65 [PMID:20399807]
 Yevenes GE *et al.* (2011) Allosteric modulation of glycine receptors. *Br J Pharmacol* **164**: 224-36 [PMID:21557733]

Ionotropic glutamate receptors

[Ion channels](#) → [Ligand-gated ion channels](#) → [Ionotropic glutamate receptors](#)

Overview: The ionotropic glutamate receptors comprise members of the NMDA (N-methyl-D-aspartate), AMPA (α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) and kainate receptor classes, named originally according to their preferred, synthetic, agonist [508, 853, 1404]. Receptor heterogeneity within each class arises from the homo-oligomeric, or hetero-oligomeric, assembly of distinct subunits into cation-selective tetramers. Each subunit of the tetrameric complex comprises an extracellular amino terminal domain (ATD), an extracellular ligand binding domain (LBD), 3 TM domains (M1, M3 and M4), a

channel lining re-entrant 'p-loop' (M2) located between M1 and M3 and an intracellular carboxy-terminal domain (CTD) [659, 757, 925, 993, 1404]. The X-ray structure of a homomeric ionotropic glutamate receptor (GluA2- see below) has been solved at 3.6 Å resolution [1298] and although providing detailed structural information, may not be representative of the subunit arrangement of, for example, the heteromeric NMDA receptors [674]. It is beyond the scope of this supplement to discuss the pharmacology of individual ionotropic glutamate receptor isoforms in detail; such information can be gleaned from [210, 266, 371,

508, 620, 621, 696, 1071, 1072, 1073, 1404, 1533]. Agents that discriminate between subunit isoforms are, where appropriate, noted in the tables and additional compounds that distinguish between receptor isoforms are indicated in the text below.

The classification of glutamate receptor subunits has been re-addressed by NC-IUPHAR [253]. The scheme developed recommends a nomenclature for ionotropic glutamate receptor subunits that is adopted here.

NMDA receptors

NMDA receptors assemble as obligate heteromers that may be drawn from GluN1, GluN2A, GluN2B, GluN2C, GluN2D, GluN3A and GluN3B subunits. Alternative splicing can generate eight isoforms of GluN1 with differing pharmacological properties. Various splice variants of GluN2B, 2C, 2D and GluN3A have also been reported. Activation of NMDA receptors containing GluN1 and GluN2 subunits requires the binding of two agonists, glutamate

to the S1 and S2 regions of the GluN2 subunit and glycine to S1 and S2 regions of the GluN1 subunit [211, 370]. The minimal requirement for efficient functional expression of NMDA receptors *in vitro* is a di-heteromeric assembly of GluN1 and at least one GluN2 subunit variant, as a dimer of heterodimers arrangement in the extracellular domain [420, 674, 925]. However, more complex tri-heteromeric assemblies, incorporating multiple subtypes

of GluN2 subunit, or GluN3 subunits, can be generated *in vitro* and occur *in vivo*. The NMDA receptor channel commonly has a high relative permeability to Ca²⁺ and is blocked, in a voltage-dependent manner, by Mg²⁺ such that at resting potentials the response is substantially inhibited.

AMPA and Kainate receptors

AMPA receptors assemble as homomers, or heteromers, that may be drawn from GluA1, GluA2, GluA3 and GluA4 subunits. Transmembrane AMPA receptor regulatory proteins (TARPs) of class I (i.e. $\gamma 2$, $\gamma 3$, $\gamma 4$ and $\gamma 8$) act, with variable stoichiometry, as auxiliary subunits to AMPA receptors and influence their trafficking, single channel conductance gating and pharmacology (reviewed in [376, 612, 956, 1395]). Functional kainate receptors can be expressed as homomers of GluK1, GluK2 or GluK3 subunits. GluK1-3 subunits are also capable of assembling into heterotetramers (e.g. GluK1/K2; [796, 1104, 1119]). Two additional kainate receptor subunits, GluK4 and GluK5, when expressed individually, form high affinity binding sites for **kainate**, but lack function, but can form heteromers when expressed with GluK1-3 subunits (e.g. GluK2/K5; reviewed in [620, 1104, 1119]). Kainate receptors may also exhibit

'metabotropic' functions [796, 1183]. As found for AMPA receptors, kainate receptors are modulated by auxiliary subunits (Neto proteins, [797, 1104]). An important function difference between AMPA and kainate receptors is that the latter require extracellular Na^+ and Cl^- for their activation [145, 1132]. RNA encoding the GluA2 subunit undergoes extensive RNA editing in which the codon encoding a p-loop glutamine residue (Q) is converted to one encoding arginine (R). This Q/R site strongly influences the biophysical properties of the receptor. Recombinant AMPA receptors lacking RNA edited GluA2 subunits are: (1) permeable to Ca^{2+} ; (2) blocked by intracellular polyamines at depolarized potentials causing inward rectification (the latter being reduced by TARPs); (3) blocked by extracellular **argiotoxin** and **joro spider toxins** and (4) demonstrate higher channel conductances than receptors con-

taining the edited form of GluA2 [603, 1248]. GluK1 and GluK2, but not other kainate receptor subunits, are similarly edited and broadly similar functional characteristics apply to kainate receptors lacking either an RNA edited GluK1, or GluK2, subunit [796, 1104]. Native AMPA and kainate receptors displaying differential channel conductances, Ca^{2+} permeabilities and sensitivity to block by intracellular polyamines have been identified [265, 603, 846]. GluA1-4 can exist as two variants generated by alternative splicing (termed 'flip' and 'flop') that differ in their desensitization kinetics and their desensitization in the presence of **cyclothiazide** which stabilises the nondesensitized state. TARPs also stabilise the nondesensitized conformation of AMPA receptors and facilitate the action of **cyclothiazide** [956]. Splice variants of GluK1-3 also exist which affects their trafficking [796, 1104].

Nomenclature	GluA1	GluA2	GluA3	GluA4
HGNC, UniProt	GRIA1 , P42261	GRIA2 , P42262	GRIA3 , P42263	GRIA4 , P48058
Agonists	(S)-5-fluorowillardiine, AMPA	(S)-5-fluorowillardiine, AMPA	(S)-5-fluorowillardiine, AMPA	(S)-5-fluorowillardiine, AMPA
Selective antagonists	ATPO, GYKI53655, GYKI53784 (active isomer, non-competitive), NBQX, tezampanel	ATPO, GYKI53655, GYKI53784 (active isomer, non-competitive), NBQX, tezampanel	ATPO, GYKI53655, GYKI53784 (active isomer, non-competitive), NBQX, tezampanel	ATPO, GYKI53655, GYKI53784 (active isomer, non-competitive), NBQX, tezampanel
Channel blockers	extracellular argiotoxin , extracellular joro spider toxin (selective for channels lacking GluA2)	extracellular argiotoxin	extracellular argiotoxin , extracellular joro spider toxin (selective for channels lacking GluA2)	extracellular argiotoxin , extracellular joro spider toxin (selective for channels lacking GluA2)
Allosteric modulators (Positive)	LY392098 (pEC ₅₀ 5.8) [962], LY404187 (pEC ₅₀ 5.2) [962], cyclothiazide (pEC ₅₀ 4.7) [962], CX516 , CX546 , IDRA-21 , LY503430 , S18986 , aniracetam , piracetam	LY404187 (pEC ₅₀ 6.8) [962], LY392098 (pEC ₅₀ 6.7) [962], cyclothiazide (pEC ₅₀ 5.7) [962], CX516 , CX546 , IDRA-21 , LY503430 , S18986 , aniracetam , piracetam	LY404187 (pEC ₅₀ 5.8) [962], LY392098 (pEC ₅₀ 5.7) [962], cyclothiazide (pEC ₅₀ 4.9) [962], CX516 , CX546 , IDRA-21 , LY503430 , S18986 , aniracetam , piracetam	LY392098 (pEC ₅₀ 6.7) [962], LY404187 (pEC ₅₀ 6.7) [962], cyclothiazide (pEC ₅₀ 5.4) [962], CX516 , CX546 , IDRA-21 , LY503430 , S18986 , aniracetam , piracetam
Labelled ligands	[³ H]AMPA (Agonist), [³ H]CNQX (Antagonist)	[³ H]AMPA (Agonist), [³ H]CNQX (Antagonist)	[³ H]AMPA, [³ H]CNQX	[³ H]AMPA (Agonist), [³ H]CNQX
Comments	Piracetam and aniracetam are examples of pyrrolidinones. Cyclothiazide , S18986 , and IDRA-21 are examples of benzothiadiazides. CX516 and CX546 are examples of benzylpiperidines. LY392098 , LY404187 and LY503430 are examples of biarylpropylsulfonamides. Also blocked by intracellular polyamines.	Piracetam and aniracetam are examples of pyrrolidinones. Cyclothiazide , S18986 , and IDRA-21 are examples of benzothiadiazides. CX516 and CX546 are examples of benzylpiperidines. LY392098 , LY404187 and LY503430 are examples of biarylpropylsulfonamides. Also blocked by intracellular polyamines.	Piracetam and aniracetam are examples of pyrrolidinones. Cyclothiazide , S18986 , and IDRA-21 are examples of benzothiadiazides. CX516 and CX546 are examples of benzylpiperidines. LY392098 , LY404187 and LY503430 are examples of biarylpropylsulfonamides. Also blocked by intracellular polyamines	Piracetam and aniracetam are examples of pyrrolidinones. Cyclothiazide , S18986 , and IDRA-21 are examples of benzothiadiazides. CX516 and CX546 are examples of benzylpiperidines. LY392098 , LY404187 and LY503430 are examples of biarylpropylsulfonamides. Also blocked by intracellular polyamines

Nomenclature	GluD1	GluD2
HGNC, UniProt	GRID1, Q9ULK0	GRID2, O43424

Nomenclature	GluK1	GluK2	GluK3	GluK4	GluK5
HGNC, UniProt	GRIK1, P39086	GRIK2, Q13002	GRIK3, Q13003	GRIK4, Q16099	GRIK5, Q16478
Agonists	dysiherbaine [1214] – Rat, SYM2081 [1095], kainate [1286], (S)-4-AHCP, (S)-5-iodowillardiine, 8-deoxy-neodysiherbaine, ATPA, domoic acid	dysiherbaine [1214] – Rat, domoic acid [176], SYM2081 [1625] – Rat, kainate [176, 1286]	SYM2081 [1211] – Rat, kainate (low potency) [1211] – Rat, dysiherbaine	SYM2081, domoic acid, dysiherbaine, kainate	SYM2081, domoic acid, dysiherbaine, kainate
Selective agonists	LY339434 [1286]	–	–	–	–
Selective antagonists	2,4-epi-neodysiherbaine, ACET, LY382884, LY466195, MSVIII-19, NS3763 (non-competitive), UBP302, UBP310	2,4-epi-neodysiherbaine	–	–	–
Allosteric modulators (Positive)	concanavalin A	concanavalin A	–	–	–
Labelled ligands	[³ H]UBP310 (Antagonist) (pK _d 7.7) [56], [³ H]SYM2081 (Agonist), [³ H]kainate (Agonist)	[³ H]kainate (Agonist) [1625] – Rat, [³ H]SYM2081 (Agonist)	[³ H]UBP310 (Antagonist) (pK _d 6.3) [56], [³ H]SYM2081 (Agonist), [³ H]kainate (Agonist)	[³ H]SYM2081 (Agonist), [³ H]kainate (Agonist)	[³ H]SYM2081 (Agonist), [³ H]kainate (Agonist)
Comments	–	Intracellular polyamines are subtype selective channel blockers (GluK3 ≫ GluK2). GluK2 mediates cold sensing in mice [169].	Domoic acid and concanavalin A are inactive at the GluK3 subunit. Intracellular polyamines are subtype selective channel blockers (GluK3 ≫ GluK2)	–	–

Nomenclature	GluN1
HGNC, UniProt	GRIN1, Q05586
Endogenous agonists	D-aspartic acid [glutamate site], D-serine [glycine site], L-aspartic acid [glutamate site], glycine [glycine site]
Agonists	(+)-HA966 [glycine site] (Partial agonist), (RS)-(tetrazol-5-yl)glycine [glutamate site], NMDA [glutamate site], homoquinolinic acid [glutamate site] (Partial agonist)
Selective antagonists	L701324 [glycine site] (pIC ₅₀ 8.7) [756] – Rat, GV196771A [glycine site] (pK _i 8.1–8.4) [231] – Rat, L689560 [glycine site] (pIC ₅₀ 8.1) [789] – Rat, 5,7-dichlorokynurenic acid [glycine site]
Labelled ligands	[³ H]MDL105519 [glycine site] (Antagonist) (pK _d ~8.5) [204] – Rat, [³ H]CGP39653 [glutamate site] (Selective Antagonist), [³ H]CGP61594 [glycine site] (Antagonist), [³ H]CGS19755 [glutamate site] (Antagonist), [³ H]CPP [glutamate site] (Selective Antagonist), [³ H]L689560 [glycine site] (Antagonist), [³ H]dizocilpine [cation channel] (Antagonist), [³ H]glycine [glycine site] (Agonist)

Nomenclature	GluN2A	GluN2B	GluN2C	GluN2D
HGNC, UniProt	<i>GRIN2A</i> , Q12879	<i>GRIN2B</i> , Q13224	<i>GRIN2C</i> , Q14957	<i>GRIN2D</i> , O15399
Endogenous agonists	D-aspartic acid [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), D-serine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A), L-aspartic acid [glutamate site] (GluN2D = GluN2B > GluN2C = GluN2A), glycine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A)	D-aspartic acid [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), D-serine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A), L-aspartic acid [glutamate site] (GluN2D = GluN2B > GluN2C = GluN2A), glycine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A)	D-aspartic acid [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), D-serine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A), L-aspartic acid [glutamate site] (GluN2D = GluN2B > GluN2C = GluN2A), glycine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A)	D-aspartic acid [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), D-serine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A), L-aspartic acid [glutamate site] (GluN2D = GluN2B > GluN2C = GluN2A), glycine [glycine site] (GluN2D > GluN2C > GluN2B > GluN2A)
Agonists	(+)-HA966 [glycine site] (Partial agonist), (RS)-(tetrazol-5-yl)glycine [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), NMDA [glutamate site] (GluN2D > GluN2C > GluN2B > GluN2A), homoquinolinic acid [glutamate site] (GluN2B ≥ GluN2A ≥ GluN2D > GluN2C; partial agonist at GluN2A and GluN2C)	(+)-HA966 [glycine site] (Partial agonist), (RS)-(tetrazol-5-yl)glycine [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), NMDA [glutamate site] (GluN2D > GluN2C > GluN2B > GluN2A), homoquinolinic acid [glutamate site] (GluN2B ≥ GluN2A ≥ GluN2D > GluN2C; partial agonist at GluN2A and GluN2C)	(RS)-(tetrazol-5-yl)glycine [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), NMDA [glutamate site] (GluN2D > GluN2C > GluN2B > GluN2A), homoquinolinic acid [glutamate site] (GluN2B ≥ GluN2A ≥ GluN2D > GluN2C; partial agonist at GluN2A and GluN2C)	(RS)-(tetrazol-5-yl)glycine [glutamate site] (GluN2D > GluN2C = GluN2B > GluN2A), NMDA [glutamate site] (GluN2D > GluN2C > GluN2B > GluN2A), homoquinolinic acid [glutamate site] (GluN2B ≥ GluN2A ≥ GluN2D > GluN2C; partial agonist at GluN2A and GluN2C)
Selective antagonists	5,7-dichlorokynurenic acid [glycine site], CGP37849 [glutamate site], GV196771A [glycine site], L689560 [glycine site], L701324 [glycine site], LY233053 [glutamate site], NVP-AAM077 [glutamate site] (GluN2A > GluN2B (human), but weakly selective for rat GluN2A versus GluN2B) [58, 393, 416, 1007], UBP141 [glutamate site] (GluN2D ≥ GluN2C > GluN2A ≥ GluN2B) [974], conantokin-G [glutamate site] (GluN2B > GluN2D = GluN2C = GluN2A), d-AP5 [glutamate site], d-CCPene [glutamate site] (GluN2A = GluN2B > GluN2C = GluN2D), selfotel [glutamate site]	5,7-dichlorokynurenic acid [glycine site], CGP37849 [glutamate site], GV196771A [glycine site], L689560 [glycine site], L701324 [glycine site], LY233053 [glutamate site], NVP-AAM077 [glutamate site] (GluN2A > GluN2B (human), but weakly selective for rat GluN2A versus GluN2B) [58, 393, 416, 1007], UBP141 [glutamate site] (GluN2D ≥ GluN2C > GluN2A ≥ GluN2B) [974], conantokin-G [glutamate site] (GluN2B > GluN2D = GluN2C = GluN2A), d-AP5 [glutamate site], d-CCPene [glutamate site] (GluN2A = GluN2B > GluN2C = GluN2D), selfotel [glutamate site]	5,7-dichlorokynurenic acid [glycine site], CGP37849 [glutamate site], GV196771A [glycine site], L689560 [glycine site], L701324 [glycine site], LY233053 [glutamate site], UBP141 [glutamate site] (GluN2D ≥ GluN2C > GluN2A ≥ GluN2B) [974], conantokin-G [glutamate site] (GluN2B > GluN2D = GluN2C = GluN2A), d-AP5 [glutamate site], d-CCPene [glutamate site] (GluN2A = GluN2B > GluN2C = GluN2D), selfotel [glutamate site]	5,7-dichlorokynurenic acid [glycine site], CGP37849 [glutamate site], GV196771A [glycine site], L689560 [glycine site], L701324 [glycine site], LY233053 [glutamate site], UBP141 [glutamate site] (GluN2D ≥ GluN2C > GluN2A ≥ GluN2B) [974], conantokin-G [glutamate site] (GluN2B > GluN2D = GluN2C = GluN2A), d-AP5 [glutamate site], d-CCPene [glutamate site] (GluN2A = GluN2B > GluN2C = GluN2D), selfotel [glutamate site]
Channel blockers	Mg²⁺ (GluN2A = GluN2B > GluN2C = GluN2D), N¹-dansyl-spermine (GluN2A = GluN2B ≫ GluN2C = GluN2D), amantadine (GluN2C = GluN2D ≥ GluN2B ≥ GluN2A), dizocilpine , ketamine , phencyclidine	Mg²⁺ (GluN2A = GluN2B > GluN2C = GluN2D), N¹-dansyl-spermine (GluN2A = GluN2B ≫ GluN2C = GluN2D), amantadine (GluN2C = GluN2D ≥ GluN2B ≥ GluN2A), dizocilpine , ketamine , phencyclidine	phencyclidine (pIC ₅₀ 7.1) [341], ketamine (pIC ₅₀ 6.2) [341], amantadine (GluN2C = GluN2D ≥ GluN2B ≥ GluN2A) (pIC ₅₀ 4.7) [341], Mg²⁺ (GluN2A = GluN2B > GluN2C = GluN2D), N¹-dansyl-spermine (GluN2A = GluN2B ≫ GluN2C = GluN2D), dizocilpine	Mg²⁺ (GluN2A = GluN2B > GluN2C = GluN2D), N¹-dansyl-spermine (GluN2A = GluN2B ≫ GluN2C = GluN2D), amantadine (GluN2C = GluN2D ≥ GluN2B ≥ GluN2A), dizocilpine , ketamine , phencyclidine

Labelled ligands	[³H]CGP39653 [glutamate site] (Antagonist), [³H]CGP61594 [glycine site] (Antagonist), [³H]CGS19755 [glutamate site] (Antagonist), [³H]CPP [glutamate site] (Antagonist), [³H]L689560 [glycine site] (Antagonist), [³H]MDL105519 [glycine site] (Antagonist), [³H]dizocilpine [cation channel] (Channel blocker), [³H]glycine [glycine site] (Agonist)	[³H]CGP39653 [glutamate site] (Antagonist), [³H]CGP61594 [glycine site] (Antagonist), [³H]CGS19755 [glutamate site] (Antagonist), [³H]CPP [glutamate site] (Antagonist), [³H]L689560 [glycine site] (Antagonist), [³H]MDL105519 [glycine site] (Antagonist), [³H]dizocilpine [cation channel] (Channel blocker), [³H]glycine [glycine site] (Agonist)	[³H]CGP39653 [glutamate site] (Antagonist), [³H]CGP61594 [glycine site] (Antagonist), [³H]CGS19755 [glutamate site] (Antagonist), [³H]CPP [glutamate site] (Antagonist), [³H]L689560 [glycine site] (Antagonist), [³H]MDL105519 [glycine site] (Antagonist), [³H]dizocilpine [cation channel] (Channel blocker), [³H]glycine [glycine site] (Agonist)	[³H]CGP39653 [glutamate site] (Antagonist), [³H]CGP61594 [glycine site] (Antagonist), [³H]CGS19755 [glutamate site] (Antagonist), [³H]CPP [glutamate site] (Antagonist), [³H]L689560 [glycine site] (Antagonist), [³H]MDL105519 [glycine site] (Antagonist), [³H]dizocilpine [cation channel] (Channel blocker), [³H]glycine [glycine site] (Agonist)
Nomenclature	GluN3A	GluN3B		
HGNC, UniProt	GRIN3A , Q8TCU5	GRIN3B , O60391		
Comments	See the main comments section below for information on the pharmacology of GluN3A and GluN3B subunits			

Comments: NMDA receptors

Potency orders unreferenced in the table are from [210, 341, 371, 758, 1073, 1404]. In addition to the glutamate and glycine binding sites documented in the table, physiologically important inhibitory modulatory sites exist for Mg²⁺, Zn²⁺, and protons [266, 508, 1404]. Voltage-independent inhibition by Zn²⁺ binding with high affinity within the ATD is highly subunit selective (GluN2A ≫ GluN2B > GluN2C ≥ GluN2D; [1073, 1404]). The receptor is also allosterically modulated, in both positive and negative directions, by endogenous neuroactive steroids in a subunit dependent manner [567, 900]. Tonic proton blockade of NMDA receptor function is alleviated by polyamines and the inclusion of exon 5 within GluN1 subunit splice variants, whereas the non-competitive antagonists [ifenprodil](#) and [traxoprodil](#) increase the fraction of receptors blocked by protons at ambient concentration. Inclusion of exon 5 also abolishes potentiation by polyamines and inhibition by Zn²⁺ that occurs through binding in the ATD [1403]. [Ifenprodil](#), [traxoprodil](#), [haloperidol](#), [felbamate](#) and

[Ro 8-4304](#) discriminate between recombinant NMDA receptors assembled from GluN1 and either GluN2A, or GluN2B, subunits by acting as selective, non-competitive, antagonists of heterooligomers incorporating GluN2B through a binding site at the ATD GluN1/GluN2B subunit interface [674]. [LY233536](#) is a competitive antagonist that also displays selectivity for GluN2B over GluN2A subunit-containing receptors. Similarly, [CGP61594](#) is a photoaffinity label that interacts selectively with receptors incorporating GluN2B versus GluN2A, GluN2D and, to a lesser extent, GluN2C subunits. [TCN-201](#) and [TCN-213](#) have been shown to block GluN2A NMDA receptors selectively by a mechanism that involves allosteric inhibition of glycine binding to the GluN1 site [112, 360, 507, 931]. In addition to influencing the pharmacological profile of the NMDA receptor, the identity of the GluN2 subunit co-assembled with GluN1 is an important determinant of biophysical properties that include sensitivity to block by Mg²⁺, single-channel conductance and maximal open probability and

channel deactivation time [266, 370, 454]. Incorporation of the GluN3A subunit into tri-heteromers containing GluN1 and GluN2 subunits is associated with decreased single-channel conductance, reduced permeability to Ca²⁺ and decreased susceptibility to block by Mg²⁺ [189, 532]. Reduced permeability to Ca²⁺ has also been observed following the inclusion of GluN3B in tri-heteromers. The expression of GluN3A, or GluN3B, with GluN1 alone forms, in *Xenopus laevis* oocytes, a cation channel with unique properties that include activation by [glycine](#) (but not [NMDA](#)), lack of permeation by Ca²⁺ and resistance to blockade by Mg²⁺ and NMDA receptor antagonists [199]. The function of heteromers composed of GluN1 and GluN3A is enhanced by Zn²⁺, or glycine site antagonists, binding to the GluN1 subunit [887]. Zn²⁺ also directly activates such complexes. The co-expression of GluN1, GluN3A and GluN3B appears to be required to form glycine-activated receptors in mammalian cell hosts [1296].

AMPA and Kainate receptors

All AMPA receptors are additionally activated by [kainate](#) (and [do-moic acid](#)) with relatively low potency, (EC₅₀ ~ 100 μM). Inclusion of TARPs within the receptor complex increases the potency and maximal effect of kainate [612, 956]. AMPA is weak partial agonist at GluK1 and at heteromeric assemblies of GluK1/GluK2, GluK1/GluK5 and GluK2/GluK5 [620]. Quinoxalinediones such as [CNQX](#) and [NBQX](#) show limited selectivity between AMPA and kainate receptors. [Tezampanel](#) also has kainate (GluK1) recep-

tor activity as has [GYKI53655](#) (GluK3 and GluK2/GluK3) [620]. [ATPO](#) is a potent competitive antagonist of AMPA receptors, has a weaker antagonist action at kainate receptors comprising GluK1 subunits, but is devoid of activity at kainate receptors formed from GluK2 or GluK2/GluK5 subunits. The pharmacological activity of [ATPO](#) resides with the (S)-enantiomer. [ACET](#) and [UBP310](#) may block GluK3, in addition to GluK1 [56, 1103]. (2S,4R)-4-methylglutamate ([SYM2081](#)) is equipotent in activating

(and desensitising) GluK1 and GluK2 receptor isoforms and, *via* the induction of desensitisation at low concentrations, has been used as a functional antagonist of kainate receptors. Both (2S,4R)-4-methylglutamate and [LY339434](#) have agonist activity at NMDA receptors. (2S,4R)-4-methylglutamate is also an inhibitor of the glutamate transporters EAAT1 and EAAT2.

Delta subunits

GluD1 and GluD2 comprise, on the basis of sequence homology, an 'orphan' class of ionotropic glutamate receptor subunit. They do not form a functional receptor when expressed solely, or in combination with other ionotropic glutamate receptor subunits, in transfected cells [1593]. However, GluD2 subunits bind **D-serine** and **glycine** and GluD2 subunits carrying the mutation A654T form a spontaneously open channel that is closed by **D-serine** [998]. Their physiological role is uncertain. It may be structural, contributing as trans-synaptic organizers [508].

Further reading on Ionotropic glutamate receptors

- Filippini A *et al.* (2017) The Good and the Bad of Glutamate Receptor RNA Editing. *Mol Neurobiol* **54**: 6795-6805 [PMID:27766534]
- Gangwar SP *et al.* (2024) Kainate receptor channel opening and gating mechanism. *Nature* **630**: 762-768 [PMID:38778115]
- Greger IH *et al.* (2017) Structural and Functional Architecture of AMPA-Type Glutamate Receptors and Their Auxiliary Proteins. *Neuron* **94**: 713-730 [PMID:28521126]
- Hackos DH *et al.* (2017) Diverse modes of NMDA receptor positive allosteric modulation: Mechanisms and consequences. *Neuropharmacology* **112**: 34-45 [PMID:27484578]
- Huettner JE. (2015) Glutamate receptor pores. *J Physiol (Lond.)* **593**: 49-59 [PMID:25556787]
- Iacobucci GJ *et al.* (2017) NMDA receptors: linking physiological output to biophysical operation. *Nat Rev Neurosci* **18**: 236-249 [PMID:28303017]
- Krieger J *et al.* (2015) Structure, Dynamics, and Allosteric Potential of Ionotropic Glutamate Receptor N-Terminal Domains. *Biophys J* **109**: 1136-48 [PMID:26255587]
- Lussier MP *et al.* (2015) Dynamic Regulation of N-Methyl-d-aspartate (NMDA) and α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic Acid (AMPA) Receptors by Posttranslational Modifications. *J Biol Chem* **290**: 28596-603 [PMID:26453298]
- Møllerud S *et al.* (2017) Lessons from crystal structures of kainate receptors. *Neuropharmacology* **112**: 16-28 [PMID:27236079]
- Segura-Covarrubias G *et al.* (2025) Structural basis of GluK2 kainate receptor activation by a partial agonist. *Nat Struct Mol Biol* **32**: 1456-1469 [PMID:40442317]
- Yuzaki M *et al.* (2017) A GluD Coming-Of-Age Story. *Trends Neurosci* **40**: 138-150 [PMID:28110935]
- Zhou HX *et al.* (2017) Advancing NMDA Receptor Physiology by Integrating Multiple Approaches. *Trends Neurosci* **40**: 129-137 [PMID:28187950]
- Zhuo M. (2017) Ionotropic glutamate receptors contribute to pain transmission and chronic pain. *Neuropharmacology* **112**: 228-234 [PMID:27543416]

IP₃ receptors

Ion channels → **Ligand-gated ion channels** → **IP₃ receptors**

Overview: The inositol 1,4,5-trisphosphate receptors (IP₃R) are ligand-gated Ca²⁺-release channels on intracellular Ca²⁺ store sites (such as the endoplasmic reticulum). They are responsible for the mobilization of intracellular Ca²⁺ stores and play an important role in intracellular Ca²⁺ signalling in a wide variety of cell types. Three different gene products (types I-III) have been isolated, which assemble as large tetrameric structures. IP₃Rs are closely associated with certain proteins: **calmodulin** (*CALM1CALM2CALM3*, P62158) and **FKBP** (and calcineurin *via* FKBP). They are phosphorylated by PKA, PKC, PKG and CaMKII.

Nomenclature	IP ₃ R1	IP ₃ R2	IP ₃ R3
HGNC, UniProt	<i>ITPR1</i> , Q14643	<i>ITPR2</i> , Q14571	<i>ITPR3</i> , Q14573
Endogenous activators	cytosolic ATP (< mM range), cytosolic Ca²⁺ Concentration range: <7.5 × 10 ⁻⁴ M, IP₃ (endogenous; nM - μ M range)	cytosolic Ca²⁺ (nM range), IP₃ (endogenous; nM - μ M range)	cytosolic Ca²⁺ (nM range), IP₃ (endogenous; nM - μ M range)
Activators	adenophostin A (pharmacological; nM range), inositol 2,4,5-trisphosphate (pharmacological; also activated by other InsP ₃ analogues)	adenophostin A (pharmacological; nM range), inositol 2,4,5-trisphosphate (pharmacological; also activated by other InsP ₃ analogues)	–
Antagonists	PIP₂ (μ M range), caffeine (mM range), decavanadate (μ M range), xestospongin C (μ M range)	decavanadate (μ M range)	decavanadate (μ M range)
Functional Characteristics	Ca ²⁺ : (P _{Ba} /P _K ~6) single-channel conductance ~70 pS (50 mM Ca ²⁺)	Ca ²⁺ : single-channel conductance ~70 pS (50 mM Ca ²⁺) ~390 pS (220 mM Cs ⁺)	Ca ²⁺ : single-channel conductance ~88 pS (55 mM Ba ²⁺)
Comments	IP ₃ R1 is also antagonised by calmodulin at high cytosolic Ca ²⁺ concentrations	–	–

Comments: The absence of a modulator of a particular isoform of receptor indicates that the action of that modulator has not been determined, not that it is without effect.

Further reading on IP₃ receptors

- Berridge MJ. (2016) The Inositol Trisphosphate/Calcium Signaling Pathway in Health and Disease. *Physiol Rev* **96**: 1261-96 [PMID:27512009]
- Gambardella J *et al.* (2020) Inositol 1,4,5-Trisphosphate Receptors in Human Disease: A Comprehensive Update. *J Clin Med* **9**: 1096 [PMID:32290556]
- Garcia MI *et al.* (2017) Cardiac inositol 1,4,5-trisphosphate receptors. *Biochim Biophys Acta* **1864**: 907-914 [PMID:27884701]
- Mak DO *et al.* (2015) Inositol 1,4,5-trisphosphate receptors in the endoplasmic reticulum: A single-channel point of view. *Cell Calcium* **58**: 67-78 [PMID:25555684]
- Prole DL *et al.* (2019) Structure and Function of IP₃ Receptors. *Cold Spring Harb Perspect Biol* **11**: [PMID:30745293]
- Rossi AM *et al.* (2018) IP₃ receptors - lessons from analyses *ex cellula*. *J Cell Sci* **132**: [PMID:30552138]
- Seo MD *et al.* (2015) Structural insights into endoplasmic reticulum stored calcium regulation by inositol 1,4,5-trisphosphate and ryanodine receptors. *Biochim Biophys Acta* **1853**: 1980-91 [PMID:25461839]
- Thillaiappan NB *et al.* (2019) IP₃ receptors and Ca²⁺ entry. *Biochim Biophys Acta Mol Cell Res* **1866**: 1092-1100 [PMID:30448464]

Nicotinic acetylcholine receptors (nACh)

Ion channels → Ligand-gated ion channels → Nicotinic acetylcholine receptors (nACh)

Overview: Nicotinic acetylcholine (ACh) receptors are members of the Cys-loop family of transmitter-gated ion channels that includes the GABA_A, strychnine-sensitive glycine and 5-HT₃ receptors [28, 951, 1281, 1360, 1528]. All nicotinic receptors are pentamers in which each of the five subunits contains 4 TM domains. Genes encoding a total of 17 subunits (α 1-10, β 1-4, γ , δ and ϵ) have been identified [662]. All subunits with the exception of α 8 (present in avian species) have been identified in mammals. All α subunits possess two tandem cysteine residues near to the site involved in acetylcholine binding, and subunits not named α lack these residues [951]. The orthosteric ligand binding site is formed by residues within at least three peptide domains on the α subunit (principal component), and three on the adjacent subunit (complementary component). Nicotinic ACh receptors contain several allosteric modulatory sites. One such site, for positive allosteric modulators (PAMs) and allosteric agonists, has been proposed to reside within an intrasubunit cavity between the 4 TM domains [456, 1579]; see also [538]. The high resolution crystal structure of the molluscan ACh binding protein, a structural homologue of the extracellular binding domain of a nicotinic receptor pentamer, in complex with several nicotinic receptor ligands (*e.g.* [191]) and the crystal structure of the extracellular domain of the α 1 subunit bound to α -bungarotoxin at 1.94Å resolution [305], has

revealed the orthosteric binding site in detail (reviewed in [196, 662, 1199, 1281]). Nicotinic receptors at the somatic neuromuscular junction of adult animals have the stoichiometry (α 1)₂ β 1 $\delta\epsilon$, whereas an extrajunctional (α 1)₂ β 1 $\gamma\delta$ receptor predominates in embryonic and denervated skeletal muscle and other pathological states. Other nicotinic receptors are assembled as combinations of α (2-6) and β (2-4) subunits. For α 2, α 3, α 4 and β 2 and β 4 subunits, pairwise combinations of α and β (*e.g.* α 3 β 4 and α 4 β 2) are sufficient to form a functional receptor *in vitro*, but far more complex isoforms may exist *in vivo* (reviewed in [467, 468, 951]). There is strong evidence that the pairwise assembly of some α and β subunits can occur with variable stoichiometry [*e.g.* (α 4)₂(β 2)₂ or (α 4)₃(β 2)₂] which influences the biophysical and pharmacological properties of the receptor [951]. α 5 and β 3 subunits lack function when expressed alone, or pairwise, but participate in the formation of functional hetero-oligomeric receptors when expressed as a third subunit with another α and β pair [*e.g.* α 4 α 5 α β 2, α 4 α β 2 β 3, α 5 α β 2, see [951] for further examples]. The α 6 subunit can form a functional receptor when co-expressed with β 4 *in vitro*, but more efficient expression ensues from incorporation of a third partner, such as β 3 [1560]. The α 7, α 8, and α 9 subunits form functional homo-oligomers, but can also combine with a second subunit to constitute a hetero-oligomeric assembly (*e.g.* α 7 β 2 and

α 9 α 10). For functional expression of the α 10 subunit, co-assembly with α 9 is necessary. The latter, along with the α 10 subunit, appears to be largely confined to cochlear and vestibular hair cells. Comprehensive listings of nicotinic receptor subunit combinations identified from recombinant expression systems, or *in vivo*, are given in [951]. In addition, numerous proteins interact with nicotinic ACh receptors modifying their assembly, trafficking to and from the cell surface, and activation by ACh (reviewed by [48, 650, 950]).

The nicotinic receptor Subcommittee of **NC-IUPHAR** has recommended a nomenclature and classification scheme for nicotinic acetylcholine (nACh) receptors based on the subunit composition of known, naturally- and/or heterologously-expressed nACh receptor subtypes [868]. Headings for this table reflect abbreviations designating nACh receptor subtypes based on the predominant α subunit contained in that receptor subtype. An asterisk following the indicated α subunit denotes that other subunits are known to, or may, assemble with the indicated α subunit to form the designated nACh receptor subtype(s). Where subunit stoichiometries within a specific nACh receptor subtype are known, numbers of a particular subunit larger than 1 are indicated by a subscript following the subunit (enclosed in parentheses- see also [253]).

Nomenclature	nicotinic acetylcholine receptor $\alpha 1$ subunit	nicotinic acetylcholine receptor $\alpha 2$ subunit	nicotinic acetylcholine receptor $\alpha 3$ subunit	nicotinic acetylcholine receptor $\alpha 4$ subunit
HGNC, UniProt	<i>CHRNA1</i> , P02708	<i>CHRNA2</i> , Q15822	<i>CHRNA3</i> , P32297	<i>CHRNA4</i> , P43681
Commonly used antagonists	($\alpha 1$) ₂ $\beta 1\gamma\delta$ and ($\alpha 1$) ₂ $\beta 1\delta\epsilon$: α-bungarotoxin > pancuronium > vecuronium > rocuroonium > tubocurarine (IC ₅₀ = 43 - 82 nM)	$\alpha 2\beta 2$: DHβE (K_B = 0.9 μ M), tubocurarine (K_B = 1.4 μ M); $\alpha 2\beta 4$: DHβE (K_B = 3.6 μ M), tubocurarine (K_B = 4.2 μ M)	$\alpha 3\beta 2$: DHβE (K_B = 1.6 μ M, IC ₅₀ = 2.0 μ M), tubocurarine (K_B = 2.4 μ M); $\alpha 3\beta 4$: DHβE (K_B = 19 μ M, IC ₅₀ = 26 μ M), tubocurarine (K_B = 2.2 μ M)	$\alpha 4\beta 2$: DHβE (K_B = 0.1 μ M; IC ₅₀ = 0.08 - 0.9 μ M), tubocurarine (K_B = 3.2 μ M, IC ₅₀ = 34 μ M); $\alpha 4\beta 4$: DHβE (K_B = 0.01 μ M, IC ₅₀ = 0.19 - 1.2 μ M), tubocurarine (K_B = 0.2 μ M, IC ₅₀ = 50 μ M)
Selective agonists	succinylcholine (selective for ($\alpha 1$) ₂ $\beta 1\gamma\delta$)	–	–	varenicline [248], rivanicline [332], TC-2559 ($\alpha 4\beta 2$) [219]
Selective antagonists	α-bungarotoxin , α-conotoxin GI , α-conotoxin MI , pancuronium , waglerin-1 (selective for ($\alpha 1$) ₂ $\beta 1\delta\epsilon$)	–	α-conotoxin AulB ($\alpha 3\beta 4$), α-conotoxin MII ($\alpha 3\beta 2$), α-conotoxin PnIA ($\alpha 3\beta 2$), α-conotoxin TxIA ($\alpha 3\beta 2$), α-conotoxin-GIC ($\alpha 3\beta 2$)	–
Channel blockers	gallamine (($\alpha 1$) ₂ $\beta 1\gamma\delta$ and ($\alpha 1$) ₂ $\beta 1\delta\epsilon$) (pIC ₅₀ ~6), mecamylamine (($\alpha 1$) ₂ $\beta 1\delta\epsilon$) (pIC ₅₀ ~5.8)	hexamethonium , mecamylamine	mecamylamine ($\alpha 3\beta 4$) (pIC ₅₀ 6.4), mecamylamine ($\alpha 3\beta 2$) (pIC ₅₀ 5.1), A-867744 ($\alpha 3\beta 4$) [906], NS1738 ($\alpha 3\beta 4$) [1389], hexamethonium ($\alpha 3\beta 4$), hexamethonium ($\alpha 3\beta 2$)	mecamylamine ($\alpha 4\beta 4$) (pIC ₅₀ 5.3–6.5), mecamylamine ($\alpha 4\beta 2$) (pIC ₅₀ 5.4–5.4), hexamethonium ($\alpha 4\beta 2$) (pIC ₅₀ 4.5–5.2), hexamethonium ($\alpha 4\beta 4$) (pIC ₅₀ 4), A-867744 ($\alpha 4\beta 2$) [906], NS1738 ($\alpha 4\beta 2$) [1389]
Allosteric modulators (Positive)	–	LY2087101 [151]	–	LY2087101 (potentiates $\alpha 4\beta 2$ and $\alpha 4\beta 4$) [151]
Selective allosteric modulators	–	–	–	NS9283 (Positive) [780]
Labelled ligands	[¹²⁵I]α-bungarotoxin (Selective Antagonist), [³H]α-bungarotoxin (Selective Antagonist)	[¹²⁵I]epibatidine (Agonist), [³H]epibatidine (Agonist), [³H]cytisine (Agonist)	[¹²⁵I]epibatidine (Agonist), [³H]epibatidine (Agonist), [¹²⁵I]epibatidine (Agonist), [³H]epibatidine (Agonist), [³H]cytisine (Agonist)	[¹²⁵I]epibatidine (Agonist), [³H]epibatidine (Agonist), [³H]cytisine (Agonist), [¹²⁵I]epibatidine (Agonist), [³H]epibatidine (Agonist), [³H]epibatidine (Agonist) – Rat, [³H]cytisine (Agonist)
Functional Characteristics	($\alpha 1$) ₂ $\beta 1\gamma\delta$: P _{Ca} /P _{Na} = 0.16 - 0.2, P _f = 2.1 - 2.9%; ($\alpha 1$) ₂ $\beta 1\delta\epsilon$: P _{Ca} /P _{Na} = 0.65 - 1.38, P _f = 4.1 - 7.2%	$\alpha 2\beta 2$: P _{Ca} /P _{Na} ~ 1.5	$\alpha 3\beta 2$: P _{Ca} /P _{Na} = 1.5; $\alpha 3\beta 4$: P _{Ca} /P _{Na} = 0.78 - 1.1, P _f = 2.7 - 4.6%	$\alpha 4\beta 2$: P _{Ca} /P _{Na} = 1.65, P _f = 2.6 - 2.9%; $\alpha 4\beta 4$: P _f = 1.5 - 3.0 %

Nomenclature	nicotinic acetylcholine receptor $\alpha 5$ subunit	nicotinic acetylcholine receptor $\alpha 6$ subunit	nicotinic acetylcholine receptor $\alpha 7$ subunit
HGNC, UniProt	CHRNA5 , P30532	CHRNA6 , Q15825	CHRNA7 , P36544
Commonly used antagonists	–	$\alpha 6/\alpha 3\beta 2\beta 3$ chimera: DHβE (IC ₅₀ = 1.1 μ M)	($\alpha 7$) ₅ : DHβE (IC ₅₀ = 8 - 20 μ M); ($\alpha 7$) ₅ : tubocurarine (IC ₅₀ = 3.1 μ M)
Selective agonists	–	–	encenicline (Partial agonist) [926, 1028], AQW051 ([125I] α -bungarotoxin binding assay) [594], 4BP-TQS (allosteric) [456], A-582941 (($\alpha 7$) ₅) [121], PHA-543613 (($\alpha 7$) ₅) [1517], PHA-709829 (($\alpha 7$) ₅) [11], PNU-282987 (($\alpha 7$) ₅) [129], bradanicline (($\alpha 7$) ₅) [521]
Selective antagonists	α-conotoxin MII , α-conotoxin PnIA , α-conotoxin TxIA , α-conotoxin-GIC	α-conotoxin MII ($\alpha 6\beta 2^*$), α-conotoxin MII [H9A, L15A] ($\alpha 6\beta 2\beta 3$), α-conotoxin PIA ($\alpha 6/\alpha 3\beta 2\beta 3$ chimera)	α-bungarotoxin (($\alpha 7$) ₅), α-conotoxin ArIB (($\alpha 7$) ₅), α-conotoxin ImI (($\alpha 7$) ₅), methyllycaconitine (($\alpha 7$) ₅)
Channel blockers	–	mecamylamine ($\alpha 6/\alpha 3\beta 2\beta 3$ chimera) (pIC ₅₀ 5), hexamethonium ($\alpha 6/\alpha 3\beta 2\beta 3$ chimera) (pIC ₅₀ 4)	mecamylamine (($\alpha 7$) ₅) (pIC ₅₀ 4.8)
Allosteric modulators (Positive)	–	–	A-867744 (($\alpha 7$) ₅ :Type 2; also blocks $\alpha 3\beta 4$ and $\alpha 4\beta 2$) [906], LY2087101 (($\alpha 7$) ₅ :Type 1) [151], NS1738 (($\alpha 7$) ₅ :Type 1; also blocks $\alpha 3\beta 4$ and $\alpha 4\beta 2$) [1389]
Selective allosteric modulators	–	–	BNC375 (Positive) (pEC ₅₀ 5.6) [1483], JNJ1930942 (Positive) [325], PNU-120596 (Positive) [593]
Labelled ligands	–	[³H]epibatidine (Agonist) – Chicken, [¹²⁵I]α-conotoxin MII (Antagonist)	[³H]epibatidine (Agonist), [³H]A-585539 (Agonist) [39], [³H]AZ11637326 (Agonist) [465], [¹²⁵I]α-bungarotoxin (Selective Antagonist) (pK _d 8.3–9.1), [³H]α-bungarotoxin (Selective Antagonist) (pK _d 8.3–9.1), [³H]methyllycaconitine (Antagonist) (pK _d 8.7) – Rat
Functional Characteristics	–	–	P _{Ca} /P _{Na} = 6.6-20, P _f = 8.8 - 11.4%

Nomenclature	nicotinic acetylcholine receptor $\alpha 8$ subunit (avian)	nicotinic acetylcholine receptor $\alpha 9$ subunit	nicotinic acetylcholine receptor $\alpha 10$ subunit
HGNC, UniProt	–	CHRNA9 , Q9UGM1	CHRNA10 , Q9GZZ6
Commonly used antagonists	($\alpha 8$) ₅ : α-bungarotoxin > atropine \geq tubocurarine \geq strychnine	($\alpha 9$) ₅ : α-bungarotoxin > methyllycaconitine > strychnine \sim tropisetron > tubocurarine ; $\alpha 9\alpha 10$: α-bungarotoxin > tropisetron = strychnine > tubocurarine	$\alpha 9\alpha 10$: α-bungarotoxin > tropisetron = strychnine > tubocurarine
Selective antagonists	–	α-bungarotoxin ($\alpha 9\alpha 10$), α-bungarotoxin (($\alpha 9$) ₅), α-conotoxin RgIA ($\alpha 9\alpha 10$), muscarine ($\alpha 9\alpha 10$), muscarine (($\alpha 9$) ₅), nicotine ($\alpha 9\alpha 10$), nicotine (($\alpha 9$) ₅), strychnine (($\alpha 9$) ₅), strychnine ($\alpha 9\alpha 10$)	α-bungarotoxin ($\alpha 9\alpha 10$), α-conotoxin RgIA ($\alpha 9\alpha 10$), muscarine ($\alpha 9\alpha 10$), nicotine ($\alpha 9\alpha 10$), strychnine ($\alpha 9\alpha 10$)
Labelled ligands	[³H]epibatidine (($\alpha 8$) ₅) (pK _d 9.7), [¹²⁵I]α-bungarotoxin (native $\alpha 8^*$) (pK _d 8.3), [³H]α-bungarotoxin (native $\alpha 8^*$) (pK _d 8.3)	[³H]methyllycaconitine (Antagonist) (pK _d 8.1), [¹²⁵I]α-bungarotoxin (Antagonist), [³H]α-bungarotoxin (Antagonist)	[³H]methyllycaconitine (Antagonist) (pK _d 8.1)
Functional Characteristics	–	($\alpha 9$) ₅ : P _{Ca} /P _{Na} = 9; $\alpha 9\alpha 10$: P _{Ca} /P _{Na} = 9, P _f = 22%	$\alpha 9\alpha 10$: P _{Ca} /P _{Na} = 9, P _f = 22%

Nomenclature	nicotinic acetylcholine receptor $\beta 1$ subunit	nicotinic acetylcholine receptor $\beta 2$ subunit	nicotinic acetylcholine receptor $\beta 3$ subunit	nicotinic acetylcholine receptor $\beta 4$ subunit	nicotinic acetylcholine receptor γ subunit	nicotinic acetylcholine receptor δ subunit	nicotinic acetylcholine receptor ϵ subunit
HGNC, UniProt	CHRN1 , P11230	CHRN2 , P17787	CHRN3 , Q05901	CHRN4 , P30926	CHRN7 , P07510	CHRN8 , Q07001	CHRN9 , Q04844
Antagonists	–	–	–	–	–	PhTX-11 (pIC ₅₀ 6.2–6.3) [1328]	–
Comments	Ligand interaction data for hetero-oligomeric receptors containing the $\beta 1$ subunit are listed under the $\alpha 1$ subunits	Ligand interaction data for hetero-oligomeric receptors containing the $\beta 2$ subunit are listed under the α subunits	Ligand interaction data for hetero-oligomeric receptors containing the $\beta 3$ subunit are listed under the α subunits	Ligand interaction data for hetero-oligomeric receptors containing the $\beta 4$ subunit are listed under the α subunits	Ligand interaction data for hetero-oligomeric receptors containing the γ subunit are listed under the $\alpha 1$ subunits	Ligand interaction data for hetero-oligomeric receptors containing the δ subunit are listed under the $\alpha 1$ subunits	Ligand interaction data for hetero-oligomeric receptors containing the ϵ subunit are listed under the $\alpha 1$ subunits

Comments: Commonly used agonists of nACh receptors that display limited discrimination in functional assays between receptor subtypes include [A-85380](#), [cytisine](#), [DMPP](#), [epibatidine](#), [nicotine](#) and the natural transmitter, ACh. A summary of their profile across differing receptors is provided in [\[468\]](#) and quantitative data across numerous assay systems are summarized in [\[628\]](#). Quantitative data presented in the table for commonly used antagonists and channel blockers for human receptors studied under voltage-clamp are from [\[164, 203, 1075, 1076, 1086, 1527\]](#). Type I PAMs increase peak agonist-evoked responses but have little, or no, effect on the rate of desensitization of $\alpha 7$ nicotinic ACh receptors whereas type II PAMs also cause a large reduction in desensitization (reviewed in [\[1513\]](#)).

Further reading on Nicotinic acetylcholine receptors (nACh)

- Auerbach A. (2015) Agonist activation of a nicotinic acetylcholine receptor. *Neuropharmacology* **96**: 150–6 [[PMID:25446670](#)]
- Bertrand D *et al.* (2015) Therapeutic Potential of $\alpha 7$ Nicotinic Acetylcholine Receptors. *Pharmacol Rev* **67**: 1025–73 [[PMID:26419447](#)]
- Bouzat C *et al.* (2018) Nicotinic acetylcholine receptors at the single-channel level. *Br J Pharmacol* **175**: 1789–1804 [[PMID:28261794](#)]
- Chatzidaki A *et al.* (2015) Allosteric modulation of nicotinic acetylcholine receptors. *Biochem Pharmacol* **97**: 408–417 [[PMID:26231943](#)]
- Corradi J *et al.* (2016) Understanding the Bases of Function and Modulation of $\alpha 7$ Nicotinic Receptors: Implications for Drug Discovery. *Mol Pharmacol* **90**: 288–99 [[PMID:27190210](#)]
- Crespi A *et al.* (2018) Proteins and chemical chaperones involved in neuronal nicotinic receptor expression and function: an update. *Br J Pharmacol* **175**: 1869–1879 [[PMID:28294298](#)]
- Dineley KT *et al.* (2015) Nicotinic ACh receptors as therapeutic targets in CNS disorders. *Trends Pharmacol Sci* **36**: 96–108 [[PMID:25639674](#)]
- Lukas RJ *et al.* (1999) International Union of Pharmacology. XX. Current status of the nomenclature for nicotinic acetylcholine receptors and their subunits. *Pharmacol Rev* **51**: 397–401 [[PMID:10353988](#)]
- Stokes C *et al.* (2015) Looking below the surface of nicotinic acetylcholine receptors. *Trends Pharmacol Sci* **36**: 514–23 [[PMID:26067101](#)]
- Wang J *et al.* (2018) Orthosteric and allosteric potentiation of heteromeric neuronal nicotinic acetylcholine receptors. *Br J Pharmacol* **175**: 1805–1821 [[PMID:28199738](#)]
- Wu J *et al.* (2016) Heteromeric $\alpha 7\beta 2$ Nicotinic Acetylcholine Receptors in the Brain. *Trends Pharmacol Sci* **37**: 562–574 [[PMID:27179601](#)]

P2X receptors

[Ion channels](#) → [Ligand-gated ion channels](#) → [P2X receptors](#)

Overview: P2X receptors (**nomenclature as agreed by the NC-IUPHAR Subcommittee on P2X Receptors** [[253, 698](#)]) have a trimeric stoichiometry [[596, 635, 686, 1009](#)] with two putative TM domains per P2X subunit, gating primarily Na⁺, K⁺ and Ca²⁺, exceptionally Cl⁻. The Nomenclature Subcommittee has recommended that for P2X receptors, structural criteria should be the initial basis for nomenclature where possible. X-ray crystallography indicates that functional P2X receptors are trimeric and three ag-

onist molecules are required to bind to a single trimeric assembly in order to activate it [[461, 520, 596, 686, 908](#)]. Native receptors may occur as either homotrimers (*e.g.* P2X1 in smooth muscle) or heterotrimers (*e.g.* P2X2:P2X3 in the nodose ganglion [[1450](#)], P2X1:P2X5 in mouse cortical astrocytes [[764](#)], and P2X2:P2X5 in mouse dorsal root ganglion, spinal cord and mid pons [[254, 1226](#)]. P2X2, P2X4 and P2X7 receptor activation can lead to influx of large cationic molecules, such as NMDG⁺, Yo-Pro, ethidium or propidium

iodide [[1098](#)]. The permeability of the P2X7 receptor is modulated by the amount of cholesterol in the plasma membrane [[983](#)]. The hemi-channel pannexin-1 was initially implicated in the action of P2X7 [[1099](#)], but not P2X2, receptors [[201](#)], but this interpretation is probably misleading [[1110](#)]. Convincing evidence now supports the view that the activated P2X7 receptor is immediately permeable to large cationic molecules, but influx proceeds at a much slower pace than that of the small cations Na⁺, K⁺, and Ca²⁺ [[317](#)].

Searchable database: <http://www.guidetopharmacology.org/>

Full Contents of ConciseGuide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.70231/full>

P2X receptors S178

Nomenclature	P2X1	P2X2	P2X3
HGNC, UniProt	P2RX1 , P51575	P2RX2 , Q9UBL9	P2RX3 , P56373
Endogenous agonists	–	ATP [615] – Rat	ATP [616]
Agonists	$\alpha\beta$ -meATP, BzATP, L- β -meATP	–	$\alpha\beta$ -meATP, BzATP
Antagonists	TNP-ATP (pIC ₅₀ ~8.9) [1437], Ip ₃ I (pIC ₅₀ ~8.5) [705], NF023 (pIC ₅₀ ~6.7) [1304], NF449 (pIC ₅₀ ~6.3) [681]	NF770 (pIC ₅₀ 7–8) [1033], NF778 (pIC ₅₀ 7–8) [1033], PSB-10211 (pIC ₅₀ ~7) [1033]	TNP-ATP (pIC ₅₀ ~8.9) [1437], AF-906 (pIC ₅₀ 8.9) [618], gefapixant (pIC ₅₀ 8.5) [618], sivopixant (pIC ₅₀ 8.4) [661], eliapixant (pIC ₅₀ 8.1) [282], camlipixant (pIC ₅₀ 7.6) [427], A317491 (pIC ₅₀ ~7.5) [623]
Selective allosteric modulators	MRS 2219 (Positive) [617]	–	–

Nomenclature	P2X4	P2X5	P2X6	P2X7
HGNC, UniProt	P2RX4 , Q99571	P2RX5 , Q93086	P2RX6 , O15547	P2RX7 , Q99572
Endogenous agonists	ATP [616]	ATP [616] – Rat	ATP [616] – Rat	ATP [616]
Agonists	$\alpha\beta$ -meATP, BzATP	–	–	–
Antagonists	BAY-1797 (pIC ₅₀ 7) [1507], PSB-12054 (pIC ₅₀ 6.7) [534], 5-BDBD (pIC ₅₀ 5–6) [618, 1033], BX-430 (pIC ₅₀ 5–6) [618, 1033], PSB-12062 (pIC ₅₀ 5–6) [618, 1033], paroxetine (pIC ₅₀ 5–6) [618, 1033]	–	–	AZ10606120 (pK _d 8.9) [941], A804598 (pIC ₅₀ ~8), brilliant blue G (pIC ₅₀ ~8) [636], A839977 (pIC ₅₀ ~7.7) [336, 338, 563], A740003 (pIC ₅₀ 7.4) [564], decavanadate (pA ₂ = 7.4) (pA ₂ 7.4) [946], A438079 (pIC ₅₀ ~6.9) [336], AZ11657312 (salt free) (pA ₂ 6.1) [53]
Selective antagonists	–	–	–	JNJ-47965567 (pK _i 7.9) [116]
Allosteric modulators (Positive)	nimodipine [1234]	–	–	GW791343 [941, 943] – Rat, LL-37 (CAMP, P49913) [1392], clemastine [1029], ivermectin [1030], polymyxin B [398]
Allosteric modulators (Negative)	amlodipine [1234]	–	–	AZ10606120 [941], GW791343 [941, 943], KN62 [434, 1263]
Selective allosteric modulators	ivermectin (Positive) (pEC ₅₀ ~6.6) [699] – Rat	–	–	chelerythrine (Negative) (pIC ₅₀ 5.2) [1263], AZ11645373 (Negative) [944, 1318]
Comments	–	–	–	Ginsenoside compounds acts as positive allosteric modulators of P2X7 [1125], however, the effects of allosteric regulators at P2X7 receptors are species-dependent.

Comments: A317491 and RO3 also block the P2X2:P2X3 heteromultimer [411, 623]. NF449, A317491 and RO3 are more than 10-fold selective for P2X1 and P2X3 receptors, respectively. Agonists listed show selectivity within recombinant P2X receptors of *ca.* one order of magnitude. A804598, A839977, A740003 and A438079 are at least 10-fold selective for P2X7 receptors and show similar affinity across human and rodent receptors [336, 338, 563].

Several P2X receptors (particularly P2X1 and P2X3) may be inhibited by desensitisation using stable agonists (*e.g.* $\alpha\beta$ -meATP); suramin and PPADS are non-selective antagonists at rat and human P2X1-3,5 and hP2X4, but not rP2X4,6,7 [163], and can also inhibit ATPase activity [258]. Ip₃I is inactive at rP2X2, an antagonist at rP2X3 (pIC₅₀ 5.6) and enhances agonist responses at rP2X4 [705]. Antagonist potency of NF023 at recombinant P2X2, P2X3

and P2X5 is two orders of magnitude lower than that at P2X1 receptors [1304]. The P2X7 receptor may be inhibited in a non-competitive manner by the protein kinase inhibitors KN62 and chelerythrine [1263], while the p38 MAP kinase inhibitor GT- γ S and the cyclic imide AZ11645373 show a species-dependent non-competitive action [337, 944, 945, 1318]. The pH-sensitive dye used in culture media, phenol red, is also reported to inhibit

P2X1 and P2X3 containing channels [706]. Some recombinant P2X receptors expressed to high density bind [³⁵S]ATPγS and [³H]αβ-meATP, although the latter can also bind to 5'-nucleotidase [942]. [³H]A317491 and [³H]A804598 have been used as high affinity antagonist radioligands for P2X3 (and P2X2/3) and P2X7 receptors, respectively [338]. Several high affinity radioligands for the P2X7 receptor have been recently synthesized, some with very promising applications in the diagnosis of inflammatory diseases of the central nervous system [387, 857, 1238, 1373, 1617].

Several P2X3 antagonists have entered clinical trials for refractory chronic cough. In 2022, **gefapixant** was approved in Japan for the management of refractory or unexplained chronic cough [911]. Oxidized ATP covalently binds un-protonated lysine residues in the vicinity of the ATP-binding site and irreversibly inhibits the P2X7 receptor. Other plasma membrane receptors exposing available lysines may also be blocked by oATP [102, 316]. The cryoelectron microscopy structures of full-length rP2X7 receptor in apo and ATP-bound states have been resolved [927]. A propor-

tion (<10%) of screened humans were found to possess full length P2X5 subunits (444 aa), which can assemble into a functional P2X5 receptor [704, 734]. An uncharged region at the N-terminus of P2X6 exerts an inhibitory effect on its assembly and export from the ER [1058]. The P2X6 subunit also lacks nine residues in the left flipper, which is a key element in agonist docking at P2X receptors [1047].

Further reading on P2X receptors

- Di Virgilio F *et al.* (2017) The P2X7 Receptor in Infection and Inflammation. *Immunity* **47**: 15-31 [PMID:28723547]
- Di Virgilio F *et al.* (2018) The Elusive P2X7 Macropore. *Trends Cell Biol* **28**: 392-404 [PMID:29439897]
- Habermacher C *et al.* (2016) Molecular structure and function of P2X receptors. *Neuropharmacology* **104**: 18-30 [PMID:26231831]
- Illes P *et al.* (2021) Update of P2X receptor properties and their pharmacology: IUPHAR Review 30. *Br J Pharmacol* **178**: 489-514 [PMID:33125712]
- Jacobson KA *et al.* (2016) Medicinal chemistry of adenosine, P2Y and P2X receptors. *Neuropharmacology* **104**: 31-49 [PMID:26686393]
- King BF. (2023) Rehabilitation of the P2X5 receptor: a re-evaluation of structure and function. *Purinergic Signal* **19**: 421-439 [PMID:36279087]
- McCarthy AE *et al.* (2019) Full-Length P2X₇ Structures Reveal How Palmitoylation Prevents Channel Desensitization. *Cell* **179**: 659-670.e13 [PMID:31587896]
- Stokes L *et al.* (2017) P2X4 Receptor Function in the Nervous System and Current Breakthroughs in Pharmacology. *Front Pharmacol* **8**: 291 [PMID:28588493]

ZAC

Ion channels → **Ligand-gated ion channels** → **ZAC**

Overview: The zinc-activated channel (ZAC, **nomenclature as agreed by the NC-IUPHAR Subcommittee for the Zinc Activated Channel**) is a member of the Cys-loop family that includes the nicotinic ACh, 5-HT₃, GABA_A and strychnine-sensitive glycine receptors [284, 574, 1402]. The channel is likely to ex-

ist as a homopentamer of 4TM subunits that form an intrinsic cation selective channel equipermeable to Na⁺, K⁺ and Cs⁺, but impermeable to Ca²⁺ and Mg²⁺ [1402]. ZAC displays constitutive activity that can be blocked by **tubocurarine**, TTFB and high concentrations of Ca²⁺ [1402]. Although denoted ZAC, the channel

is more potently activated by H⁺ and Cu²⁺, with greater and lesser efficacy than Zn²⁺, respectively [1402]. Orthologs of the human ZACN gene are present in a wide range of mammalian genomes, but notably not in the mouse or rat genomes. [284, 574].

Nomenclature	ZAC
HGNC, UniProt	ZACN, Q401N2
Endogenous agonists	H ⁺ [1402], Cu ²⁺ [1402], Zn ²⁺ [284, 1402]
Antagonists	tubocurarine (pIC ₅₀ 5.2) [284], Ca ²⁺ (pIC ₅₀ 2) [1402]
Allosteric modulators	TTFB (Antagonist) (pIC ₅₀ 5.5) [886]
Functional Characteristics	Outwardly rectifying current (both constitutive and evoked by Zn ²⁺)

Comments: A ZACN gene does not appear to exist in the mouse or rat genomes [284]. Although tabulated as an antagonist, it is possible that **tubocurarine** acts as a channel blocker. Antagonism by Ca²⁺ is voltage-independent. ZAC is not activated (at 1 mM) by transition metals including Fe²⁺, Co²⁺, Ni²⁺, Cd²⁺, or Al³⁺ [1402]. The concentration response relationship to Cu²⁺ is biphasic, with concentrations exceeding 30 μM being associated with reduced activation [1402]. The N-(thiazol-2-yl)-benzamide analog TTFB has been identified as a moderately potent negative allosteric modulator of ZAC. TTFB displays negligible activity at representatives of the GABA_A, glycine, 5-HT₃ and nicotinic ACh receptors, and thus it constitutes a potential pharmacological tool for ZAC.

Further reading on ZAC

- Collingridge GL *et al.* (2009) A nomenclature for ligand-gated ion channels. *Neuropharmacology* **56**: 2-5 [PMID:18655795]
- Madjroh N *et al.* (2021) Delineation of the functional properties exhibited by the Zinc-Activated Channel (ZAC) and its high-frequency Thr¹²⁸Ala variant (rs2257020) in *Xenopus* oocytes. *Pharmacol Res* **169**: 105653 [PMID:33962015]
- Madjroh N *et al.* (2021) Probing the molecular basis for signal transduction through the Zinc-Activated Channel (ZAC). *Biochem Pharmacol* **193**: 114781 [PMID:34560053]
- Madjroh N *et al.* (2021) Discovery and functional characterization of N-(thiazol-2-yl)-benzamide analogs as the first class of selective antagonists of the Zinc-Activated Channel (ZAC). *Biochem Pharmacol* **193**: 114782 [PMID:34560054]
- Peralta FA *et al.* (2016) Zinc as Allosteric Ion Channel Modulator: Ionotropic Receptors as Metalloproteins. *Int J Mol Sci* **17**: [PMID:27384555]
- Trattinig SM *et al.* (2016) Copper and protons directly activate the zinc-activated channel. *Biochem Pharmacol* **103**: 109-17 [PMID:26872532]

Voltage-gated ion channels

[Ion channels](#) → [Voltage-gated ion channels](#)

Overview: The voltage-gated ion channels and their structural relatives comprise a superfamily encoded by at least 143 genes in the human genome and are therefore one of the largest superfamilies of signal transduction proteins, following the G protein-coupled receptors and the protein kinases in number [185]. In addition to their prominence in signal transduction, these ion channels are also among the most common drug targets. As for

other large protein superfamilies, understanding the molecular relationships among family members, developing a unified, rational nomenclature for the ion channel families and subfamilies, and assigning physiological functions and pharmacological significance to each family member has been an important challenge. Some of the ion channels placed under the 'Voltage-gated' umbrella are not in fact gated by voltage, but for the reasons

mentioned above it is useful to consider them within this superfamily. The inwardly rectifying potassium channels, two-pore domain potassium channels (K2P), ryanodine receptors (RyR) and transient receptor potential channels (TRP) are those that are NOT voltage-gated.

CatSper and Two-Pore channels (TPC)

[Ion channels](#) → [Voltage-gated ion channels](#) → [CatSper and Two-Pore channels \(TPC\)](#)

Overview: CatSper channels (CatSper1-4, **nomenclature as agreed by NC-IUPHAR** [246]) are putative 6TM, voltage-gated, alkalization-activated calcium permeant channels that are presumed to assemble as a tetramer of α -like subunits and mediate the current I_{CatSper} [707]. In mammals, CatSper subunits are structurally most closely related to individual domains of voltage-activated calcium channels (Ca_v) [1172]. CatSper1 [1172], CatSper2 [1153] and CatSper3 and 4 [639, 851, 1144],

in common with a putative 2TM auxiliary CatSper β protein [841] and two putative 1TM associated CatSper γ and CatSper δ proteins [237, 1468], are restricted to the testis and localised to the principle piece of sperm tail. The novel cross-species CatSper channel inhibitor, RU1968, has been proposed as a useful tool to aid characterisation of native CatSper channels [1173].

Two-pore channels (TPCs) are structurally related to CatSper3, Ca_v s and Na_v s. TPCs have a 2x6TM structure with twice the num-

ber of TMs of CatSper3 and half that of Ca_v s. There are three animal TPCs (TPC1-TPC3). Humans have TPC1 and TPC2, but not TPC3. TPC1 and TPC2 are localized in endosomes and lysosomes [170]. TPC3 is also found on the plasma membrane and forms a voltage-activated, non-inactivating Na^+ channel [173]. All the three TPCs are Na^+ -selective under whole-cell or whole-organelle patch clamp recording [174, 175, 1534]. The channels may also conduct Ca^{2+} [972].

Nomenclature	CatSper1	CatSper2	CatSper3	CatSper4
HGNC, UniProt	CATSPER1 , Q8NEC5	CATSPER2 , Q96P56	CATSPER3 , Q86XQ3	CATSPER4 , Q7RTX7
Activators	CatSper1 is constitutively active, weakly facilitated by membrane depolarisation, strongly augmented by intracellular alkalinisation. In human, but not mouse, progesterone (EC ₅₀ ~ 8 nM) also potentiates the CatSper current (I _{CatSper}). [836, 1330]	–	–	–
Channel blockers	ruthenium red (Inhibition) (pIC ₅₀ 5) [707] – Mouse, HC-056456 (pIC ₅₀ 4.7) [181], Cd²⁺ (Inhibition) (pIC ₅₀ 3.7) [707] – Mouse, Ni²⁺ (Inhibition) (pIC ₅₀ 3.5) [707] – Mouse	–	–	–
Selective channel blockers	NNC55-0396 (Inhibition) (pIC ₅₀ 5.7) [-80mV – 80mV] [836, 1330], mibefradil (Inhibition) (pIC ₅₀ 4.4–4.5) [1330]	–	–	–
Functional Characteristics	Calcium selective ion channel (Ba ²⁺ >Ca ²⁺ >>Mg ²⁺ >>Na ⁺); quasilinear monovalent cation current in the absence of extracellular divalent cations; alkalinization shifts the voltage-dependence of activation towards negative potentials [V ₁ @ pH 6.0 = +87 mV (mouse); V ₁ @ pH 7.5 = +11mV (mouse) or pH 7.4 = +85 mV (human)]; required for I _{CatSper} and male fertility (mouse and human)	Required for I _{CatSper} and male fertility (mouse and human)	Required for I _{CatSper} and male fertility (mouse)	Required for I _{CatSper} and male fertility (mouse)

Nomenclature	TPC1	TPC2
HGNC, UniProt	TPCN1 , Q9ULQ1	TPCN2 , Q8NHX9
Activators	phosphatidyl (3,5) inositol bisphosphate (pEC ₅₀ 6.5) [174]	phosphatidyl (3,5) inositol bisphosphate (pEC ₅₀ 6.4) [1484]
Selective activators	–	TPC2-A1-N (pEC ₅₀ 5.1) [442]
Channel blockers	verapamil (Inhibition) (pIC ₅₀ 4.6) [174], Cd²⁺ (Inhibition) (pIC ₅₀ 3.7) [174]	verapamil (Inhibition) (pIC ₅₀ 5) [1484]
Functional Characteristics	Organelle voltage-gated Na ⁺ -selective channel (Na ⁺ >>K ⁺ >>Ca ²⁺); Required for the generation of action potential-like long depolarization in lysosomes. Voltage-dependence of activation is sensitive to luminal pH (determined from lysosomal recordings). $\psi_{1/2}$ @ pH4.6 = +91 mV; $\psi_{1/2}$ @ pH6.5 = +2.6 mV. Maximum activity requires PI(3,5)P2 and reduced [ATP], or depletion of extracellular amino acids.	Organelle voltage-independent Na ⁺ -selective channel (Na ⁺ >>K ⁺ >>Ca ²⁺). Sensitive to the levels of PI(3,5)P2. Activated by decreases in [ATP] or depletion of extracellular amino acids

Comments: CatSper channel subunits expressed singly, or in combination, fail to functionally express in heterologous expression systems [1153, 1172]. The properties of CatSper1 tabulated above are derived from whole cell voltage-clamp recordings comparing currents endogenous to spermatozoa isolated from the *corpus epididymis* of wild-type and *Catsper1*^(-/-) mice [707] and also mature human sperm [836, 1330]. I_{CatSper} is also undetectable in the spermatozoa of *Catsper2*^(-/-), *Catsper3*^(-/-), *Catsper4*^(-/-), or CatSper δ ^(-/-) mice, and CatSper 1 associates with CatSper 2, 3, 4, β , γ , and δ [237, 841, 1144]. Moreover, targeted disruption of *Catsper1*, 2, 3, 4, or δ genes results in an identical phenotype in which spermatozoa fail to exhibit the hyperactive movement (whip-like flagellar beats) necessary for penetration of the egg *cumulus* and *zona pellucida* and subsequent fertilization. Such disruptions are associated with a deficit in alkalization and depolarization-evoked Ca²⁺ entry into spermatozoa [182, 237, 1144]. Thus, it is likely that the CatSper pore is formed by a heterotetramer of CatSper1-4 [1144] in association with the auxiliary subunits (β , γ , δ) that are also essential for function [237]. CatSper channels are required for the

increase in intracellular Ca²⁺ concentration in sperm evoked by egg *zona pellucida* glycoproteins [1534]. Mouse and human sperm swim against the fluid flow and Ca²⁺ signaling through CatSper is required for the rheotaxis [949]. *In vivo*, CatSper1-null spermatozoa cannot ascend the female reproductive tracts efficiently [238, 552]. It has been shown that CatSper channels form four linear Ca²⁺ signaling domains along the flagella, which orchestrate capacitation-associated tyrosine phosphorylation [238]. The driving force for Ca²⁺ entry is principally determined by a mildly outwardly rectifying K⁺ channel (KSper) that, like CatSper, is activated by intracellular alkalization [1000]. Mouse KSper is encoded by *mSlo3*, a protein detected only in testis [915, 1000, 1598]. In human sperm, such alkalization may result from the activation of H_v1, a proton channel [837]. Mutations in CatSper are associated with syndromic and non-syndromic male infertility [540]. In human ejaculated spermatozoa, progesterone (<50 nM) potentiates the CatSper current by a non-genomic mechanism and acts synergistically with intracellular alkalisation [836, 1330]. Sperm cells from infertile patients with a deletion in

CatSper2 gene lack I_{CatSper} and the progesterone response [1290]. In addition, certain prostaglandins (e.g. PGF_{1 α} , PGE₁) also potentiate CatSper mediated currents [836, 1330]. In human sperm, CatSper channels are also activated by various small molecules including endocrine disrupting chemicals and proposed as a polymodal sensor [149, 149]. TPCs are the major Na⁺ conductance in lysosomes; knocking out TPC1 and TPC2 eliminates the Na⁺ conductance and renders the organelle's membrane potential insensitive to changes in [Na⁺] (31). The channels are regulated by luminal pH [174], PI(3,5)P2 [1484], intracellular ATP and extracellular amino acids [175]. TPCs are also involved in the NAADP-activated Ca²⁺ release from lysosomal Ca²⁺ stores [170, 972]. Mice lacking TPCs are viable but have phenotypes including compromised lysosomal pH stability, reduced physical endurance [175], resistance to Ebola viral infection [1215] and fatty liver [476]. No major human disease-associated TPC mutation has been reported.

Further reading on CatSper and Two-Pore channels (TPC)

Clapham DE *et al.* (2005) International Union of Pharmacology. L. Nomenclature and structure-function relationships of CatSper and two-pore channels. *Pharmacol Rev* **57**: 451-4 [PMID:16382101]

Grimm C *et al.* (2017) Two-Pore Channels: Catalyzers of Endolysosomal Transport and Function. *Front Pharmacol* **8**: 45 [PMID:28223936]

Kintzer AF *et al.* (2018) On the structure and mechanism of two-pore channels. *FEBS J* **285**: 233-243 [PMID:28656706]

Cyclic nucleotide-regulated channels (CNG)

Ion channels → Voltage-gated ion channels → Cyclic nucleotide-regulated channels (CNG)

Overview: Cyclic nucleotide-gated (CNG) channels are responsible for signalling in the primary sensory cells of the vertebrate visual and olfactory systems. CNG channels are voltage-independent cation channels formed as tetramers. Each subunit has 6TM, with the pore-forming domain between TM5 and TM6. CNG channels were first found in rod photoreceptors [400, 685], where light signals through rhodopsin and transducin to stimulate phosphodiesterase and reduce intracellular cyclic GMP level. This results in a closure of CNG channels and a reduced 'dark current'. Similar channels were found in the cilia of olfactory neurons [994] and the pineal gland [342]. The cyclic nucleotides bind

to a domain in the C terminus of the subunit protein: other channels directly binding cyclic nucleotides include hyperpolarisation-activated, cyclic nucleotide-gated channels (HCN), ether-a-go-go and certain plant potassium channels.

The **HCN channels** are cation channels that are activated by hyperpolarisation at voltages negative to ~-50 mV. The cyclic nucleotides cyclic AMP and cyclic GMP directly bind to the cyclic nucleotide-binding domain of HCN channels and shift their activation curves to more positive voltages, thereby enhancing channel activity. HCN channels underlie pacemaker currents found in many excitable cells including cardiac cells and neu-

rons [322, 1074]. In native cells, these currents have a variety of names, such as I_h, I_q and I_r. The four known HCN channels have six transmembrane domains and form tetramers. It is believed that the channels can form heteromers with each other, as has been shown for HCN1 and HCN4 [35]. High resolution structural studies of CNG and HCN channels has provided insight into the gating processes of these channels [778, 779, 808]. **A standardised nomenclature for CNG and HCN channels has been proposed by the NC-IUPHAR Subcommittee on voltage-gated ion channels [556].**

Nomenclature	CNGA1	CNGA2	CNGA3	CNGB1	CNGB3
HGNC, UniProt	CNGA1, P29973	CNGA2, Q16280	CNGA3, Q16281	CNGB1, Q14028	CNGB3, Q9N-QW8
Activators	cyclic GMP (EC ₅₀ ~ 30 μM) ≫ cyclic AMP	cyclic GMP > cyclic AMP (EC ₅₀ ~ 1 μM)	cyclic GMP (EC ₅₀ ~ 30 μM) ≫ cyclic AMP	–	–
Channel blockers	dequalinium (Antagonist) (pIC ₅₀ 6.7) [0mV] [1189], L-(cis)-diltiazem (Antagonist) (pK _i 4) [-80mV – 80mV] [213]	dequalinium (Antagonist) (pIC ₅₀ 5.6) [0mV] [1188]	L-(cis)-diltiazem (high affinity binding requires presence of CNGB subunits)	–	L-(cis)-diltiazem (Antagonist) (pIC ₅₀ 5.5) [0mV] [443] – Mouse
Functional Characteristics	γ = 25-30 pS P _{Ca} /P _{Na} = 3.1	γ = 35 pS P _{Ca} /P _{Na} = 6.8	γ = 40 pS P _{Ca} /P _{Na} = 10.9	–	–
Comments	–	–	–	L-(cis)-diltiazem acts as a channel blocker when CNGB1 is co-expressed with CNGA1.	–

Nomenclature	HCN1	HCN2	HCN3	HCN4
HGNC, UniProt	HCN1, O60741	HCN2, Q9UL51	HCN3, Q9P1Z3	HCN4, Q9Y3Q4
Activators	cyclic AMP > cyclic GMP (both weak)	cyclic AMP > cyclic GMP	–	cyclic AMP > cyclic GMP
Channel blockers	MEL57A (pEC ₅₀ 6.5) [-80mV] [303] – Mouse, ivabradine (Antagonist) (pIC ₅₀ 5.7) [1314], ZD7288 (Antagonist) (pIC ₅₀ 4.7) [1313], EC18 (pEC ₅₀ 4.7) [303] – Mouse, Cs ⁺ (Antagonist) (pIC ₅₀ 3.7) [-40mV] [1313]	ivabradine (Antagonist) (pIC ₅₀ 5.6) [1314] – Mouse, clonidine (Antagonist) (pIC ₅₀ 5.1) [-40mV] [721] – Mouse, MEL57A (pEC ₅₀ 4.9) [303] – Mouse, EC18 (pEC ₅₀ 4.7) [303] – Mouse, ZD7288 (Antagonist) (pIC ₅₀ 4.4) [1313], Cs ⁺ (Antagonist) (pIC ₅₀ 3.7) [-40mV] [1313]	ivabradine (Antagonist) (pIC ₅₀ 5.7) [1314], ZD7288 (Antagonist) (pIC ₅₀ 4.5) [1313], Cs ⁺ (Antagonist) (pIC ₅₀ 3.8) [-40mV] [1313]	ivabradine (Antagonist) (pIC ₅₀ 5.7) [1314], EC18 (pEC ₅₀ 5.4) [-80mV] [303], clonidine (Antagonist) (pIC ₅₀ 5) [-40mV] [721] – Mouse, ZD7288 (Antagonist) (pIC ₅₀ 4.7) [1313], MEL57A (pEC ₅₀ 4.1) [303], Cs ⁺ (Antagonist) (pIC ₅₀ 3.8) [-40mV] [1313]

Comments: CNGA1, CNGA2 and CNGA3 express functional channels as homomers. Three additional subunits *CNGA4* (Q8IV77), *CNGB1* (Q14028) and *CNGB3* (Q9NQW8) do not, and are referred to as auxiliary subunits. The subunit composition of the native channels is believed to be as follows. Rod: CNGA1₃/

CNGB1a [87]; Cone: CNGA3₃/CNGB3₁ [1618] ; Olfactory neurons: CNGA2₂/CNGA4/CNGB1b [1100, 1505, 1615, 1616, 1622]. HCN channels are permeable to both Na⁺ and K⁺ ions, with a Na⁺/K⁺ permeability ratio of about 0.2. Functionally, they differ from each other in terms of time constant of activation with

HCN1 the fastest, HCN4 the slowest and HCN2 and HCN3 intermediate. The compounds ZD7288 [140] and ivabradine [162] have proven useful in identifying and studying functional HCN channels in native cells. Zatebradine and clobradine are also useful blocking agents.

Further reading on Cyclic nucleotide-regulated channels (CNG)

Barret DCA *et al.* (2022) The structure of cyclic nucleotide-gated channels in rod and cone photoreceptors. *Trends Neurosci* **45**: 763-776 [PMID:35934530]
 Gerhardt MJ *et al.* (2023) Biology, Pathobiology and Gene Therapy of CNG Channel-Related Retinopathies. *Biomedicines* **11**: [PMID:36830806]
 Hennis K *et al.* (2022) Paradigm shift: new concepts for HCN4 function in cardiac pacemaking. *Pflugers Arch* **474**: 649-663 [PMID:35556164]
 Napolitano LMR *et al.* (2021) CNG channel structure, function, and gating: a tale of conformational flexibility. *Pflugers Arch* **473**: 1423-1435 [PMID:34357442]

Santoro B *et al.* (2020) Hyperpolarization-Activated Cyclic Nucleotide-Gated Channels as Drug Targets for Neurological Disorders. *Annu Rev Pharmacol Toxicol* **60**: 109-131 [PMID:31914897]
 Sartiani L *et al.* (2017) The Hyperpolarization-Activated Cyclic Nucleotide-Gated Channels: from Biophysics to Pharmacology of a Unique Family of Ion Channels. *Pharmacol Rev* **69**: 354-395 [PMID:28878030]

Potassium channels

Ion channels → Voltage-gated ion channels → Potassium channels

Overview: Activation of potassium channels regulates excitability and can control the shape of the action potential waveform. They are present in all cells within the body and can influence processes as diverse as cognition, muscle contraction and hormone secretion. Potassium channels are subdivided into families,

based on their structural and functional properties. The largest family consists of potassium channels that activated by membrane depolarization, with other families consisting of channels that are either activated by a rise of intracellular calcium ions or are constitutively active. A standardised nomenclature for potas-

sium channels has been proposed by the **NC-IUPHAR Subcommittee** on potassium channels [460, 494, 754, 1503], which has placed cloned channels into groups based on gene family and structure of channels that exhibit 6, 4 or 2 transmembrane domains (TM).

Calcium- and sodium-activated potassium channels (K_{Ca} , K_{Na})

Ion channels → Voltage-gated ion channels → Potassium channels → Calcium- and sodium-activated potassium channels (K_{Ca} , K_{Na})

Overview: Calcium- and sodium- activated potassium channels are members of the 6TM family of K channels which comprises the voltage-gated K_V subfamilies, including the KCNQ subfamily, the EAG subfamily (which includes hERG channels), the Ca^{2+} -activated Slo subfamily (actually with 6 or 7TM) and the Ca^{2+} - and Na^+ -activated SK subfamily (**nomenclature as agreed by the NC-IUPHAR Subcommittee on Calcium- and sodium-activated potassium channels** [658]). As for the 2TM family, the pore-forming a subunits form tetramers and heteromeric channels may be formed within subfamilies (e.g. $K_V1.1$ with $K_V1.2$; KCNQ2 with KCNQ3).

Nomenclature	$K_{Ca}1.1$
HGNC, UniProt	<i>KCNMA1</i> , Q12791
Activators	dehydrosoyasaponin I (pEC ₅₀ ~7) [453], 16,17-EpDPE (pEC ₅₀ 6.6) [1479], GoSlo-SR-5-69 (pEC ₅₀ 6.6) [1193], flindokalner (pEC ₅₀ 6.4–6.5) [629], magnolol (pEC ₅₀ ~6) [1530], BC5 (pEC ₅₀ 5.6) [1604], BMS-191011 (pEC ₅₀ ~5.6) [1186], NS1619 (pEC ₅₀ 5.4) [784, 1049], NS004 (pEC ₅₀ 4.5–5), NS11021 (pEC ₅₀ 4.7–5) [109], NS19504 (pEC ₅₀ 5) [999], mefenamic acid (pEC ₅₀ 4) [1372]
Channel blockers	BmTx2 (pIC ₅₀ 9.5) [124, 1185], BmTx1 (pIC ₅₀ 9.2) [1185], slotoxin (blocks channels with $\beta 1$ auxiliary subunit) (pIC ₅₀ 8.8) [432], paxilline (pK _i 8.7) [0mV] [1221] – Mouse, Penitrem A (pIC ₅₀ 8.2) [52], kaliotoxin (pIC ₅₀ 7.7) [259], BmP09 (pIC ₅₀ 7.6) [1566], natrin (pIC ₅₀ 7.5) [1474], ChTX-Lq2 (pIC ₅₀ 7.4) [866], Lqh Tx 15-1 (pIC ₅₀ 7.3) [913], butantoxin (pIC ₅₀ 7.3) [1035], martentoxin (pIC ₅₀ 7.1) [1369], verruculogen (pIC ₅₀ ~7) [968], charybdotoxin (pIC ₅₀ 6.6) [428], iberiotoxin (pIC ₅₀ 6.6) [422], clotrimazole (pIC ₅₀ 5.5) [1531], tetrandrine (pIC ₅₀ 5) [1466]
Functional Characteristics	Maxi K_{Ca}

Nomenclature	$K_{Ca}2.1$	$K_{Ca}2.2$	$K_{Ca}2.3$
HGNC, UniProt	<i>KCNN1</i> , Q92952	<i>KCNN2</i> , Q9H2S1	<i>KCNN3</i> , Q9UGI6
Activators	EBIO (Agonist) Concentration range: $2 \times 10^{-3}M$ [-80mV] [1092, 1500], NS309 (Agonist) Concentration range: $3 \times 10^{-8}M$ - $1 \times 10^{-7}M$ [-90mV] [1327, 1500]	NS309 (Agonist) (pEC ₅₀ 6.2) Concentration range: $3 \times 10^{-8}M$ - $1 \times 10^{-7}M$ [1091, 1327, 1500], NS13001 (Agonist) (pEC ₅₀ 5.7) [682] – Mouse, rimituzalcap (pIC ₅₀ ~5.3) [1609], CyPPA (Agonist) (pEC ₅₀ 4.9) [571], EBIO (Agonist) (pEC ₅₀ 3.3) [1091, 1500]	NS13001 (Agonist) (pEC ₅₀ 6.8) [682] – Mouse, CyPPA (Agonist) (pEC ₅₀ 5.3) [571], EBIO (Agonist) (pEC ₅₀ 3.8) [1500, 1519], NS309 (Agonist) Concentration range: $3 \times 10^{-8}M$ [1327, 1500]

Inhibitors	UCL1684 (pIC ₅₀ 9.1) [1326, 1500], apamin (pIC ₅₀ 8.4) [760, 1255, 1316, 1326], CM-TPMF (pEC ₅₀ 7.6) [572]	UCL1684 (pIC ₅₀ 10.1) [386, 996, 1500], apamin (pK _d 9.4) [619], Lei-Dab7 (pIC ₅₀ 8.4) [1256], AP14145 (pIC ₅₀ 6) [323, 1279], AP30663 (pIC ₅₀ 5.6) [562]	UCL1684 (pIC ₅₀ 8–9) [386, 1500]
Channel blockers	NS8593 (pIC ₅₀ 6.4) [0mV] [1325], AP30663 (pIC ₅₀ 6) [108, 562], tetraethylammonium (pIC ₅₀ 2.7) [1500]	NS8593 (pIC ₅₀ 6.4) [1325], AP30663 (pIC ₅₀ 5.6) [108, 562], tetraethylammonium (pIC ₅₀ 2.7) [1500]	apamin (Antagonist) (pIC ₅₀ 8.6) [78, 570, 760] – Rat, AP14145 (pIC ₅₀ 6) [1279], AP30663 (pIC ₅₀ 6) [108, 562], tetraethylammonium (pIC ₅₀ 2.7) [1500]
Functional Characteristics	SK _{Ca}	SK _{Ca}	SK _{Ca}
Comments	The rat isoform does not form functional channels when expressed alone in cell lines. N- or C-terminal chimeric constructs permit functional channels that are insensitive to apamin [1500]. Heteromeric channels are formed between K _{Ca} 2.1 and 2.2 subunits that show intermediate sensitivity to apamin [242].	Cryo-EM-derived structures of human K _{Ca} 2.2-K _{Ca} 3.1 chimeric channels have been deposited to the Protein Data Bank (PDB) with IDs 9O48 (Ca ₂₊ bound state), 9O51 (Ca ₂₊ free state), 9O53 (inhibitor bound state), 9O50 (activator bound state) and 9O52 (apamin bound state).	–

Nomenclature	K _{Ca} 3.1	K _{Na} 1.1	K _{Na} 1.2	K _{Ca} 5.1
HGNC, UniProt	KCNN4, O15554	KCNT1, Q5JUK3	KCNT2, Q6UVM3	KCNUI, A8MYU2
Activators	NS309 (Agonist) (pEC ₅₀ 8) [-90mV] [1327, 1500], SKA-121 (Agonist) (pEC ₅₀ 7) [250], SKA-111 (Agonist) (pEC ₅₀ 6.9) [250], EBIO (Agonist) (pEC ₅₀ 4.1–4.5) [-100mV – -50mV] [1092, 1350, 1500]	bithionol (Agonist) (pEC ₅₀ 5–6) [1557] – Rat, niclosamide (Agonist) (pEC ₅₀ 5.5) [122], loxapine (Agonist) (pEC ₅₀ 5.4) [122]	niflumic acid (Agonist) (pEC ₅₀ 8.7) [271, 433]	–
Gating inhibitors	–	bepiridil (pIC ₅₀ 5–6) [1557] – Rat	–	–
Channel blockers	maurotoxin (Inhibition) (pIC ₅₀ 9) [184], charybdotoxin (pIC ₅₀ 7.6–8.7) [630, 649]	PRX-2904 (pIC ₅₀ 7.4) [474], CPK20 (pIC ₅₀ 6.7) [601], VU0948578 (pIC ₅₀ 6.2) [1155], VU0948578 (pIC ₅₀ 6.2) [1155], quinidine (pIC ₅₀ 4) [115, 1557] – Rat	CPK20 (pIC ₅₀ 5.9) [601], compound 43 (pIC ₅₀ 5.6) [1154], Ba ²⁺ (Inhibition) (pIC ₅₀ 3) [115], quinidine (Inhibition) Concentration range: 1 × 10 ⁻³ M [115] – Rat	VU0546110 (pIC ₅₀ 5.9) [875], quinidine Concentration range: 2 × 10 ⁻⁵ M [1367, 1526] – Mouse
Selective channel blockers	DHP-103 (pIC ₅₀ 8.2) [1056], NS6180 (Inhibition) (pIC ₅₀ 8.1) [1324], TRAM-34 (Inhibition) (pK _d 7.6–8) [1532], senicapoc (Inhibition) (pIC ₅₀ 8) [1315]	–	–	–
Functional Characteristics	IK _{Ca} ; K _{Ca} 3.1 may form heteromers with K _{Ca} 2.1 [539].	K _{Na}	K _{Na}	Sperm pH-regulated K ⁺ current, K _{SPER}
Comments	Cryo-EM-derived structures of human K _{Ca} 2.2-K _{Ca} 3.1 chimeric channels have been deposited to the Protein Data Bank (PDB) with IDs 9O48 (Ca ₂₊ bound state), 9O51 (Ca ₂₊ free state), 9O53 (inhibitor bound state), 9O50 (activator bound state) and 9O52 (apamin bound state).	–	–	–

Further reading on Calcium- and sodium-activated potassium channels (K_{Ca} , K_{Na})Brown BM *et al.* (2020) Pharmacology of Small- and Intermediate-Conductance Calcium-Activated Potassium Channels. *Annu Rev Pharmacol Toxicol* **60**: 219-240 [PMID:31337271]Dopico AM *et al.* (2018) Calcium- and voltage-gated BK channels in vascular smooth muscle. *Pflugers Arch* **470**: 1271-1289 [PMID:29748711]Kaczmarek LK *et al.* (2017) International Union of Basic and Clinical Pharmacology. C. Nomenclature and Properties of Calcium-Activated and Sodium-Activated Potassium Channels. *Pharmacol Rev* **69**: 1-11 [PMID:28267675]Kshatri AS *et al.* (2018) Physiological Roles and Therapeutic Potential of Ca^{2+} Activated Potassium Channels in the Nervous System. *Front Mol Neurosci* **11**: 258 [PMID:30104956]

Inwardly rectifying potassium channels (K_{IR})

Ion channels → Voltage-gated ion channels → Potassium channels → Inwardly rectifying potassium channels (K_{IR})**Overview:** The 2TM domain family of K channels are also known as the inward-rectifier K channel family. This family includes the strong inward-rectifier K channels (K_{IR} 2.x) that are constitutively active, the G protein-activated inward-rectifier K channels (K_{IR} 3.x) and the ATP-sensitive K channels (K_{IR} 6.x, which combine with sulphonylurea receptors (SUR1-3)). The pore-forming α subunits form tetramers, and heteromeric channels may be formed within subfamilies (e.g. K_{IR} 3.2 with K_{IR} 3.3).

Nomenclature	K_{IR} 1.1
HGNC, UniProt	KCNJ1 , P48048
Ion Selectivity and Conductance	NH_4^+ [62pS] > K^+ [38. pS] > Tl^+ [21pS] > Rb^+ [15pS] (Rat) [229, 551]
Channel blockers	tertiapin-Q (Inhibition) (pIC_{50} 8.9) [641], Ba^{2+} (Antagonist) (pIC_{50} 2.3–4.2) Concentration range: $1 \times 10^{-4}M$ [voltage dependent 0mV – -100mV] [551, 1623] – Rat, Cs^+ (Antagonist) (pIC_{50} 2.9) [voltage dependent -120mV] [1623] – Rat
Functional Characteristics	K_{IR} 1.1 is weakly inwardly rectifying, as compared to classical (strong) inward rectifiers.

	K_{IR} 2.1	K_{IR} 2.2	K_{IR} 2.3	K_{IR} 2.4
Nomenclature	K_{IR} 2.1	K_{IR} 2.2	K_{IR} 2.3	K_{IR} 2.4
HGNC, UniProt	KCNJ2 , P63252	KCNJ12 , Q14500	KCNJ4 , P48050	KCNJ14 , Q9UNX9
Endogenous activators	PIP_2 (Agonist) Concentration range: $1 \times 10^{-5}M$ – $5 \times 10^{-5}M$ [-30mV] [581, 1167, 1303] – Mouse	–	–	–
Endogenous inhibitors	–	Intracellular Mg^{2+} (pIC_{50} 5) [40mV] [1554]	–	Intracellular Mg^{2+}
Gating inhibitors	–	Ba^{2+} (Antagonist) Concentration range: $5 \times 10^{-5}M$ [-150mV – -50mV] [1354] – Mouse, Cs^+ (Antagonist) Concentration range: $5 \times 10^{-6}M$ – $5 \times 10^{-5}M$ [-150mV – -50mV] [1354] – Mouse	–	–

Endogenous channel blockers	spermine (Antagonist) (pK_d 9.1) [<i>voltage dependent</i> 40mV] [609, 1559] – Mouse, spermidine (Antagonist) (pK_d 8.1) [<i>voltage dependent</i> 40mV] [1559] – Mouse, putrescine (Antagonist) (pK_d 5.1) [<i>voltage dependent</i> 40mV] [609, 1559] – Mouse, Intracellular Mg²⁺ (Antagonist) (pK_d 4.8) [<i>voltage dependent</i> 40mV] [1559] – Mouse	–	Intracellular Mg²⁺ (Antagonist) (pK_d 5) [<i>voltage dependent</i> 50mV] [854], putrescine (Antagonist) Concentration range: $5 \times 10^{-5}M$ – $1 \times 10^{-3}M$ [-80mV – 80mV] [854], spermidine (Antagonist) Concentration range: $2.5 \times 10^{-5}M$ – $1 \times 10^{-3}M$ [-80mV – 80mV] [854], spermine (Antagonist) Concentration range: $5 \times 10^{-5}M$ – $1 \times 10^{-3}M$ [-80mV – 80mV] [854]	–
Channel blockers	Ba²⁺ (Antagonist) (pK_d 3.9–5.6) Concentration range: $1 \times 10^{-6}M$ – $1 \times 10^{-4}M$ [<i>voltage dependent</i> 0mV – -80mV] [25] – Mouse, Cs⁺ (Antagonist) (pK_d 1.3–4) Concentration range: $3 \times 10^{-5}M$ – $3 \times 10^{-4}M$ [<i>voltage dependent</i> 0mV – -102mV] [8] – Mouse	–	Ba²⁺ (Antagonist) (pIC_{50} 5) Concentration range: $3 \times 10^{-6}M$ – $5 \times 10^{-4}M$ [-60mV] [898, 1136, 1368], Cs⁺ (Antagonist) (pK_i 1.3–4.5) Concentration range: $3 \times 10^{-6}M$ – $3 \times 10^{-4}M$ [0mV – -130mV] [898]	Cs⁺ (Antagonist) (pK_d 3–4.1) [<i>voltage dependent</i> -60mV – -100mV] [590], Ba²⁺ (Antagonist) (pK_d 3.3) [<i>voltage dependent</i> 0mV] [590]
Functional Characteristics	IK ₁ in heart, ‘strong’ inward-rectifier current	IK ₁ in heart, ‘strong’ inward-rectifier current	IK ₁ in heart, ‘strong’ inward-rectifier current	IK ₁ in heart, ‘strong’ inward-rectifier current
Comments	K _{ir} 2.1 is also inhibited by intracellular polyamines	K _{ir} 2.2 is also inhibited by intracellular polyamines	K _{ir} 2.3 is also inhibited by intracellular polyamines	K _{ir} 2.4 is also inhibited by intracellular polyamines

Nomenclature	K_{ir} 3.1	K_{ir} 3.2	K_{ir} 3.3	K_{ir} 3.4
HGNC, UniProt	KCNJ3, P48549	KCNJ6, P48051	KCNJ9, Q92806	KCNJ5, P48544
Endogenous activators	PIP₂ (Agonist) (pK_d 6.3) Concentration range: $5 \times 10^{-5}M$ [physiological voltage] [581]	PIP₂ (Agonist) (pK_d 6.3) Concentration range: $5 \times 10^{-5}M$ [physiological voltage] [581]	PIP₂ [542]	PIP₂ [90, 542]
Gating inhibitors	–	pimozide (Antagonist) (pEC_{50} 5.5) [-70mV] [724] – Mouse	–	–
Channel blockers	tertiapin-Q (Antagonist) (pIC_{50} 7.9) [640], Ba²⁺ (Antagonist) (pIC_{50} 4.7) [281] – Rat	desipramine (Antagonist) (pIC_{50} 4.4) [-70mV] [725] – Mouse	–	tertiapin-Q (Antagonist) (pIC_{50} 7.9) [640]
Functional Characteristics	G protein-activated inward-rectifier current	G protein-activated inward-rectifier current	G protein-activated inward-rectifier current	G protein-activated inward-rectifier current
Comments	K _{ir} 3.1 is also activated by G _{βγ} . K _{ir} 3.1 is not functional alone. The functional expression of K _{ir} 3.1 in <i>Xenopus oocytes</i> requires coassembly with the endogenous <i>Xenopus</i> K _{ir} 3.5 subunit. The major functional assembly in the heart is the K _{ir} 3.1/3.4 heteromultimer, while in the brain it is K _{ir} 3.1/3.2, K _{ir} 3.1/3.3 and K _{ir} 3.2/3.3.	K _{ir} 3.2 is also activated by G _{βγ} . K _{ir} 3.2 forms functional heteromers with K _{ir} 3.1/3.3.	K _{ir} 3.3 is also activated by G _{βγ}	K _{ir} 3.4 is also activated by G _{βγ}

Nomenclature	K_{ir} 4.1	K_{ir} 4.2	K_{ir} 5.1
HGNC, UniProt	KCNJ10 , P78508	KCNJ15 , Q99712	KCNJ16 , Q9NPI9
Channel blockers	Ba^{2+} (Antagonist) Concentration range: $3 \times 10^{-6}M$ - $1 \times 10^{-3}M$ [-160mV – 60mV] [731 , 1357 , 1364] – Rat, Cs^{+} (Antagonist) Concentration range: $3 \times 10^{-5}M$ - $3 \times 10^{-4}M$ [-160mV – 50mV] [1357] – Rat	Ba^{2+} (Antagonist) Concentration range: $1 \times 10^{-5}M$ - $1 \times 10^{-4}M$ [-120mV – 100mV] [1090] – Mouse, Cs^{+} (Antagonist) Concentration range: $1 \times 10^{-5}M$ - $1 \times 10^{-4}M$ [-120mV – 100mV] [1090] – Mouse	Ba^{2+} (Antagonist) Concentration range: $3 \times 10^{-3}M$ [-120mV – 20mV] [1363] – Rat
Functional Characteristics	Inward-rectifier current	Inward-rectifier current	Weakly inwardly rectifying

Nomenclature	K_{ir} 6.1	K_{ir} 6.2	K_{ir} 7.1
HGNC, UniProt	KCNJ8 , Q15842	KCNJ11 , Q14654	KCNJ13 , O60928
Associated subunits	SUR1, SUR2A, SUR2B	SUR1, SUR2A, SUR2B	–
Activators	cromakalim , diazoxide (Agonist) Concentration range: $2 \times 10^{-4}M$ [-60mV] [1552] – Mouse, minoxidil , nicorandil (Agonist) Concentration range: $3 \times 10^{-4}M$ [-60mV – 60mV] [1552] – Mouse	diazoxide (Agonist) (pEC_{50} 4.2) [physiological voltage] [597] – Mouse, cromakalim (Agonist) Concentration range: $3 \times 10^{-5}M$ [-60mV] [598] – Mouse, minoxidil , nicorandil	–
Inhibitors	glibenclamide , tolbutamide	glibenclamide , tolbutamide	–
Channel blockers	–	–	Ba^{2+} (Antagonist) (pK_i 3.2) [voltage dependent -100mV] [339 , 742 , 759 , 1082], Cs^{+} (Antagonist) (pK_i 1.6) [voltage dependent -100mV] [339 , 742 , 1082]
Functional Characteristics	ATP-sensitive, inward-rectifier current	ATP-sensitive, inward-rectifier current	Inward-rectifier current

Two-pore domain potassium channels (K_{2P})

Ion channels → Voltage-gated ion channels → Potassium channels → Two-pore domain potassium channels (K_{2P})

Overview: The 4TM family of K channels mediate many of the background potassium currents observed in native cells. They are open across the physiological voltage-range and are regulated by a wide array of neurotransmitters and biochemical mediators. The pore-forming α -subunit contains two pore loop (P) domains and two subunits assemble to form one ion conduction pathway

lined by four P domains. It is important to note that single channels do not have two pores but that each subunit has two P domains in its primary sequence; hence the name two-pore domain, or K_{2P} channels (and not two-pore channels). Some of the K_{2P} subunits can form heterodimers across subfamilies (*e.g.* $K_{2P}3.1$ with $K_{2P}9.1$). The nomenclature of 4TM K channels in the liter-

ature is still a mixture of IUPHAR and common names. The suggested division into subfamilies, described in the [More detailed introduction online](#), is based on similarities in both structural and functional properties within subfamilies and this explains the “common abbreviation” nomenclature in the tables below.

Nomenclature	K_{2p}1.1	K_{2p}2.1	K_{2p}3.1	K_{2p}4.1
Common abbreviation	TWIK1	TREK1	TASK1	TRAAK1
HGNC, UniProt	KCNK1, O00180	KCNK2, O95069	KCNK3, O14649	KCNK4, Q9NYG8
Endogenous activators	–	arachidonic acid (studied at 1-10 µM) (pEC ₅₀ 5) [1085]	–	arachidonic acid (studied at 1-10 µM) [402]
Activators	–	GI-530159 (pEC ₅₀ 6.1) [859], ONO-2920632 (pEC ₅₀ 5.6) [1567], BL-1249 (pEC ₅₀ 5.3) [1134], LPS2336 (pEC ₅₀ 4.9) [146], chloroform (studied at 1-5 mM) Concentration range: 8 × 10 ⁻³ M [1084], halothane (studied at 1-5 mM) [1084], isoflurane (studied at 1-5 mM) [1084]	halothane (studied at 1-10 mM) [774]	riluzole (studied at 1-100 µM) [352]
Inhibitors	magnolol (pIC ₅₀ 5.2) [1473]	ONO-TR-772 (pIC ₅₀ 7.8) [1362] – Rat, Cpd8l (pIC ₅₀ 6.1) [842] – Rat, norfluoxetine (pIC ₅₀ 5.1) [694]	KU124 (pIC ₅₀ 5.8) [350]	–
Channel blockers	–	–	R-(+)-methanandamide (pIC ₅₀ ~6.2) [894], anandamide (pIC ₅₀ ~6.2) [894]	–
Functional Characteristics	Background current	Background current	Background current	Background current
Comments	K _{2p} 1.1 is inhibited by acid pH _o external acidification with a pK _a ~6.7 [1127]. K _{2p} 1 forms heterodimers with K _{2p} 3 and K _{2p} 9 [1128].	K _{2p} 2.1 is also activated by membrane stretch, heat and acid pH _i [893, 895]. K _{2p} 2 can heterodimerize with K _{2p} 4 [126] and K _{2p} 10 [802].	Knock-out of the <i>kcnk3</i> gene leads to a prolonged QT interval in mice [293] and disrupted development of the adrenal cortex [526]. K _{2p} 3.1 is inhibited by acid pH _o with a pK _a of 6.4 [855]. K _{2p} 3 forms heterodimers with K _{2p} 1 [1128] and K _{2p} 9 [269].	K _{2p} 4 is activated by membrane stretch [892], and increased temperature (~12 to 20-fold between 17 and 40°C [667]) and can heterodimerize with K _{2p} 2 [126].

Nomenclature	K_{2p}5.1	K_{2p}6.1	K_{2p}7.1	K_{2p}9.1	K_{2p}10.1
Common abbreviation	TASK2	TWIK2	–	TASK3	TREK2
HGNC, UniProt	KCNK5 , O95279	KCNK6 , Q9Y257	KCNK7 , Q9Y2U2	KCNK9 , Q9NPC2	KCNK10 , P57789
Endogenous activators	–	–	–	–	arachidonic acid (studied at 1-10 μ M) [798]
Activators	–	–	dapagliflozin (pEC ₅₀ 4) [876]	halothane (studied at 1-5 mM) [1359]	Nb-Activator-67 (pEC ₅₀ 7) [1156], ONO-2920632 (pEC ₅₀ 6.5) [1567], Nb-Activator-76 (pEC ₅₀ 6.4) [1156], GI-530159 [859], halothane (studied at 1-5 mM) [798]
Inhibitors	–	–	–	A-769662 (effective at 300 μ M) (pIC ₅₀ 3.5) [1212], R-(+)-methanandamide (studied at 1-10 μ M) [1158], anandamide (studied at 1-10 μ M) [1158], benzimidazole 991 (effective concentration range 3-30 μ M) [1212]	ONO-TR-772 (pIC ₅₀ 6.8) [1362], Nb-Inhibitor-61 (pIC ₅₀ 6.2) [1156]
Functional Characteristics	Background current	Unknown	Unknown	Background current	Background current
Comments	K _{2p} 5.1 is activated by alkaline pH _o [1175]. Knockout of the <i>kcnk5</i> gene in mice is associated with metabolic acidosis, hyponatremia and hypotension due to impaired bicarbonate handling in the kidney [1494], as well as deafness [190]. The T108P mutation is associated with Balkan Endemic Nephropathy in humans [1396].	–	–	K _{2p} 9.1 is also inhibited by acid pH _o with a pK _a of ~6 [1158]. Imprinting of the <i>KCNK9</i> gene is associated with Birk Barel syndrome [77]. K _{2p} 9 can form heterodimers with K _{2p} 1 [1128] or K _{2p} 3 [269].	K _{2p} 10.1 is also activated by membrane stretch [798] and can heterodimerize with K _{2p} 2 [802].

Nomenclature	K _{2p} 12.1	K _{2p} 13.1	K _{2p} 15.1	K _{2p} 16.1	K _{2p} 17.1	K _{2p} 18.1
Common abbreviation	THIK2	THIK1	TASK5	TALK1	TALK2	TRESK
HGNC, UniProt	KCNK12 , Q9HB15	KCNK13 , Q9HB14	KCNK15 , Q9H427	KCNK16 , Q96T55	KCNK17 , Q96T54	KCNK18 , Q7Z418
Endogenous inhibitors	–	–	–	–	–	arachidonic acid (studied at 10–50 µM) [1223]
Inhibitors	–	halothane (studied at ~5 mM) [127]	–	–	–	–
Functional Characteristics	Does not function as a homodimer [1157] but can form a functional heterodimer with K _{2p} 13 [127].	Background current	Unknown	Background current	Background current	Background current
Comments	–	Forms a heterodimer with K _{2p} 12 [127].	–	K _{2p} 16.1 current is increased by alkaline pH _o with a pK _a of 7.8 [668].	K _{2p} 17.1 current is increased by alkaline pH _o with a pK _a of 8.8 [668].	A frame-shift mutation (F139WfsX24) in the <i>KCNK18</i> gene, is associated with migraine with aura in humans [763]. Sodium butyrate is a TRESK channel activator [550].

Comments: The K_{2p}6, K_{2p}7.1, K_{2p}15.1 and K_{2p}12.1 subtypes, when expressed in isolation, are nonfunctional. All 4TM channels are insensitive to the classical potassium channel blockers tetraethylammonium and fampidine, but are blocked to varying degrees by Ba²⁺ ions.

Further reading on Two-pore domain potassium channels (K_{2p})

- Gada K *et al.* (2019) Two-pore domain potassium channels: emerging targets for novel analgesic drugs: IUPHAR Review 26. *Br J Pharmacol* **176**: 256–266 [PMID:30325008]
- Mathie A *et al.* (2021) Two-Pore Domain Potassium Channels as Drug Targets: Anesthesia and Beyond. *Annu Rev Pharmacol Toxicol* **61**: 401–420 [PMID:32679007]
- Jin X *et al.* (2020) Targeting Two-Pore Channels: Current Progress and Future Challenges. *Trends Pharmacol Sci* **41**: 582–594 [PMID:32679067]

Voltage-gated potassium channels (K_v)

[Ion channels](#) → [Voltage-gated ion channels](#) → [Potassium channels](#) → [Voltage-gated potassium channels \(K_v\)](#)

Overview: The 6TM family of K channels comprises the voltage-gated K_v subfamilies, the EAG subfamily (which includes hERG channels), the Ca²⁺-activated Slo subfamily (actually with 7TM, termed BK) and the Ca²⁺-activated SK subfamily. These channels possess a pore-forming α subunit that comprises tetramers of identical subunits (homomeric) or of different subunits (heteromeric). Heteromeric channels can only be formed within subfamilies (e.g.

K_v1.1 with K_v1.2; K_v7.2 with K_v7.3). The pharmacology largely reflects the subunit composition of the functional channel.

K_v7 channels K_v7.1–K_v7.5 (KCNQ1–5) K⁺ channels are voltage-gated K⁺ channels with major roles in neurons, muscle cells and epithelia where they underlie physiologically important K⁺ currents, such as the neuronal M-current and the cardiac IKs. Genetic deficiencies in all five KCNQ genes result in human excitability

disorders, including epilepsy, autism spectrum disorders, cardiac arrhythmias and deafness. Thanks to the recent knowledge of the structure and function of human KCNQ-encoded proteins, these channels are increasingly used as drug targets for treating diseases [1, 587, 651, 1434].

Nomenclature	K_v1.1	K_v1.2	K_v1.3	K_v1.4	K_v1.5	K_v1.6	K_v1.7
HGNC, UniProt	KCNA1, Q09470	KCNA2, P16389	KCNA3, P22001	KCNA4, P22459	KCNA5, P22460	KCNA6, P17658	KCNA7, Q96RP8
Associated subunits	K _v 1.2, K _v 1.4, K _v β1 and K _v β2 [251]	K _v 1.1, K _v 1.4, K _v β1 and K _v β2 [251]	K _v 1.1, K _v 1.2, K _v 1.4, K _v 1.6, K _v β1 and K _v β2 [251]	K _v 1.1, K _v 1.2, K _v β1 and K _v β2 [251]	K _v β1 and K _v β2	K _v β1 and K _v β2	K _v β1 and K _v β2
Channel blockers	α-dendrotoxin (pEC ₅₀ 7.7–9) [481, 592] – Rat, margatoxin (Inhibition) (pIC ₅₀ 8.4) [88], tetraethylammonium (Inhibition) (pK _d 3.5) [481] – Mouse	margatoxin (Inhibition) (pIC ₅₀ 11.2) [88], α-dendrotoxin (pIC ₅₀ 7.8–9.4) [481, 592] – Rat, noxiustoxin (pK _d 8.7) [481] – Rat, κM-conotoxin RIIIK (pIC ₅₀ 6.6) [0mV] [396]	margatoxin (pIC ₅₀ 10–10.3) [431, 445], noxiustoxin (pK _d 9) [481] – Mouse, maurotoxin (pIC ₅₀ 6.8) [1181], tetraethylammonium (pK _d 2) [481] – Mouse	fampridine (pIC ₅₀ 1.9) [1331] – Rat	fampridine (pIC ₅₀ 4.3) [390]	α-dendrotoxin (pIC ₅₀ 7.7) [485], tetraethylammonium (pIC ₅₀ 2.2) [485]	noxiustoxin (pIC ₅₀ 7.7) [664] – Mouse, fampridine (pIC ₅₀ 3.6) [664] – Mouse
Selective channel blockers	–	–	correolide (pIC ₅₀ 7.1) [392]	–	–	–	–
Functional Characteristics	K _v	K _v	K _v	K _A	K _v	K _v	K _v
Comments	–	–	Resistant to dendrotoxins	Resistant to dendrotoxins	Resistant to external TEA	–	–

Nomenclature	K_v1.8	K_v2.1	K_v2.2	K_v3.1	K_v3.2	K_v3.3	K_v3.4
HGNC, UniProt	KCNA10, Q16322	KCNB1, Q14721	KCNB2, Q92953	KCNC1, P48547	KCNC2, Q96PR1	KCNC3, Q14003	KCNC4, Q03721
Associated subunits	K _v β1 and K _v β2	K _v 5.1, K _v 6.1–6.4, K _v 8.1–8.2 and K _v 9.1–9.3	K _v 5.1, K _v 6.1–6.4, K _v 8.1–8.2 and K _v 9.1–9.3	–	–	–	MiRP2 is an associated subunit for K _v 3.4
Gating inhibitors	–	RY796 (pIC ₅₀ 6.6) [535], RY785 (pIC ₅₀ 5.9) [535], GxTx-1E (pIC ₅₀ 2) [536]	GxTx-1E (pK _d 8.6) [536], RY796 (pIC ₅₀ 7.1) [535], RY785 (pIC ₅₀ 6.7) [535]	–	–	–	–
Channel blockers	fampridine (pIC ₅₀ 2.8) [769]	tetraethylammonium (Pore blocker) (pIC ₅₀ 2) [524] – Rat	fampridine (pIC ₅₀ 2.8) [1236], tetraethylammonium (pIC ₅₀ 2.6) [1236]	fampridine (pIC ₅₀ 4.5) [481] – Mouse, tetraethylammonium (pIC ₅₀ 3.7) [481] – Mouse	fampridine (pIC ₅₀ 4.6) [823] – Rat, tetraethylammonium (pIC ₅₀ 4.2) [823] – Rat	tetraethylammonium (pIC ₅₀ 3.9) [1422] – Rat	tetraethylammonium (pIC ₅₀ 3.5) [1174, 1245] – Rat
Selective channel blockers	–	–	–	–	–	–	sea anemone toxin BDS-I (pIC ₅₀ 7.3) [329] – Rat
Functional Characteristics	K _v	K _v	–	K _v	K _v	K _A	K _A

Nomenclature	K _v 4.1	K _v 4.2	K _v 4.3
HGNC, UniProt	KCND1, Q9NSA2	KCND2, Q9NZV8	KCND3, Q9UK17
Associated subunits	KChIP 1-4, DPP6, DPP10	KChIP 1-4, DPP6, DPP10, K _v β1, NCS-1, Na _v β1	KChIP 1-4, DPP6 and DPP10, MinK, MiRPs
Channel blockers	fampridine (pIC ₅₀ 2) [604]	–	–
Functional Characteristics	K _A	K _A	K _A

Nomenclature	K _v 5.1	K _v 6.1	K _v 6.2	K _v 6.3	K _v 6.4
HGNC, UniProt	KCNF1, Q9H3M0	KCNG1, Q9UIX4	KCNG2, Q9UJ96	KCNG3, Q8TAE7	KCNG4, Q8TDN1

Nomenclature	K _v 7.1	K _v 7.2	K _v 7.3	K _v 7.4	K _v 7.5
HGNC, UniProt	KCNQ1, P51787	KCNQ2, O43526	KCNQ3, O43525	KCNQ4, P56696	KCNQ5, Q9NR82
Activators	ML277 (pEC ₅₀ 6.6) [924]	–	gabapentin (pEC ₅₀ 8.3) [909], retigabine (pEC ₅₀ 6.2) [1371]	retigabine (pEC ₅₀ 5.2) [1371]	retigabine (pEC ₅₀ 5) [353]
Selective activators	–	–	–	retigabine derivative 10g (pEC ₅₀ 6) [1477]	gabapentin (pEC ₅₀ 8.7) [909], retigabine derivative 10g (pEC ₅₀ 6) [1477]
Inhibitors	XE991 (pK _d 6.1) [1471], linopirdine (pIC ₅₀ 4.4) [1043] – Mouse	XE991 (pIC ₅₀ 6.2) [1472], linopirdine (pIC ₅₀ 5.3) [1472]	linopirdine (pIC ₅₀ 5.4) [1472] – Rat	XE991 (pIC ₅₀ 5.3) [1299], linopirdine (pIC ₅₀ 4.9) [1299]	linopirdine (pK _d 4.8) [795]
Sub/family-selective inhibitors	–	–	–	–	XE991 (pIC ₅₀ 4.2) [1244]
Channel blockers	JNJ303 (pIC ₅₀ 7.2) [1399]	tetraethylammonium (pIC ₅₀ 3.5–3.9) [499, 1510]	–	tetraethylammonium (pIC ₅₀ 1.3) [70]	–
Allosteric modulators	–	retigabine (Activation) (pEC ₅₀ 5.6) [814, 1371]	–	–	–
Functional Characteristics	cardiac IK _S	M current as a heteromer between K _v 7.2 and K _v 7.3	M current as heteromeric K _v 7.2/K _v 7.3 or K _v 7.3/K _v 7.5	–	M current as heteromeric K _v 7.3/K _v 7.5

Comments	Polyunsaturated fatty acids (PUFA) activate K _v 7.1 (KCNQ1) [772]. The PUFA analogue DHA-glycine is a selective activator of IKS with pEC ₅₀ 5.2 [133]. A single binding site for ML277 was identified, localized to a pocket lined by the S4-S5 linker, S5 and S6 helices of two separate subunits [878, 1512]. 3D structures have revealed the structural basis of hKCNQ1 modulation and gating [1336, 1337]. ML277 induces an upward movement of the S4-S5 linker and opening of the activation gate without affecting the C-terminal domain [878].	The cryo-EM structure of K _v 7.2 channel in its apo state and in complex with two activators, ztz240 or retigabine. While ztz240 binds to the voltage sensor domain, retigabine interacts with the pore domain [814]. Other K _v 7.2 cryo-EM structures have been determined in complex with additional activators such as Ebio1, Ebio2, cannabidiol (CBD), PIP ₂ and HN37 revealing the binding of two CBD, one PIP ₂ , and two HN37 molecules in each K _v 7.2 subunit [806, 877, 1610] and with a blocker Ebio3 [806].	Gentisic acid (GA) is found to be selective for K _v 7.3 versus other K _v 7 homomers; it requires S5 residue K _v 7.3-W265 for K _v 7.3 channel activation [2].	Two PIP ₂ molecules are identified in the open-state structure of K _v 7.4 (KCNQ4), which act as a bridge to couple the voltage-sensing domain (VSD) and pore domain (PD) of KCNQ4 leading to the channel opening [812, 1619]. In K _v 7.4, retigabine nestles in each fenestration, whereas linopirdine resides in a cytosolic cavity underneath the channel's inner gate [812].	The cryo-EM structure of human KCNQ5-calmodulin (CaM) complex in the apo, PIP ₂ -bound, and both PIP ₂ - and the activator HN37-bound states in either a closed or an open conformation was solved [1565].
----------	--	--	--	--	--

Nomenclature	K _v 8.1	K _v 8.2	K _v 9.1	K _v 9.2	K _v 9.3	K _v 10.1	K _v 10.2
HGNC, UniProt	KCNV1, Q6PIU1	KCNV2, Q8TDN2	KCNS1, Q96KK3	KCNS2, Q9ULS6	KCNS3, Q9BQ31	KCNH1, O95259	KCNH5, Q8NCM2

Nomenclature	K _v 11.1	K _v 11.2	K _v 11.3	K _v 12.1	K _v 12.2	K _v 12.3
HGNC, UniProt	KCNH2, Q12809	KCNH6, Q9H252	KCNH7, Q9NS40	KCNH8, Q96L42	KCNH3, Q9ULD8	KCNH4, Q9UQ05
Associated subunits	minK (KCNE1) and MiRP1 (KCNE2)	minK (KCNE1)	minK (KCNE1)	minK (KCNE1)	minK (KCNE1) and MiRP2 (KCNE3)	–
Channel blockers	astemizole (pIC ₅₀ 9) [1629], terfenadine (pIC ₅₀ 7.3) [1162], disopyramide (Inhibition) (pIC ₅₀ 4) [695]	–	–	–	–	–
Sub/family-selective channel blockers	E4031 (pIC ₅₀ 8.1) [1628]	–	–	–	–	–
Selective channel blockers	dofetilide (Inhibition) (pK _i 8.2) [1282], ibutilide (pIC ₅₀ 7.6–8) [695, 1106]	–	–	–	–	–
Functional Characteristics	cardiac I _{K_R}	–	–	–	–	–
Comments	RPR260243 is an activator of K _v 11.1 [670].	–	–	–	–	–

Further reading on Potassium channels

- Aguilar-Bryan L *et al.* (1998) Toward understanding the assembly and structure of KATP channels. *Physiol Rev* **78**: 227-45 [PMID:9457174]
- Borsotto M *et al.* (2015) Targeting two-pore domain K(+) channels TREK-1 and TASK-3 for the treatment of depression: a new therapeutic concept. *Br J Pharmacol* **172**: 771-84 [PMID:25263033]
- Chang PC *et al.* (2015) SK channels and ventricular arrhythmias in heart failure. *Trends Cardiovasc Med* **25**: 508-14 [PMID:25743622]
- Decher N *et al.* (2017) Stretch-activated potassium currents in the heart: Focus on TREK-1 and arrhythmias. *Prog Biophys Mol Biol* **130**: 223-232 [PMID:28526352]
- Feliciangeli S *et al.* (2015) The family of K2P channels: salient structural and functional properties. *J Physiol (Lond.)* **593**: 2587-603 [PMID:25530075]
- Foster MN *et al.* (2016) KATP Channels in the Cardiovascular System. *Physiol Rev* **96**: 177-252 [PMID:26660852]
- Gada K *et al.* (2019) Two-pore domain potassium channels: emerging targets for novel analgesic drugs: IUPHAR Review 26. *Br J Pharmacol* **176**: 256-266 [PMID:30325008]
- Goldstein SA *et al.* (2005) International Union of Pharmacology. LV. Nomenclature and molecular relationships of two-P potassium channels. *Pharmacol Rev* **57**: 527-40 [PMID:16382106]
- Greene DL *et al.* (2017) Modulation of Kv7 channels and excitability in the brain. *Cell Mol Life Sci* **74**: 495-508 [PMID:27645822]
- Gutman GA *et al.* (2003) International Union of Pharmacology. XLI. Compendium of voltage-gated ion channels: potassium channels. *Pharmacol Rev* **55**: 583-6 [PMID:14657415]
- Kaczmarek LK *et al.* (2017) International Union of Basic and Clinical Pharmacology. C. Nomenclature and Properties of Calcium-Activated and Sodium-Activated Potassium Channels. *Pharmacol Rev* **69**: 1-11 [PMID:28267675]
- Kubo Y *et al.* (2005) International Union of Pharmacology. LIV. Nomenclature and molecular relationships of inwardly rectifying potassium channels. *Pharmacol Rev* **57**: 509-26 [PMID:16382105]
- Latorre R *et al.* (2017) Molecular Determinants of BK Channel Functional Diversity and Functioning. *Physiol Rev* **97**: 39-87 [PMID:27807200]
- Niemeyer MI *et al.* (2016) Gating, Regulation, and Structure in K2P K+ Channels: In Varietate Concordia? *Mol Pharmacol* **90**: 309-17 [PMID:27268784]
- Poveda JA *et al.* (2017) Towards understanding the molecular basis of ion channel modulation by lipids: Mechanistic models and current paradigms. *Biochim Biophys Acta* **1859**: 1507-1516 [PMID:28408206]
- Rifkin RA *et al.* (2017) G Protein-Gated Potassium Channels: A Link to Drug Addiction. *Trends Pharmacol Sci* **38**: 378-392 [PMID:28188005]
- Taylor KC *et al.* (2017) Regulation of KCNQ/Kv7 family voltage-gated K+ channels by lipids. *Biochim Biophys Acta* **1859**: 586-597 [PMID:27818172]
- Vivier D *et al.* (2016) Perspectives on the Two-Pore Domain Potassium Channel TREK-1 (TWIK-Related K(+) Channel 1). A Novel Therapeutic Target? *J Med Chem* **59**: 5149-57 [PMID:26588045]
- Wei AD *et al.* (2005) International Union of Pharmacology. LII. Nomenclature and molecular relationships of calcium-activated potassium channels. *Pharmacol Rev* **57**: 463-72 [PMID:16382103]
- Yang KC *et al.* (2016) Mechanisms contributing to myocardial potassium channel diversity, regulation and remodeling. *Trends Cardiovasc Med* **26**: 209-18 [PMID:26391345]

Ryanodine receptors (RyR)

Ion channels → **Voltage-gated ion channels** → **Ryanodine receptors (RyR)**

Overview: The ryanodine receptors (RyRs) are found on intracellular Ca²⁺ storage/release organelles. The family of RyR genes encodes three highly related Ca²⁺ release channels: RyR1, RyR2 and RyR3, which assemble as large tetrameric structures. These RyR channels are ubiquitously expressed in many types of cells and participate in a variety of important Ca²⁺ signaling phenom-

ena (neurotransmission, secretion, *etc.*). In addition to the three mammalian isoforms described below, various nonmammalian isoforms of the ryanodine receptor have been identified [1343]. The function of the ryanodine receptor channels may also be influenced by closely associated proteins such as the tacrolimus (FK506)-binding protein, calmodulin [1553], triadin, calseques-

trin, junctin and sorcin, and by protein kinases and phosphatases. Recent studies solving the structure of the ryanodine receptor have shed light on the structural basis of ryanodine receptor function [see, for example, Samsó (2017) [1220] and Meissner (2017) [937]].

Nomenclature	RyR1	RyR2	RyR3
HGNC, UniProt	RYR1 , P21817	RYR2 , Q92736	RYR3 , Q15413
Endogenous activators	cytosolic ATP (endogenous; mM range), luminal Ca²⁺ (endogenous), cytosolic Ca²⁺ (endogenous; μ M range)	cytosolic ATP (endogenous; mM range), cytosolic Ca²⁺ (endogenous; μ M range), luminal Ca²⁺ (endogenous)	cytosolic ATP (endogenous; mM range), cytosolic Ca²⁺ (endogenous; μ M range)
Activators	caffeine (pharmacological; mM range), ryanodine (pharmacological; nM - μ M range), suramin (pharmacological; μ M range)	caffeine (pharmacological; mM range), ryanodine (pharmacological; nM - μ M range), suramin (pharmacological; μ M range)	caffeine (pharmacological; mM range), ryanodine (pharmacological; nM - μ M range)
Endogenous antagonists	cytosolic Ca²⁺ Concentration range: $>1 \times 10^{-4}$ M, cytosolic Mg²⁺ (mM range)	cytosolic Ca²⁺ Concentration range: $>1 \times 10^{-3}$ M, cytosolic Mg²⁺ (mM range)	cytosolic Ca²⁺ Concentration range: $>1 \times 10^{-3}$ M, cytosolic Mg²⁺ (mM range)
Antagonists	dantrolene	–	dantrolene
Channel blockers	procaine , ruthenium red , ryanodine Concentration range: $>1 \times 10^{-4}$ M	procaine , ruthenium red , ryanodine Concentration range: $>1 \times 10^{-4}$ M	ruthenium red
Functional Characteristics	Ca²⁺ : (P_{Ca}/P_K) single-channel conductance: 90 pS (50 mM Ca²⁺), 770 pS (200 mM K⁺)	Ca²⁺ : (P_{Ca}/P_K) single-channel conductance: 90 pS (50 mM Ca²⁺), 720 pS (210 mM K⁺)	Ca²⁺ : (P_{Ca}/P_K) single-channel conductance: 140 pS (50 mM Ca²⁺), 777 pS (250 mM K⁺)
Comments	RyR1 is also activated by depolarisation <i>via</i> DHP receptor, calmodulin at low cytosolic Ca²⁺ concentrations, CaM kinase and PKA; antagonised by calmodulin at high cytosolic Ca²⁺ concentrations	RyR2 is also activated by CaM kinase and PKA; antagonised by calmodulin at high cytosolic Ca²⁺ concentrations	RyR3 is also activated by calmodulin at low cytosolic Ca²⁺ concentrations; antagonised by calmodulin at high cytosolic Ca²⁺ concentrations

Comments: The modulators of channel function included in this table are those most commonly used to identify ryanodine-sensitive [Ca²⁺](#) release pathways. Numerous other modulators of ryanodine receptor/channel function can be found in the reviews listed below. The absence of a modulator of a particular isoform of receptor indicates that the action of that modulator has not been determined, not that it is without effect. The potential role of cyclic ADP ribose as an endogenous regulator of ryanodine receptor channels is controversial. A region of RyR likely to be involved in ion translocation and selection has been identified [[423](#), [1613](#)].

Further reading on Ryanodine receptors (RyR)

- Dulhunty AF *et al.* (2017) Physiology and Pharmacology of Ryanodine Receptor Calcium Release Channels. *Adv Pharmacol* **79**: 287-324 [[PMID:28528672](#)]
- Meissner G. (2017) The structural basis of ryanodine receptor ion channel function. *J Gen Physiol* **149**: 1065-1089 [[PMID:29122978](#)]
- O'Brien F *et al.* (2015) The ryanodine receptor provides high throughput [Ca²⁺](#)-release but is precisely regulated by networks of associated proteins: a focus on proteins relevant to phosphorylation. *Biochem Soc Trans* **43**: 426-33 [[PMID:26009186](#)]
- Samsó M. (2017) A guide to the 3D structure of the ryanodine receptor type 1 by cryoEM. *Protein Sci* **26**: 52-68 [[PMID:27671094](#)]
- Van Petegem F. (2015) Ryanodine receptors: allosteric ion channel giants. *J Mol Biol* **427**: 31-53 [[PMID:25134758](#)]
- Zalk R *et al.* (2017) [Ca²⁺](#) Release Channels Join the 'Resolution Revolution'. *Trends Biochem Sci* **42**: 543-555 [[PMID:28499500](#)]

Transient Receptor Potential channels (TRP)

Ion channels → Voltage-gated ion channels → Transient Receptor Potential channels (TRP)

Overview:

The TRP superfamily of channels (**nomenclature as agreed by NC-IUPHAR** [247, 1529]), whose founder member is the *Drosophila* Trp channel, exists in mammals as six families; TRPC, TRPM, TRPV, TRPA, TRPP and TRPML based on amino acid homologies. TRP subunits contain six putative TM domains and assemble as homo- or hetero-tetramers to form cation selective channels with diverse modes of activation and varied permeation properties (reviewed by [1065]). Established, or potential, physiological functions of the individual members of the TRP families are discussed in detail in the recommended reviews and in a number of books [367, 610, 1018, 1631]. The established, or potential, involvement of TRP channels in disease [1592] is reviewed

in [711, 1016], [1020] and [727], together with a special edition of *Biochimica et Biophysica Acta* on the subject [1016]. Additional disease related reviews, for pain [971], stroke [1603], sensation and inflammation [1423], itch [183], and airway disease [470, 1508], are available. The pharmacology of most TRP channels has been advanced in recent years. Broad spectrum agents are listed in the tables along with more selective, or recently recognised, ligands that are flagged by the inclusion of a primary reference. See Rubaiy (2019) for a review of pharmacological tools for TRPC1/C4/C5 channels [1195]. Most TRP channels are regulated by phosphoinositides such as $\text{PI}(4,5)\text{P}_2$ although the effects reported are often complex, occasionally contradictory, and likely

to be dependent upon experimental conditions, such as intracellular ATP levels (reviewed by [1021, 1184, 1440]). Such regulation is generally not included in the tables. When thermosensitivity is mentioned, it refers specifically to a high Q10 of gating, often in the range of 10-30, but does not necessarily imply that the channel's function is to act as a 'hot' or 'cold' sensor. In general, the search for TRP activators has led to many claims for temperature sensing, mechanosensation, and lipid sensing. All proteins are of course sensitive to energies of binding, mechanical force, and temperature, but the issue is whether the proposed input is within a physiologically relevant range resulting in a response.

TRPA (ankyrin) family

TRPA1 is the sole mammalian member of this group (reviewed by [430]). TRPA1 activation of sensory neurons contribute to nociception [653, 932, 1321]. Pungent chemicals such as mustard oil (AITC), **allicin**, and **cinnamaldehyde** activate TRPA1 by modification of free thiol groups of cysteine side chains, especially those located in its amino terminus [92, 547, 883, 885]. Alkenals with α , β -unsaturated bonds, such as propenal (**acrolein**), butenal (**croty-**

ldehyde), and **2-pentenal** can react with free thiols *via* Michael addition and can activate TRPA1. However, potency appears to weaken as carbon chain length increases [43, 92]. Covalent modification leads to sustained activation of TRPA1. Chemicals including **carvacrol**, menthol, and local anesthetics reversibly activate TRPA1 by non-covalent binding [675, 790, 1540, 1541]. TRPA1 is not mechanosensitive under physiological conditions, but can be

activated by cold temperatures [302, 676]. The electron cryo-EM structure of TRPA1 [1087] indicates that it is a 6-TM homotetramer. Each subunit of the channel contains two short 'pore helices' pointing into the ion selectivity filter, which is big enough to allow permeation of partially hydrated Ca^{2+} ions.

Nomenclature	TRPA1
HGNC, UniProt	TRPA1, O75762
Chemical activators	Isothiocyanates (covalent) and 1,4-dihydropyridines (non-covalent)
Oxidative stress compounds	4-oxo-nonenal: pEC ₅₀ 5.7, H ₂ O ₂ : pEC ₅₀ 3.6, hypochlorite: EC ₅₀ 11 ppm (human) and 7 ppm (mouse) (Mouse) [41, 110, 1227]
Physical activators	Cooling (<17°C) (disputed) [653, 989, 1420]
Activators	polygodial (pEC ₅₀ 6.4) [374], acrolein (Agonist) (pEC ₅₀ 5.3) [physiological voltage] [92], allicin (Agonist) (pEC ₅₀ 5.1) [physiological voltage] [93], Δ ⁹ -tetrahydrocannabinol (Agonist) (pEC ₅₀ 4.9) [-60mV] [653], nicotine (non-covalent) (pEC ₅₀ 4.8) [-75mV] [1358], thymol (non-covalent) (pEC ₅₀ 4.7) Concentration range: 6.2 × 10 ⁻⁶ M-2.5 × 10 ⁻⁵ M [786], URB597 (Agonist) (pEC ₅₀ 4.6) [1015], (-)-menthol (Partial agonist) (pEC ₅₀ 4-4.5) [675, 1535], allyl isothiocyanate (pEC ₅₀ 4.2) [547], cinnamaldehyde (Agonist) (pEC ₅₀ 4.2) [physiological voltage] [72] – Mouse, formalin (covalent. This level of activity is also observed for rat TRPA1) (pEC ₅₀ 3.4) [885, 932] – Mouse, icilin (Agonist) Concentration range: 1 × 10 ⁻⁴ M [physiological voltage] [1321] – Mouse
Selective activators	JT010 (pEC ₅₀ 9.2) [1355], PF-4840154 (This compound has similar activity at rat and mouse TRPA1) (pEC ₅₀ 7.6) [1205], chlorobenzylidene malononitrile (covalent) (pEC ₅₀ 6.7) [153], ASP7663 (pEC ₅₀ 6.3) [728]
Channel blockers	AP18 (Inhibition) (pIC ₅₀ 5.5) [1109], ruthenium red (Inhibition) (pIC ₅₀ 5.5) [-80mV] [989] – Mouse
Selective channel blockers	GDC-0334 (Inhibition) (pIC ₅₀ 8.8) [71], AM-0902 (Antagonist) (pIC ₅₀ 7.7) [1233], A-967079 (Inhibition) (pIC ₅₀ 7.2) [209]
Functional Characteristics	γ = 87-100 pS; conducts mono- and di-valent cations non-selectively (P _{Ca} /P _{Na} = 0.84); outward rectification; activated by elevated intracellular Ca ²⁺
Comments	miRNA-711 is a selective activator of TRPA1 (pEC ₅₀ ~5.0) [504]. GRC 17536 (structure not disclosed) is a TRPA1 antagonist with potential as an anti-tussive therapeutic [980]. Some pathogen-derived molecules activate human TRPA1, such as lipopolysaccharide (LPS) [938], indole (pEC ₅₀ 4.1) [241] and indole-3-carboxyaldehyde (pEC ₅₀ 4.1) [1568].

TRPC (canonical) family

Members of the TRPC subfamily (reviewed by [7, 36, 99, 120, 414, 709, 1083, 1143]) fall into the subgroups outlined below. TRPC2 is a pseudogene in humans. It is generally accepted that all TRPC channels are activated downstream of G_{q/11}-coupled receptors, or receptor tyrosine kinases (reviewed by [1129, 1405, 1529]). A comprehensive listing of G protein-coupled receptors that activate TRPC channels is given in [7]. Hetero-oligomeric complexes of TRPC channels and their association with proteins to form signal-

ling complexes are detailed in [36] and [710]. TRPC channels have frequently been proposed to act as store-operated channels (SOCs) (or components of multimeric complexes that form SOCs), activated by depletion of intracellular calcium stores (reviewed by [36, 95, 223, 224, 1055, 1094, 1135, 1217, 1591]). However, the weight of the evidence is that they are not directly gated by conventional store-operated mechanisms, as established for STIM-gated Orai channels. TRPC channels are not mechanically gated in physio-

logically relevant ranges of force. All members of the TRPC family are blocked by 2-APB and SKF96365 [517, 518]. Activation of TRPC channels by lipids is discussed by [99]. Important progress has been recently made in TRPC pharmacology [136, 268, 426, 697, 958, 1195, 1259]. TRPC channels regulate a variety of physiological functions and are implicated in many human diseases [100, 137, 217, 436, 633, 848, 1312, 1341, 1467, 1486].

TRPC1/C4/C5 subgroup

TRPC1 alone may not form a functional ion channel [320]. The structures of the apo and antagonist-bound states of TRPC1/TRPC4 heteromeric channels have been resolved by cryo-EM [1521]. TRPC4/C5 may be distinguished from other TRP channels by their potentiation by micromolar concentrations of La³⁺. TRPC2 is a pseudogene in humans, but in other mammals appears to be an ion channel localized to microvilli of the vomeronasal organ. It is required for normal sexual behavior in response to pheromones in mice. It may also function in the main olfactory epithelia in mice [826, 1053, 1054, 1574, 1580, 1585, 1639].

TRPC3/C6/C7 subgroup

All members are activated by diacylglycerol independent of protein kinase C stimulation [518].

	TRPC1	TRPC2	TRPC3	TRPC4
Nomenclature	TRPC1	TRPC2	TRPC3	TRPC4
HGNC, UniProt	TRPC1, P48995	TRPC2, –	TRPC3, Q13507	TRPC4, Q9UBN4
Chemical activators	NO-mediated cysteine S-nitrosylation	Diacylglycerol (SAG, OAG, DOG): strongly inhibited by Ca ²⁺ /CaM once activated by DAG [1305]	diacylglycerols	NO-mediated cysteine S-nitrosylation, potentiation by extracellular protons
Physical activators	membrane stretch (likely direct)	DAG kinase; regulates DAG concentration in vomeronasal sensory neurons	–	–
Endogenous activators	–	Intracellular Ca ²⁺	–	–
Activators	–	DOG (Agonist) Concentration range: 1 × 10 ⁻⁴ M [-80mV] [865] – Mouse, SAG (Agonist) Concentration range: 1 × 10 ⁻⁴ M [-80mV] [865] – Mouse	pyrazolopyrimidine 4n (pEC ₅₀ 7.7) [1150], GSK1702934A (Agonist) (pEC ₅₀ 7.1) [1548]	(-)-englerin A (Agonist) (pEC ₅₀ 7.9) [18], tonantzitlone (pEC ₅₀ 6.9) [1198], La ³⁺ (μM range)
Channel blockers	2-APB (Antagonist) [-70mV] [1329], Gd ³⁺ (Antagonist) Concentration range: 2 × 10 ⁻⁵ M [-70mV] [1636], La ³⁺ (Antagonist) Concentration range: 1 × 10 ⁻⁴ M [-70mV] [1329]	2-APB (Antagonist) Concentration range: 5 × 10 ⁻⁵ M [-70mV – 80mV] [865] – Mouse, U73122 (Antagonist) Concentration range: 1 × 10 ⁻⁵ M – Mouse	GSK2833503A (pIC ₅₀ 7.7) [80mV] [1252], GSK417651A (Antagonist) (pIC ₅₀ 7.1) [1495], Gd ³⁺ (Antagonist) (pEC ₅₀ 7) [-60mV] [502], SAR7334 (pIC ₅₀ 6.6) [891], BTP2 (Antagonist) (pIC ₅₀ 6.5) [-80mV] [523], Pyr3 (pIC ₅₀ 6.2) [715], Pyr10 (Antagonist) (pIC ₅₀ 6.1) [1235], norgestimate (pK _i 5.5) [947], La ³⁺ (Antagonist) (pIC ₅₀ 5.4) [-60mV] [502], clemizole (pIC ₅₀ 5) [1176], 2-APB (Antagonist) (pIC ₅₀ 5) [physiological voltage] [824], Ni ²⁺ , SKF96365	HC-070 (Antagonist) (pIC ₅₀ 7.3) [655], ML204 (pIC ₅₀ 5.5) [952], M084 (Inhibition) (pIC ₅₀ 5.3) [1632], clemizole (pIC ₅₀ 5.2) [1176], La ³⁺ (mM range), SKF96365, niflumic acid (Antagonist) Concentration range: 3 × 10 ⁻⁵ M [-60mV] [1459] – Mouse
Functional Characteristics	It is not yet clear that TRPC1 forms a homomer. It does form heteromers with TRPC4 and TRPC5	γ = 42 pS linear single channel conductance in 150 mM symmetrical Na ⁺ in vomeronasal sensory neurons. P _{Ca} /P _{Na} = 2.7; permeant to Na ⁺ , Cs ⁺ , Ca ²⁺ , but not NMDG [1053, 1580]	γ = 66 pS; conducts mono and divalent cations non-selectively (P _{Ca} /P _{Na} = 1.6); monovalent cation current suppressed by extracellular Ca ²⁺ ; dual (inward and outward) rectification	γ = 30 -41 pS, conducts mono and divalent cations non-selectively (P _{Ca} /P _{Na} = 1.1 - 7.7); dual (inward and outward) rectification

Nomenclature	TRPC5	TRPC6	TRPC7
HGNC, UniProt	TRPC5, Q9UL62	TRPC6, Q9Y210	TRPC7, Q9HCX4
Chemical activators	NO-mediated cysteine S-nitrosylation (disputed), potentiation by extracellular protons	Diacylglycerols	diacylglycerols
Physical activators	Membrane stretch	Membrane stretch	–
Endogenous activators	intracellular Ca ²⁺ (at negative potentials) (pEC ₅₀ 6.2), lysophosphatidylcholine	20-HETE, arachidonic acid, lysophosphatidylcholine	–
Activators	(-)-englerin A (Agonist) (pEC ₅₀ 8.1) [18], tonantzitolone (pEC ₅₀ 7.1) [1198], BTD (pEC ₅₀ 5.8) [98], riluzole (pEC ₅₀ 5) [1177], methylprednisolone (pEC ₅₀ 4.9) [98], rosiglitazone (pEC ₅₀ 4.5) [897], Gd ³⁺ Concentration range: 1 × 10 ⁻⁴ M, La ³⁺ (µM range), Pb ²⁺ Concentration range: 5 × 10 ⁻⁶ M, genistein (independent of tyrosine kinase inhibition) [1522]	AM-0883 (Agonist) (pEC ₅₀ 7.3) [68], GSK1702934A (Agonist) (pIC ₅₀ 6.4) [1548], pyrazolopyrimidine 4n (pEC ₅₀ 5.9) [1150], OptoBI-1 (photoswitch activation; concentration range: 1-2x10 ⁻⁵ M) [1388], OptoDARg (photoswitch activation; concentration range: 3x10 ⁻⁵ M) [822], flufenamate, hyp 9 [800], hyperforin [801]	pyrazolopyrimidine 4n (pIC ₅₀ 6.1) [1150], Op-toBI-1 (photoswitch activation; concentration range: 1-2x10 ⁻⁵ M) [1388]
Selective activators	AM237 (pEC ₅₀ 7.7) [957]	–	–
Channel blockers	Pico145 (Inhibition) (pIC ₅₀ 8.9) [1197], HC-070 (Antagonist) (pIC ₅₀ 8) [655], AM12 (Inhibition) (pIC ₅₀ 6.6) [1001], GFB-8438 (Inhibition) (pIC ₅₀ 6.5) [1587], galangin (pK _i 6.3) [1001], clemizole (pIC ₅₀ 6) [1176], KB-R7943 (Inhibition) (pIC ₅₀ 5.9) [736], M084 (Inhibition) (pIC ₅₀ 5.1) [1632], ML204 (pIC ₅₀ ~5) [952], 2-APB (Antagonist) (pIC ₅₀ 4.7) [-80mV] [1546], La ³⁺ (Antagonist) Concentration range: 5 × 10 ⁻³ M [-60mV] [654] – Mouse	AM-1473 (Antagonist) (pIC ₅₀ 9.7) [68], GSK2833503A (pIC ₅₀ 8.5) [80mV] [1252], GSK2332255B (Antagonist) (pIC ₅₀ 8.4) [1252], SAR7334 (pIC ₅₀ 8) [891], BTDM (Inhibition) (pIC ₅₀ 8) [1366], DS88790512 (Inhibition) (pIC ₅₀ ~7.9) [976], BI 749327 (Antagonist) (pIC ₅₀ 7.9) [827], SH045 (pIC ₅₀ 7.2) [500], larixyl acetate (Inhibition) (pIC ₅₀ 7) [1414], GSK417651A (Antagonist) (pIC ₅₀ 6.4) [1495], Pyrazolo-pyrimidine 14a (Inhibition) (pIC ₅₀ ~6) [324], clemizole (pIC ₅₀ 5.9) [1176], Gd ³⁺ (Antagonist) (pIC ₅₀ 5.7) [-60mV] [599] – Mouse, SKF96365 (Antagonist) (pIC ₅₀ 5.4) [-60mV] [599] – Mouse, norgestimate (pIC ₅₀ 5.3) [947], La ³⁺ (pIC ₅₀ ~5.2), amiloride (Antagonist) (pIC ₅₀ 3.9) [-60mV] [599] – Mouse, Cd ²⁺ (Antagonist) (pIC ₅₀ 3.6) [-60mV] [599] – Mouse, 2-APB, ACAA, GsMTx-4, Extracellular H ⁺ , KB-R7943, ML9	SH045 (pIC ₅₀ ~7.7) [500], SAR7334 (pIC ₅₀ 6.7) [891], BI 749327 (Antagonist) (pIC ₅₀ 6.3) [827], larixyl acetate (Inhibition) (pIC ₅₀ ~6.3) [1414], 2-APB, La ³⁺ (Antagonist) Concentration range: 1 × 10 ⁻⁴ M [-60mV] [1045] – Mouse, SKF96365 (Antagonist) Concentration range: 2.5 × 10 ⁻⁵ M [-60mV] [1045] – Mouse, amiloride
Selective channel blockers	AC1903 (Inhibition) (pIC ₅₀ 4.8) [1627]	–	–
Functional Characteristics	γ = 41-63 pS; conducts mono- and di-valent cations non-selectively (P _{Ca} /P _{Na} = 1.8 - 9.5); dual rectification (inward and outward) as a homomer, outwardly rectifying when expressed with TRPC1 or TRPC4	γ = 28-37 pS; conducts mono and divalent cations with a preference for divalents (P _{Ca} /P _{Na} = 4.5-5.0); monovalent cation current suppressed by extracellular Ca ²⁺ and Mg ²⁺ , dual rectification (inward and outward), or inward rectification	γ = 25-75 pS; conducts mono and divalent cations with a preference for divalents (P _{Ca} /P _{CS} = 5.9); modest outward rectification (monovalent cation current recorded in the absence of extracellular divalents); monovalent cation current suppressed by extracellular Ca ²⁺ and Mg ²⁺

TRPM (melastatin) family

Members of the TRPM subfamily (reviewed by [408, 517, 1094, 1620]) fall into the five subgroups outlined below.

TRPM1/M3 subgroup

In darkness, glutamate released by the photoreceptors and ON-bipolar cells binds to the metabotropic glutamate receptor 6, leading to activation of Go. This results in the closure of TRPM1. When the photoreceptors are stimulated by light, glutamate release is reduced, and TRPM1 channels are more active, resulting in cell membrane depolarization. Human TRPM1 mutations are associated with congenital stationary night blindness (CSNB),

whose patients lack rod function. TRPM1 is also found in melanocytes. Isoforms of TRPM1 may present in melanocytes, melanoma, brain, and retina. In melanoma cells, TRPM1 is prevalent in highly dynamic intracellular vesicular structures [602, 1037]. TRPM3 (reviewed by [1040]) exists as multiple splice variants which differ significantly in their biophysical properties. TRPM3 is expressed in somatosensory neurons and may be important in

development of heat hyperalgesia during inflammation (see review [1375]). TRPM3 is frequently coexpressed with TRPA1 and TRPV1 in these neurons. TRPM3 is expressed in pancreatic beta cells as well as brain, pituitary gland, eye, kidney, and adipose tissue [1039, 1374]. TRPM3 may contribute to the detection of noxious heat [1447].

TRPM2

TRPM2 is activated under conditions of oxidative stress (respiratory burst of phagocytic cells). The direct activators are calcium, adenosine diphosphate ribose (ADPR) [1413] and cyclic ADPR (cADPR) [1588]. As for many ion channels, PI(4,5)P2 must also be present [1574]. Numerous splice variants of TRPM2 exist which differ in their activation mechanisms [345]. Recent studies have reported structures of human (hs) TRPM2, which demonstrate two ADPR binding sites in hsTRPM2, one in the N-terminal MHR1/2 domain and the other in the C-terminal NUDT9-H do-

main. In addition, one Ca²⁺ binding site in the intracellular S2-S3 loop is revealed and proposed to mediate Ca²⁺ binding that induces conformational changes leading the ADPR-bound closed channel to open [588, 1476]. Meanwhile, a quadruple-residue motif (979FGQI982) was identified as the ion selectivity filter and a gate to control ion permeation in hsTRPM2 [1589]. TRPM2 is involved in warmth sensation [1252], and contributes to several diseases [107]. TRPM2 interacts with extra synaptic NMDA receptors (NMDAR) and enhances NMDAR activity in ischemic stroke

[1637]. Activation of TRPM2 in macrophages promotes atherosclerosis [1612, 1638]. Moreover, silica nanoparticles induce lung inflammation in mice *via* ROS/PARP/TRPM2 signaling-mediated lysosome impairment and autophagy dysfunction [1478]. Recent studies have designed various compounds for their potential to selectively inhibit the TRPM2 channel, including ACA derivatives A23, and 2,3-dihydroquinazolin-4(1H)-one derivatives [1606, 1608].

TRPM4/5 subgroup

TRPM4 and TRPM5 have the distinction within all TRP channels of being impermeable to Ca²⁺ [1529]. A splice variant of TRPM4 (*i.e.* TRPM4b) and TRPM5 are molecular candidates for endogenous calcium-activated cation (CAN) channels [488]. TRPM4 is active in the late phase of repolarization of the cardiac ventric-

ular action potential. TRPM4 deletion or knockout enhances beta adrenergic-mediated inotropy [922]. Mutations are associated with conduction defects [614, 922, 1307]. TRPM4 has been shown to be an important regulator of Ca²⁺ entry in to mast cells [1426] and dendritic cell migration [76]. TRPM5 in taste receptor

cells of the tongue appears essential for the transduction of sweet, amino acid and bitter stimuli [825] TRPM5 contributes to the slow afterdepolarization of layer 5 neurons in mouse prefrontal cortex [792]. Both TRPM4 and TRPM5 are required for transduction of taste stimuli [356].

TRPM6/7 subgroup

TRPM6 and 7 combine channel and enzymatic activities ('chanzymes') [245]. These channels have the unusual property of permeation by divalent (Ca²⁺, Mg²⁺, Zn²⁺) and monovalent cations, high single channel conductances, but overall extremely small inward conductance when expressed to the plasma membrane. They are inhibited by internal Mg²⁺ at ~0.6 mM, around the free

level of Mg²⁺ in cells. Whether they contribute to Mg²⁺ homeostasis is a contentious issue. PIP2 is required for TRPM6 and TRPM7 activation [1202, 1536]. When either gene is deleted in mice, the result is embryonic lethality [638, 1524]. The C-terminal kinase region of TRPM6 and TRPM7 is cleaved under unknown stimuli, and the kinase phosphorylates nuclear histones [740, 741].

TRPM7 is responsible for oxidant-induced Zn²⁺ release from intracellular vesicles [6] and contributes to intestinal mineral absorption essential for postnatal survival [961]. The putative metal transporter proteins CNNM1-4 interact with TRPM7 and regulate TRPM7 channel activity [69, 730].

TRPM8

Is a channel activated by cooling and pharmacological agents

evoking a 'cool' sensation and participates in the thermosensa-

tion of cold temperatures [94, 249, 315] reviewed by [723, 850,

902, 1441]. Direct chemical agonists include menthol and **icilin** [1545]. Besides, **linalool** can promote ERK phosphorylation in human dermal microvascular endothelial cells, down-regulate intracellular ATP levels, and activate TRPM8 [97]. Recent studies have found that TRPM8 has typical S4-S5 connectomes with clear selective filters and exowell rings [791], and have identified cryo-electron microscopy structures of mouse TRPM8 in closed, intermediate, and open states along the ligand- and PIP₂-dependent gated pathways [1575]. Moreover, the last 36 amino acids at

the carboxyl terminal of TRPM8 are key protein sequences for TRPM8's temperature-sensitive function [270]. TRPM8 deficiency reduced the expression of S100A9 and increased the expression of HNF4 α in the liver of mice, which reduced inflammation and fibrosis progression in mice with liver fibrosis, and helped to alleviate the symptoms of bile duct disease [844]. Channel deficiency also shortens the time of hypersensitivity reactions in migraine mouse models by promoting the recovery of normal sensitivity [26]. A cyclic peptide DeC-1.2 was designed to inhibit ligand acti-

vation of TRPM8 but not cold activation, which can eliminate the side effects of cold dysalgesia in oxaliplatin-treated mice without changing body temperature [17]. Analysis of clinical data shows that TRPM8-specific blockers WS12 can reduce tumor growth in colorectal cancer xenografted mice by reducing transcription and activation of Wnt signaling regulators and β -catenin and its target oncogenes, such as C-Myc and Cyclin D1 [1067].

Nomenclature	TRPM1	TRPM2
HGNC, UniProt	TRPM1, Q7Z4N2	TRPM2, O94759
Chemical activators	–	Agents producing reactive oxygen (e.g. H ₂ O ₂) and nitrogen (e.g. GEA 3162) species
Physical activators	–	Heat ~ 35°C
Endogenous activators	pregnenolone sulphate [766]	intracellular cADPR (Agonist) (pEC ₅₀ 5) [-80mV – -60mV] [96, 729, 1390], intracellular ADP ribose (Agonist) (pEC ₅₀ 3.9–4.4) [-80mV] [1105], intracellular Ca²⁺ (perhaps <i>via</i> calmodulin), H₂O₂ (Agonist) Concentration range: 5 × 10 ⁻⁷ M–5 × 10 ⁻⁵ M [physiological voltage] [410, 514, 738, 1291, 1502], membrane PIP₂ [1397], arachidonic acid (Potentiation) Concentration range: 1 × 10 ⁻⁵ M–3 × 10 ⁻⁵ M [physiological voltage] [514]
Activators	–	GEA 3162
Endogenous channel blockers	Zn²⁺ (pIC ₅₀ 6)	Zn²⁺ (pIC ₅₀ 6), extracellular H⁺
Channel blockers	–	2-APB (Antagonist) (pIC ₅₀ 6.1) [-60mV] [1391], ACAA (Antagonist) (pIC ₅₀ 5.8) [physiological voltage] [737], clotrimazole (Antagonist) Concentration range: 3 × 10 ⁻⁶ M–3 × 10 ⁻⁵ M [-60mV – -15mV] [544], econazole (Antagonist) Concentration range: 3 × 10 ⁻⁶ M–3 × 10 ⁻⁵ M [-60mV – -15mV] [544], flufenamic acid (Antagonist) Concentration range: 5 × 10 ⁻⁵ M–1 × 10 ⁻³ M [-60mV – -50mV] [543, 1391], miconazole (Antagonist) Concentration range: 1 × 10 ⁻⁵ M [-60mV] [1391]
Functional Characteristics	Conducts mono- and di-valent cations non-selectively, dual rectification (inward and outward)	γ = 52–60 pS at negative potentials, 76 pS at positive potentials; conducts mono- and di-valent cations non-selectively (P _{Ca} /P _{Na} = 0.6–0.7); non-rectifying; inactivation at negative potentials; activated by oxidative stress probably <i>via</i> PARP-1, PARP inhibitors reduce activation by oxidative stress, activation inhibited by suppression of APDR formation by glycohydrolase inhibitors.
Comments	–	Additional endogenous activators include 2'-deoxy-ADPR, 3'-deoxy-ADPR, 2'-phospho-ADPR, 2-F-ADPR and ADP-ribose-2'-phosphate (ADPRP) [409, 1398]. 8-Br-cADPR acts as a gating inhibitor [729].

Nomenclature	TRPM3	TRPM4
HGNC, UniProt	TRPM3, Q9HCF6	TRPM4, Q8TD43
Other channel blockers	–	Intracellular nucleotides including ATP , ADP , adenosine 5'-monophosphate and AMP-PNP with an IC ₅₀ range of 1.3–1.9 μM
Physical activators	heat (Q ₁₀ = 7.2 between 15 - 25°C), hypotonic cell swelling [479, 1447, 1448]	Membrane depolarization (V _{0.5} = -20 mV to +60 mV dependent upon conditions) in the presence of elevated [Ca ²⁺] _i , heat (Q ₁₀ = 8.5 @ +25 mV between 15 and 25°C)
Endogenous activators	pregnenolone sulphate (pEC ₅₀ 4.9) [1452], sphingosine (Agonist) (pEC ₅₀ 4.9) [physiological voltage] [480], dihydrosphingosine (Agonist) (pEC ₅₀ 4.7) [480], epipregnanolone sulphate [896]	intracellular Ca²⁺ (Agonist) (pEC ₅₀ 3.9–6.3) [-100mV – 100mV] [1019, 1023, 1024, 1356]
Activators	CIM0216 (pEC ₅₀ 6.1) [529, 1375], nifedipine , pentafluoro-trityl clotrimazole analogue 29a (Agonist) [660]	BTP2 (Agonist) (pEC ₅₀ 8.1) [-80mV] [1356], decavanadate (Agonist) (pEC ₅₀ 5.7) [-100mV] [1023]
Gating inhibitors	2-APB (Antagonist) (pIC ₅₀ 4) [1546]	flufenamic acid (Antagonist) (pIC ₅₀ 5.6) [100mV] [1413] – Mouse, clotrimazole (Antagonist) Concentration range: 1 × 10 ⁻⁶ M–1 × 10 ⁻⁵ M [1027]
Endogenous channel blockers	Mg²⁺ (Antagonist) (pIC ₅₀ 2) [1038] – Mouse, extracellular Na⁺ (TRPM3α2 only)	–
Channel blockers	isosakuranetin (pIC ₅₀ 7.3) [1322], primidone (pIC ₅₀ 6.2) [751], maprotiline (pIC ₅₀ 5.8) [751], (S)-liquiritigenin (pIC ₅₀ 5.2) [1322], diclofenac (pIC ₅₀ 5.2) [1345], naringenin (pIC ₅₀ 5.2) [1322, 1323], Gd³⁺ (Antagonist) (pIC ₅₀ 4) [479, 785], La³⁺ (Antagonist) (pIC ₅₀ 4) [479, 785], chloroform (Antagonist) (pIC ₅₀ 3.8) [690], halothane (Antagonist) (pIC ₅₀ 3.3) [690]	NBA (pIC ₅₀ 6.4) [1317], LBA (pIC ₅₀ 5.8) [1317], CBA (Antagonist) (pIC ₅₀ 5.7) [1066], meclofenamic acid (pIC ₅₀ 5.5) [1421], 9-phenanthrol (pIC ₅₀ 4.6–4.8) [471], spermine (Antagonist) (pIC ₅₀ 4.2) [100mV] [1025], adenosine (pIC ₅₀ 3.2)
Functional Characteristics	TRPM3 ₁₃₂₅ : γ = 83 pS (Na ⁺ current), 65 pS (Ca ²⁺ current); conducts mono and di-valent cations non-selectively (P _{Ca} /P _{Na} = 1.6) TRPM3α1: selective for monovalent cations (P _{Ca} /P _{Cs} ~0.1); TRPM3α2: conducts mono- and di-valent cations non-selectively (P _{Ca} /P _{Cs} = 1–10); In- and outwardly rectifying currents by co-application of pregnenolone sulphate and clotrimazole or single application of CIM0216 [529, 1446]. TRPM3α1 is activated by clotrimazole but not by pregnenolone sulphate [528].	γ = 23 pS (within the range 60 to +60 mV); permeable to monovalent cations; impermeable to Ca ²⁺ ; strong outward rectification; slow activation at positive potentials, rapid deactivation at negative potentials, deactivation blocked by decavanadate
Comments	G protein βγ subunits can act as endogenous inhibitors of TRPM3 channel activity [63, 309, 1152].	–

Nomenclature	TRPM5	TRPM6
HGNC, UniProt	TRPM5, Q9NZQ8	TRPM6, Q9BX84
EC number	–	2.7.11.1
Other chemical activators	–	constitutively active, activated by reduction of intracellular Mg ²⁺
Physical activators	membrane depolarization (V ₁ = 0 to + 120 mV dependent upon conditions), heat (Q ₁₀ = 10.3 @ -75 mV between 15 and 25°C)	–
Endogenous activators	intracellular Ca ²⁺ (Agonist) (pEC ₅₀ 4.5–6.2) [-80mV – 80mV] [557, 840, 1413] – Mouse	extracellular H ⁺ (Potentiation), intracellular Mg ²⁺
Activators	compound 39 (Agonist) (pEC ₅₀ 7.5) [1206]	2-APB (Agonist) (pEC ₅₀ 3.4–3.7) [-120mV – 100mV] [807]
Endogenous channel blockers	–	Mg ²⁺ (inward current mediated by monovalent cations is blocked) (pIC ₅₀ 5.5–6), Ca ²⁺ (inward current mediated by monovalent cations is blocked) (pIC ₅₀ 5.3–5.3)
Channel blockers	flufenamic acid (pIC ₅₀ 4.6), intracellular spermine (pIC ₅₀ 4.4), Extracellular H ⁺ (pIC ₅₀ 3.2)	ruthenium red (pIC ₅₀ 7) [voltage dependent -120mV]
Allosteric modulators	APV207095A (Potentiation) (pEC ₅₀ 5) [1436], APV207094A (Potentiation) (pEC ₅₀ 4.4) [1436], APV207010A (Potentiation) (pEC ₅₀ 4.4) [1436], APV206690A (Potentiation) (pEC ₅₀ 4) [1436]	–
Functional Characteristics	γ = 15–25 pS; conducts monovalent cations selectively (P _{Ca} /P _{Na} = 0.05); strong outward rectification; slow activation at positive potentials, rapid inactivation at negative potentials; activated and subsequently desensitized by [Ca ²⁺] _i	γ = 40–87 pS; permeable to mono- and di-valent cations with a preference for divalents (Mg ²⁺ > Ca ²⁺ ; P _{Ca} /P _{Na} = 6.9), conductance sequence Zn ²⁺ > Ba ²⁺ > Mg ²⁺ = Ca ²⁺ = Mn ²⁺ > Sr ²⁺ > Cd ²⁺ > Ni ²⁺ ; strong outward rectification abolished by removal of extracellular divalents, inhibited by intracellular Mg ²⁺ (IC ₅₀ = 0.5 mM) and ATP
Comments	TRPM5 is not blocked by ATP. APV206512A and APV206513A are TRPM5 blockers, with IC ₅₀ s of 15 μM [1436]. Steviol glycosides (sweet-tasting organic molecules) act as positive modulators of TRPM5 activity [1111].	–

Nomenclature	TRPM7	TRPM8
HGNC, UniProt	TRPM7, Q96QT4	TRPM8, Q7Z2W7
EC number	2.7.11.1	–
Chemical activators	–	agonist activities are temperature dependent and potentiated by cooling
Physical activators	–	depolarization ($V_{0.5} \sim +50$ mV at 15°C), cooling (< 22–26°C)
Endogenous activators	Extracellular H^+ (Potentiation) (pEC_{50} 4.5) [634]	–
Activators	2-APB Concentration range: $>1 \times 10^{-3}M$ [807] – Mouse, naltrexone [558]	icilin (Agonist) (pEC_{50} 6.7–6.9) [physiological voltage] [40, 101] – Mouse, tacrolimus (Agonist) (pEC_{50} 4.8) [49], (-)-menthol (inhibited by intracellular Ca^{2+}) (pEC_{50} 4.6) [-120mV – 160mV] [1439]
Selective activators	–	acoltremone (Full agonist) (pEC_{50} 4.9) [physiological voltage] [879, 1267] – Rat
Selective antagonists	–	KPR-5714 (pIC_{50} 7.6) [995]
Channel blockers	sphingosine (Inhibition) (pIC_{50} 6.2) [-100mV – 100mV] [1149] – Mouse, fingolimod (Inhibition) (pIC_{50} 6.1) [-100mV – 100mV] [1149] – Mouse, spermine (Inhibition) (pK_i 5.6) [-110mV – 80mV] [735] – Rat, 2-APB (Inhibition) (pIC_{50} 3.8) [-100mV – 100mV] [807] – Mouse, carvacrol (Inhibition) (pIC_{50} 3.5) [-100mV – 100mV] [1080] – Mouse, Mg^{2+} (Antagonist) (pIC_{50} 2.5) [80mV] [988] – Mouse, La^{3+} (Antagonist) Concentration range: $2 \times 10^{-3}M$ [-100mV – 100mV] [1203] – Mouse	BCTC (Antagonist) (pIC_{50} 6.1) [physiological voltage] [101] – Mouse, scutellarein (pIC_{50} 5.8) [1222], 2-APB (Antagonist) (pIC_{50} 4.9–5.1) [100mV – -100mV] [578, 1002] – Mouse, capsazepine (Antagonist) (pIC_{50} 4.7) [physiological voltage] [101] – Mouse
Selective channel blockers	–	PF-05105679 (Antagonist) (pIC_{50} 7) [voltage dependent] [44]
Functional Characteristics	$\gamma = 40$ –105 pS at negative and positive potentials respectively; conducts mono- and di-valent cations with a preference for monovalents ($P_{Ca}/P_{Na} = 0.34$); conductance sequence $Ni^{2+} > Zn^{2+} > Ba^{2+} = Mg^{2+} > Ca^{2+} = Mn^{2+} > Sr^{2+} > Cd^{2+}$; outward rectification, decreased by removal of extracellular divalent cations; inhibited by intracellular Mg^{2+} , Ba^{2+} , Sr^{2+} , Zn^{2+} , Mn^{2+} and Mg .ATP (disputed); activated by and intracellular alkalinization; sensitive to osmotic gradients	$\gamma = 40$ –83 pS at positive potentials; conducts mono- and di-valent cations non-selectively ($P_{Ca}/P_{Na} = 1.0$ –3.3); pronounced outward rectification; demonstrates desensitization to chemical agonists and adaptation to a cold stimulus in the presence of Ca^{2+} ; modulated by lysophospholipids and PUFAs
Comments	2-APB acts as a channel blocker in the μM range. Recent study shows cAMP inhibits TRPM7-mediated Ca^{2+} influx [152]. Waixenicin-A specifically inhibits TRPM7 [1634].	Cannabidiol and Δ^9 -tetrahydrocannabinol are examples of cannabinoid activators. TRPM8 is insensitive to ruthenium red. Icilin requires intracellular Ca^{2+} for full agonist activity.

TRPML (mucolipin) family

The TRPML family [264, 1140, 1146, 1544, 1596] consists of three mammalian members (TRPML1–3). TRPML channels are probably restricted to intracellular vesicles and mutations in the gene (*MCOLN1*) encoding TRPML1 (mucolipin-1) cause the neurodegenerative disorder mucopolipidosis type IV (MLIV) in man. TRP-

ML1 is a cation selective ion channel that is important for sorting/transport of endosomes in the late endocytotic pathway and specifically, fission from late endosome-lysosome hybrid vesicles and lysosomal exocytosis [1219]. TRPML2 and TRPML3 show increased channel activity in low luminal sodium and/or increased

luminal pH, and are activated by similar small molecules [206, 477, 1306]. A naturally occurring gain of function mutation in TRPML3 (*i.e.* A419P) results in the varitint waddler (*Va*) mouse phenotype (reviewed by [1022, 1146]).

Nomenclature	TRPML1	TRPML2	TRPML3
HGNC, UniProt	MCOLN1 , Q9GZU1	MCOLN2 , Q8IZK6	MCOLN3 , Q8TDD5
Endogenous activators	phosphatidyl (3,5) inositol bisphosphate (Also activates other TRPMLs) (pEC ₅₀ 7.3) [334]	–	–
Activators	ML SA1 (pEC ₅₀ 7.3) [-140mV] [1264], MK6-83 (pEC ₅₀ 7) [208], SF-22 (pEC ₅₀ 6.3) [-200mV] [208], ML-SA5 (pEC ₅₀ 5.3) [1586], SF-51 (pEC ₅₀ 4.5) [1264], ML1-SA1 [1306] TRPML1 ^{Va} : Constitutively active, current potentiated by extracellular acidification (equivalent to intralysosomal acidification)	ML SA1 Concentration range: 1×10^{-5} M [-140mV] [1264], phosphatidyl (3,5) inositol bisphosphate Concentration range: 1×10^{-6} M [-140mV] [334] TRPML2 ^{Va} : Constitutively active, current potentiated by extracellular acidification (equivalent to intralysosomal acidification)	SF-11 (pEC ₅₀ 6.6) [478], EVP-21 (pEC ₅₀ 4.3) [1170], ML SA1 Concentration range: 1×10^{-5} M [-140mV] [1264], phosphatidyl (3,5) inositol bisphosphate Concentration range: 1×10^{-6} M [-140mV] [334]
Selective activators	–	ML2-SA1 (Agonist) (pEC ₅₀ 1.2) [1131]	ML3-SA1 (pEC ₅₀ 5.1) [1306] – Mouse
Channel blockers	estradiol 3-methyl ether (pIC ₅₀ 6.7) [1200], PRU-12 (Inhibition) (pIC ₅₀ 6.6) [1200], estradiol 3-methyl ether (pIC ₅₀ 6.2) [1200]	–	Gd³⁺ (Antagonist) (pIC ₅₀ 4.7) [-80mV] [990] – Mouse
Functional Characteristics	TRPML1 ^{Va} : $\gamma = 40$ pS and 76-86 pS at very negative holding potentials with Fe ²⁺ and monovalent cations as charge carriers, respectively; conducts Na ⁺ \cong K ⁺ > Cs ⁺ and divalent cations (Ba ²⁺ > Mn ²⁺ > Fe ²⁺ > Ca ²⁺ > Mg ²⁺ > Ni ²⁺ > Co ²⁺ > Cd ²⁺ > Zn ²⁺ >> Cu ²⁺); monovalent cation flux suppressed by divalent cations (<i>e.g.</i> Ca ²⁺ , Fe ²⁺); inwardly rectifying	Conducts Na ⁺ ; monovalent cation flux suppressed by divalent cations; inwardly rectifying	TRPML3 ^{Va} : $\gamma = 49$ pS at very negative holding potentials with monovalent cations as charge carrier; conducts Na ⁺ > K ⁺ > Cs ⁺ with maintained current in the presence of Na ⁺ , conducts Ca ²⁺ and Mg ²⁺ , but not Fe ²⁺ , impermeable to protons; inwardly rectifying Wild type TRPML3: $\gamma = 59$ pS at negative holding potentials with monovalent cations as charge carrier; conducts Na ⁺ > K ⁺ > Cs ⁺ and Ca ²⁺ (P _{Ca} /P _K \cong 350), slowly inactivates in the continued presence of Na ⁺ within the extracellular (extracytosolic) solution; outwardly rectifying
Comments	TRPML1 current is potentiated by acidic pH and sphingosine [1264].	TRPML2 current is inhibited by intralysosomal acidification [1131].	Current is activated by Na ⁺ -free extracellular (extracytosolic) solution, and is inhibited by extracellular acidification (equivalent to intra-lysosomal acidification). Channel blockers include the ML-SI series of compounds (<i>e.g.</i> ML-S11; concentration range 5×10^{-5} M; -120mV) [940].

TRPP (polycystin) family

The TRPP family (reviewed by [306, 308, 452, 555, 1520]) or PKD2 family is comprised of PKD2 (PC2), PKD2L1 (PC2L1), PKD2L2 (PC2L2), which have been renamed TRPP1, TRPP2 and TRPP3, respectively [1529]. It should also be noted that the nomenclature

of PC2 was TRPP2 in old literature. However, PC2 has been uniformed to be called TRPP2 [516]. PKD2 family channels are clearly distinct from the PKD1 family, whose function is unknown. PKD1 and PKD2 form a hetero-oligomeric complex with a 1:3

ratio. [1333]. Although still being sorted out, TRPP family members appear to be 6TM spanning nonselective cation channels.

	TRPP1	TRPP2	TRPP3
Nomenclature	TRPP1	TRPP2	TRPP3
HGNC, UniProt	PKD2 , Q13563	PKD2L1 , Q9P0L9	PKD2L2 , Q9NZM6
Activators	–	Calmidazolium (in primary cilia): 10 μ M	–
Channel blockers	–	phenamil (pIC ₅₀ 6.9), benzamil (pIC ₅₀ 6), ethylisopropylamiloride (pIC ₅₀ 5), amiloride (pIC ₅₀ 3.8), Gd³⁺ Concentration range: 1 \times 10 ⁻⁴ M [-50mV] [218], La³⁺ Concentration range: 1 \times 10 ⁻⁴ M [-50mV] [218], flufenamate	–
Functional Characteristics	TRPP1 (PKD2) forms a cation channel (as a homomer consisting of 4 PKD2 subunits or as a heteromer combining 3 PKD2 subunits with one PKD1 subunit) that is expressed on primary cilia of kidney epithelial cells [847]. In kidney epithelial cells TRPP1 is only functional in the ciliary membrane, but not in the plasma membrane. In oocyte overexpression TRPP1 forms functional homomeric and heteromeric channels. Gain of function mutations in TRPP1 in either the S4-S5 linker (F604P) or in the lower gate (L677A, N681A) result in constitutively active channels [50, 495, 1492]. TRPP1 prefers monovalent cations over divalent cations in the order of K ⁺ >Na ⁺ >Ca ²⁺ (permeability 1:0.4:0.025), showing low selectivity for Ca ²⁺ . The conductance of TRPP1 varies depending on the ion (K ⁺ : 144 pS, Na ⁺ : 89 pS, Ca ²⁺ : 4pS) [847]. TRPP1 homomeric channel produces a larger conductance of 82 pS than the PC-1/TRPP1 heteromeric channel (79.5 pS) with higher absolute open probability (TRPP1 homomeric channel: 0.58, PC-1/TRPP1 heteromeric channel: 0.08) in primary cilia [495]. Specific activators or channel blockers of TRPP1 remain unknown.	TRPP2 is a nonselective cation channel functionally expressed on primary cilia and/or the plasma membrane depending on cell type. It can form a functional channel with PC1-L1 on primary cilia of retinal pigmented epithelial cells. TRPP2 (PKD2L1) displays calcium dependent activation. Calcium accumulation due to prolonged channel activity may lead to outward-moving Ca ²⁺ ions within the pore to close the channel [292]. TRPP2 permeates K ⁺ , Na ⁺ and Ca ²⁺ with the single-channel conductance of 189 pS for K ⁺ , 156 for Na ⁺ , and 53 pS for Ca ²⁺ , respectively [373]. PKD2L1 forms a heteromeric channel with PC-1L3 (PKD1L3) that may be activated by intracellular Ca ²⁺ [1332].	TRPP3 is not fully characterized yet. One report suggests the single channel activity of PKD2L2 in HEK293 cells as a 25 pS conductance [1344], but these recordings have not been confirmed.
Comments	Several studies have reported that TRPP1 forms heteromeric ion channels with other TRP channels such as TRPM3 and TRPM4, but the physiological significance of these potential heteromers remains unclear [407, 717]. TRPP1 has also been reported to function as a heteromeric channel with PC-1-L1 (PKD1L1) in the embryonic node, but the biophysical characteristics of this heteromeric channel have not yet been characterized [684].	–	–

TRPV (vanilloid) family

Members of the TRPV family (reviewed by [1427]) can broadly be divided into the non-selective cation channels, TRPV1-4 and the more calcium selective channels TRPV5 and TRPV6.

TRPV1-V4 subfamily

TRPV1 is involved in the development of thermal hyperalgesia following inflammation and may contribute to the detection of noxious heat (reviewed by [1118, 1308, 1351]). Numerous splice variants of TRPV1 have been described, some of which modulate the activity of TRPV1, or act in a dominant-negative man-

ner when co-expressed with TRPV1 [1247]. The pharmacology of TRPV1 channels is discussed in detail in [490] and [1445]. TRPV2 is probably not a thermosensor in man [1077], but has recently been implicated in innate immunity [834]. Functional TRPV2 expression is described in placental trophoblast cells of

mouse [286]. TRPV3 and TRPV4 are both thermosensitive. There are claims that TRPV4 is also mechanosensitive, but this has not been established to be within a physiological range in a native environment [177, 820].

TRPV5/V6 subfamily

TRPV5 and TRPV6 are highly expressed in placenta, bone, and kidney. Under physiological conditions, TRPV5 and TRPV6 are calcium selective channels involved in the absorption and reabsorption of calcium across intestinal and kidney tubule epithelia (reviewed by [287, 389, 987, 1518]). TRPV6 is reported to play a key role in calcium transport in the mouse placenta [1516].

Nomenclature	TRPV1	TRPV2
HGNC, UniProt	TRPV1, Q8NER1	TRPV2, Q9Y5S1
Other chemical activators	NO-mediated cysteine S-nitrosylation	–
Physical activators	depolarization ($V_{0.5} \sim 0$ mV at 35°C), noxious heat (> 43°C at pH 7.4)	–
Endogenous activators	extracellular H^+ (at 37°C) (pEC ₅₀ 5.4), 12S-HPETE (Agonist) (pEC ₅₀ 5.1) [-60mV] [595] – Rat, anandamide (pEC ₅₀ 5) [14], LTB ₄ (Agonist) (pEC ₅₀ 4.9) [-60mV] [595] – Rat, 5S-HETE	–
Activators	resiniferatoxin (Agonist) (pEC ₅₀ 8.4) [physiological voltage] [1287], capsaicin (Agonist) (pEC ₅₀ 7.5) [-100mV – 160mV] [1439], RhTx (pEC ₅₀ 6.3) [1562], piperine (Agonist) (pEC ₅₀ 4.4–5) [-70mV] [933], camphor, diphenylboronic anhydride, phenylacetylirvanil [47]	2-APB (pEC ₅₀ 5) [1004, 1148] – Rat, Δ^9 -tetrahydrocannabinol (pEC ₅₀ 4.8) [1148] – Rat, cannabidiol (pEC ₅₀ 4.5) [1148], probenecid (pEC ₅₀ 4.5) [73] – Rat, 2-APB (Agonist) (pEC ₅₀ 3.8–3.9) [physiological voltage] [578, 656] – Mouse, diphenylboronic anhydride (Agonist) Concentration range: 1×10^{-4} M [-80mV] [239, 656] – Mouse
Selective activators	olvaniil (Agonist) (pEC ₅₀ 7.7) [physiological voltage] [1287], DkTx (pEC ₅₀ 6.6) [physiological voltage] [135] – Rat	–
Channel blockers	5'-iodoresiniferatoxin (pIC ₅₀ 8.4), 6-iodo-nordihydrocapsaicin (pIC ₅₀ 8), AMG 9810 (Inhibition) (pIC ₅₀ 7.8) [physiological voltage] [437], BCTC (Antagonist) (pIC ₅₀ 7.5) [200], capsazepine (Antagonist) (pIC ₅₀ 7.4) [-60mV] [930], ruthenium red (pIC ₅₀ 6.7–7)	ruthenium red (pIC ₅₀ 6.2), tranilast (Inhibition) (pIC ₅₀ 4–5) [1010], SKF96365 (pIC ₅₀ 4) [656], TRIM (Inhibition) Concentration range: 5×10^{-4} M [656] – Mouse
Selective channel blockers	AMG517 (pIC ₅₀ 9) [128], AMG628 (pIC ₅₀ 8.4) [1470] – Rat, A425619 (pIC ₅₀ 8.3) [366], A778317 (pIC ₅₀ 8.3) [117], SB366791 (pIC ₅₀ 8.2) [492], JYL1421 (Antagonist) (pIC ₅₀ 8) [1489] – Rat, JNJ17203212 (Antagonist) (pIC ₅₀ 7.8) [physiological voltage] [1347], SB452533 (Antagonist) (p _B 7.7), SB705498 (Antagonist) (pIC ₅₀ 7.1) [491]	SET2 (pIC ₅₀ 6.3) [193], loratadine (Inhibition) (pIC ₅₀ 5.5) [1418]
Allosteric modulators (Positive)	s-RhTx (pEC ₅₀ 6.1) [1605], MRS-1477 [683]	–

Labelled ligands	[³H]A778317 (Channel blocker) (pK _d 8.5) [117], [¹²⁵ I] resiniferatoxin (Channel blocker, Antagonist) (pIC ₅₀ 8.4) [-50mV] [1453] – Rat, [³ H] resiniferatoxin (Activator)	–
Functional Characteristics	γ = 35 pS at - 60 mV; 77 pS at + 60 mV, conducts mono and di-valent cations with a selectivity for divalents (P _{Ca} /P _{Na} = 9.6); voltage- and time- dependent outward rectification; potentiated by ethanol; activated/potentiated/upregulated by PKC stimulation; extracellular acidification facilitates activation by PKC; desensitisation inhibited by PKA; inhibited by Ca ²⁺ / calmodulin; cooling reduces vanilloid-evoked currents; may be tonically active at body temperature	Conducts mono- and di-valent cations (P _{Ca} /P _{Na} = 0.9-2.9); dual (inward and outward) rectification; current increases upon repetitive activation by heat; translocates to cell surface in response to IGF-1 to induce a constitutively active conductance, translocates to the cell surface in response to membrane stretch. Cannabidiol sensitizes TRPV2 channels to activation by 2APB [458].
Nomenclature	TRPV3	TRPV4
HGNC, UniProt	TRPV3 , Q8NET8	TRPV4 , Q9HBA0
Other chemical activators	NO-mediated cysteine S-nitrosylation	Epoxyeicosatrienoic acids and NO-mediated cysteine S-nitrosylation
Physical activators	depolarization (V _{0.5} ~ +80 mV, reduced to more negative values following heat stimuli), heat (23°C - 39°C, temperature threshold reduces with repeated heat challenge)	Constitutively active, heat (> 24°C - 32°C), mechanical stimuli
Activators	incensole acetate (pEC ₅₀ 4.8) [978] – Mouse, 2-APB (Full agonist) (pEC ₅₀ 4.6) [-80mV – 80mV] [240] – Mouse, diphenylboronic anhydride (Full agonist) (pEC ₅₀ 4.1–4.2) [voltage dependent -80mV – 80mV] [239] – Mouse, thymol (Full agonist) (pEC ₅₀ 3.3) [1541] – Mouse, eugenol (Full agonist) (pEC ₅₀ 2.5) [1541] – Mouse, camphor (Full agonist) (pEC ₅₀ 2) [970] – Mouse, (-)-menthol (pEC ₅₀ 1.7) [-80mV – 80mV] [884] – Mouse, carvacrol (Full agonist) Concentration range: 5 × 10 ⁻⁴ M [-80mV – 80mV] [1541] – Mouse	phorbol 12-myristate 13-acetate (Agonist) (pEC ₅₀ 7.9) [physiological voltage] [1539], quinazolin-4(3H) derivative 36 (pEC ₅₀ 7.2) [57], curcumin (pEC ₅₀ 5.2) [1257], arachidonic acid (pEC ₅₀ 5) [1496] – Mouse, puerarin (pEC ₅₀ 4.8) [1626], vildagliptin (pEC ₅₀ 3) [424]
Selective activators	KS0365 (pEC ₅₀ 5.3) [890], 6-tert-butyl-m-cresol (pEC ₅₀ 3.4) [1443] – Mouse	GSK1016790A (pEC ₅₀ 8.7) [physiological voltage] [1387], 4α-PDH (pEC ₅₀ 7.1) [physiological voltage] [716] – Mouse, 4α-PDD (Agonist) (pEC ₅₀ 6.5) [1539], RN1747 (pEC ₅₀ 6.1) [physiological voltage] [1435], bisandrographolide (Agonist) (pEC ₅₀ 6) [-60mV] [1292] – Mouse
Inhibitors	–	GSK2798745 (pIC ₅₀ 8.8) [155], GSK-Bz derivative 2b (pIC ₅₀ 7.7) [16], paracetamol (pIC ₅₀ 6.5) [992], cimifugin (pIC ₅₀ 5.8) [1555], propofol (pIC ₅₀ 4.4) [1464], Cro-tamiton [714]
Channel blockers	forsythoside B (Inhibition) (pIC ₅₀ 6.2) [1607], ruthenium red (Inhibition) (pIC ₅₀ 6) [1096] – Mouse, diphenyltetrahydrofuran (Antagonist) (pIC ₅₀ 5–5.2) [-80mV – 80mV] [239] – Mouse, Citrusinine II (Inhibition) (pIC ₅₀ 4.9) [505], osthole (Inhibition) (pIC ₅₀ 4.4) [1340]	NSC151066 (Inhibition) (pIC ₅₀ 6.8) [331], ruthenium red (Inhibition) (pIC ₅₀ 6.7) [489] – Rat, Gd³⁺ , La³⁺
Selective channel blockers	Trpvicin (Inhibition) (pIC ₅₀ 6.4) [385], Isochlorogenic acid B (Inhibition) (pIC ₅₀ 6.1) [1145], Isochlorogenic acid A (Inhibition) (pIC ₅₀ 5.6) [1145], verbascoside (Inhibition) (pIC ₅₀ 4.8) [1338]	HC067047 (Inhibition) (pIC ₅₀ 7.3) [-40mV] [382], RN-9893 (Antagonist) (pIC ₅₀ 6.2) [1504], RN1734 (Inhibition) (pIC ₅₀ 5.6) [physiological voltage] [1435]
Functional Characteristics	γ = 197 pS at = +40 to +80 mV, 48 pS at negative potentials; conducts mono- and di-valent cations; outward rectification; potentiated by arachidonic acid	γ = ~60 pS at -60 mV, ~90-100 pS at +60 mV; conducts mono- and di-valent cations with a preference for divalents (P _{Ca} /P _{Na} =6-10); dual (inward and outward) rectification; potentiated by intracellular Ca ²⁺ via Ca ²⁺ / calmodulin; inhibited by elevated intracellular Ca ²⁺ via an unknown mechanism (IC ₅₀ = 0.4 μM)

Nomenclature	TRPV5	TRPV6
HGNC, UniProt	TRPV5, Q9NQA5	TRPV6, Q9H1D0
Other channel blockers	$Pb^{2+} = Cu^{2+} = Gd^{3+} > Cd^{2+} > Zn^{2+} > La^{3+} > Co^{2+} > Fe^{2+}$	–
Activators	constitutively active (with strong buffering of intracellular Ca^{2+})	acetaldehyde (pEC ₅₀ 6.7) [935], ethanol (pEC ₅₀ 0.8) [935], 2-APB (Potentiation) constitutively active (with strong buffering of intracellular Ca^{2+})
Inhibitors	gentamicin (pIC ₅₀ 6) [1419], tetrahydrocannabivarin (pIC ₅₀ 5.4) [622], oxoglaucine (pIC ₅₀ 4.8) [1621]	GSK3527497 (pIC ₅₀ 7.9) [154], SOR-C13 (pIC ₅₀ 7.8) [144], TRPV6 inhibitor cis-22 a (pIC ₅₀ 6.5) [1280], tetrahydrocannabivarin (pIC ₅₀ 5.1) [622]
Channel blockers	ruthenium red (pIC ₅₀ 6.9), Mg^{2+}	ruthenium red (Antagonist) (pIC ₅₀ 5) [-80mV] [554] – Mouse, Cd^{2+} , La^{3+} , Mg^{2+}
Functional Characteristics	$\gamma = 59\text{--}78$ pS for monovalent ions at negative potentials, conducts mono- and di-valents with high selectivity for divalents ($P_{Ca}/P_{Na} > 107$); voltage- and time- dependent inward rectification; inhibited by intracellular Ca^{2+} promoting fast inactivation and slow down-regulation; feedback inhibition by Ca^{2+} reduced by calcium binding protein 80-K-H; inhibited by extracellular and intracellular acidosis; upregulated by 1,25-dihydrovitamin D3	$\gamma = 58\text{--}79$ pS for monovalent ions at negative potentials, conducts mono- and di-valents with high selectivity for divalents ($P_{Ca}/P_{Na} > 130$); voltage- and time-dependent inward rectification; inhibited by intracellular Ca^{2+} promoting fast and slow inactivation; gated by voltage-dependent channel blockade by intracellular Mg^{2+} ; slow inactivation due to Ca^{2+} -dependent calmodulin binding; phosphorylation by PKC inhibits Ca^{2+} -calmodulin binding and slow inactivation; upregulated by 1,25-dihydroxyvitamin D3

Comments:**TRPA (ankyrin) family**

Agents activating TRPA1 in a covalent manner are thiol reactive electrophiles that bind to cysteine and lysine residues within the cytoplasmic domain of the channel [547, 882]. TRPA1 is activated

by a wide range of endogenous and exogenous compounds and only a few representative examples are mentioned in the table: an exhaustive listing can be found in [75]. In addition, TRPA1 is

potently activated by intracellular zinc ($EC_{50} = 8$ nM) [42, 577]. A gain-of-function mutation in TRPA1 was found to cause familial episodic pain syndrome [746].

TRPM (melastatin) family

Ca^{2+} activates all splice variants of TRPM2, but other activators listed are effective only at the full length isoform [345]. Inhibition of TRPM2 by clotrimazole, miconazole, econazole, flufenamic acid is largely irreversible. Co-application of pregnenolone sulphate and clotrimazole caused TRPM3 currents to acquire an inwardly rectifying component at negative voltages, resulting in a biphasic conductance-voltage relationship. Evidence was presented that the inward current might reflect the permeation of cations through the opening of a non-canonical pore [1446]. TRPM3 activity is impaired in chronic fatigue syndrome/myalgic encephalomyelitis

patients suggesting changes in intracellular Ca^{2+} concentration, which may impact natural killer cellular functions [167]. TRPM4 exists as multiple splice variants: data listed are for TRPM4b. The sensitivity of TRPM4b and TRPM5 to activation by $[Ca^{2+}]_i$ demonstrates a pronounced and time-dependent reduction following excision of inside-out membrane patches [1413]. The $V_{0.5}$ for activation of TRPM4 and TRPM5 demonstrates a pronounced negative shift with increasing temperature. Activation of TRPM8 by depolarization is strongly temperature-dependent via a channel-closing rate that decreases with decreasing temperature. The $V_{0.5}$ is shifted

in the hyperpolarizing direction both by decreasing temperature and by exogenous agonists, such as (-)-menthol [1439] whereas antagonists produce depolarizing shifts in $V_{0.5}$ [901]. The $V_{0.5}$ for the native channel is far more positive than that of heterologously expressed TRPM8 [901]. It should be noted that (-)-menthol and structurally related compounds can elicit release of Ca^{2+} from the endoplasmic reticulum independent of activation of TRPM8 [889]. Intracellular pH modulates activation of TRPM8 by cold and icilin, but not (-)-menthol [40].

TRPML (mucolipin) family

Data in the table are for TRPML proteins mutated (*i.e.* TRPML1^{Va}, TRPML2^{Va} and TRPML3^{Va}) at loci equivalent to TRPML3 A419P to allow plasma membrane expression when expressed in HEK-293 cells and subsequent characterisation by patch-clamp recording [333, 475, 702, 990, 1542]. Data for wild type TRPML3 are also

tabulated [702, 703, 990, 1542]. It should be noted that alternative methodologies, particularly in the case of TRPML1, have resulted in channels with differing biophysical characteristics (reviewed by [1140]). Initial functional characteristics of TRPML channels are performed on their Va mutations of TRPMLs at loci

equivalent to TRPML3 A419P. Current pharmacological characterization of channel activators and blockers are conducted on wild-type channel proteins using endolysosomal patch-clamp [208, 334, 1131, 1264].

TRPP (polycystin) family

Data in the table are extracted from [272, 308] and [1269]. Broadly similar single channel conductance, mono- and di-valent cation selectivity and sensitivity to blockers are observed for TRPP2 co-expressed with TRPP1 [307]. Ca²⁺, Ba²⁺ and Sr²⁺ permeate TRPP3, but reduce inward currents carried by Na⁺. Mg²⁺ is largely impermeant and exerts a voltage dependent inhibition that increases with hyperpolarization.

TRPV (vanilloid) family

Activation of TRPV1 by depolarisation is strongly temperature-dependent via a channel opening rate that increases with increasing temperature. The V_{0.5} is shifted in the hyperpolarizing direction both by increasing temperature and by exogenous agonists [1439]. TRPV3 channel dysfunction caused by genetic gain-of-function mutations is implicated in the pathogenesis of skin inflammation, dermatitis, and chronic itch. In rodents, a spontaneous gain-of-function mutation of the TRPV3 gene causes the development of skin lesions with pruritus and dermatitis [51, 830]. In contrast to other thermoTRP channels, TRPV3 sensitizes rather than desensitizes, upon repeated stimulation with either heat or agonists [240, 839, 1543]. The sensitivity of TRPV4 to heat, but not 4 α -PDD is lost upon patch excision. TRPV4 is activated by anandamide and arachidonic acid following P450

epoxygenase-dependent metabolism to 5,6-epoxyeicosatrienoic acid (reviewed by [1026]). Activation of TRPV4 by cell swelling, but not heat, or phorbol esters, is mediated via the formation of epoxyeicosatrienoic acids. Phorbol esters bind directly to TRPV4. Different TRPV4 mutations lead to a broad spectrum of dominant skeletal dysplasias [739, 1182] and spinal muscular atrophies and hereditary motor and sensory neuropathies [59, 311]. Similar mutations were also found in patients with Charcot-Marie-Tooth disease type 2C [768]. TRPV5 preferentially conducts Ca²⁺ under physiological conditions, but in the absence of extracellular Ca²⁺, conducts monovalent cations. Single channel conductances listed for TRPV5 and TRPV6 were determined in divalent cation-free extracellular solution. Ca²⁺-induced inactivation occurs at hyperpolarized potentials when Ca²⁺ is present extracellularly. Sin-

gle channel events cannot be resolved (probably due to greatly reduced conductance) in the presence of extracellular divalent cations. Measurements of P_{Ca}/P_{Na} for TRPV5 and TRPV6 are dependent upon ionic conditions due to anomalous mole fraction behaviour. Blockade of TRPV5 and TRPV6 by extracellular Mg²⁺ is voltage-dependent. Intracellular Mg²⁺ also exerts a voltage dependent block that is alleviated by hyperpolarization and contributes to the time-dependent activation and deactivation of TRPV6 mediated monovalent cation currents. TRPV5 and TRPV6 differ in their kinetics of Ca²⁺-dependent inactivation and recovery from inactivation. TRPV5 and TRPV6 function as homo- and hetero-tetramers.

Further reading on Transient Receptor Potential channels (TRP)

- Aghazadeh Tabrizi M *et al.* (2017) Medicinal Chemistry, Pharmacology, and Clinical Implications of TRPV1 Receptor Antagonists. *Med Res Rev* **37**: 936-983 [PMID:27976413]
- Basso L *et al.* (2017) Transient Receptor Potential Channels in neuropathic pain. *Curr Opin Pharmacol* **32**: 9-15 [PMID:27835802]
- Ciarlo MG *et al.* (2017) Lipids as central modulators of sensory TRP channels. *Biochim Biophys Acta* **1859**: 1615-1628 [PMID:28432033]
- Clapham DE *et al.* (2003) International Union of Pharmacology. XLIII. Compendium of voltage-gated ion channels: transient receptor potential channels. *Pharmacol Rev* **55**: 591-6 [PMID:14657417]
- Diaz-Franulic I *et al.* (2016) Allosterism and Structure in Thermally Activated Transient Receptor Potential Channels. *Annu Rev Biophys* **45**: 371-98 [PMID:27297398]
- Emir TLR. (2017) Neurobiology of TRP Channels *Neurobiology of TRP Channels*[PMID:29356487]
- Grayson TH *et al.* (2017) Transient receptor potential canonical type 3 channels: Interactions, role and relevance - A vascular focus. *Pharmacol Ther* **174**: 79-96 [PMID:28223224]
- Nilius B *et al.* (2014) Mammalian transient receptor potential (TRP) cation channels. Preface. *Handb Exp Pharmacol* **223**: v - vi [PMID:25296415]
- Rubaiy HN. (2019) Treasure troves of pharmacological tools to study transient receptor potential canonical 1/4/5 channels. *Br J Pharmacol* **176**: 832-846 [PMID:30656647]
- Wu LJ *et al.* (2010) International Union of Basic and Clinical Pharmacology. LXXXVI. Current progress in the mammalian TRP ion channel family. *Pharmacol Rev* **62**: 381-404 [PMID:20716668]
- Zhu MX. (2011) Various TRP Channels (CRC Press/Taylor & Francis)[PMID:22593967]
- Zierler S *et al.* (2017) TRPM channels as potential therapeutic targets against pro-inflammatory diseases. *Cell Calcium* **67**: 105-115 [PMID:28549569]

Voltage-gated calcium channels (Ca_v)

Ion channels → Voltage-gated ion channels → Voltage-gated calcium channels (Ca_v)

Overview: Ca²⁺ channels are voltage-gated ion channels present in the membrane of most excitable cells. The nomenclature for Ca²⁺ channels was proposed by [372] and **approved by the NC-IUPHAR Subcommittee on Ca²⁺ channels** [188]. Most Ca²⁺ channels form hetero-oligomeric complexes. The α1 subunit is pore-forming and provides the binding site(s) for practically all agonists and antagonists. The 10 cloned α1-subunits can be grouped into three families: (1) the high-voltage activated dihydropyridine-sensitive (L-type, Ca_v1.x) channels; (2) the

high- to moderate-voltage activated dihydropyridine-insensitive (Ca_v2.x) channels and (3) the low-voltage-activated (T-type, Ca_v3.x) channels. Each α1 subunit has four homologous repeats (I-IV), each repeat having six transmembrane domains (S1-S6) forming a voltage-sensing domain (VSD, S1-S4) coupled to a pore-forming module (S5, S6 and their connecting linker that contains the selectivity filter. Voltage-dependent gating is driven by voltage-induced transmembrane movements of the S4-helix enabled by conserved positive charges interacting with negative

counter-charges within the VSD [186]. All of the α1-subunit genes give rise to alternatively spliced products. At least for high-voltage activated channels, it is likely that native channels comprise co-assemblies of α1, β and α2-δ subunits. CACHD1 is an α2δ-like protein that modulates Ca_v3 channel activity [257]. The γ subunits have not been proven to associate with channels other than the α1s skeletal muscle Ca_v1.1 channel. The α2-δ1 and α2-δ2 subunits bind **gabapentin** and **pregabalin** [221].

Nomenclature	Ca _v 1.1	Ca _v 1.2	Ca _v 1.3	Ca _v 1.4
HGNC, UniProt	<i>CACNA1S</i> , Q13698	<i>CACNA1C</i> , Q13936	<i>CACNA1D</i> , Q01668	<i>CACNA1F</i> , O60840
Activators	FPL64176 (pEC ₅₀ ~7.8), (-)-(S)-BayK8644 (pEC ₅₀ ~7.8)	(-)-(S)-BayK8644 (pEC ₅₀ ~7.8), FPL64176 Concentration range: 1 × 10 ⁻⁷ M-1 × 10 ⁻⁶ M [669, 843]	FPL64176 (pEC ₅₀ ~7.8), (-)-(S)-BayK8644 (pEC ₅₀ ~7.8)	(-)-(S)-BayK8644 (pEC ₅₀ ~7.8)
Gating inhibitors	nifedipine (Antagonist) (pIC ₅₀ 6.3) [voltage dependent -90mV] [765] – Rat, nimodipine (Antagonist) (pIC ₅₀ ~6) [-70mV]	amlodipine (pIC ₅₀ 9.3) [589] – Rabbit, isradipine (Antagonist) (pIC ₅₀ 8.8) [1059], nifedipine (Antagonist) (pIC ₅₀ 8.1–8.7) [-40mV] [1102, 1117] – Rat, isradipine (Antagonist) (pIC ₅₀ 8.5) [1059], nimodipine (Antagonist) (pIC ₅₀ 6.8) [-80mV] [1547] – Rat	nitrendipine (Inhibition) (pIC ₅₀ 8.4) [1283], isradipine (dopamine neuron-like activity; splice variant-dependent) (pIC ₅₀ 7.8–8.2) [1059], nifedipine (Antagonist) (pIC ₅₀ 7.7) [1283], nimodipine (Antagonist) (pIC ₅₀ 5.7–6.6) [-80mV – -40mV] [1210, 1547] – Rat	nifedipine (Antagonist) (pIC ₅₀ 6) [-100mV] [934], nimodipine (Antagonist) (pIC ₅₀ ~6) [-70mV], nitrendipine (Antagonist) (pIC ₅₀ ~6) [-70mV]
Channel blockers	verapamil (Antagonist) (pIC ₅₀ ~5) [100mV] [1461] – Rabbit, diltiazem (Antagonist) (pIC ₅₀ 4.2) [-100mV] [1461] – Rabbit	verapamil (Antagonist) (pIC ₅₀ 5.3–6.5) [645] – Rat, diltiazem (Antagonist) (pIC ₅₀ 6.3) [666] – Ferret	verapamil (Antagonist) (pIC ₅₀ 3.7) [-70mV] [1370] – Mouse, diltiazem (pIC ₅₀ 3.5) [-70mV] [1370] – Mouse	diltiazem (pIC ₅₀ 4) [-80mV] [91] – Mouse, verapamil Concentration range: 1 × 10 ⁻⁴ M [-80mV] [91] – Mouse
Sub/family-selective channel blockers	–	calciseptine (Antagonist) (pIC ₅₀ 7.1–7.8) [86, 291, 939, 1498]	–	–
Functional Characteristics	L-type calcium current: High voltage-activated, very slow voltage dependent inactivation	L-type calcium current: High voltage-activated, voltage- and calcium-dependent inactivation	L-type calcium current: more negative activation voltage range than Ca _v 1.2, calcium-dependent inactivation	L-type calcium current: More negative activation voltage range than Ca _v 1.2, no/weak calcium-dependent inactivation

Comments	Serves primarily as voltage-sensor for excitation contraction coupling in skeletal muscle. The adaptor protein STAC3 binds to Ca _v 1.1 α1 and is essential for transmitting its conformational changes to ryanodine receptor-mediated excitation contraction coupling [1133].	Amlodipine, isradipine, nifedipine and nimodipine are examples of dihydropyridine calcium channel antagonists. Verapamil is a phenylalkylamine calcium channel blocker. Diltiazem is an example of a benzothiazepine calcium channel blocker. Inhibition by dihydropyridines (e.g. nifedipine or isradipine) is voltage-dependent with a higher apparent affinity at more depolarised potentials [1352]; phenylalkylamines and diltiazem exhibit strong use-dependence with a higher apparent affinity at higher stimulation frequencies. Calciseptin is a potent inhibitor of Ca _v 1.2 but not of Ca _v 1.3 [939].	Ca _v 1.3 is about 5-10 fold less sensitive to dihydropyridine antagonists [1352].	Ca _v 1.4 is less sensitive to dihydropyridine antagonists than other Ca _v 1 channels
Nomenclature	Ca _v 2.1	Ca _v 2.2	Ca _v 2.3	
HGNC, UniProt	CACNA1A, O00555	CACNA1B, Q00975	CACNA1E, Q15878	
Gating inhibitors	–	NP118809 (pIC ₅₀ 7) [-80mV] [1594] – Rat	–	
Selective gating inhibitors	ω-agatoxin IVA (Antagonist) (pIC ₅₀ 7–8.7) [-100mV – -90mV] [142, 960] – Rat, ω-agatoxin IVB (Antagonist) (pK _d 8.5) [-80mV] [12] – Rat	C2230 (pIC ₅₀ 6) [-50mV] [1365]	SNX482 (Antagonist) (pIC ₅₀ 7.5–8) [physiological voltage] [1005]	
Channel blockers	–	–	Ni ²⁺ (Antagonist) (pIC ₅₀ 4.6) [-90mV] [1515]	
Sub/family-selective channel blockers	ω-conotoxin MVIIC (Antagonist) (pIC ₅₀ 8.2–9.2) Concentration range: 2 × 10 ⁻⁶ M–5 × 10 ⁻⁶ M [physiological voltage] [803] – Rat	ω-conotoxin GVIA (Antagonist) (pIC ₅₀ 10.4) [-80mV] [803] – Rat, ω-conotoxin MVIIC (Antagonist) (pIC ₅₀ 6.1–8.5) [-80mV] [546, 803, 929] – Rat	–	
Functional Characteristics	P/Q-type calcium current: High voltage-activated, moderate voltage-dependent inactivation	N-type calcium current: High voltage-activated, moderate voltage-dependent inactivation	R-type calcium current: Moderate voltage-activated, fast voltage-dependent inactivation	
Comments	All three Ca _v 2.x types directly contribute towards triggering neurotransmitter release at fast synapses in the mammalian nervous system. In many cell types, P- and Q-current components cannot be adequately separated and many researchers in the field have therefore adopted the terminology 'P/Q-type' current when referring to either component. Both of these physiologically defined current types are conducted by alternative forms of Ca _v 2.1 [815]. Ziconotide (a synthetic peptide equivalent to omega-Conotoxin-MVIIA) has been approved for the treatment of chronic pain [1514].	–	–	

Nomenclature	Ca_v3.1	Ca_v3.2	Ca_v3.3
HGNC, UniProt	CACNA1G, O43497	CACNA1H, O95180	CACNA1I, Q9P0X4
Gating inhibitors	kurtoxin (Antagonist) (pIC ₅₀ 7.3–7.8) [-90mV] [236 , 1273] – Rat	kurtoxin (Antagonist) (pIC ₅₀ 7.3–7.6) [-90mV] [236 , 1273] – Rat	–
Channel blockers	ACT-709478 (Inhibition) (pIC ₅₀ 8.2) [114], ulixacaltamide (Pore blocker) (pIC ₅₀ 7.3) [-80mV] [1406], TTA-A2 (Pore blocker) (pIC ₅₀ 7) [-75mV] [413], mibefradil (Antagonist) (pIC ₅₀ 6–6.6) [-110mV – -100mV] [914], Ni²⁺ (Antagonist) (pIC ₅₀ 3.6–3.8) [voltage dependent -90mV] [782] – Rat	TTA-A2 (Pore blocker) (pIC ₅₀ 8) [-75mV] [413], ACT-709478 (Inhibition) (pIC ₅₀ 7.7) [114], mibefradil (Pore blocker) (pIC ₅₀ 5.9–7.2) [-110mV – -80mV] [914], and derivatives pimozide (Pore blocker) (pIC ₅₀ 6.8) [679], ulixacaltamide (Pore blocker) (pIC ₅₀ 6.8) [-75mV] [1406], efonidipine (Pore blocker) (pIC ₅₀ 6.4) [788], ABT-639 (Pore blocker) (pIC ₅₀ 5.7) [-70mV] [46 , 625], Ni²⁺ (Pore blocker) (pIC ₅₀ 4.9–5.2) [voltage dependent -90mV] [782]	ACT-709478 (Inhibition) (pIC ₅₀ 8.1) [114], TTA-A2 (Pore blocker) (pIC ₅₀ 7.5) [-75mV] [413], ulixacaltamide (Pore blocker) (pIC ₅₀ 7) [-75mV] [1406], mibefradil (Antagonist) (pIC ₅₀ 5.8) [-110mV] [914], Ni²⁺ (Antagonist) (pIC ₅₀ 3.7–4.1) [voltage dependent -90mV] [782] – Rat
Functional Characteristics	T-type calcium current: Low voltage-activated, fast voltage-dependent inactivation	T-type calcium current: Low voltage-activated, fast voltage-dependent inactivation	T-type calcium current: Low voltage-activated, moderate voltage-dependent inactivation

Further reading on Voltage-gated calcium channels (Ca_v)

- Catterall WA. (2023) Voltage gated sodium and calcium channels: Discovery, structure, function, and Pharmacology. *Channels (Austin)* **17**: 2281714 [[PMID:37983307](#)]
- Catterall WA *et al.* (2005) International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev* **57**: 411-25 [[PMID:16382099](#)]
- Catterall WA *et al.* (2015) Deciphering voltage-gated Na(+) and Ca(2+) channels by studying prokaryotic ancestors. *Trends Biochem Sci* **40**: 526-34 [[PMID:26254514](#)]
- Chen Z *et al.* (2023) EMC chaperone-Ca_v structure reveals an ion channel assembly intermediate. *Nature* **619**: 410-419 [[PMID:37196677](#)]
- Dolphin AC. (2018) Voltage-gated calcium channels: their discovery, function and importance as drug targets. *Brain Neurosci Adv* **2**: [[PMID:30320224](#)]
- Filippini L *et al.* (2023) Ca_v 1.3-selective inhibitors of voltage-gated L-type Ca²⁺ channels: Fact or (still) fiction? *Br J Pharmacol* **180**: 1289-1303 [[PMID:36788128](#)]
- Huang J *et al.* (2017) Regulation of voltage gated calcium channels by GPCRs and post-translational modification. *Curr Opin Pharmacol* **32**: 1-8 [[PMID:27768908](#)]
- Nanou E *et al.* (2018) Calcium Channels, Synaptic Plasticity, and Neuropsychiatric Disease. *Neuron* **98**: 466-481 [[PMID:29723500](#)]
- Papa A *et al.* (2022) Adrenergic Regulation of Calcium Channels in the Heart. *Annu Rev Physiol* **84**: 285-306 [[PMID:34752709](#)]
- Pelizzari S *et al.* (2024) Ca_v1.1 voltage-sensing domain III exclusively controls skeletal muscle excitation-contraction coupling. *Nat Commun* **15**: 7440 [[PMID:39198449](#)]
- Rougier JS *et al.* (2016) Cardiac voltage-gated calcium channel macromolecular complexes. *Biochim Biophys Acta* **1863**: 1806-12 [[PMID:26707467](#)]
- Weiss N *et al.* (2024) The T-type calcium channelosome. *Pflugers Arch* **476**: 163-177 [[PMID:38036777](#)]
- Yao X *et al.* (2024) Structural biology of voltage-gated calcium channels. *Channels (Austin)* **18**: 2290807 [[PMID:38062897](#)]
- Zamponi GW. (2016) Targeting voltage-gated calcium channels in neurological and psychiatric diseases. *Nat Rev Drug Discov* **15**: 19-34 [[PMID:26542451](#)]
- Zamponi GW *et al.* (2015) The Physiology, Pathology, and Pharmacology of Voltage-Gated Calcium Channels and Their Future Therapeutic Potential. *Pharmacol Rev* **67**: 821-70 [[PMID:26362469](#)]

Voltage-gated proton channel (H_v1)

[Ion channels](#) → [Voltage-gated ion channels](#) → [Voltage-gated proton channel \(H_v1\)](#)

Overview: The voltage-gated proton channel (provisionally denoted H_v1) is a putative 4TM proton-selective channel gated by membrane depolarization and which is sensitive to the transmembrane pH gradient [[178](#), [294](#), [295](#), [1165](#), [1225](#)]. The structure of H_v1 is homologous to the voltage-sensing domain (VSD) of the superfamily of voltage-gated ion channels (*i.e.* segments S1 to S4) and contains

no discernable pore region [[1165](#), [1225](#)]. Proton flux through H_v1 is instead most likely mediated by a hydrogen-bonded chain [[296](#), [991](#)] formed in a crevice of the protein when the voltage-sensing S4 helix moves in response to a change in transmembrane potential [[1164](#), [1523](#)]. Proton selective conduction requires an aspartate residue at the center of the pore [[202](#), [986](#), [1294](#)]. Both selectivity and

conduction may result from obligatory protonation by each conducted proton [[297](#), [349](#)]. H_v1 expresses largely as a dimer mediated by intracellular C-terminal coiled-coil interactions [[811](#)] but individual promoters nonetheless support gated H⁺ flux *via* separate conduction pathways [[726](#), [787](#), [1108](#), [1393](#)]. Within dimeric structures, the two protomers do not function independently, but dis-

Searchable database: <http://www.guidetopharmacology.org/>

Full Contents of ConciseGuide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.70231/full>

Voltage-gated proton channel (Hv1) S215

play co-operative interactions during gating resulting in increased voltage sensitivity, but slower activation, of the dimeric, *versus* monomeric, complexes [462, 1394]. The otopetrin proteins appear to form proton-selective ion channels and to date 3 subtypes have been identified in eukaryotes; otopetrin 1 [1163, 1409], otopetrin 2 [856] and otopetrin 3 [591].

Nomenclature	H _v 1
HGNC, UniProt	HVCN1, Q96D96
Channel blockers	Zn ²⁺ (pIC ₅₀ ~5.7–6.3), Cd ²⁺ (pIC ₅₀ ~5)
Functional Characteristics	Activated by membrane depolarization mediating macroscopic currents with time-, voltage- and pH-dependence; outwardly rectifying; voltage dependent kinetics with relatively slow current activation sensitive to extracellular pH and temperature, relatively fast deactivation; voltage threshold for current activation determined by pH gradient ($\Delta\text{pH} = \text{pH}_o - \text{pH}_i$) across the membrane

Comments: The voltage threshold (V_{thr}) for activation of H_v1 is not fixed but is set by the pH gradient across the membrane such that V_{thr} is positive to the Nernst potential for H⁺, which ensures that only outwardly directed flux of H⁺ occurs under physiological conditions [178, 227, 294, 295]. Phosphorylation of H_v1 within the N-terminal domain by PKC enhances the gating of the channel [298, 299, 984]. Tabulated IC₅₀ values for Zn²⁺ and Cd²⁺

are for heterologously expressed human and mouse H_v1 [1165, 1225]. Zn²⁺ is not a conventional pore blocker, but is coordinated by two, or more, external protonation sites involving histamine residues [1165]. Zn²⁺ binding may occur at the dimer interface between pairs of histamine residues from both monomers where it may interfere with channel opening [985]. Mouse knockout studies [365, 973, 1166] support the view that H_v1 participates

in both charge compensation and pH regulation in granulocytes during the respiratory burst of NADPH oxidase-dependent reactive oxygen species production that assists in the clearance of bacterial pathogens [300, 531, 673, 1243]. Additional physiological functions of H_v1 are reviewed by [178].

Further reading on Voltage-gated proton channel (H_v1)

Castillo K *et al.* (2015) Voltage-gated proton (H(v)1) channels, a singular voltage sensing domain. *FEBS Lett* **589**: 3471-8 [PMID:26296320]
 DeCoursey TE. (2018) Gating currents indicate complex gating of voltage-gated proton channels. *Proc Natl Acad Sci USA* **115**: 9057-9059 [PMID:30135099]
 DeCoursey TE. (2018) Voltage and pH sensing by the voltage-gated proton channel, H_v1. *J R Soc Interface* **15**: [PMID:29643227]

Fernández A *et al.* (2016) Pharmacological Modulation of Proton Channel Hv1 in Cancer Therapy: Future Perspectives. *Mol Pharmacol* **90**: 385-402 [PMID:27260771]
 Okamura Y *et al.* (2015) Gating mechanisms of voltage-gated proton channels. *Annu Rev Biochem* **84**: 685-709 [PMID:26034892]

Voltage-gated sodium channels (Na_v)

Ion channels → **Voltage-gated ion channels** → **Voltage-gated sodium channels (Na_v)**

Overview: Sodium channels are voltage-gated sodium-selective ion channels present in the membrane of most excitable cells. Sodium channels comprise one pore-forming α subunit, which may be associated with either one or two β subunits [611]. α -Subunits consist of four homologous domains (I-IV), each containing six transmembrane segments (S1-S6) and a pore-forming loop. The positively charged fourth transmembrane segment (S4) acts as a voltage sensor and is involved in channel gating. The crystal structure of the bacterial NavAb channel has revealed a number of novel structural features compared to earlier potassium chan-

nel structures including a short selectivity filter with ion selectivity determined by interactions with glutamate side chains [1088]. Interestingly, the pore region is penetrated by fatty acyl chains that extend into the central cavity which may allow the entry of small, hydrophobic pore-blocking drugs [1088]. Auxiliary β 1, β 2, β 3 and β 4 subunits consist of a large extracellular N-terminal domain, a single transmembrane segment and a shorter cytoplasmic domain. Pharmacological targeting of voltage-gated sodium channels has long been a cornerstone of clinical treatment for a range of conditions. Classical sodium channel blockers, many of

which act by occluding the central pore, are widely used as local anesthetics, antiarrhythmic agents, and anticonvulsants [412]. More recently, suzetrigine, a highly selective Na_v1.8 inhibitor, received FDA approval for the treatment of acute post-operative pain [652, 1499].

The nomenclature for sodium channels was proposed by Goldin *et al.*, (2000) [459] and approved by the NC-IUPHAR Subcommittee on sodium channels (Catterall *et al.*, 2005, [185]).

Nomenclature	Na _v 1.1	Na _v 1.2	Na _v 1.3	Na _v 1.4	Na _v 1.5
HGNC, UniProt	SCN1A, P35498	SCN2A, Q99250	SCN3A, Q9NY46	SCN4A, P35499	SCN5A, Q14524
Sub/family-selective activators	batrachotoxin, veratridine	batrachotoxin (Agonist) (pK _d 9.1) [physiological voltage] [831] – Rat, veratridine (Partial agonist) (pK _d 5.2) [physiological voltage] [187] – Rat	batrachotoxin, veratridine	batrachotoxin (Full agonist) Concentration range: 5 × 10 ⁻⁶ M [-100mV] [1481] – Rat, veratridine (Partial agonist) Concentration range: 2 × 10 ⁻⁴ M [-100mV] [1481] – Rat	batrachotoxin (Full agonist) (pK _d 7.6) [physiological voltage] [1262] – Rat, veratridine (Partial agonist) (pEC ₅₀ 6.3) [-30mV] [1465] – Rat
Channel blockers	tetrodotoxin (Pore blocker) (pK _d 8) [-100mV] [1293] – Rat, cannabidiol (pIC ₅₀ 5.5) [-70mV] [449], cannabidiol (pIC ₅₀ 4.9) [-100mV] [449]	cannabidiol (pIC ₅₀ 5.5) [-45mV] [449]	cannabidiol (pIC ₅₀ 5.5) [-45mV] [449]	cannabidiol (pIC ₅₀ 5.7) [-45mV] [449], cannabidiol (pIC ₅₀ 5) [-110mV] [446]	cannabidiol (pIC ₅₀ 5.4) [-60mV] [449]
Sub/family-selective channel blockers	ICA-121431 (pIC ₅₀ 7.7) [928], Hm1a [1062] – Rat, saxitoxin (Pore blocker)	saxitoxin (Pore blocker) (pIC ₅₀ 8.8) [-120mV] [150] – Rat, tetrodotoxin (Pore blocker) (pIC ₅₀ 8) [-120mV] [150] – Rat, XPC-5462 (pIC ₅₀ 8) [-45mV] [463], lacosamide (Antagonist) (pIC ₅₀ 4.5) [-80mV] [3] – Rat	tetrodotoxin (Pore blocker) (pIC ₅₀ 8.4) [220], ICA-121431 (pIC ₅₀ 7.7) [928], saxitoxin (Pore blocker)	saxitoxin (Pore blocker) (pIC ₅₀ 8.4) [-100mV] [1101] – Rat, tetrodotoxin (Pore blocker) (pIC ₅₀ 7.6) [-120mV] [192], μ-conotoxin GIIIA (Pore blocker) (pIC ₅₀ 5.9) [-100mV] [192]	tetrodotoxin (Pore blocker) (pK _d 5.8) [-80mV] [260, 1597] – Rat
Functional Characteristics	Activation V _{0.5} = -20 mV. Fast inactivation (τ = 0.7 ms for peak sodium current).	Activation V _{0.5} = -24 mV. Fast inactivation (τ = 0.8 ms for peak sodium current).	Activation V _{0.5} = -24 mV. Fast inactivation (0.8 ms)	Activation V _{0.5} = -30 mV. Fast inactivation (0.6 ms)	Activation V _{0.5} = -26 mV. Fast inactivation (τ = 1 ms for peak sodium current).
Comments	–	The anti-sense oligonucleotide <i>elsunersen</i> is designed to reduce expression of gain-of-function SCN2A, as a proposed therapy for SCN2A-driven early-onset developmental and epileptic encephalopathy (DEE) [809].	–	–	–

Nomenclature	Na _v 1.6	Na _v 1.7	Na _v 1.8	Na _v 1.9
HGNC, UniProt	SCN8A, Q9UQD0	SCN9A, Q15858	SCN10A, Q9Y5Y9	SCN11A, Q9UI33
Sub/family-selective activators	batrachotoxin, veratridine	batrachotoxin, veratridine	–	–
Channel blockers	cannabidiol (pIC ₅₀ 5.5) [-45mV] [449]	cannabidiol (pIC ₅₀ 5.5–6.7) [449, 583], cannabigerol (pIC ₅₀ 5.3) [-90mV] [448], amorolfine (pIC ₅₀ 5.2) [-80mV] [450], ANP-230 (At resting state) (pIC ₅₀ 5.2) [665], cannabinalol (pIC ₅₀ 5) [-90mV] [447], cannabigerol (pIC ₅₀ 4.7) [-110mV] [448], cannabinalol (pIC ₅₀ 4.5) [-100mV] [447], amorolfine (pIC ₅₀ 4) [-100mV] [450]	cannabidiol (pIC ₅₀ 5.7) [451] – Mouse, cannabigerol (pIC ₅₀ 5.3) [451], ANP-230 (AT resting state) (pIC ₅₀ 4.9) [665], cannabinalol (pIC ₅₀ 4.8) [451] – Mouse	ANP-230 (At resting state) (pIC ₅₀ 5.2) [665]
Sub/family-selective channel blockers	tetrodotoxin (Pore blocker) (pIC ₅₀ 9) [-130mV] [321] – Rat, XPC-5462 (pIC ₅₀ 8) [-45mV] [463], XPC-7224 (pIC ₅₀ 7.1) [-45mV] [463], saxitoxin (Pore blocker)	PF-05089771 (pIC ₅₀ 7.5–8) [34, 642], tetrodotoxin (Pore blocker) (pIC ₅₀ 7.6) [-100mV] [720], PF-04856264 (pIC ₅₀ 7.6) [928], saxitoxin (Pore blocker) (pIC ₅₀ 6.2) [1458]	suzetrigine (pIC ₅₀ 9.6) [643, 1417], A-887826 (pIC ₅₀ 8) [643, 1611], VX-150 (pIC ₅₀ 7.8) [1417], LTGO-33 (pIC ₅₀ 7.5–7.6) [455, 643], PF-01247324 (Pore blocker) (pIC ₅₀ 6.7) [voltage dependent] [1089], tetrodotoxin (Pore blocker) (pIC ₅₀ 4.2) [-60mV] [20] – Rat	tetrodotoxin (Pore blocker) (pIC ₅₀ 4.4) [-120mV] [267] – Rat
Functional Characteristics	Activation V _{0.5} = -29 mV. Fast inactivation (1 ms)	Activation V _{0.5} = -27 mV. Fast inactivation (0.5 ms)	Activation V _{0.5} = -16 mV. Inactivation (6 ms)	Activation V _{0.5} = -32 mV. Slow inactivation (16 ms)
Comments	Na _v 1.6 is the target of the potential anti-epileptic channel blocker zandatrigrine (NBI-921352) [646, 809].	–	–	–

Comments: Sodium channels are also blocked by local anaesthetic agents, antiarrhythmic drugs and antiepileptic drugs. In general, these drugs are not highly selective among channel subtypes. There are two clear functional fingerprints for distinguishing different subtypes. These are sensitivity to tetrodotoxin (Na_v1.5, Na_v1.8 and Na_v1.9 are much less sensitive to block) and rate of fast inactivation (Na_v1.8 and particularly Na_v1.9 inactivate more slowly). All sodium channels also have a slow inactivation process that is engaged during long depolarizations (>100 ms) or repetitive trains of stimuli. All sodium channel subtypes are blocked by intracellular QX-314.

Further reading on Voltage-gated sodium channels (Na_v)

- Alsoum M *et al.* (2025) Voltage-gated sodium channels in excitable cells as drug targets. *Nat Rev Drug Discov* **24**: 358-378 [PMID:39901031]
- Catterall WA *et al.* (2005) International Union of Pharmacology. XLVII. Nomenclature and structure-function relationships of voltage-gated sodium channels. *Pharmacol Rev* **57**: 397-409 [PMID:16382098]
- Catterall WA *et al.* (2017) The chemical basis for electrical signaling. *Nat Chem Biol* **13**: 455-463 [PMID:28406893]
- Deuis JR *et al.* (2017) The pharmacology of voltage-gated sodium channel activators. *Neuropharmacology* **127**: 87-108 [PMID:28416444]
- Ghovanloo MR *et al.* (2022) Cannabidiol and Sodium Channel Pharmacology: General Overview, Mechanism, and Clinical Implications. *Neuroscientist* **28**: 318-334 [PMID:34027742]
- Jiang D *et al.* (2018) Structural basis for gating pore current in periodic paralysis. *Nature* **557**: 590-594 [PMID:29769724]
- Kanellopoulos AH *et al.* (2016) Voltage-gated sodium channels and pain-related disorders. *Clin Sci* **130**: 2257-2265 [PMID:27815510]
- Shen H *et al.* (2018) Structural basis for the modulation of voltage-gated sodium channels by animal toxins. *Science* **362**: [PMID:30049784]
- Terragni B *et al.* (2018) Post-translational dysfunctions in channelopathies of the nervous system. *Neuropharmacology* **132**: 31-42 [PMID:28571716]
- Waxman SG. (2023) Targeting a Peripheral Sodium Channel to Treat Pain. *N Engl J Med* **389**: 466-469 [PMID:37530829]

Comments on Voltage-gated ion channels: The voltage-dependent anion channels (VDACs) plays a key role in regulating metabolic and energetic flux across the outer mitochondrial membrane. It is involved in the transport of ATP, ADP, pyruvate, malate, and other metabolites, and thus communicates extensively with enzymes from metabolic pathways. They are a class of porin ion channel located on the outer mitochondrial membrane. VDAC1, VDAC2 and VDAC3 are involved in the regula-

tion of apoptosis, cell metabolism, mitochondrial apoptosis, and spermatogenesis [888, 1271].

The calcium homeostasis modulator (CALHM) ion channels are apparently voltage- and extracellular Ca^{2+} -gated, and constitute a novel ion channel family that is widely expressed in the brain and taste buds throughout vertebrates and in sensory neurons and body wall muscles in *C. elegans*. In humans, the CALHM family encompasses six paralogs, some of which function as non-

selective channels that are permeable to large substances such as ATP. CALHM channels are thought to play important roles in neuronal excitability, neurotransmission of tastes, and muscle cell function. The voltage- and extracellular Ca^{2+} -dependent gating mechanisms, structural features that define the gate and ion permeation pathway and additional physiological roles, remain to be discovered [881].

Other ion channels

[Ion channels](#) → [Other ion channels](#)

Overview: A number of ion channels in the human genome do not fit readily into the classification of either ligand-gated or voltage-gated ion channels. These are identified below.

Aquaporins

[Ion channels](#) → [Other ion channels](#) → [Aquaporins](#)

Overview: Aquaporins and aquaglyceroporins are membrane channels that allow the permeation of water and certain other small solutes across the cell membrane, or in the case of AQP6, AQP11 and AQP12A, intracellular membranes, such as vesicles and the endoplasmic reticulum membrane [745]. Since the isolation and cloning of the first aquaporin (AQP1) [1137], 12 additional mammalian members of the family have been identified, although little is known about the functional properties of one of these (AQP12A; Q8IXF9) and it is thus not tabulated. The other 12 aquaporins can be broadly divided into three families: orthodox aquaporins (AQP0,-1,-2,-4,-5, -6 and -8) permeable mainly to water, but for some additional solutes [285]; aquaglyceroporins (AQP3,-7 -9 and -10), additionally permeable to glycerol and for some isoforms urea [712], and superaquaporins (AQP11 and

12) located within cells [608]. Some aquaporins also conduct ammonia and/or H_2O_2 giving rise to the terms 'ammoniaporins' ('aquaammoniaporins') and 'peroxiporins', respectively. Aquaporins are impermeable to protons and other inorganic and organic cations, with the possible exception of AQP1, although this is controversial [712]. One or more members of this family of proteins have been found to be expressed in almost all tissues of the body [reviewed in Yang (2017) [1556]]. AQP1s are involved in numerous processes that include systemic water homeostasis, adipocyte metabolism, brain oedema, cell migration and fluid secretion by epithelia. Loss of function mutations of some human AQP1s, or their disruption by autoantibodies further underscore their importance [reviewed by Verkman *et al.* (2014) [1429], Kitchen *et al.* (2105) [712]].

Functional AQP1s exist as homotetramers that are the water conducting units wherein individual AQP subunits (each a protomer) have six TM helices and two half helices that constitute a seventh 'pseudotransmembrane domain' that surrounds a narrow water conducting channel [745]. In addition to the four pores contributed by the protomers, an additional hydrophobic pore exists within the center of the complex [745] that may mediate the transport through AQP1. Although numerous small molecule inhibitors of aquaporins, particularly AQP1, have been reported primarily from *Xenopus* oocyte swelling assays, the activity of most has subsequently been disputed upon retesting using assays of water transport that are less prone to various artifacts [377] and they are therefore excluded from the tables [see Tradtrantip *et al.* (2017) [1400] for a review].

Nomenclature	AQP0	AQP1	AQP2	AQP3
HGNC, UniProt	MIP, P30301	AQP1, P29972	AQP2, P41181	AQP3, Q92482
Endogenous activator	AQP0 is gated by calmodulin [712]	cGMP (see comment)	–	–
Permeability	water (rat single channel permeability $0.25 \times 10^{-14} \text{cm}^3 \text{s}^{-1}$) (Rat) [1558]	water (rat single channel permeability $6.0 \times 10^{-14} \text{cm}^3 \text{s}^{-1}$), ammonia, H_2O_2 [444, 1558]	water (rat single channel permeability $3.3 \times 10^{-14} \text{cm}^3 \text{s}^{-1}$) [880]	water (rat single channel permeability $2.1 \times 10^{-14} \text{cm}^3 \text{s}^{-1}$), glycerol, ammonia, H_2O_2 [119, 444, 1558]
Inhibitors	Hg^{2+}	compound 1 (pIC_{50} 5.1) [1249], Ag^+ , Hg^{2+} , pCMBS	Hg^{2+}	Auphen (pIC_{50} 6.1) [916], Audien (pIC_{50} 4.8) [916], Hg^{2+}
Channel blockers	–	5-Hydroxymethyl-2-furfural (Inhibition) (pIC_{50} 6.4) [232]	–	–
Comments	–	Permeability to H_2O_2 has been demonstrated for rat, but not human, AQP1 [119]. Numerous small molecule inhibitors of AQP1 have been proposed, but re-evaluation indicates that they have no significant effect upon water permeability at concentrations in excess of their originally reported IC_{50} values [435]. A fifth pore located at the central axis of the tetrameric complex has, controversially, been described as a cation conductance activated by cGMP and phosphorylation by protein kinases A and C. Evidence in support and against this proposal is discussed in detail by Kitchen <i>et al.</i> (2015) [712].	–	AQP3 is also inhibited by acid pH: permeability to urea is controversial [712].

Nomenclature	AQP4	AQP5
HGNC, UniProt	AQP4, P55087	AQP5, P55064
Permeability	water (rat single channel permeability $24 \times 10^{-14} \text{cm}^3 \text{s}^{-1}$) [1558]	water (rat single channel permeability $5.0 \times 10^{-14} \text{cm}^3 \text{s}^{-1}$), H_2O_2 [663]
Inhibitors	–	Hg^{2+}
Comments	AQP4 is inhibited by PKC activation (although this is probably due to phosphorylation-dependent protein localisation rather than inhibition of the channel <i>per se</i>), but not by HgCl_2 . AQP4 is predicted to be permeable to NO [1490]. Recent work suggests that the membrane trafficking of AQP4 could be an alternative target to pore-blockers [713].	AQP5 may conduct CO_2 [444].

Nomenclature	AQP6	AQP7	AQP8	AQP9	AQP10
HGNC, UniProt	AQP6 , Q13520	AQP7 , O14520	AQP8 , O94778	AQP9 , O43315	AQP10 , Q96PS8
Permeability	water (zero, or very low basal, permeability is enhanced by low pH and in mouse and rat by Hg ²⁺), glycerol, ammonia, urea, anions [444 , 561 , 712 , 1159]	water (high), glycerol, ammonia, urea [444 , 606]	water (mouse single channel permeability 8.2 x 10 ⁻¹⁴ cm ³ s ⁻¹), ammonia, H ₂ O ₂ [119 , 444 , 712 , 880]	water (low), glycerol, ammonia, urea, H ₂ O ₂ , monocarboxylates [444 , 560 , 1159 , 1497]	water (low), glycerol, urea [607]
Activators	Hg ²⁺	–	–	–	–
Inhibitors	–	Auphen (Effective at 15 μM), Hg ²⁺	Hg ²⁺	Hg ²⁺ , phloretin	Hg ²⁺
Comments	AQP6 is an intracellular channel that localises to acid secreting intercalated cells of the renal collecting ducts. Notably, AQP6 is activated by Hg ²⁺ and by low pH and is unusually permeable to anions (with the permeability sequence NO ₃ ⁻ >I ⁻ >Br ⁻ >Cl ⁻ >F ⁻) as well as water, both through the monomeric pore [712 , 1159].	AQP7 also transports silicon [435].	Permeability to urea is controversial, but might be explained by differences between mouse and human caused by a pore-lining amino acid residue that differs between species [712].	AQP9 may transport silicon [435].	It is not known if AQP10 is permeable to ammonia. Permeability to silicon has been described [435].

Further reading on Aquaporins

- Abir-Awan M *et al.* (2019) Inhibitors of Mammalian Aquaporin Water Channels. *Int J Mol Sci* **20**: [[PMID:30934923](#)]
- Agre P. (2006) The aquaporin water channels. *Proc Am Thorac Soc* **3**: 5-13 [[PMID:16493146](#)]
- Beitz E *et al.* (2015) Challenges and achievements in the therapeutic modulation of aquaporin functionality. *Pharmacol Ther* **155**: 22-35 [[PMID:26277280](#)]
- Bill RM. (2024) Drugging aquaporins. *Biochim Biophys Acta Biomembr* **1866**: 184164 [[PMID:37146744](#)]
- Carbrey JM *et al.* (2009) Discovery of the aquaporins and development of the field. *Handb Exp Pharmacol* 3-28 [[PMID:19096770](#)]
- Geng X *et al.* (2017) Transport Characteristics of Aquaporins. *Adv Exp Med Biol* **969**: 51-62 [[PMID:28258565](#)]
- Kitchen P *et al.* (2015) Beyond water homeostasis: Diverse functional roles of mammalian aquaporins. *Biochim Biophys Acta* **1850**: 2410-21 [[PMID:26365508](#)]
- Tesse A *et al.* (2018) Aquaporins as Targets of Dietary Bioactive Phytocompounds. *Front Mol Biosci* **5**: 30 [[PMID:29721498](#)]
- Tradtrantip L *et al.* (2017) Aquaporin-Targeted Therapeutics: State-of-the-Field. *Adv Exp Med Biol* **969**: 239-250 [[PMID:28258578](#)]
- Yang B. (2017) *In Advances in Experimental Medicine and Biology* Edited by Yang B: Springer: 1-276 [ISBN: 9789402410570]

Chloride channels

[Ion channels](#) → [Other ion channels](#) → [Chloride channels](#)

Overview: Chloride channels are a functionally and structurally diverse group of anion selective channels involved in processes including the regulation of the excitability of neurones, skeletal, cardiac and smooth muscle, cell volume regulation, transepithelial salt transport, the acidification of internal and extracellular compartments, the cell cycle and apoptosis (reviewed in [[354](#)]).

Excluding the transmittergated GABA_A and glycine receptors (see separate tables), well characterised chloride channels can be classified as certain members of the voltage-sensitive ClC subfamily, calcium-activated channels, high (maxi) conductance channels, the cystic fibrosis transmembrane conductance regulator (CFTR) and volume regulated channels [[1430](#)]. No official recommen-

dation exists regarding the classification of chloride channels. Functional chloride channels that have been cloned from, or characterised within, mammalian tissues are listed with the exception of several classes of intracellular channels (*e.g.* CLIC) that are reviewed by in [[362](#)].

Searchable database: <http://www.guidetopharmacology.org/>

Full Contents of ConciseGuide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.70231/full>

Chloride channels S221

CIC family

Ion channels → Other ion channels → Chloride channels → CIC family

Overview: The mammalian CIC family (reviewed in [10, 212, 354, 357, 632]) contains 9 members that fall, on the basis of sequence homology, into three groups; CIC-1, CIC-2, hCIC-Ka (rCIC-K1) and hCIC-Kb (rCIC-K2); CIC-3 to CIC-5, and CIC-6 and -7. CIC-1 and CIC-2 are plasma membrane chloride channels. CIC-Ka and CIC-Kb are also plasma membrane channels (largely expressed in the kidney and inner ear) when associated with barttin (*BSND*, *Q8WZ55*), a 320 amino acid 2TM protein [378]. The localisation of the remaining members of the CIC family is likely to be predominantly intracellular *in vivo*, although they may traffic to the plasma membrane in overexpression systems. Numer-

ous recent reports indicate that CIC-4, CIC-5, CIC-6 and CIC-7 (and by inference CIC-3) function as Cl⁻/H⁺ antiporters (secondary active transport), rather than classical Cl⁻ channels [472, 793, 1003, 1113, 1232]; reviewed in [10, 1142]). It has recently been reported that the activity of CIC-5 as a Cl⁻/H⁺ exchanger is important for renal endocytosis [1034]. Alternative splicing increases the structural diversity within the CIC family. The crystal structure of two bacterial CIC proteins has been described [358] and a eukaryotic CIC transporter (CmCLC) has recently been described at 3.5 Å resolution [394]. Each CIC subunit, with a complex topology of 18 intramembrane segments, contributes a single pore

to a dimeric 'double-barrelled' CIC channel that contains two independently gated pores, confirming the predictions of previous functional and structural investigations (reviewed in [212, 357, 632, 1142]). As found for CIC-4, CIC-5, CIC-6 and CIC-7, the prokaryotic CIC homologue (CIC-ec1) and CmCLC function as H⁺/Cl⁻ antiporters, rather than as ion channels [9, 394]. The generation of monomers from dimeric CIC-ec1 has firmly established that each CIC subunit is a functional unit for transport and that cross-subunit interaction is not required for Cl⁻/H⁺ exchange in CIC transporters [1180].

Nomenclature	CIC-1	CIC-2
HGNC, UniProt	CLCN1 , P35523	CLCN2 , P51788
Endogenous activators	–	arachidonic acid
Activators	–	lubiprostone, omeprazole
Inhibitors	NMD670 (pEC ₅₀ 5.8) [1285]	–
Channel blockers	9-anthroic acid, S-(-)CPB, S-(-)CPP, Cd ²⁺ , Zn ²⁺ , fenofibric acid, niflumic acid	GaTx2 (pK _d 10.8) [voltage dependent -100mV], Cd ²⁺ , NPPB, Zn ²⁺ , diphenylamine-2-carboxylic acid
Functional Characteristics	γ = 1-1.5 pS; voltage-activated (depolarization) (by fast gating of single protopores and a slower common gate allowing both pores to open simultaneously); inwardly rectifying; incomplete deactivation upon repolarization, ATP binding to cytoplasmic cystathionine β-synthase related (CBS) domains inhibits CIC-1 (by closure of the common gate), depending on its redox status	γ = 2-3 pS; voltage-activated by membrane hyperpolarization by fast protopore and slow cooperative gating; channels only open negative to E _{Cl} resulting in steady-state inward rectification; voltage dependence modulated by permeant anions; activated by cell swelling, PKA, and weak extracellular acidosis; potentiated by SGK1; inhibited by phosphorylation by p34(cdc2)/cyclin B; cell surface expression and activity increased by association with Hsp90
Comments	CIC-1 is constitutively active	CIC-2 is also activated by amidation

Nomenclature	ClC-Ka	ClC-Kb	ClC-3	ClC-4
HGNC, UniProt	CLCNKA, P51800	CLCNKB, P51801	CLCN3, P51790	CLCN4, P51793
Activators	niflumic acid (pEC ₅₀ 3–5)	niflumic acid (pEC ₅₀ 3–5)	–	–
Channel blockers	3-phenyl-CPP , DIDS , niflumic acid	3-phenyl-CPP , DIDS	phloretin (pIC ₅₀ 4.5)	Zn²⁺ (pIC ₅₀ 4.3) [1063], Cd²⁺ (pIC ₅₀ 4.2) [1063]
Functional Characteristics	γ = 26 pS; linear current-voltage relationship except at very negative potentials; no time dependence; inhibited by extracellular protons (pK = 7.1); potentiated by extracellular Ca ²⁺	Bidirectional rectification; no time dependence; inhibited by extracellular protons; potentiated by extracellular Ca ²⁺	Cl ⁻ /H ⁺ antiporter [923]; pronounced outward rectification; slow activation, fast deactivation; activity enhanced by CaM kinase II; inhibited by intracellular Ins(3,4,5,6)P4 and extracellular acidosis	Cl ⁻ /H ⁺ antiporter (2Cl ⁻ :1H ⁺) [30, 1113, 1232]; extreme outward rectification; voltage-dependent gating with midpoint of activation at +73 mV [1057]; rapid activation and deactivation; inhibited by extracellular acidosis; non-hydrolytic nucleotide binding required for full activity
Comments	ClC-Ka is constitutively active (when co-expressed with barttin), and can be blocked by benzofuran derivatives	ClC-Kb is constitutively active (when co-expressed with barttin), and can be blocked by benzofuran derivatives	insensitive to the channel blockers DIDS , NPPB and tamoxifen (10 μM)	–

Nomenclature	ClC-5	ClC-6	ClC-7
HGNC, UniProt	CLCN5, P51795	CLCN6, P51797	CLCN7, P51798
Channel blockers	–	DIDS (pIC ₅₀ 3)	DIDS (pIC ₅₀ 4.4) [1246], NSS818 (pIC ₅₀ 4.3) [1246], NPPB (pIC ₅₀ 3.8) [1246]
Functional Characteristics	Cl ⁻ /H ⁺ antiporter (2Cl ⁻ :1H ⁺) [1113, 1232, 1288, 1635]; extreme outward rectification; voltage-dependent gating with midpoint of activation of 116.0 mV; rapid activation and deactivation; potentiated and inhibited by intracellular and extracellular acidosis, respectively; ATP binding to cytoplasmic cystathionine β-synthase related (CBS) domains activates ClC-5	Cl ⁻ /H ⁺ antiporter (2Cl ⁻ :1H ⁺) [1003]; outward rectification, rapid activation and deactivation	Cl ⁻ /H ⁺ antiporter (2Cl ⁻ :1H ⁺) [472, 793, 1246]; strong outward rectification; voltage-dependent gating with a threshold more positive than ~ +20 mV; very slow activation and deactivation
Comments	Insensitive to the channel blockers DIDS (1 mM), diphenylamine-2-carboxylic acid (1 mM), 9-anthroic acid (2 mM), NPPB (0.5 mM) and niflumic acid (1 mM)	–	active when co-expressed with Ostm1

Comments: ClC channels display the permeability sequence Cl⁻ > Br⁻ > I⁻ (at physiological pH). ClC-1 has significant opening probability at resting membrane potential, accounting for 75% of the membrane conductance at rest in skeletal muscle, and is important for stabilization of the membrane potential. [S\(-\)CPP](#), [9-anthroic acid](#) and [niflumic acid](#) act intracellularly and exhibit a strongly voltage-dependent block with strong inhibition at negative voltages and relief of block at depolarized potentials ([818] and reviewed in [1141]). Inhibition of ClC-2 by the peptide [GaTx2](#), from *Leiurus quinquestriatus herbareus* venom, is likely

to occur through inhibition of channel gating, rather than direct open channel blockade [1383]. Although ClC-2 can be activated by cell swelling, it does not correspond to the VRAC channel (see below). Alternative potential physiological functions for ClC-2 are reviewed in [1126]. Functional expression of human ClC-Ka and ClC-Kb requires the presence of barttin [378, 1239] reviewed in [384]. The properties of ClC-Ka/barttin and ClC-Kb/barttin tabulated are those observed in mammalian expression systems: in oocytes the channels display time- and voltage-dependent gating. The rodent homologue (ClC-K1) of ClC-Ka demonstrates

limited expression as a homomer, but its function is enhanced by barttin which increases both channel opening probability in the physiological range of potentials [378, 404, 1239] reviewed in [384]. ClC-Ka is approximately 5 to 6-fold more sensitive to block by [3-phenyl-CPP](#) and [DIDS](#) than ClC-Kb, while newly synthesized benzofuran derivatives showed the same blocking affinity (<10 μM) on both ClC-K isoforms [819]. The biophysical and pharmacological properties of ClC-3, and the relationship of the protein to the endogenous volume-regulated anion channel(s) VRAC [31, 486] are controversial and further complicated by the

possibility that CIC-3 may function as both a Cl⁻/H⁺ exchanger and an ion channel [31, 1113, 1485]. The functional properties tabulated are those most consistent with the close structural relationship between CIC-3, CIC-4 and CIC-5. Activation of heterologously expressed CIC-3 by cell swelling in response to hypotonic

solutions is disputed, as are many other aspects of its regulation. Dependent upon the predominant extracellular anion (*e.g.* SCN⁻ versus Cl⁻), CIC-4 can operate in two transport modes: a slippage mode in which behaves as an ion channel and an exchanger mode in which unitary transport rate is 10-fold lower [30]. Sim-

ilar findings have been made for CIC-5 [1595]. CIC-7 associates with a β subunit, Ostm1, which increases the stability of the former [770] and is essential for its function [793].

CFTR

Ion channels → **Other ion channels** → **Chloride channels** → **CFTR**

Overview: CFTR is a member of the ABC transporter superfamily, but, uniquely, it is an ion channel, allowing electrodiffusion of Cl⁻ and HCO₃⁻. It is activated by phosphorylation, mainly by PKA on its regulatory domain (R domain). Conserved nucleotide binding domains (NBD1 and NBD2) couple ATP binding and hydrolysis to gate opening and closing, respectively [262]. CFTR is expressed apically in polarized epithelial cells in various organs where it controls volume and pH of fluid secretions as well as mucin unfolding and release [904]. CFTR transcripts are present in secretory and ionocyte cells in airway epithelia [1048, 1130], crypt enterocytes, goblet and CFTR-high expressing cells in the intestine [113, 180], pancreatic duct cells [397], intra- and extra-

hepatic cholangiocytes 33318612 [1432] and others. Mutations in the CFTR gene cause the genetic disease cystic fibrosis (CF) [1272]. The most common mutation, F508del, is present in at least one gene copy in ~80% of patients worldwide, but there are ~1000 different variants known to cause CF. Mutations affect CFTR biogenesis (folding, maturation, trafficking, metabolic stability) and/or ion-channel function. Vertex Pharmaceuticals developed small-molecule CFTR modulator drugs that improve biogenesis ("correctors") or open probability ("potentiators") of defective CFTR variants. Triple combination therapies, including two correctors and one potentiator (*e.g.* Trikafta®: **elexacaftor**, **tezacaftor**, **ivacaftor**), are standard of care for patients carrying at

least one copy of the F508del variant. Patients carrying mutations only affecting ion-channel function ("gating mutations" *e.g.* G551D) are treated with ivacaftor (potentiator) alone. Cryo-EM structures of Trikafta-bound F508del-E1371Q-CFTR reveal that all three compounds bind at the protein-membrane interface, in shallow pockets on CFTR's surface [401].

While low/absent CFTR activity causes CF, over-activation of CFTR (due to bacterial toxins such as cholera toxin) results in secretory diarrhoeas, causing large intestinal loss of fluid and alkali [288]. No inhibitors have been approved yet for emergency treatment of secretory diarrhoeas.

Nomenclature	CFTR
HGNC, UniProt	CFTR, P13569
Activators	ivacaftor (Potentiation) (pEC ₅₀ 9.3) [261], ivacaftor (Potentiation) (pEC ₅₀ 8.5) [496], elexacaftor (Correction) (pEC ₅₀ 6.4–7.2) [4, 689], GLPG1837 (Potentiation) (pEC ₅₀ 6.6) [1569], deutivacaftor (Potentiation) (pEC ₅₀ 6.6) [515], tezacaftor (Correction) (pEC ₅₀ 6.6) [1194], felodipine (Potentiation) (pEC ₅₀ 5.7) [1093], vanzacaftor (Corrector) [688]
Inhibitors	(R)-BPO-27 (pIC ₅₀ 8.4) [1297], CFTR_{inh}-172 (pIC ₅₀ 7.1) [425], GaTx1 (pIC ₅₀ 7) [419]
Channel blockers	GlyH-101 (Pore blocker) (pIC ₅₀ 5.4) [voltage dependent] [979], glibenclamide (Pore blocker) (pK _i 4.7) [voltage dependent] [1266]
Functional Characteristics	$\gamma = 6-9$ pS; permeability ratio sequence: SCN ⁻ > NO ₃ ⁻ > Br ⁻ > Cl ⁻ > I ⁻ > formate ⁻ > F ⁻ ; conductance ratio sequence: Cl ⁻ > NO ₃ ⁻ > Br ⁻ ~ formate ⁻ > SCN ⁻ > I ⁻ . Slight outward rectification, only observed in intact cells, is probably due to cytosolic anions resulting in voltage-dependent channel block [835]. Activated by binding of and phosphorylation by PKA [948]. Opening of the channel gate follows ATP binding at two nucleotide binding domains (NBD1 and NBD2), and formation of an intramolecular NBD1/NBD2 tight dimer [1428]; closing is coupled to ATP hydrolysis and dimer dissociation [263]. Positively regulated by PKC [197] and PKGII (in enterocytes) [1250, 1416].
Comments	UCCF-339 , UCCF-029 , apigenin and genistein are examples of flavones. UCCF-853 and NS004 are examples of benzimidazolones. CBIQ is an example of a benzquinoline. Felodipine and nimodipine are examples of 1,4-dihydropyridines. Phenylglycine-01 is an example of a phenylglycine. Sulfonamide-01 is an example of a sulfonamide. Malonic acid hydrazide conjugates are also CFTR channel blockers (see Verkman and Galietta, 2009 [1430])

Comments: Cystic fibrosis mouse models have been useful in studies of the intestinal symptoms of CF (*e.g.* [1361]). However, they do not develop severe lung symptoms typically observed in humans with CF [487]. CF pigs [1319] exhibit respiratory defects and mucus accumulation similar to humans [1311]. More recently, several ferret (*Mustela putorius furo*) CF models have also been developed, by providing *in utero* treatment with CFTR modulators to reduce mortality in the foetal and neonatal period [380, 1339].

Further reading on CFTR

Fiedorczuk K *et al.* (2022) Mechanism of CFTR correction by type I folding correctors. *Cell* **185**: 158-168.e11 [PMID:34995514] Levring J *et al.* (2023) CFTR function, pathology and pharmacology at single-molecule resolution. *Nature* **616**: 606-614 [PMID:36949202]

Calcium activated chloride channel (CaCC)

[Ion channels](#) → [Other ion channels](#) → [Chloride channels](#) → [Calcium activated chloride channel \(CaCC\)](#)

Overview: Chloride channels activated by intracellular Ca^{2+} (CaCC) are widely expressed in excitable and non-excitable cells where they perform diverse functions [519]. CaCCs are activated by a rise in intracellular free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$), typically following activation of G_q protein coupled receptors (G_q -PCR). This section centres on CaCC channels encoded by the TMEM16A gene (HUGO gene nomenclature: Anoctamin 1). The

TMEM16 family consists of 10 paralogs (TMEM16A-K; Anoctamin 1-10). The TMEM16A and TMEM16B genes (*ANO1* and *ANO2*) encode for CaCCs, while the other members function as lipid scramblases or have combined scramblase and non-selective ion channel function [13, 395, 522, 1116, 1242]. TMEM16A has a broad tissue distribution and a variety of established cellular roles, while the main physiological role for TMEM16B identified

thus far is in olfaction [319, 776]. Alternative splicing regulates the voltage- and Ca^{2+} -dependence of TMEM16A and such post-transcriptional process may be tissue-specific and contribute to functional diversity [399]. TMEM16A is a potential drug target for a variety of conditions spanning from respiratory to vascular (see “Comments” section for further detail).

Nomenclature	CaCC
HGNC, UniProt	ANO1 , Q5XXA6
Endogenous activators	intracellular Ca^{2+}
Activators	PAM_16A (pEC ₅₀ 8.4) [21], E_{act} [1431]
Endogenous inhibitors	Ins(3,4,5,6)P₄
Inhibitors	MONNA (pIC ₅₀ 7.1) [130, 1042, 1407], T16Ainh-A01 (pIC ₅₀ ~6) [130, 1114], crofelemer (pIC ₅₀ 5.2) [1401], 9-anthroic acid , CaCCinh-A01 [130, 289], DCDPC , DIDS , NPPB , SITS , flufenamic acid , fluoxetine , mibefradil , niflumic acid , tannic acid
Selective inhibitors	TM_{inh}-23 (pIC ₅₀ 7.5) [1407], Ani9 (pIC ₅₀ 7) [1253]
Functional Characteristics	$\gamma = 0.5\text{-}5$ pS; permeability sequence, $\text{SCN}^- > \text{NO}_3^- > \text{I}^- > \text{Br}^- > \text{Cl}^- > \text{F}^-$; relative permeability of $\text{SCN}^-:\text{Cl}^- \sim 8$. $\text{I}^-:\text{Cl}^- \sim 3$, aspartate: $\text{Cl}^- \sim 0.15$, outward rectification (decreased by increasing $[\text{Ca}^{2+}]_i$); sensitivity to activation by $[\text{Ca}^{2+}]_i$ decreased at hyperpolarized potentials; slow activation at positive potentials (accelerated by increasing $[\text{Ca}^{2+}]_i$); rapid deactivation at negative potentials, deactivation kinetics modulated by anions binding to an external site; modulated by redox status
Comments	TMEM16A has been reported to regulate a wide variety of physiological processes ranging from sensory transduction to smooth muscle contraction and epithelial secretion [582]. A CaCC (TMEM16A) potentiator compound (GDC-6988/ETD002, undisclosed structure; acquired by Roche from Enterprise Therapeutics) entered Phase 1 clinical evaluation as a novel approach that has potential to provide benefit to all patients with cystic fibrosis (mentioned in [1120]). Up-regulating chloride transport <i>via</i> CaCC is proposed to mitigate the effect of loss of chloride transport <i>via</i> CFTR in CF. See Enterprise Therapeutics' reports of CaCC potentiator ETX001 for more detailed background information [273, 278].

Comments: Therapeutic indications: TMEM16A regulates a wide variety of physiological processes ranging from nociception to smooth muscle contraction and epithelial secretion [582]. In airway epithelial cells, TMEM16A expression is enhanced by inflammatory stimuli that also lead to goblet cell metaplasia and increased mucus secretion. Thus, pharmacological modulation of TMEM16A could help to improve mucociliary clearance in cystic fibrosis (CF) and chronic obstructive respiratory disease [274, 421].

TMEM16A activation might also be of potential therapeutic benefit in Sjögren's syndrome. Given the established role of TMEM16A role in smooth muscle and cerebral pericyte contraction, pharmacological inhibitors of TMEM16A are expected to promote vasodilation and could have therapeutic use in a range of diseases of altered vessel tone including stroke, hypertension and vascular dementia [23, 440]. Other suggested therapeutic indications in which TMEM16A could be targeted include neuropathic pain, given the

identified role of TMEM16A in dorsal root ganglion (DRG) neurons [838]. TMEM16A is also expressed in interstitial cells of Cajal and may be involved in the control of gastric emptying and motility [243]; thus, TMEM16A modulators could be used to modulate gastric contractility. TMEM16A was also reported in the smooth muscle of the bladder, uterus, and internal urethral sphincter; therefore, TMEM16A modulators could have a role in the treatment of overactive bladder and urinary incontinence [391] and to

foster myometrium relaxation [1525]. TMEM16A is overexpressed in a variety of cancer types and TMEM16A inhibitors have been suggested as a potential anticancer treatment [580, 1151].

Pharmacology: A CaCC (TMEM16A) activator compound (GDC-6988/ETD002, undisclosed structure; acquired by Roche from Enterprise Therapeutics) entered Phase 1 clinical evaluation as a novel approach that has potential to provide benefit to all patients with cystic fibrosis (CF) (mentioned in [1120]). Up-regulating chloride transport *via* CaCC is proposed to mitigate the effect of loss of Cl⁻ transport *via* CFTR in CF. See Enterprise Therapeutics' reports of CaCC potentiator ETX001 for more detailed background information [273, 278]. The compound Eact has been reported to directly activate TMEM16A [1431]. However, recent data have challenged this direct activation of TMEM16A and indicate that Eact induces an increase in [Ca²⁺]_i through an agonist effect on TRPV1 and TRPV4 [438, 845]. Many of the list-

ed TMEM16A channel blockers are recognised as being unselective [22, 130, 439]. Ani9 demonstrates selectivity for TMEM16A versus TMEM16B [1253] and, in contrast to several of the other inhibitors, (i) has no effects on intracellular Ca²⁺ signalling in human airway epithelial cells [276] and (ii) does not affect a range of other ion current types [22]. Ani9 possess an imine/hydrazone group that might render the compound unstable and potentially reactive *in vivo*. TMinh-23 is one the most potent TMEM16A inhibitors reported thus far (IC₅₀ ~30 nM) [244, 1407]. However, the selectivity of this compound is not fully defined, and the compound might be metabolically unstable *in vivo* due to potential hydrolysis of one or both amide groups present in this molecule. TMEM16A is inhibited by the therapeutic drugs tamsulosin (IC₅₀ ~7 μM) [810]. The anthelmintic niclosamide was also reported to inhibit TMEM16A [959], but further studies demonstrated that niclosamide can act both as an inhibitor or an activator depend-

ing on the membrane potential and [Ca²⁺]_i [276, 687, 816]. A range of natural compounds including gallotannins [997], evo-diamine and rutecarpine [1614], theaflavin [1268] and allicin [67] are non-selective, low potency TMEM16A inhibitors. CaCC currents in CFPAC-1 have also been reported to be inhibited by Ins(3,4,5,6)P4 [553]. Complex effects on native CaCC and cloned TMEM16A and TMEM16B currents have been observed for a range of compounds including extracellular niflumic acid; DCD-PC and 9-anthroic acid (but not DIDS) enhancing and inhibiting inwardly and outwardly directed currents in a manner dependent upon [Ca²⁺]_i and membrane potential [226, 1121, 1209, 1353] (and [777] for summary). Blockade of CaCC currents by niflumic acid, DIDS and 9-anthroic acid is voltage-dependent, whereas block by NPPB is voltage-independent [519]. Considerable cross-over in pharmacology with large conductance Ca²⁺-activated K⁺ channels also exists (see [473] for overview).

Maxi chloride channel

[Ion channels](#) → [Other ion channels](#) → [Chloride channels](#) → [Maxi chloride channel](#)

Overview: Maxi Cl⁻ channels are high conductance, anion selective, channels initially characterised in skeletal muscle and subsequently found in many cell types including neurones, glia, cardiac muscle, lymphocytes, secreting and absorbing epithelia, macula densa cells of the kidney and human placenta syncytiotrophoblasts [1208]. The physiological significance of the maxi Cl⁻ chan-

nel is uncertain, but roles in cell volume regulation and apoptosis have been claimed. Evidence suggests a role for maxi Cl⁻ channels as a conductive pathway in the swelling-induced release of ATP from mouse mammary C127i cells that may be important for autocrine and paracrine signalling by purines [355, 1207]. A similar channel mediates ATP release from macula densa cells within the

thick ascending of the loop of Henle in response to changes in luminal NaCl concentration [106]. A family of human high conductance Cl⁻ channels (TTYH1-3) that resemble Maxi Cl⁻ channels has been cloned [1346], but alternatively, Maxi Cl⁻ channels have also been suggested to correspond to the voltage-dependent anion channel, VDAC, expressed at the plasma membrane [65, 1044].

Nomenclature	Maxi Cl ⁻
Activators	cytosolic GTPγS, extracellular chlorpromazine, extracellular tamoxifen, extracellular toremifene, extracellular triflupromazine
Endogenous channel blockers	intracellular arachidonic acid
Channel blockers	DIDS (pIC ₅₀ 4.4) [1246], extracellular Zn ²⁺ (pIC ₅₀ 4.3) [1063], NPPB (pIC ₅₀ 3.8) [1246], extracellular Gd ³⁺ , SITS, diphenylamine-2-carboxylic acid
Functional Characteristics	γ = 280-430 pS (main state); permeability sequence, I > Br > Cl > F > gluconate (P _{Cl} /P _{Cl} = ~1.5); ATP is a voltage dependent permeant blocker of single channel activity (P _{ATP} /P _{Cl} = 0.08-0.1); channel activity increased by patch-excision; channel opening probability (at steady-state) maximal within approximately ± 20 mV of 0 mV, opening probability decreased at more negative and (commonly) positive potentials yielding a bell-shaped curve; channel conductance and opening probability regulated by annexin 6
Comments	Maxi Cl ⁻ is also activated by G protein-coupled receptors and cell swelling. Tamoxifen and toremifene are examples of triphenylethylene anti-oestrogens

Comments: Differing ionic conditions may contribute to variable estimates of γ reported in the literature. Inhibition by arachidonic acid (and cis-unsaturated fatty acids) is voltage-independent, occurs at an intracellular site, and involves both channel shut down (K_d = 4-5 μM) and a reduction of γ (K_d = 13-14 μM). Blockade of channel activity by SITS, DIDS, Gd³⁺ and

arachidonic acid is paralleled by decreased swelling-induced release of ATP [355, 1207]. Channel activation by anti-oestrogens in whole cell recordings requires the presence of intracellular nucleotides and is prevented by pre-treatment with 17β-estradiol, bucladesine, or intracellular dialysis with GDPβS [318]. Activation by tamoxifen is suppressed by low concentrations of oka-

daic acid, suggesting that a dephosphorylation event by protein phosphatase PP2A occurs in the activation pathway [318]. In contrast, 17β-estradiol and tamoxifen appear to directly inhibit the maxi Cl⁻ channel of human placenta reconstituted into giant liposomes and recorded in excised patches [1178].

Volume regulated chloride channels (VRAC)

Ion channels → **Other ion channels** → **Chloride channels** → **Volume regulated chloride channels (VRAC)**

Overview: Volume activated chloride channels (also termed VSOAC, volume-sensitive organic osmolyte/anion channel; VRC, volume regulated channel and VSOR, volume expansion-sensing outwardly rectifying anion channel) participate in regulatory volume decrease (RVD) in response to cell swelling. VRAC may also be important for several other processes including the regulation

of membrane excitability, transcellular Cl⁻ transport, angiogenesis, cell proliferation, necrosis, apoptosis, glutamate release from astrocytes, **insulin** (*INS*, P01308) release from pancreatic β cells and resistance to the anti-cancer drug, **cisplatin** (reviewed by [111, 981, 1017, 1046]). VRAC may not be a single entity, but may instead represent a number of different channels that are

expressed to a variable extent in different tissues and are differentially activated by cell swelling. In addition to ClC-3 expression products (see above) several former VRAC candidates including *MDR1* (ABCB1 P-glycoprotein), Icln, Band 3 anion exchanger and phospholemman are also no longer considered likely to fulfil this function (see reviews [1017, 1224]).

Nomenclature	VRAC
Activators	GTPγS
Endogenous channel blockers	intracellular Mg ²⁺ , arachidonic acid
Channel blockers	1,9-dideoxyforskolin, 9-anthroic acid, DCPIB, DIDS, IAA-94, NPPB, NS3728, carbenoxolone, clomiphene, diBA-(5)-C4, gossypol, mefloquine, mibefradil, nafoxidine, nordihydroguaiaretic acid, quinidine, quinine, tamoxifen
Functional Characteristics	γ = 10-20 pS (negative potentials), 50-90 pS (positive potentials); permeability sequence SCN ⁻ > I ⁻ > NO ₃ ⁻ > Br ⁻ > Cl ⁻ > F ⁻ > gluconate; outward rectification due to voltage dependence of γ; inactivates at positive potentials in many, but not all, cell types; time dependent inactivation at positive potentials; intracellular ionic strength modulates sensitivity to cell swelling and rate of channel activation; rate of swelling-induced activation is modulated by intracellular ATP concentration; ATP dependence is independent of hydrolysis and modulated by rate of cell swelling; inhibited by increased intracellular free Mg ²⁺ concentration; swelling induced activation of several intracellular signalling cascades may be permissive of, but not essential to, the activation of VRAC including: the Rho-Rho kinase-MLCK; Ras-Raf-MEK-ERK; PIK3-NOX-H ₂ O ₂ and Src-PLCγ-Ca ²⁺ pathways; regulation by PKCα required for optimal activity; cholesterol depletion enhances activity; activated by direct stretch of β1-integrin
Comments	VRAC is also activated by cell swelling and low intracellular ionic strength. VRAC is also blocked by chromones, extracellular nucleotides and nucleoside analogues

Comments: In addition to conducting monovalent anions, in many cell types the activation of VRAC by a hypotonic stimulus can allow the efflux of organic osmolytes such as amino acids and polyols that may contribute to RVD.

Comments on Chloride channels: Other chloride channels

In addition to some intracellular chloride channels that are not considered here, plasma membrane channels other than those listed have been functionally described. Many cells and tissues contain outwardly rectifying chloride channels (ORCC) that may correspond to VRAC active under isotonic conditions. A **cyclic AMP**-activated Cl⁻ channel that does not correspond to CFTR has been described in intestinal Paneth cells [1408]. A Cl channel

activated by **cyclic GMP** with a dependence on raised intracellular Ca²⁺ has been recorded in various vascular smooth muscle cells types, which has a pharmacology and biophysical characteristics very different from the 'conventional' CaCC [920, 1122]. It has been proposed that **bestrophin-3** (*BEST3*, Q8N1M1) is an essential component of the **cyclic GMP**-activated channel [921]. A proton-activated, outwardly rectifying anion channel has also

been described [767].

The chloride intracellular channel proteins (**CLICs**) are non-canonical ion channels with six homologs, distinct from most ion channels in that they have both soluble and integral membrane forms. The physiological role of CLICs appears to be maintenance of intracellular membranes, which is associated with tubulogenesis but may involve other substructures [493].

Further reading on Chloride channels

Adkins GB *et al.* (2015) Potential role of cardiac chloride channels and transporters as novel therapeutic targets. *Pharmacol Ther* **145**: 67-75 [PMID:25160469]

Huang F *et al.* (2012) International Union of Basic and Clinical Pharmacology. LXXXV: calcium-activated chloride channels. *Pharmacol Rev* **64**: 1-15 [PMID:22090471]

Searchable database: <http://www.guidetopharmacology.org/>

Full Contents of ConciseGuide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.70231/full>

Volume regulated chloride channels (VRAC) S227

- Kamaleddin MA. (2018) Molecular, biophysical, and pharmacological properties of calcium-activated chloride channels. *J Cell Physiol* **233**: 787-798 [PMID:28121009]
- Kunzelmann K. (2015) TMEM16, LRRC8A, bestrophin: chloride channels controlled by Ca(2+) and cell volume. *Trends Biochem Sci* **40**: 535-43 [PMID:26254230]
- Pedersen SF *et al.* (2016) Biophysics and Physiology of the Volume-Regulated Anion Channel (VRAC)/Volume-Sensitive Outwardly Rectifying Anion Channel (VSOR). *Pflugers Arch* **468**: 371-83 [PMID:26739710]
- Peretti M *et al.* (2015) Chloride channels in cancer: Focus on chloride intracellular channel 1 and 4 (CLIC1 AND CLIC4) proteins in tumor development and as novel therapeutic targets. *Biochim Biophys Acta* **1848**: 2523-31 [PMID:25546839]
- Sabirov RZ *et al.* (2016) The properties, functions, and pathophysiology of maxi-anion channels. *Pflugers Arch* **468**: 405-20 [PMID:26733413]
- Zegarra-Moran O *et al.* (2017) CFTR pharmacology. *Cell Mol Life Sci* **74**: 117-128 [PMID:27704174]

Connexins and Pannexins

[Ion channels](#) → [Other ion channels](#) → [Connexins and Pannexins](#)

Overview: Gap junctions are essential for many physiological processes including cardiac and smooth muscle contraction, regulation of neuronal excitability and epithelial electrolyte transport [161, 255, 381]. Gap junction channels allow the passive diffusion of molecules of up to 1,000 Daltons which can include nutrients, metabolites and second messengers (such as IP₃) as well as cations and anions. 21 connexin genes and 3 pannexin genes which are structurally related to the invertebrate innexin genes, code for gap junction proteins in humans. Each connexin

gap junction comprises 2 hemichannels or 'connexons' which are themselves formed from 6 connexin molecules. The various connexins have been observed to combine into both homomeric and heteromeric combinations, each of which may exhibit different functional properties. It is also suggested that individual hemichannels formed by a number of different connexins might be functional in at least some cells [537]. Connexins have a common topology, with four α -helical transmembrane domains, two extracellular loops, a cytoplasmic loop, and N- and C-termini

located on the cytoplasmic membrane face. In mice, the most abundant connexins in electrical synapses in the brain seem to be Cx36, Cx45 and Cx57 [1300]. Mutations in connexin genes are associated with the occurrence of a number of pathologies, such as peripheral neuropathies, cardiovascular diseases and hereditary deafness. The pannexin genes Px1 and Px2 are widely expressed in the mammalian brain [1442]. Like the connexins, at least some of the pannexins can form hemichannels [161, 1099].

Nomenclature	Cx23	Cx25	Cx26	Cx30	Cx30.2	Cx30.3	Cx31
HGNC, UniProt	GJE1 , A6NN92	GJB7 , Q6PEY0	GJB2 , P29033	GJB6 , O95452	GJC3 , Q8NFK1	GJB4 , Q9NTQ9	GJB3 , O75712
Endogenous inhibitors	extracellular Ca ²⁺ (blocked by raising external Ca ²⁺)						
Inhibitors	carbenoxolone, flufenamic acid, octanol						

Nomenclature	Cx31.1	Cx31.9	Cx32	Cx36	Cx37	Cx40	Cx40.1
HGNC, UniProt	GJB5 , O95377	GJD3 , Q8N144	GJB1 , P08034	GJD2 , Q9UKL4	GJA4 , P35212	GJA5 , P36382	GJD4 , Q96KN9
Endogenous inhibitors	extracellular Ca ²⁺ (blocked by raising external Ca ²⁺)						
Inhibitors	carbenoxolone, flufenamic acid, octanol						

Nomenclature	Cx43	Cx45	Cx46	Cx47	Cx50	Cx59	Cx62
HGNC, UniProt	GJA1, P17302	GJC1, P36383	GJA3, Q9Y6H8	GJC2, Q5T442	GJA8, P48165	GJA9, P57773	GJA10, Q969M2
Endogenous inhibitors	extracellular Ca ²⁺ (blocked by raising external Ca ²⁺)						
Inhibitors	carbenoxolone, flufenamic acid, octanol						

Nomenclature	Px1	Px2	Px3
HGNC, UniProt	PANX1, Q96RD7	PANX2, Q96RD6	PANX3, Q96QZ0
Inhibitors	EG-2184 (pI _{C₅₀ 7.3) [761], carbenoxolone, flufenamic acid (little block by flufenamic acid)}	carbenoxolone , flufenamic acid (little block by flufenamic acid)	carbenoxolone , flufenamic acid (little block by flufenamic acid)
Comments	Electrophysiological studies demonstrate that endogenously expressed hPx1 forms intercellular channels with distinct voltage-dependent properties [1068]. Channel function is unaffected by raising external Ca ²⁺ .	Unaffected by raising external Ca ²⁺	Unaffected by raising external Ca ²⁺

Comments: Connexins are most commonly named according to their molecular weights, so, for example, Cx23 is the connexin protein of 23 kDa. This can cause confusion when comparing between species- for example, the mouse connexin Cx57 is orthologous to the human connexin Cx62. No natural toxin or specific inhibitor of junctional channels has been identified yet however two compounds often used experimentally to block connexins are [carbenoxolone](#) and [flufenamic acid](#) [1216]. At least some pannexin hemichannels are more sensitive to [carbenoxolone](#) than connexins but much less sensitive to [flufenamic acid](#) [160]. It has been suggested that 2-aminoethoxydiphenyl borate ([2-APB](#)) may be a more effective blocker of some connexin channel subtypes (Cx26, Cx30, Cx36, Cx40, Cx45, Cx50) compared to others (Cx32, Cx43, Cx46, [66]).

Further reading on Connexins and Pannexins

- Decrock E *et al.* (2015) Connexin and pannexin signaling pathways, an architectural blueprint for CNS physiology and pathology? *Cell Mol Life Sci* **72**: 2823-51 [PMID:26118660]
- Esseltine JL *et al.* (2016) Next-Generation Connexin and Pannexin Cell Biology. *Trends Cell Biol* **26**: 944-955 [PMID:27339936]
- Freund-Michel V *et al.* (2016) Expression and role of connexin-based gap junctions in pulmonary inflammatory diseases. *Pharmacol Ther* **164**: 105-19 [PMID:27126473]
- Harris AL. (2018) Electrical coupling and its channels. *J Gen Physiol* **150**: 1606-1639 [PMID:30389716]
- Navis KE *et al.* (2020) Pannexin 1 Channels as a Therapeutic Target: Structure, Inhibition, and Outlook. *ACS Chem Neurosci* **11**: 2163-2172 [PMID:32639715]
- Palacios-Prado N *et al.* (2022) Endogenous pannexin1 channels form functional intercellular cell-cell channels with characteristic voltage-dependent properties. *Proc Natl Acad Sci U S A* **119**: e2202104119 [PMID:35486697]
- Sáez JC *et al.* (2015) Regulation of pannexin and connexin channels and their functional role in skeletal muscles. *Cell Mol Life Sci* **72**: 2929-35 [PMID:26084874]
- Willebrords J *et al.* (2017) Inhibitors of connexin and pannexin channels as potential therapeutics. *Pharmacol Ther* **180**: 144-160 [PMID:28720428]

Piezo channels

Ion channels → **Other ion channels** → **Piezo channels**

Overview: PIEZO proteins are the pore-forming subunits of trimeric ion channels that open rapidly in response to mechanical stimuli such as membrane stretch, allowing positively charged ions such as the calcium ion to flow into the cell to change cellular structure and function. The proteins are eukaryotic, not prokaryotic, and lack sequence or structural homology to other proteins. There are two main types: In humans, there is PIEZO1 of 2521 amino acids and PIEZO2 of 2752 amino acids, which

can form homomeric PIEZO1 and PIEZO2 channels. Membrane stretching causes complex changes in the channels that include a reduced inward curvature, an expansion of the N-terminal transmembrane helical units and an opening of the central ion pore region comprising the last two C-terminal helices. When there is sustained membrane stretching, the channels may adapt by inactivating at various rates depending on the cell type and context. There are many roles of the channels in diverse organ systems

including cardiovascular, gastrointestinal, haematological, hepatobiliary, immune, integumental, musculoskeletal, neuronal, reproductive, respiratory and urinary systems as well as adipose tissue and cancers. The expression and functions of PIEZO1 are broad and diverse across many cell types and organs, whereas PIEZO2 is more restricted.

Nomenclature	Piezo1	Piezo2
HGNC, UniProt	<i>PIEZO1</i> , Q92508	<i>PIEZO2</i> , Q9H515
Endogenous modulators	Cholesterol, 7-ketocholesterol, phosphatidylserine, phosphatidylinositol 4,5-bisphosphate, ceramide, docosahexaenoic acid, eicosapentaenoic acid, margaric acid, protons, amyloid beta (1-40) peptide	Phosphatidic acid, palmitoyl lysophosphatidic acid
Physical activators	Membrane stretch, cell indentation, fluid shear stress, cell traction force, substrate stiffening	Cell indentation
Selective activators	Yoda2b (pEC ₅₀ 6.9) [867] – Mouse, Yoda2 (pEC ₅₀ 6.8) [1081], Yaddle1 (pEC ₅₀ 6.4) [464], Yoda1 (pEC ₅₀ 4.6) [1349], Jedi1 (pEC ₅₀ 3.8) [1487] – Mouse, Jedi2 (pEC ₅₀ 3.8) [1487] – Mouse	–
Inhibitors	salvianolic acid B (pIC ₅₀ 5.9) [1070], Dooku1 (pIC ₅₀ 5.9) [379], Dooku1 (pIC ₅₀ 5.8) [379], escin (pIC ₅₀ 5.8) [1488], GsMTx-4 (pIC ₅₀ 5.4) [64], benzbromarone (pIC ₅₀ 5.4) [817], Gd³⁺ (@ 30 μM) [256], propofol (@ 50 μM) [1581], ruthenium red (@ 30 μM) [256]	GsMTx-4 (pIC ₅₀ 5.3) [29], Gd³⁺ (@ 30 μM) [256], propofol (@ 50 μM) [1581], ruthenium red (@ 30 μM) [256]
Functional Characteristics	Activation and deactivation, a few milliseconds; Inactivation, time constant 15 ms at -80 mV ranging to seconds or minutes and slower at positive voltages; Unitary conductance, 29 pS; Ion selectivity, monovalent and divalent cations (PCa/PCs 1.21) - Mouse (Mm).	Activation and deactivation, a few milliseconds; Inactivation, time constant 7 ms at -80 mV and slower at positive voltages; Unitary conductance, 24 pS; Ion selectivity, monovalent and divalent cations (PCa/PCs 1.34) - Mouse (Mm).
Comments	No small molecule or toxin binding sites are confirmed on the channel, although an interaction site for Yoda1 has been suggested. The evidence for selective activation is greatest for Yoda1 [379, 1480], but none of the activators is guaranteed to be selective and the selectivity achieved will likely depend on the concentration of the activator and the biological context in which it is administered. Yoda1 has low aqueous solubility, which may be a factor behind the range of pEC ₅₀ s observed for it [1081]. Analogues such as Yoda2 have better aqueous solubility. Yoda1 , Yoda2 , Yoda2b , Yaddle1 and Dooku1 are chemically related molecules (Yoda1 analogues). OB-1 is thought to inhibit the channels <i>via</i> stomatin like 3 (STOML3 ; Q8TAV4), with its effect taking up to 3 hours to occur [1509].	No small molecule or toxin binding sites are confirmed on the channel.

Further reading on Piezo channels

- Cheng CW *et al.* (2025) *PIEZO1* variant implications for biological understanding and human health. *Open Biol* **15**: 240345 [PMID:40628291]
- Coste B *et al.* (2010) Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. *Science* **330**: 55-60 [PMID:20813920]
- Douguet D *et al.* (2019) Mammalian Mechanoelectrical Transduction: Structure and Function of Force-Gated Ion Channels. *Cell* **179**: 340-354 [PMID:31585078]
- Guo YR *et al.* (2017) Structure-based membrane dome mechanism for Piezo mechanosensitivity. *Elife* **6**: [PMID:29231809]
- Kinsella JA *et al.* (2024) Pharmacology of PIEZO1 channels. *Br J Pharmacol* **181**: 4714-4732 [PMID:39402010]
- Mulhall EM *et al.* (2023) Direct observation of the conformational states of PIEZO1. *Nature* **620**: 1117-1125 [PMID:37587339]
- Smith KA *et al.* (2025) Regulation of PIEZO1 channel force sensitivity by interblade handshaking. *Sci Adv* **11**: eadt7046 [PMID:40512861]
- Wang L *et al.* (2019) Structure and mechanogating of the mammalian tactile channel PIEZO2. *Nature* **573**: 225-229 [PMID:31435011]
- Xiao B. (2024) Mechanisms of mechanotransduction and physiological roles of PIEZO channels. *Nat Rev Mol Cell Biol* **25**: 886-903 [PMID:39251883]
- Yang X *et al.* (2022) Structure deformation and curvature sensing of PIEZO1 in lipid membranes. *Nature* **604**: 377-383 [PMID:35388220]
- Zhou Z *et al.* (2023) Mechanisms of PIEZO Channel Inactivation. *Int J Mol Sci* **24**: [PMID:37762415]

Sodium leak channel, non-selective (NALCN)

[Ion channels](#) → [Other ion channels](#) → [Sodium leak channel, non-selective \(NALCN\)](#)

Overview: The sodium leak channel, non selective (NALCN) is a member of the family of four-domain voltage-gated cation channels that include voltage-gated sodium (Na_v) and calcium (Ca_v) channels [781, 1582]. It possesses distinctive ion selectivity and pharmacological properties compared to these latter ion channels [235, 1415]. NALCN, which is insensitive to tetrodotoxin (10 μM), has been proposed to mediate the tetrodotoxin-resistant and voltage-insensitive Na^+ leak current ($I_{\text{L-Na}}$) observed in many types of neurone [861]. However, whether NALCN is constitutively active has been challenged [139, 363,

1348]. NALCN is widely distributed within the central nervous system and is also expressed in the heart and pancreas specifically, in rodents, within the islets of Langerhans [781, 861]. There is now strong functional and structural evidence indicating that NALCN forms a channelosome with obligatory auxiliary subunits UNC79, UNC80 and FAM155A (also known as NALF1) [141, 235, 671, 753, 1624]. NALCN is the pore-forming α subunit, UNC79 and UNC80 are massive HEAT-repeat proteins that form an intertwined anti-parallel superhelical assembly that docks intracellularly onto NALCN. FAM155A forms an extracellular dome

that shields extracellular access pathways to the selective filter of NALCN. There is also increasing evidence suggesting that the NALCN-UNC79-UNC80-FAM155A channelosome is modulated by additional auxiliary subunits including G proteins [1112] and neuronal SNARE complex proteins [1415]. However, there remain many areas of uncertainty surrounding NALCN function. It is worth noting that there is currently no NALCN-specific pharmacology. Inhibitors include multivalent cations (Gd^{3+} , Ca^{2+} , Mg^{2+} , Ba^{2+} , Zn^{2+}) and small molecules (verapamil, 2-APB, DPBA, fluvastatin, L-703,606) [235, 501, 752, 1241].

Nomenclature	NALCN
HGNC, UniProt	NALCN, Q8IZF0
Activators	Constitutively active (Lu <i>et al.</i> , 2007), or activated downstream of Src family tyrosine kinases (SFKs) (Lu <i>et al.</i> , 2009; Swayne <i>et al.</i> , 2009) [235, 861, 862, 863, 1348]
Channel blockers	Gd^{3+} (pIC_{50} 5.6), Ca^{2+} (pIC_{50} 3.9) [235], Cd^{2+} (pIC_{50} 3.8), Co^{2+} (pIC_{50} 3.6), verapamil (pIC_{50} 3.4)
Functional Characteristics	$\gamma = 27$ pS [753], $P_{\text{Na}}/P_{\text{Cs}} = 2.5$, $P_{\text{Na}}/P_{\text{Li}} = 1.0$, $P_{\text{Na}}/P_{\text{K}} = 1.7$ [235], activation $V_{1/2}$ of around +60 mV [1415]

Comments: It has been previously suggested that NALCN function is modulated by different signaling molecules/proteins including NK1, neurotensin, M3R, G proteins, Src tyrosine kinases and CaSR [862, 862, 862, 863, 1348, 1348, 1469]. There is an expansive body of evidence indicating that NALCN forms a complex with UNC79, UNC80 and FAM155 [969]. Clinical reports indicate that NALCN variants cause severe neurodevelopmental disorders including CLIFAHDD (congenital contractures of the limbs and face, hypotonia, and developmental delay), and IHPRF1 (infantile hypotonia with psychomotor retardation and characteristic facies-1) and -2 [24, 230, 657].

Further reading on Sodium leak channel, non-selective (NALCN)

- Lu TZ *et al.* (2012) NALCN: a regulator of pacemaker activity. *Mol Neurobiol* **45**: 415-23 [PMID:22476981]
- Monteil A *et al.* (2024) New insights into the physiology and pathophysiology of the atypical sodium leak channel NALCN. *Physiol Rev* **104**: 399-472 [PMID:37615954]
- Philippart F *et al.* (2018) G_{1/o} protein-coupled receptors in dopamine neurons inhibit the sodium leak channel NALCN. *Elife* **7**: [PMID:30556810]

Orai channels

[Ion channels](#) → [Other ion channels](#) → [Store-operated ion channels](#) → [Orai channels](#)

Overview: Orai channels are pore forming proteins which underlie calcium release-activated calcium (CRAC) channels. In numerous cell types, calcium influx is predominantly governed by store-operated calcium channels (SOCs). The process of store-operated calcium entry (SOCE) is orchestrated through the concerted interaction of two essential molecular components: the pore-forming Orai proteins (Orai1-3) and the endoplasmic reticulum calcium-sensing stromal interaction molecules (STIM1 and STIM2) [1196].

Nomenclature	Orai1	Orai2	Orai3
HGNC, UniProt	ORAI1 , Q96D31	ORAI2 , Q96SN7	ORAI3 , Q9BRQ5
Selective channel blockers	zegocractin (pIC ₅₀ 6.9) [157, 1425], JPIII [89], compound 4k [821]	–	–
Comments	ORAI1 is the gene that encodes the prototypical CRAC store-operated Ca ²⁺ entry (SOCE) channel. CRAC channels are activated by depletion of Ca ²⁺ in the endoplasmic reticulum that results from antigen-induced activation of a range of immunoreceptors (including TCRs, BCRs, Fcγ and Fcε receptors, chemokine GPCRs and some innate pattern-recognition receptors). A constitutively active Orai1 G98S mutant has been identified in patients with tubular aggregate myopathy.	–	–

Comments: The pathophysiological effect of functional CRAC channel deficiency can be caused by loss-of-function mutations in [ORAI1](#) or STIM molecules. Such CRAC channelopathies are characterised by impaired immune cell function and have been identified as an underlying cause of primary immunodeficiency with predominant features that resemble severe combined immunodeficiency disease.

[ORAI1](#) and [ORAI2](#) proteins form heteromeric complexes that constitute the pore of Ca²⁺ release-activated Ca²⁺ (CRAC) channels. Mice with double [Orai1/Orai2](#) knockout have severely impaired T cell function. [NCT04195347](#) is a phase I/II trial evaluating CM4620 (Auxora™), a CRAC channel inhibitor, for treating asparaginase-associated pancreatitis in pediatric and young adult leukemia/lymphoma

patients. The study focuses on determining CM4620's safety, tolerability, and optimal dosing, while exploring its ability to prevent severe complications such as pancreatic necrosis, pseudocyst formation, and SIRS. Administered intravenously over four days soon after symptom onset, CM4620 is being tested as a targeted approach to limit calcium overload-driven pancreatic injury and inflammation, with results expected after 2026 [157].

Further reading on Orai channels

- Lewis RS. (2020) Store-Operated Calcium Channels: From Function to Structure and Back Again. *Cold Spring Harb Perspect Biol* **12**: [PMID:31570335]
- Rubaiy HN. (2023) ORAI Calcium Channels: Regulation, Function, Pharmacology, and Therapeutic Targets. *Pharmaceuticals (Basel)* **16**: [PMID:37259313]

References

- Abbott GW. (2020) [32655402]
- Abbott GW *et al.* (2024) [39390220]
- Abdelsayed M *et al.* (2013) [24065921]
- Abela AR *et al.* (2018) Patent number: WO2018107100A1.
- Abi-Dargham A *et al.* (1993) [8419547]
- Abiria SA *et al.* (2017) [28696294]
- Abramowitz J *et al.* (2009) [18940894]
- Abrams CJ *et al.* (1996) [8799888]
- Accardi A *et al.* (2004) [14985752]
- Accardi A *et al.* (2010) [20188062]
- Acker BA *et al.* (2008) [18490160]
- Adams ME *et al.* (1993) [8232218]
- Agostinelli E *et al.* (2022) [35163502]
- Ahern GP. (2003) [12761211]
- Ahrens J *et al.* (2009) [19204413]
- Ai C *et al.* (2023) [36696767]
- Aierken A *et al.* (2021) [34658162]
- Akbulut Y *et al.* (2015) [25707820]
- Akopian AN *et al.* (2000) [10923674]
- Akopian AN *et al.* (1996) [8538791]
- Al-Hosni R *et al.* (2025) *BJP*
- Al-Hosni R *et al.* (2025) [39829151]
- Al-Hosni R *et al.* (2022) [35811176]
- Al-Sayed MD *et al.* (2013) [24075186]
- Alagem N *et al.* (2001) [11454958]
- Alarcón-Alarcón D *et al.* (2022) [36272975]
- Albaugh PA *et al.* (2002) [12408715]
- Albuquerque EX *et al.* (2009) [19126755]
- Alcaino C *et al.* (2017) [28085630]
- Alekov AK *et al.* (2009) [19364886]
- Alekov AK *et al.* (2008) [19036336]
- Alexander SP *et al.* (2015) [26650442]
- Alexander SP *et al.* (2017) [29055039]
- Alexandrou AJ *et al.* (2016) [27050761]
- Altomare C *et al.* (2003) [12702747]
- Ambudkar IS *et al.* (2007) [17486362]
- Anand D *et al.* (2022) [35276025]
- Anand P *et al.* (2020) [32452762]
- Anderson DJ *et al.* (2008) [17959745]
- Andersson DA *et al.* (2004) [15190109]
- Andersson DA *et al.* (2008) [18322093]
- Andersson DA *et al.* (2009) [19416844]
- André E *et al.* (2008) [18568077]
- Andrews MD *et al.* (2015) [25893043]
- Andrey F *et al.* (2005) [16085050]
- Antunes FTT *et al.* (2024) [38342359]
- Appendino G *et al.* (2005) [15356216]
- Araud T *et al.* (2010) [20346921]
- Arcas JM *et al.* (2019) [30545944]
- Arif Pavel M *et al.* (2016) [27071085]
- Asakawa M *et al.* (2006) [16858425]
- Asano S *et al.* (2012) [22580348]
- AstraZeneca. astrazeneca.com. Accessed on 12/09/2014.
- Atack JR. (2008) [18482097]
- Atack JR. (2010) [21309116]
- Atlason PT *et al.* (2010) [20837679]
- Atobe M *et al.* (2019) [30629441]
- Auberson YP *et al.* (2002) [11909726]
- Auer-Grumbach M *et al.* (2010) [20037588]
- Babinski K *et al.* (2000) [10842183]
- Babinski K *et al.* (1999) [9886053]
- Baconguis I *et al.* (2014) [24507937]
- Badheka D *et al.* (2017) [28829742]
- Bae C *et al.* (2011) [21696149]
- Bahamonde MI *et al.* (2003) [12794078]
- Bai D *et al.* (2006) [16985167]
- Bai X *et al.* (2023) [36574498]
- Bai Y *et al.* (2020) [32149605]
- Bai Z *et al.* (2021) [34928937]
- Bal M *et al.* (2008) [18786918]
- Balestrini A *et al.* (2017) [33620419]
- Bandell M *et al.* (2004) [15046718]
- Bang S *et al.* (2007) [17850966]
- Bang-Andersen B *et al.* (2011) [21486038]
- Baraldi PG *et al.* (2010) [20356305]
- Barbet G *et al.* (2008) [18758465]
- Barel O *et al.* (2008) [18678320]
- Barford ET *et al.* (2001) [11245600]
- Barnard EA *et al.* (1998) [9647870]
- Barnes NM *et al.* (2009) [18761359]
- Baron A *et al.* (2013) [23624383]
- Baron A *et al.* (2015) [25613302]
- Baron A *et al.* (2001) [11457851]
- Baron A *et al.* (2008) [18256271]
- Barrera NP *et al.* (2005) [16116092]
- Barrere C *et al.* (2020) *Archives of Cardiovascular Diseases Supplements* 12: 259
- Barret DCA *et al.* (2022) [34969975]
- Bartok A *et al.* (2014) [24878374]
- Bartoli F *et al.* (2020) [31906693]
- Baukowitz T *et al.* (1998) [9804555]
- Baumann L *et al.* (2004) [14744918]
- Bautista DM *et al.* (2006) [16564016]
- Bautista DM *et al.* (2005) [16103371]
- Bautista DM *et al.* (2007) [17538622]
- Bavencoffe A *et al.* (2017) [28900918]
- Beck A *et al.* (2006) [16585058]
- Becker V *et al.* (2021) [33655414]
- Beckmann H *et al.* (2017) [28807145]
- Beech DJ. (2012) [21624095]
- Beech DJ. (2013) [23412755]
- Behrendt HJ *et al.* (2004) [14757700]
- Beigi RD *et al.* (2003) [14522842]
- Belelli D *et al.* (1995) [8848005]
- Belelli D *et al.* (2009) [19828786]
- Belelli D *et al.* (2005) [15959466]
- Bell PD *et al.* (2003) [12655045]
- Belrose JC *et al.* (2018) [29671419]
- Bentzen BH *et al.* (2020) [32477117]
- Bentzen BH *et al.* (2007) [17636045]
- Bessac BF *et al.* (2008) [18398506]
- Best L *et al.* (2010) [21099297]
- Bettini E *et al.* (2010) [20810618]
- Beumer J *et al.* (2021) [32958874]
- Bezençon O *et al.* (2017) [29116786]
- Bhattacharjee A *et al.* (2003) [14684870]
- Bhattacharya A *et al.* (2013) [23889535]
- Bianchi BR *et al.* (2007) [17660385]
- Bianchi MT *et al.* (2003) [14645489]
- Bienert GP *et al.* (2014) [24060746]
- Birnbaumer L. (2009) [19281310]
- Bitner RS *et al.* (2007) [17898229]
- Biton B *et al.* (2012) [22171093]
- Bize V *et al.* (2007) [17670907]
- Blanc E *et al.* (1998) [9730813]
- Blanchard MG *et al.* (2012) [21943094]
- Blin S *et al.* (2016) [27035965]
- Blin S *et al.* (2014) [25148687]
- Blum CA *et al.* (2010) [20307063]
- Bodnar AL *et al.* (2005) [15715459]
- Bodtker DM *et al.* (2015) [26013995]
- Bogdanović R *et al.* (2012) [21956615]
- Boggula VR *et al.* (2018) [29885352]
- Bohannon BM *et al.* (2020) [32207683]
- Bohlen CJ *et al.* (2011) [22094702]
- Bohlen CJ *et al.* (2010) [20510930]
- Bon RS *et al.* (2013) [23763262]
- Bon RS *et al.* (2022) [34499525]
- Bonin RP *et al.* (2008) [18201756]
- Boone AN *et al.* (2014) [24945283]
- BoSmith RE *et al.* (1993) [7693281]
- Bouasse M *et al.* (2019) [31409833]
- Bourinet E *et al.* (1999) [10321243]
- Bourque CW. (2008) [18509340]
- Bowen CV *et al.* (2013) [23554944]
- Bowie D. (2010) [19822544]
- Boyer R *et al.* (2025) [40427633]
- Brady CA *et al.* (2001) [11489465]
- Brams M *et al.* (2020) [31990273]
- Brenker C *et al.* (2012) [22354039]
- Bricelj VM *et al.* (2005) [15815630]
- Broad LM *et al.* (2006) [16738207]
- Broertjes J *et al.* (2019) [30615643]
- Brône B *et al.* (2008) [18501939]
- Brooks CA *et al.* (2019) [31532662]
- Brooks CA *et al.* (2019) [31413810]
- Brown N *et al.* (2002) [12145096]
- Bruen C *et al.* (2021) [33939666]
- Brunner FS *et al.* (2017) [28320947]
- Brüss M *et al.* (2000) [11111833]
- Bruzzone R *et al.* (2005) [15715654]
- Bruzzone R *et al.* (2003) [14597722]
- Bucchi A *et al.* (2002) [12084770]
- Buell G *et al.* (1996) [8598206]
- Buisson B *et al.* (1996) [8987816]
- Buta A *et al.* (2015) [25974655]
- Butterworth MB *et al.* (2008) [18713729]
- Cabanas H *et al.* (2018) [30134818]
- Cadiou H *et al.* (2007) [18045919]
- Cai W *et al.* (2024) [38467901]
- Calcraft PJ *et al.* (2009) [19387438]
- Canessa CM *et al.* (1994) [7810611]
- Canessa CM *et al.* (1994) [8107805]
- Cang C *et al.* (2014) [25256615]
- Cang C *et al.* (2014) [24776928]
- Cang C *et al.* (2013) [23394946]
- Cantrell BE *et al.* (1996) [8809152]
- Cao E *et al.* (2013) [24305161]
- Capasso M *et al.* (2011) [20961760]
- Capogna M *et al.* (2011) [21145601]
- Carlos Dos Reis D *et al.* (2023) [36967909]
- Carlson AE *et al.* (2009) [19718436]
- Carlson AE *et al.* (2005) [16036917]
- Carstens E *et al.* (2014) [24830011]
- Castle NA *et al.* (2003) [12527813]
- Catterall WA *et al.* (2005) [16382098]
- Catterall WA *et al.* (2020) [31537174]
- Catterall WA *et al.* (1981) [6114956]
- Catterall WA *et al.* (2005) [16382099]
- Cavara NA *et al.* (2008) [18654865]

190. Cazals Y *et al.* (2015) [26549439]
 191. Celie PH *et al.* (2004) [15046723]
 192. Chahine M *et al.* (1994) [8058462]
 193. Chai H *et al.* (2019) [30620187]
 194. Chang SS *et al.* (1996) [8589714]
 195. Changeux JP. (2010) [20055696]
 196. Changeux JP *et al.* (2008) [18262468]
 197. Chappe V *et al.* (2003) [12588899]
 198. Chassagnon IR *et al.* (2017) [28320941]
 199. Chatterton JE *et al.* (2002) [11823786]
 200. Chaudhari SS *et al.* (2013) [24055075]
 201. Chaumont S *et al.* (2008) [18689682]
 202. Chaves G *et al.* (2016) [26866814]
 203. Chavez-Noriega LE *et al.* (1997) [8996215]
 204. Chazot PL *et al.* (1998) [9721050]
 205. Chebib M. (2004) [15566397]
 206. Chen CC *et al.* (2017) [28732201]
 207. Chen CC *et al.* (1998) [9707631]
 208. Chen CC *et al.* (2014) [25119295]
 209. Chen J *et al.* (2011) [21402443]
 210. Chen PE *et al.* (2008) [17962328]
 211. Chen PE *et al.* (2006) [16474411]
 212. Chen TY. (2005) [15709979]
 213. Chen TY *et al.* (1993) [7682292]
 214. Chen X *et al.* (2005) [15955877]
 215. Chen X *et al.* (2006) [16505147]
 216. Chen X *et al.* (2010) [20114056]
 217. Chen X *et al.* (2020) [32872338]
 218. Chen XZ *et al.* (1999) [10517637]
 219. Chen Y *et al.* (2003) [12871651]
 220. Chen YH *et al.* (2000) [11122339]
 221. Chen Z *et al.* (2023) [36973510]
 222. Chen ZW *et al.* (2007) [17083446]
 223. Cheng KT *et al.* (2011) [21290310]
 224. Cheng KT *et al.* (2013) [23890115]
 225. Cheng YR *et al.* (2018) [29793480]
 226. Cherian OL *et al.* (2015) [25620774]
 227. Cherny VV *et al.* (1999) [10578017]
 228. Chien YC *et al.* (2025) [40264173]
 229. Choe H *et al.* (2000) [10736307]
 230. Chong JX *et al.* (2015) [25683120]
 231. Chopra B *et al.* (2000) [10780999]
 232. Chow PH *et al.* (2020) [32434851]
 233. Chu XP *et al.* (2006) [16707785]
 234. Chu XP *et al.* (2004) [15470133]
 235. Chua HC *et al.* (2020) [32494638]
 236. Chuang RS *et al.* (1998) [10196582]
 237. Chung JJ *et al.* (2011) [21224844]
 238. Chung JJ *et al.* (2014) [24813608]
 239. Chung MK *et al.* (2005) [15722340]
 240. Chung MK *et al.* (2004) [15175387]
 241. Chung S *et al.* (2022) [34817438]
 242. Church TW *et al.* (2015) [25421315]
 243. Cil O *et al.* (2019) [31299174]
 244. Cil O *et al.* (2021) [33836171]
 245. Clapham DE. (2003) [14654832]
 246. Clapham DE *et al.* (2005) [16382101]
 247. Clapham DE *et al.* (2003) [14657417]
 248. Coe JW *et al.* (2005) [16171993]
 249. Colburn RW *et al.* (2007) [17481392]
 250. Coleman N *et al.* (2014) [24958817]
 251. Coleman SK *et al.* (1999) [10428084]
 252. Collier DM *et al.* (2011) [21149458]
 253. Collingridge GL *et al.* (2009) [18655795]
 254. Compan V *et al.* (2012) [22442090]
 255. Connors BW *et al.* (2004) [15217338]
 256. Coste B *et al.* (2010) [20813920]
 257. Cottrell GS *et al.* (2018) [30181139]
 258. Crack BE *et al.* (1994) [7889301]
 259. Crest M *et al.* (1992) [1730708]
 260. Cribbs LL *et al.* (1990) [2175715]
 261. Csanády L *et al.* (2019) [31205003]
 262. Csanády L *et al.* (2019) [30516439]
 263. Csanády L *et al.* (2010) [19966305]
 264. Cuajungco MP *et al.* (2016) [26336837]
 265. Cull-Candy S *et al.* (2006) [16713244]
 266. Cull-Candy SG *et al.* (2004) [15494561]
 267. Cummins TR *et al.* (1999) [10594087]
 268. Curcic S *et al.* (2019) [30974125]
 269. Czirájk G *et al.* (2002) [11733509]
 270. Díaz-Franulic I *et al.* (2020) [32747539]
 271. Dai L *et al.* (2010) [20176855]
 272. Dai XQ *et al.* (2007) [17804601]
 273. Danahay H *et al.* (2020) [32821878]
 274. Danahay H *et al.* (2020) [32235608]
 275. Danahay H *et al.* (2025) [39561495]
 276. Danahay H *et al.* (2023) [36950016]
 277. Danahay H *et al.* (2025) [38851923]
 278. Danahay HL *et al.* (2020) [31898911]
 279. Daniel C *et al.* (2009) [19909284]
 280. Das P *et al.* (2003) [14625088]
 281. Dascal N *et al.* (1993) [8234283]
 282. Davenport AJ *et al.* (2021) [34615939]
 283. Davies PA *et al.* (1999) [9950429]
 284. Davies PA *et al.* (2003) [12381728]
 285. Day RE *et al.* (2014) [24090884]
 286. De Clercq K *et al.* (2021) [33884443]
 287. de Groot T *et al.* (2008) [18596722]
 288. de Jonge HR *et al.* (2020) [33113586]
 289. De La Fuente R *et al.* (2008) [18083779]
 290. de Weille JR *et al.* (1998) [9744806]
 291. de Weille JR *et al.* (1991) [1848702]
 292. DeCaen PG *et al.* (2016) [27348301]
 293. Decher N *et al.* (2011) [21865850]
 294. DeCoursey TE. (2008) [18463791]
 295. DeCoursey TE. (2008) [18801839]
 296. DeCoursey TE. (2003) [12663866]
 297. DeCoursey TE. (2017) [29023793]
 298. DeCoursey TE. (2010) [20134026]
 299. DeCoursey TE *et al.* (2000) [10823889]
 300. DeCoursey TE *et al.* (2003) [12673252]
 301. Deeb TZ *et al.* (2009) [19131665]
 302. del Camino D *et al.* (2010) [21068322]
 303. Del Lungo M *et al.* (2012) [22091830]
 304. Dellaunay A *et al.* (2012) [22829666]
 305. Dellisanti CD *et al.* (2007) [17643119]
 306. Delmas P. (2005) [15889307]
 307. Delmas P *et al.* (2004) [14766803]
 308. Delmas P *et al.* (2004) [15336986]
 309. Deemba S *et al.* (2017) [28826482]
 310. Demir R *et al.* (2009) [19307742]
 311. Deng HX *et al.* (2010) [20037587]
 312. Deval E *et al.* (2015) [25724084]
 313. Deval E *et al.* (2011) [21508231]
 314. Deval E *et al.* (2008) [18923424]
 315. Dhaka A *et al.* (2007) [17481391]
 316. Di Virgilio F. (2003) [14522840]
 317. Di Virgilio F *et al.* (2018) [29439897]
 318. Diaz M *et al.* (2001) [11579158]
 319. Dibattista M *et al.* (2024) [38677213]
 320. Dietrich A *et al.* (2014) [25268281]
 321. Dietrich PS *et al.* (1998) [9603190]
 322. DiFrancesco D. (1993) [7682405]
 323. Diness JG *et al.* (2017) [29018164]
 324. Ding M *et al.* (2018) [29859875]
 325. Dinko T *et al.* (2011) [21084390]
 326. Diochot S *et al.* (2004) [15044953]
 327. Diochot S *et al.* (2012) [23034652]
 328. Diochot S *et al.* (2007) [17113616]
 329. Diochot S *et al.* (1998) [9506974]
 330. Do HQ *et al.* (2023) [37026993]
 331. Doñate-Macian P *et al.* (2022) [32959389]
 332. Dogruer D *et al.* (2004) [14698195]
 333. Dong XP *et al.* (2008) [18794901]
 334. Do XP *et al.* (2010) [20802798]
 335. Donier E *et al.* (2008) [18662336]
 336. Donnelly-Roberts DL *et al.* (2007) [17471177]
 337. Donnelly-Roberts DL *et al.* (2004) [14634045]
 338. Donnelly-Roberts DL *et al.* (2009) [19558545]
 339. Döring F *et al.* (1998) [9786970]
 340. Doyle DA *et al.* (1998) [9525859]
 341. Dravid SM *et al.* (2007) [17303642]
 342. Dryer SE *et al.* (1991) [1719422]
 343. Du J *et al.* (2015) [26344198]
 344. Du J *et al.* (2014) [24889629]
 345. Du J *et al.* (2009) [19372375]
 346. Dubé GR *et al.* (2005) [16061325]
 347. Dubin AE *et al.* (1999) [10521471]
 348. Duc C *et al.* (1994) [7806569]
 349. Dudev T *et al.* (2015) [25955978]
 350. Dumani A *et al.* (2025) [40635742]
 351. Dunlop J *et al.* (2008) [18356919]
 352. Duprat F *et al.* (2000) [10779373]
 353. Dupuis DS *et al.* (2002) [11890900]
 354. Duran C *et al.* (2010) [19827947]
 355. Dutta AK *et al.* (2002) [12154180]
 356. Dutta Banik D *et al.* (2018) [29311301]
 357. Dutzler R. (2007) [17452037]
 358. Dutzler R *et al.* (2002) [11796999]
 359. Edelheit O *et al.* (2014) [24571549]
 360. Edman S *et al.* (2012) [22579927]
 361. Edwards E *et al.* (1996) [8891244]
 362. Edwards JC *et al.* (2010) [20100480]
 363. Egan JM *et al.* (2018) [29892559]
 364. Eichler SA *et al.* (2008) [19210758]
 365. El Chemaly A *et al.* (2010) [20026664]
 366. El Kouhen R *et al.* (2005) [15837819]
 367. Emir TLR. (2017) *In Neurobiology of TRP Channels (2nd Ed.)* Edited by Emir TLR: CRC Press/Taylor & Francis: [PMID:29356469]
 368. Enuka Y *et al.* (2012) [22207244]
 369. Er SY *et al.* (2017) [28327374]
 370. Erreger K *et al.* (2004) [15701057]
 371. Erreger K *et al.* (2007) [17622578]
 372. Ertel EA *et al.* (2000) [10774722]
 373. Esarte Palomero O *et al.* (2023) [36763973]
 374. Escalera J *et al.* (2008) [18550530]
 375. Escoubas P *et al.* (2000) [10829030]
 376. Esteban JA. (2008) [18026130]
 377. Esteva-Font C *et al.* (2016) [26993802]
 378. Estévez R *et al.* (2001) [11734858]
 379. Evans EL *et al.* (2018) [29498036]
 380. Evans IA *et al.* (2024) [38646935]
 381. Evans WH *et al.* (2002) [12126230]
 382. Everaerts W *et al.* (2010) [20956320]
 383. Fabjan J *et al.* (2020) [33519364]
 384. Fahlke C *et al.* (2010) [21423394]

385. Fan J *et al.* (2023) [36302896]
 386. Fanger CM *et al.* (2001) [11278890]
 387. Fantoni ER *et al.* (2017) [28374288]
 388. Farrant M *et al.* (2005) [15738957]
 389. Fecher-Trost C *et al.* (2014) [24756713]
 390. Fedida D *et al.* (1993) [8508531]
 391. Fedigan S *et al.* (2017) [28733893]
 392. Felix JP *et al.* (1999) [10213593]
 393. Feng B *et al.* (2004) [14718249]
 394. Feng L *et al.* (2010) [20929736]
 395. Feng Z *et al.* (2024) [38749289]
 396. Ferber M *et al.* (2004) [15208025]
 397. Fernández Á *et al.* (2024) [38908487]
 398. Ferrari D *et al.* (2004) [15383600]
 399. Ferrera L *et al.* (2009) [19819874]
 400. Fessenko EE *et al.* (1985) [2578616]
 401. Fiedorczuk K *et al.* (2022) [36264792]
 402. Fink M *et al.* (1998) [9628867]
 403. Finol-Urdaneta RK *et al.* (2023) [37260417]
 404. Fischer M *et al.* (2010) [20538786]
 405. Fisher JL. (2009) [18585399]
 406. Fisher JL. (2002) [12021393]
 407. Flannery RJ *et al.* (2015) [26290373]
 408. Fleig A *et al.* (2004) [15530641]
 409. Fliegert R *et al.* (2017) [28671679]
 410. Fonfria E *et al.* (2004) [15302683]
 411. Ford AP *et al.* (2006) [16465177]
 412. Fouda MA *et al.* (2022) [35436004]
 413. Francois A *et al.* (2013) [23257507]
 414. Freichel M *et al.* (2005) [15975974]
 415. Friese MA *et al.* (2007) [17994101]
 416. Frizelle PA *et al.* (2006) [16778008]
 417. Frølund B *et al.* (2002) [12171573]
 418. Fucile S *et al.* (2000) [11144365]
 419. Fuller MD *et al.* (2007) [17951250]
 420. Furukawa H *et al.* (2005) [16281028]
 421. Galletta LJV. (2022) [35364521]
 422. Galvez A *et al.* (1990) [1694175]
 423. Gao L *et al.* (2000) [10920015]
 424. Gao P *et al.* (2020) [31735085]
 425. Gao X *et al.* (2024) [39107303]
 426. Gao YY *et al.* (2021) [33763843]
 427. Garceau D *et al.* (2019) [30902655]
 428. Garcia ML *et al.* (1995) [7543240]
 429. García-Añoveros J *et al.* (1997) [9037075]
 430. García-Añoveros J *et al.* (2007) [17217068]
 431. García-Calvo M *et al.* (1993) [8360176]
 432. Garcia-Valdes J *et al.* (2001) [11576530]
 433. Garg P *et al.* (2012) [22851714]
 434. Gargett CE *et al.* (1997) [9113369]
 435. Garneau AP *et al.* (2015) [26313002]
 436. Gaunt HJ *et al.* (2016) [27289383]
 437. Gavva NR *et al.* (2005) [15615864]
 438. Genovese M *et al.* (2019) [31622498]
 439. Genovese M *et al.* (2023) [36444690]
 440. Genovese M *et al.* (2024) [38788257]
 441. Gentzsch M *et al.* (2020) [33201937]
 442. Gerndt S *et al.* (2020) [32167471]
 443. Gerstner A *et al.* (2000) [10662822]
 444. Geyer RR *et al.* (2013) [23485707]
 445. Ghanshani S *et al.* (2000) [10961988]
 446. Ghovanloo MR *et al.* (2021) [33836525]
 447. Ghovanloo MR *et al.* (2024) [38263462]
 448. Ghovanloo MR *et al.* (2022) [35297036]
 449. Ghovanloo MR *et al.* (2018) [30219789]
 450. Ghovanloo MR *et al.* (2025) [40090585]
 451. Ghovanloo MR *et al.* (2025) [39835903]
 452. Giamarchi A *et al.* (2006) [16880824]
 453. Giangiacomo KM *et al.* (1998) [9758866]
 454. Gielen M *et al.* (2009) [19404260]
 455. Gilchrist JM *et al.* (2024) [38195157]
 456. Gill JK *et al.* (2011) [21436053]
 457. Giraldez T *et al.* (2012) [22573384]
 458. Gochman A *et al.* (2023) [36747846]
 459. Goldin AL *et al.* (2000) [11144347]
 460. Goldstein SA *et al.* (2005) [16382106]
 461. Gonzales EB *et al.* (2009) [19641589]
 462. Gonzalez C *et al.* (2010) [20023639]
 463. Goodchild SJ *et al.* (2024) [38359277]
 464. Goon S *et al.* (2024) [38716967]
 465. Gordon JC *et al.* (2010) [20674564]
 466. Gornati D *et al.* (2021) [34097384]
 467. Gotti C *et al.* (2009) [19481063]
 468. Gotti C *et al.* (2006) [16876883]
 469. Gründer S *et al.* (2023) [36479972]
 470. Grace MS *et al.* (2014) [24286227]
 471. Grand T *et al.* (2008) [18297105]
 472. Graves AR *et al.* (2008) [18449189]
 473. Greenwood IA *et al.* (2007) [17150263]
 474. Griffin AM *et al.* (2021) [33859800]
 475. Grimm C *et al.* (2007) [18048323]
 476. Grimm C *et al.* (2014) [25144390]
 477. Grimm C *et al.* (2012) [22753890]
 478. Grimm C *et al.* (2010) [20189104]
 479. Grimm C *et al.* (2003) [12672799]
 480. Grimm C *et al.* (2005) [15550678]
 481. Grissmer S *et al.* (1994) [7517498]
 482. Gründer S. (2020) *In The Oxford Handbook of Neuronal Ion Channels* Oxford University Press:
 483. Gründer S *et al.* (2000) [10852210]
 484. Gründer S *et al.* (2015) [25585135]
 485. Grupe A *et al.* (1990) [2347305]
 486. Guan YY *et al.* (2006) [16697056]
 487. Guilbault C *et al.* (2007) [16888286]
 488. Guinamard R *et al.* (2011) [21290294]
 489. Guler AD *et al.* (2002) [12151520]
 490. Gunthorpe MJ *et al.* (2009) [19063991]
 491. Gunthorpe MJ *et al.* (2007) [17392405]
 492. Gunthorpe MJ *et al.* (2004) [14654105]
 493. Gururaja Rao S *et al.* (2018) [30040212]
 494. Gutman GA *et al.* (2005) [16382104]
 495. Ha K *et al.* (2020) [33164752]
 496. Hadida S *et al.* (2014) [25441013]
 497. Hadingham KL *et al.* (1996) [8632757]
 498. Hadingham KL *et al.* (1995) [8719414]
 499. Hadley JK *et al.* (2000) [10711337]
 500. Häfner S *et al.* (2018) [29522264]
 501. Hahn S *et al.* (2020) [32436268]
 502. Halaszovich CR *et al.* (2000) [10970899]
 503. Haloi N *et al.* (2025) [40215320]
 504. Han Q *et al.* (2018) [30033153]
 505. Han Y *et al.* (2021) [33501656]
 506. Hanna MC *et al.* (2000) [10854267]
 507. Hansen KB *et al.* (2012) [22553026]
 508. Hansen KB *et al.* (2021) [34753794]
 509. Hanukoglu A. (1991) [1939532]
 510. Hanukoglu A *et al.* (2008) [18634878]
 511. Hanukoglu A *et al.* (2018) [29702750]
 512. Hanukoglu I *et al.* (2017) [28130590]
 513. Hanukoglu I *et al.* (2016) [26772908]
 514. Hara Y *et al.* (2002) [11804595]
 515. Harbeson SL *et al.* (2017) [28611092]
 516. Harding SD *et al.* (2018) [29149325]
 517. Harteneck C. (2005) [15843919]
 518. Harteneck C *et al.* (2011) [20932261]
 519. Hartzell C *et al.* (2005) [15709976]
 520. Hattori M *et al.* (2012) [22535247]
 521. Hauser TA *et al.* (2009) [19482012]
 522. Hawn MB *et al.* (2021) [34488544]
 523. He LP *et al.* (2005) [15647288]
 524. He Y *et al.* (2006) [16478442]
 525. Heckel A *et al.* (2017) Patent number: WO2017028927A1.
 526. Heitzmann D *et al.* (2008) [18034154]
 527. Hejazi N *et al.* (2006) [16332990]
 528. Held K *et al.* (2022) [32780479]
 529. Held K *et al.* (2015) [25733887]
 530. Hemmings HC *et al.* (2005) [16126282]
 531. Henderson LM *et al.* (1987) [2825632]
 532. Henson MA *et al.* (2010) [20097255]
 533. Herd MB *et al.* (2007) [17531325]
 534. Hernandez-Olmos V *et al.* (2012) [23075067]
 535. Herrington J *et al.* (2011) [21948463]
 536. Herrington J *et al.* (2006) [16567526]
 537. Hervé JC *et al.* (2007) [17507078]
 538. Hibbs RE *et al.* (2011) [21572436]
 539. Higham J *et al.* (2019) [31048549]
 540. Hildebrand MS *et al.* (2010) [20648059]
 541. Hilf RJ *et al.* (2009) [19646860]
 542. Hilgemann DW *et al.* (1996) [8688080]
 543. Hill K *et al.* (2004) [15275834]
 544. Hill K *et al.* (2004) [15549272]
 545. Hille B. (1986) [2452140]
 546. Hillyard DR *et al.* (1992) [1352986]
 547. Hinman A *et al.* (2006) [17164327]
 548. Hirata T *et al.* (2007) [17652911]
 549. Hirzel K *et al.* (2006) [17114051]
 550. Ho IHT *et al.* (2025) [39542827]
 551. Ho K *et al.* (1993) [7680431]
 552. Ho K *et al.* (2009) [19210926]
 553. Ho MW *et al.* (1997) [9142840]
 554. Hoenderop JG *et al.* (2001) [11744752]
 555. Hofherr A *et al.* (2011) [21290302]
 556. Hofmann F *et al.* (2005) [16382102]
 557. Hofmann T *et al.* (2003) [12842017]
 558. Hofmann T *et al.* (2014) [24633576]
 559. Holbrook JD *et al.* (2009) [19012743]
 560. Holm LM *et al.* (2005) [15988592]
 561. Holm LM *et al.* (2004) [14985982]
 562. Holst AG *et al.* (2024) [38092897]
 563. Honore P *et al.* (2009) [19464323]
 564. Honore P *et al.* (2006) [16982702]
 565. Hope AG *et al.* (1993) [7683998]
 566. Hope AG *et al.* (1996) [8818349]
 567. Horak M *et al.* (2006) [16257494]
 568. Horisberger JD *et al.* (2004) [14988660]
 569. Hosie AM *et al.* (2007) [17560657]
 570. Hosseini R *et al.* (2001) [11533126]
 571. Hougaard C *et al.* (2007) [17486140]
 572. Hougaard C *et al.* (2012) [22046005]
 573. Houser A *et al.* (2025) [39667931]
 574. Houtani T *et al.* (2005) [16083862]
 575. Howard RJ *et al.* (2011) [21730162]
 576. Hoyer D *et al.* (1994) [7938165]
 577. Hu H *et al.* (2009) [19202543]
 578. Hu HZ *et al.* (2004) [15194687]
 579. Hu XQ *et al.* (2008) [18187416]
 580. Hu Y *et al.* (2025) [40356890]
 581. Huang CL *et al.* (1998) [9486652]

582. Huang F *et al.* (2012) [22090471]
 583. Huang J *et al.* (2023) [37330538]
 584. Huang X *et al.* (2015) [26416729]
 585. Huang X *et al.* (2017) [28479061]
 586. Huang X *et al.* (2017) [27991902]
 587. Huang Y *et al.* (2023) [37976835]
 588. Huang Y *et al.* (2019) [31513012]
 589. Hughes AD *et al.* (1993) [8388295]
 590. Hughes BA *et al.* (2000) [10942728]
 591. Hurlle B *et al.* (2003) [12651873]
 592. Hurst RS *et al.* (1991) [1921987]
 593. Hurst RS *et al.* (2005) [15858066]
 594. Hurth K *et al.* ACS Med Chem Meeting Abstracts, March 2014. Accessed on 06/04/2014.
 595. Hwang SW *et al.* (2000) [10823958]
 596. Illes P *et al.* (2021) [33125712]
 597. Inagaki N *et al.* (1995) [7502040]
 598. Inagaki N *et al.* (1996) [8630239]
 599. Inoue R *et al.* (2001) [11179201]
 600. Introini B *et al.* (2005) [39232129]
 601. Iraci N *et al.* (2024) [38782404]
 602. Irie S *et al.* (2014) [24756714]
 603. Isaac JT *et al.* (2007) [17582328]
 604. Isbrandt D *et al.* (2000) [10729221]
 605. Isenberg KE *et al.* (1993) [7509203]
 606. Ishibashi K *et al.* (1997) [9252401]
 607. Ishibashi K *et al.* (2002) [12084581]
 608. Ishibashi K *et al.* (2014) [24189537]
 609. Ishihara K *et al.* (1996) [8866861]
 610. Islam MS. (2011) [21290328]
 611. Isom LL. (2001) [11486343]
 612. Jackson AC *et al.* (2011) [21521608]
 613. Jacob TC *et al.* (2008) [18382465]
 614. Jacobs G *et al.* (2015) [25600961]
 615. Jacobson KA *et al.* (2009) [18600475]
 616. Jacobson KA *et al.* (2002) [12213051]
 617. Jacobson KA *et al.* (1998) [9632352]
 618. Jacobson KA *et al.* (2016) [26686393]
 619. Jäger H *et al.* (2000) [10713270]
 620. Jane DE *et al.* (2009) [18793656]
 621. Jane DE *et al.* (2000) *In Handbook of Experimental Pharmacology, Pharmacology of Ionic Channel Function: Activators and Inhibitors* Edited by Endo M, Kurachi Y, Mishina M: Springer: 415-478
 622. Janssens A *et al.* (2018) [30170189]
 623. Jarvis MF *et al.* (2002) [12482951]
 624. Jarvis MF *et al.* (2009) [18657557]
 625. Jarvis MF *et al.* (2014) [24726441]
 626. Jasti J *et al.* (2007) [17882215]
 627. Jensen AA *et al.* (2008) [18597859]
 628. Jensen AA *et al.* (2005) [16033252]
 629. Jensen BS. (2002) [12481191]
 630. Jensen BS *et al.* (1998) [9730970]
 631. Jensen TN *et al.* (2006) [16647053]
 632. Jentsch TJ. (2008) [18307107]
 633. Jeon J *et al.* (2020) [33490083]
 634. Jiang J *et al.* (2005) [16009728]
 635. Jiang LH *et al.* (2003) [14523092]
 636. Jiang LH *et al.* (2000) [10860929]
 637. Jiang Q *et al.* (2010) [20580786]
 638. Jin J *et al.* (2008) [18974357]
 639. Jin JL *et al.* (2005) [16107607]
 640. Jin W *et al.* (1999) [10572004]
 641. Jin W *et al.* (1999) [10572003]
 642. Jo S *et al.* (2020) [32193331]
 643. Jo S *et al.* (2025) [40136042]
 644. Joeres N *et al.* (2016) [27277303]
 645. Johnson BD *et al.* (1996) [8913371]
 646. Johnson JP *et al.* (2022) [35234610]
 647. Johnston GA. (2005) [15974965]
 648. Johnston GA *et al.* (2010) [20963487]
 649. Joiner WJ *et al.* (1997) [9380751]
 650. Jones AK *et al.* (2010) [20674046]
 651. Jones F *et al.* (2021) [33860384]
 652. Jones J *et al.* (2023) [37530822]
 653. Jordt SE *et al.* (2004) [14712238]
 654. Jung S *et al.* (2003) [12456670]
 655. Just S *et al.* (2018) [29385160]
 656. Juvin V *et al.* (2007) [17673572]
 657. Köroğlu Ç *et al.* (2013) [23749988]
 658. Kaczmarek LK *et al.* (2017) [28267675]
 659. Kaczor AA *et al.* (2010) [20491632]
 660. Kahler JP *et al.* (2023) [36762958]
 661. Kai H *et al.* (2021) [34587541]
 662. Kalamida D *et al.* (2007) [17651090]
 663. Kaldenhoff R *et al.* (2014) [24141139]
 664. Kalman K *et al.* (1998) [9488722]
 665. Kamei T *et al.* (2024) [38776832]
 666. Kanaya S *et al.* (1984) [6694106]
 667. Kang D *et al.* (2005) [15677687]
 668. Kang D *et al.* (2004) [14985088]
 669. Kang J *et al.* (2012) [22353878]
 670. Kang J *et al.* (2005) [15548764]
 671. Kang Y *et al.* (2022) [35550517]
 672. Kapeller J *et al.* (2011) [21192076]
 673. Kapus A *et al.* (1993) [8270911]
 674. Karakas E *et al.* (2011) [21677647]
 675. Karashima Y *et al.* (2007) [17855602]
 676. Karashima Y *et al.* (2009) [19144922]
 677. Karim N *et al.* (2013) [23385381]
 678. Karnovsky AM *et al.* (2003) [14597179]
 679. Kasanami Y *et al.* (2022) [36075145]
 680. Kashlan OB *et al.* (2012) [22405998]
 681. Kassack MU *et al.* (2004) [15072843]
 682. Kasumu AW *et al.* (2012) [23102227]
 683. Kaszas K *et al.* (2012) [22005042]
 684. Katoh TA *et al.* (2023) [36603091]
 685. Kaupp UB *et al.* (1989) [2481236]
 686. Kawate T *et al.* (2009) [19641588]
 687. Kaye R *et al.* (2025) [40491382]
 688. Keating C *et al.* (2025) [39756424]
 689. Keating D *et al.* (2018) [30334692]
 690. Kelemen B *et al.* (2020) [31987857]
 691. Kellenberger S *et al.* (2003) [14500741]
 692. Kellenberger S *et al.* (2015) [25287517]
 693. Kelley SP *et al.* (2003) [12867984]
 694. Kennard LE *et al.* (2005) [15685212]
 695. Keserü GM. (2003) [12873512]
 696. Kew JN *et al.* (2005) [15731895]
 697. Khairatkar-Joshi N *et al.* (2015) [26030081]
 698. Khakh BS *et al.* (2001) [11171941]
 699. Khakh BS *et al.* (1999) [10460235]
 700. Khom S *et al.* (2006) [16272224]
 701. Khom S *et al.* (2010) [20718240]
 702. Kim HJ *et al.* (2007) [17962195]
 703. Kim HJ *et al.* (2008) [18369318]
 704. King BF. (2023) [36279087]
 705. King BF *et al.* (1999) [10556935]
 706. King BF *et al.* (2005) [15778739]
 707. Kirichok Y *et al.* (2006) [16467839]
 708. Kirsch J. (2006) [16807723]
 709. Kiselyov K *et al.* (2009) [19273053]
 710. Kiselyov K *et al.* (2007) [17217079]
 711. Kiselyov K *et al.* (2007) [17138610]
 712. Kitchen P *et al.* (2015) [26365508]
 713. Kitchen P *et al.* (2020) [32413299]
 714. Kittaka H *et al.* (2017) [28612138]
 715. Kiyonaka S *et al.* (2009) [19289841]
 716. Klausen TK *et al.* (2009) [19361196]
 717. Kleene SJ *et al.* (2019) [30883612]
 718. Kleyman TR *et al.* (2020) [31721612]
 719. Kleyman TR *et al.* (2018) [29120692]
 720. Klugbauer N *et al.* (1995) [7720699]
 721. Knaus A *et al.* (2007) [17261653]
 722. Knepper MA *et al.* (2015) [25830425]
 723. Knowlton WM *et al.* (2011) [20932257]
 724. Kobayashi T *et al.* (2000) [10780978]
 725. Kobayashi T *et al.* (2004) [15150531]
 726. Koch HP *et al.* (2008) [18583477]
 727. Koivisto AP *et al.* (2022) [34526696]
 728. Kojima R *et al.* (2014) [24291101]
 729. Kolisek M *et al.* (2005) [15808509]
 730. Kollwe A *et al.* (2021) [34766907]
 731. Konstas AA *et al.* (2003) [12456399]
 732. Korpi ER *et al.* (2007) [17591542]
 733. Korpi ER *et al.* (2002) [12126658]
 734. Kotnis S *et al.* (2010) [20223879]
 735. Kozak JA *et al.* (2002) [12149283]
 736. Kraft R. (2007) [17658472]
 737. Kraft R *et al.* (2006) [16604090]
 738. Kraft R *et al.* (2004) [14512294]
 739. Krakow D *et al.* (2009) [19232556]
 740. Krapivinsky G *et al.* (2014) [24855944]
 741. Krapivinsky G *et al.* (2017) [28784805]
 742. Krapivinsky G *et al.* (1998) [9620703]
 743. Krauson AJ *et al.* (2018) [29782492]
 744. Krehan D *et al.* (2003) [14604650]
 745. Kreida S *et al.* (2015) [26342685]
 746. Kremeyer B *et al.* (2010) [20547126]
 747. Kreple CJ *et al.* (2014) [24952644]
 748. Krishek BJ *et al.* (1998) [9508826]
 749. Kristensson C *et al.* (2022) [35227647]
 750. Krosgaard-Larsen P *et al.* (2002) [12469353]
 751. Krügel U *et al.* (2017) [28106668]
 752. Kschonsak M *et al.* (2020) [32698188]
 753. Kschonsak M *et al.* (2022) [34929720]
 754. Kubo Y *et al.* (2005) [16382105]
 755. Kuduk SD *et al.* (2010) [22778804]
 756. Kulagowski JJ *et al.* (1994) [8182696]
 757. Kumar J *et al.* (2013) [22974439]
 758. Kuner T *et al.* (1996) [8642401]
 759. Kusaka S *et al.* (2001) [11179389]
 760. Kuzmenkov AI *et al.* (2022) [36188602]
 761. Kwok CHT *et al.* (2024) [39048565]
 762. Ladefoged LK *et al.* (2018) [30257860]
 763. Lafrenière RG *et al.* (2010) [20871611]
 764. Lalo U *et al.* (2008) [18495881]
 765. Lamb GD *et al.* (1987) [2451745]
 766. Lambert S *et al.* (2011) [21278253]
 767. Lambert S *et al.* (2005) [15961423]
 768. Landouré G *et al.* (2010) [20037586]
 769. Lang R *et al.* (2000) [10836990]
 770. Lange PF *et al.* (2006) [16525474]
 771. Lankiewicz S *et al.* (1998) [9463477]
 772. Larsson JE *et al.* (2020) [32595524]
 773. Laube B *et al.* (2002) [12413807]
 774. Lazarenko RM *et al.* (2010) [20519544]
 775. Leacock S *et al.* (2018) [29445326]
 776. Leblanc N *et al.* (2015) [26064450]

777. Leblanc N *et al.* (2005) [16091780]
 778. Lee CH *et al.* (2017) [28086084]
 779. Lee CH *et al.* (2019) [31787376]
 780. Lee CH *et al.* (2011) [21763685]
 781. Lee JH *et al.* (1999) [10094463]
 782. Lee JH *et al.* (1999) [10585925]
 783. Lee JYP *et al.* (2018) [29134638]
 784. Lee K *et al.* (1995) [7589189]
 785. Lee N *et al.* (2003) [12672827]
 786. Lee SP *et al.* (2008) [18334983]
 787. Lee SY *et al.* (2008) [18509058]
 788. Lee TS *et al.* (2006) [16899990]
 789. Leeson PD *et al.* (1992) [1534584]
 790. Leffler A *et al.* (2011) [21861907]
 791. Lei X *et al.* (2022) [36526620]
 792. Lei YT *et al.* (2014) [25237295]
 793. Leisle L *et al.* (2011) [21527911]
 794. Lemmens-Gruber R *et al.* (2023) [37175488]
 795. Lerche C *et al.* (2000) [10787416]
 796. Lerma J. (2006) [16361114]
 797. Lerma J. (2011) [21709676]
 798. Lesage F *et al.* (2000) [10880510]
 799. Leslie TK *et al.* (2023) [36183245]
 800. Leuner K *et al.* (2010) [20008516]
 801. Leuner K *et al.* (2007) [17666455]
 802. Levitz J *et al.* (2016) [27035963]
 803. Lewis RJ *et al.* (2000) [10938268]
 804. Li GD *et al.* (2009) [19282280]
 805. Li GD *et al.* (2006) [17093081]
 806. Li J *et al.* (2025) [39814994]
 807. Li M *et al.* (2006) [16636202]
 808. Li M *et al.* (2017) [28099415]
 809. Li N *et al.* (2025) [40808385]
 810. Li S *et al.* (2025) [39739807]
 811. Li SJ *et al.* (2010) [20147290]
 812. Li T *et al.* (2021) [33238160]
 813. Li WG *et al.* (2010) [21143836]
 814. Li X *et al.* (2021) [32884139]
 815. Li Z *et al.* (2024) [38443561]
 816. Liang P *et al.* (2024) [38814250]
 817. Liang P *et al.* (2024) [38033286]
 818. Liantonio A *et al.* (2007) [17128287]
 819. Liantonio A *et al.* (2008) [18216243]
 820. Liao M *et al.* (2013) [24305160]
 821. Liardo E *et al.* (2024) [39232360]
 822. Lichtenegger M *et al.* (2018) [29556099]
 823. Lien CC *et al.* (2002) [11790809]
 824. Lievreumont JP *et al.* (2005) [15933213]
 825. Liman ER. (2007) [17217064]
 826. Liman *et al.* (2007) Edited by W. B. Liedtke and S. Heller:
 827. Lin BL *et al.* (2019) [31028142]
 828. Lin SH *et al.* (2015) [25828470]
 829. Lin SH *et al.* (2015) [25582292]
 830. Lin Z *et al.* (2012) [22405088]
 831. Linford NJ *et al.* (1998) [9811906]
 832. Lingueglia E *et al.* (1997) [9368048]
 833. Lingueglia E *et al.* (1993) [8382172]
 834. Link TM *et al.* (2010) [20118928]
 835. Linsdell P. (2017) [27699452]
 836. Lishko PV *et al.* (2011) [21412339]
 837. Lishko PV *et al.* (2010) [20679352]
 838. Liu B *et al.* (2010) [20335661]
 839. Liu B *et al.* (2011) [22006988]
 840. Liu D *et al.* (2003) [14657398]
 841. Liu J *et al.* (2007) [17478420]
 842. Liu K *et al.* (2025) [40040241]
 843. Liu L *et al.* (2003) [12842134]
 844. Liu Q *et al.* (2022) [35525986]
 845. Liu S *et al.* (2016) [26756551]
 846. Liu SJ *et al.* (2007) [17275103]
 847. Liu X *et al.* (2018) [29443690]
 848. Liu X *et al.* (2020) [31936014]
 849. Liu Y *et al.* (2021) [33564124]
 850. Liu Y *et al.* (2011) [21290296]
 851. Lobley A *et al.* (2003) [12932298]
 852. Lochner M *et al.* (2010) [20409468]
 853. Lodge D. (2009) [18765242]
 854. Lopatin AN *et al.* (1994) [7969496]
 855. Lopes CM *et al.* (2000) [10748056]
 856. Lopez IA *et al.* (2019) [31092032]
 857. Lord B *et al.* (2015) [26386289]
 858. Lou J *et al.* (2022) [35221714]
 859. Loucif AJC *et al.* (2018) [29150838]
 860. Lozovaya N *et al.* (2005) [16107637]
 861. Lu B *et al.* (2007) [17448995]
 862. Lu B *et al.* (2009) [19092807]
 863. Lu B *et al.* (2010) [21040849]
 864. Lu M *et al.* (2008) [18326490]
 865. Lucas P *et al.* (2003) [14642279]
 866. Lucchesi K *et al.* (1989) [14775548]
 867. Ludlow MJ *et al.* (2025) *iScience*
 868. Lukas RJ *et al.* (1999) [10353988]
 869. Lummis SC *et al.* (2011) [21775477]
 870. Luscher B *et al.* (2011) [21555068]
 871. Lynagh T *et al.* (2018) [30061402]
 872. Lynagh T *et al.* (2017) [28949138]
 873. Lynch JW. (2009) [18721822]
 874. Lynch JW. (2004) [15383648]
 875. Lyon M *et al.* (2023) [36649421]
 876. Müller ME *et al.* (2024) [38891889]
 877. Ma D *et al.* (2023) [37857637]
 878. Ma D *et al.* (2022) [36763058]
 879. Ma S *et al.* (2008) [18930858]
 880. Ma T *et al.* (1997) [9388476]
 881. Ma Z *et al.* (2016) [26603282]
 882. Macpherson LJ *et al.* (2007) [17237762]
 883. Macpherson LJ *et al.* (2005) [15916949]
 884. Macpherson LJ *et al.* (2006) [16829128]
 885. Macpherson LJ *et al.* (2007) [17942735]
 886. Madjroh N *et al.* (2021) [34560054]
 887. Madry C *et al.* (2008) [18711142]
 888. Magri A *et al.* (2018) [29682501]
 889. Mahieu F *et al.* (2007) [17142461]
 890. Maier M *et al.* (2022) [35916168]
 891. Maier M *et al.* (2015) [25847402]
 892. Maingret F *et al.* (1999) [9880510]
 893. Maingret F *et al.* (2000) [10835347]
 894. Maingret F *et al.* (2001) [11226154]
 895. Maingret F *et al.* (1999) [10480871]
 896. Majeed Y *et al.* (2010) [20735426]
 897. Majeed Y *et al.* (2011) [21406603]
 898. Makhina EN *et al.* (1994) [8051145]
 899. Maksud G *et al.* (2009) [19383091]
 900. Malayev A *et al.* (2002) [11861317]
 901. Mälkiä A *et al.* (2007) [17317754]
 902. Mälkiä A *et al.* (2011) [20932258]
 903. Mall MA. (2020) [32732328]
 904. Mall MA *et al.* (2024) [39117676]
 905. Malosio ML *et al.* (1991) [1651228]
 906. Malysz J *et al.* (2009) [19389923]
 907. Mamet J *et al.* (2002) [12486159]
 908. Mansoor SE *et al.* (2016) [27626375]
 909. Manville RW *et al.* (2018) [30021858]
 910. Maricq AV *et al.* (1991) [1718042]
 911. Markham A. (2022) [35347635]
 912. Marra S *et al.* (2016) [26772186]
 913. Marshall DL *et al.* (1994) [7533951]
 914. Martin RL *et al.* (2000) [10991994]
 915. Martínez-López P *et al.* (2009) [19338774]
 916. Martins AP *et al.* (2012) [22624030]
 917. Mascia MP *et al.* (2000) [10908659]
 918. Masiulis S *et al.* (2019) [30602790]
 919. Matalon S *et al.* (2015) [26432872]
 920. Matchkov VV *et al.* (2004) [14718479]
 921. Matchkov VV *et al.* (2008) [18776041]
 922. Mathar I *et al.* (2014) [24226423]
 923. Matsuda JJ *et al.* (2008) [17977943]
 924. Mattmann ME *et al.* (2012) [22910039]
 925. Mayer ML. (2006) [16554805]
 926. Mazurov AA *et al.* (2011) [21919481]
 927. McCarthy AE *et al.* (2019) [31587896]
 928. McCormack K *et al.* (2013) [23818614]
 929. McDouough SI *et al.* (1996) [8786437]
 930. McIntyre P *et al.* (2001) [11226139]
 931. McKay S *et al.* (2012) [22022974]
 932. McNamara CR *et al.* (2007) [17686976]
 933. McNamara FN *et al.* (2005) [15685214]
 934. McRory JE *et al.* (2004) [14973233]
 935. Meena AS *et al.* (2022) [35705057]
 936. Meier JC *et al.* (2005) [15895087]
 937. Meissner G. (2017) [29122978]
 938. Meseguer V *et al.* (2014) [24445575]
 939. Mesirca P *et al.* (2024) [38167790]
 940. Miao Y *et al.* (2015) [26027738]
 941. Michel AD *et al.* (2008) [18071294]
 942. Michel AD *et al.* (1995) [8548175]
 943. Michel AD *et al.* (2008) [18660826]
 944. Michel AD *et al.* (2009) [19309360]
 945. Michel AD *et al.* (2006) [17031382]
 946. Michel AD *et al.* (2006) [16487507]
 947. Miehe S *et al.* (2012) [22530015]
 948. Mihályi C *et al.* (2024) [39495914]
 949. Miki K *et al.* (2013) [23453951]
 950. Millar NS. (2008) [18246096]
 951. Millar NS *et al.* (2009) [18723036]
 952. Miller M *et al.* (2011) [21795696]
 953. Miller PS *et al.* (2005) [15905212]
 954. Miller PS *et al.* (2005) [16144831]
 955. Miller PS *et al.* (2010) [20096941]
 956. Milstein AD *et al.* (2008) [18514334]
 957. Minard A *et al.* (2019) [31277085]
 958. Minard A *et al.* (2018) [29865154]
 959. Miner K *et al.* (2019) [30837866]
 960. Mintz IM *et al.* (1992) [1311418]
 961. Mittermeier L *et al.* (2019) [30770447]
 962. Miu P *et al.* (2001) [11406188]
 963. Miyake A *et al.* (1995) [7565620]
 964. Mochizuki S *et al.* (1999) [10204690]
 965. Mochizuki S *et al.* (2000) [10884508]
 966. Mody I *et al.* (2004) [15331240]
 967. Möhler H. (2007) [17394533]
 968. Molinari EJ *et al.* (2000) [10666507]
 969. Monteil A *et al.* (2024) [37615954]
 970. Moqrich A *et al.* (2005) [15746429]
 971. Moran MM *et al.* (2018) [28924972]
 972. Morgan AJ *et al.* (2014) [24277557]
 973. Morgan D *et al.* (2009) [19805063]
 974. Morley RM *et al.* (2005) [15801853]
 975. Moss SJ *et al.* (2001) [11283747]

976. Motoyama K *et al.* (2018) [29752182]
 977. Moura Barbosa AJ *et al.* (2010) [20724042]
 978. Moussaieff A *et al.* (2008) [18492727]
 979. Muanprasat C *et al.* (2004) [15277574]
 980. Mukhopadhyay I *et al.* (2014) [24819048]
 981. Mulligan SJ *et al.* (2006) [17047222]
 982. Munro G *et al.* (2016) [26663905]
 983. Murrell-Lagnado RD *et al.* (2022) [35776329]
 984. Musset B *et al.* (2010) [20037153]
 985. Musset B *et al.* (2010) [20231140]
 986. Musset B *et al.* (2011) [22020278]
 987. Na T *et al.* (2014) [24756712]
 988. Nadler MJ *et al.* (2001) [11385574]
 989. Nagata K *et al.* (2005) [15843607]
 990. Nagata K *et al.* (2008) [18162548]
 991. Nagle JF *et al.* (1978) [272644]
 992. Nakagawa F *et al.* (2020) [32051870]
 993. Nakagawa T. (2010) [21080238]
 994. Nakamura T *et al.* (1987) [3027574]
 995. Nakanishi O *et al.* (2020) [32102918]
 996. Nam YW *et al.* (2025) [40246884]
 997. Namkung W *et al.* (2010) [20581223]
 998. Naur P *et al.* (2007) [17715062]
 999. Nausch B *et al.* (2014) [24951278]
 1000. Navarro B *et al.* (2007) [17460039]
 1001. Naylor J *et al.* (2016) [26565375]
 1002. Nazirog'lu M *et al.* (2012) [21964764]
 1003. Neagoe I *et al.* (2010) [20466723]
 1004. Neeper MP *et al.* (2007) [17395593]
 1005. Newcomb R *et al.* (1998) [9799496]
 1006. Newman AS *et al.* (2013) [23594147]
 1007. Neyton J *et al.* (2006) [16452656]
 1008. Ng CK *et al.* (2011) [21428815]
 1009. Nicke A *et al.* (1998) [9606184]
 1010. Nie L *et al.* (1997) [9292225]
 1011. Niesler B. (2011) [21345729]
 1012. Niesler B *et al.* (2003) [12801637]
 1013. Niesler B *et al.* (2008) [18466097]
 1014. Niesler B *et al.* (2007) [17392525]
 1015. Niforatos W *et al.* (2007) [17314320]
 1016. Nilius B. (2007) [17368864]
 1017. Nilius B *et al.* (2003) [12558550]
 1018. Nilius B *et al.* (2014) [25296415]
 1019. Nilius B *et al.* (2006) [16424899]
 1020. Nilius B *et al.* (2010) [20127491]
 1021. Nilius B *et al.* (2008) [18923420]
 1022. Nilius B *et al.* (2007) [17237345]
 1023. Nilius B *et al.* (2004) [15331675]
 1024. Nilius B *et al.* (2005) [15590641]
 1025. Nilius B *et al.* (2004) [14758478]
 1026. Nilius B *et al.* (2004) [14707014]
 1027. Nina DuUllrich. (2005) In *TRPM4 and TRPM5: Functional characterisation and comparison of two novel Ca²⁺-activated cation channels of the TRPM subfamily* Faculteit Geneeskunde, Dept. Moleculaire Celbiologie, KU Leuven: 1028. No authors listed. (2004) [15293871]
 1029. Nörenberg W *et al.* (2011) [21262970]
 1030. Nörenberg W *et al.* (2012) [22506590]
 1031. Noreng S *et al.* (2018) [30251954]
 1032. Noreng S *et al.* (2020) [32729833]
 1033. North RA *et al.* (2013) [23253448]
 1034. Novarino G *et al.* (2010) [20430975]
 1035. Novello JC *et al.* (1999) [10082164]
 1036. Nury H *et al.* (2011) [21248852]
 1037. Oancea E *et al.* (2009) [19436059]
 1038. Oberwinkler J *et al.* (2005) [15824111]
 1039. Oberwinkler J *et al.* (2014) [24756716]
 1040. Oberwinkler J *et al.* (2007) [17217062]
 1041. Oertel J *et al.* (2007) [17145751]
 1042. Oh SJ *et al.* (2013) [23997117]
 1043. Ohya S *et al.* (2003) [12690036]
 1044. Okada SF *et al.* (2004) [15477379]
 1045. Okada T *et al.* (1999) [10488066]
 1046. Okada Y *et al.* (2009) [19171657]
 1047. Oken AC *et al.* (2022) [35784697]
 1048. Okuda K *et al.* (2021) [33321047]
 1049. Olesen SP *et al.* (1994) [8137869]
 1050. Olsen RW *et al.* (2011) [21194017]
 1051. Olsen RW *et al.* (2008) [18790874]
 1052. Olsen RW *et al.* (2009) [18760291]
 1053. Omura M *et al.* (2014) [25001287]
 1054. Omura M *et al.* (2015) [25701815]
 1055. Ong HL *et al.* (2017) [28900914]
 1056. Ong ST *et al.* (2015) [40294255]
 1057. Orhan G *et al.* (2011) [21354396]
 1058. Ormond SJ *et al.* (2006) [16452399]
 1059. Ortner NJ *et al.* (2017) [28592699]
 1060. Osmakov DI *et al.* (2020) [32733241]
 1061. Osmakov DI *et al.* (2013) [23001332]
 1062. Osteen JD *et al.* (2016) [27281198]
 1063. Osteen JD *et al.* (2008) [18658230]
 1064. Overington JP *et al.* (2006) [17139284]
 1065. Owsianik G *et al.* (2006) [16460288]
 1066. Ozhatil LC *et al.* (2018) [29579323]
 1067. Pagano E *et al.* (2023) [36168728]
 1068. Palacios-Prado N *et al.* (2022) [35486697]
 1069. Palmer LG *et al.* (2012) [22038262]
 1070. Pan X *et al.* (2022) [35194776]
 1071. Paoletti P. (2011) [21395862]
 1072. Paoletti P *et al.* (2013) [23686171]
 1073. Paoletti P *et al.* (2007) [17088105]
 1074. Pape HC. (1996) [8815797]
 1075. Papke RL *et al.* (2008) [18448138]
 1076. Papke RL *et al.* (2001) [11303054]
 1077. Park U *et al.* (2011) [21832173]
 1078. Parker RM *et al.* (1996) [8994113]
 1079. Parker RM *et al.* (1996) [8936343]
 1080. Parnas M *et al.* (2009) [19135721]
 1081. Parsonage G *et al.* (2023) [36457143]
 1082. Partiseti M *et al.* (1998) [9738472]
 1083. Patel A *et al.* (2010) [20490539]
 1084. Patel AJ *et al.* (1999) [10321245]
 1085. Patel AJ *et al.* (1998) [9687497]
 1086. Paul M *et al.* (2002) [11867382]
 1087. Paulsen CE *et al.* (2015) [25855297]
 1088. Payandeh J *et al.* (2011) [21743477]
 1089. Payne CE *et al.* (2015) [25625641]
 1090. Pearson WL *et al.* (1999) [9882736]
 1091. Pedarzani P *et al.* (2005) [16239218]
 1092. Pedarzani P *et al.* (2001) [11134030]
 1093. Pedemonte N *et al.* (2007) [17452495]
 1094. Pedersen SF *et al.* (2005) [16098585]
 1095. Pedregal C *et al.* (2000) [10821708]
 1096. Peier AM *et al.* (2002) [12016205]
 1097. Peigneur S *et al.* (2012) [22972919]
 1098. Pelegrin P *et al.* (2009) [19212823]
 1099. Pelegrin P *et al.* (2007) [17121814]
 1100. Peng C *et al.* (2004) [15134637]
 1101. Penzotti JL *et al.* (2001) [11159437]
 1102. Pérez-Vizcaíno F *et al.* (1993) [7506331]
 1103. Perrais D *et al.* (2009) [18761361]
 1104. Perrais D *et al.* (2010) [20850188]
 1105. Perraud AL *et al.* (2001) [11385575]
 1106. Perry M *et al.* (2004) [15266014]
 1107. Peters JA *et al.* (2005) [16194573]
 1108. Petheo GL *et al.* (2010) [21124855]
 1109. Petrus M *et al.* (2007) [18086313]
 1110. Peverini L *et al.* (2018) [30108481]
 1111. Philippaert K *et al.* (2017) [28361903]
 1112. Philippart F *et al.* (2018) [30556810]
 1113. Picollo A *et al.* (2005) [16034421]
 1114. Piechowicz KA *et al.* (2016) [26796863]
 1115. Pierandrei S *et al.* (2021) [33916525]
 1116. Pifferi S *et al.* (2009) [19475416]
 1117. Pignier C *et al.* (2000) [11045961]
 1118. Pingle SC *et al.* (2007) [17217056]
 1119. Pinheiro P *et al.* (2006) [16847640]
 1120. Pinto MC *et al.* (2021) [34326672]
 1121. Piper AS *et al.* (2002) [11850506]
 1122. Piper AS *et al.* (2004) [14724180]
 1123. Pirahanchi Y *et al.* (2021) [30725773]
 1124. Pitt SJ *et al.* (2008) [18987182]
 1125. Piyasirananda W *et al.* (2021) [33334897]
 1126. Planells-Cases R *et al.* (2009) [19708126]
 1127. Plant LD *et al.* (2010) [20498050]
 1128. Plant LD *et al.* (2012) [23169818]
 1129. Plant TD *et al.* (2003) [12765689]
 1130. Plasschaert LW *et al.* (2018) [30069046]
 1131. Plesch E *et al.* (2018) [30479274]
 1132. Pledsted AJ. (2011) [21713670]
 1133. Polster A *et al.* (2015) [25548159]
 1134. Pope L *et al.* (2018) [30089357]
 1135. Potier M *et al.* (2008) [18536932]
 1136. Preisig-Müller R *et al.* (2002) [12032359]
 1137. Preston GM *et al.* (1992) [1373524]
 1138. Price MP *et al.* (1996) [8626462]
 1139. Pritchett DB *et al.* (1989) [2551039]
 1140. Puertollano R *et al.* (2002) [19158345]
 1141. Pusch M *et al.* (2002) [12512775]
 1142. Pusch M *et al.* (2006) [16179405]
 1143. Putney JW. (2005) [16133266]
 1144. Qi H *et al.* (2007) [17227845]
 1145. Qi H *et al.* (2022) [35256942]
 1146. Qian F *et al.* (2005) [15971078]
 1147. Qiang M *et al.* (2018) [30042215]
 1148. Qin N *et al.* (2008) [18550765]
 1149. Qin X *et al.* (2013) [23145923]
 1150. Qu C *et al.* (2017) [28395140]
 1151. Qu Z *et al.* (2014) [24639373]
 1152. Quallo T *et al.* (2017) [28826490]
 1153. Quill TA *et al.* (2001) [11675491]
 1154. Qunies AM *et al.* (2023) [37812884]
 1155. Qunies AM *et al.* (2024) [38893312]
 1156. Rödström KEJ *et al.* (2024) [38755204]
 1157. Rajan S *et al.* (2001) [11060316]
 1158. Rajan S *et al.* (2000) [10747866]
 1159. Rambow J *et al.* (2014) [25225485]
 1160. Ramerstorfer J *et al.* (2010) [20303942]
 1161. Ramerstorfer J *et al.* (2011) [21248110]
 1162. Rampe D *et al.* (1997) [9395068]
 1163. Ramsey IS *et al.* (2018) [29483132]
 1164. Ramsey IS *et al.* (2010) [20543828]
 1165. Ramsey IS *et al.* (2006) [16554753]
 1166. Ramsey IS *et al.* (2009) [19372380]
 1167. Rapedius M *et al.* (2005) [15980413]
 1168. Rash LD. (2017) [28528673]
 1169. Rauh R *et al.* (2017) [27941075]
 1170. Rautenberg S *et al.* (2023) [35902436]

1171. Redd MA *et al.* (2021) [34264749]
 1172. Ren D *et al.* (2001) [11595941]
 1173. Rennhack A *et al.* (2018) [29723408]
 1174. Rettig J *et al.* (1992) [1378392]
 1175. Reyes R *et al.* (1998) [9812978]
 1176. Richter JM *et al.* (2014) [25140002]
 1177. Richter JM *et al.* (2014) [24117252]
 1178. Riquelme G. (2009) [19604577]
 1179. Roberts A *et al.* (2020) [32102919]
 1180. Robertson JL *et al.* (2010) [21048711]
 1181. Rochat H *et al.* (1998) [9792177]
 1182. Rock MJ *et al.* (2008) [18587396]
 1183. Rodríguez-Moreno A *et al.* (2007) [17981346]
 1184. Rohacs T. (2009) [19376575]
 1185. Romi-Lebrun R *et al.* (1997) [9354615]
 1186. Romine JL *et al.* (2007) [1726205]
 1187. Rook ML *et al.* (2021) [32306405]
 1188. Rosenbaum T *et al.* (2004) [14981138]
 1189. Rosenbaum T *et al.* (2008) [12508052]
 1190. Rossier BC *et al.* (2015) [25540145]
 1191. Rossier BC *et al.* (2009) [18928407]
 1192. Rotin D *et al.* (2011) [20972579]
 1193. Roy S *et al.* (2014) [24561672]
 1194. Ruah SSH *et al.* (2009) Patent number: US20090131492A1.
 1195. Rubaiy HN. (2019) [30656647]
 1196. Rubaiy HN. (2023) [37259313]
 1197. Rubaiy HN *et al.* (2017) [28325835]
 1198. Rubaiy HN *et al.* (2018) [29859013]
 1199. Rucktooa P *et al.* (2009) [19576182]
 1200. Rühl P *et al.* (2021) [33859333]
 1201. Rundström N *et al.* (1994) [8090751]
 1202. Runnels LW *et al.* (2002) [11941371]
 1203. Runnels LW *et al.* (2001) [11161216]
 1204. Rüschi D *et al.* (2007) [17360702]
 1205. Ryckmans T *et al.* (2011) [21741838]
 1206. Sabat M *et al.* (2022) [36402081]
 1207. Sabirov RZ *et al.* (2001) [11524456]
 1208. Sabirov RZ *et al.* (2009) [19340557]
 1209. Sabirova RKh *et al.* (1972) [4642630]
 1210. Safa P *et al.* (2001) [11514547]
 1211. Sagot E *et al.* (2008) [18578478]
 1212. Said EA *et al.* (2024) [39362502]
 1213. Sakai H *et al.* (1999) [10457052]
 1214. Sakai R *et al.* (2001) [11160654]
 1215. Sakurai Y *et al.* (2015) [25722412]
 1216. Salameh A *et al.* (2005) [16216217]
 1217. Salido GM *et al.* (2009) [19061922]
 1218. Salinas M *et al.* (2006) [16284080]
 1219. Samie M *et al.* (2013) [23993788]
 1220. Samsó M. (2017) [27671094]
 1221. Sanchez M *et al.* (1996) [8938726]
 1222. Sanechika S *et al.* (2021) [33877504]
 1223. Sano Y *et al.* (2003) [12754259]
 1224. Sardini A *et al.* (2003) [14729152]
 1225. Sasaki M *et al.* (2006) [16556803]
 1226. Saul A *et al.* (2013) [24391538]
 1227. Sawada Y *et al.* (2008) [18364033]
 1228. Saxena A *et al.* (2002) [12107247]
 1229. Saxena A *et al.* (1998) [9813171]
 1230. Saxena NC *et al.* (1997) [9263639]
 1231. Schaefer L *et al.* (2000) [10767424]
 1232. Scheel O *et al.* (2005) [16034422]
 1233. Schenkel LB *et al.* (2016) [26942860]
 1234. Schiller IC *et al.* (2022) [35335209]
 1235. Schleifer H *et al.* (2012) [22862290]
 1236. Schmalz F *et al.* (1998) [9612272]
 1237. Schmidt A *et al.* (2017) [29025967]
 1238. Schmitt S *et al.* (2023) [36674884]
 1239. Scholl U *et al.* (2006) [16849430]
 1240. Scholze P *et al.* (1996) [8813598]
 1241. Schott K *et al.* (2024) [38787877]
 1242. Schreiber R *et al.* (2024) [38642428]
 1243. Schrenzel J *et al.* (1998) [9565037]
 1244. Schroeder BC *et al.* (2000) [10816588]
 1245. Schröter KH *et al.* (1991) [1840526]
 1246. Schulz P *et al.* (2010) [20830208]
 1247. Schumacher MA *et al.* (2010) [20515731]
 1248. Seeburg PH *et al.* (2003) [12850211]
 1249. Seeliger D *et al.* (2013) [23113556]
 1250. Seidler U *et al.* (1997) [9423183]
 1251. Semyanov A *et al.* (2004) [15111008]
 1252. Seo K *et al.* (2014) [24453217]
 1253. Seo Y *et al.* (2016) [27219012]
 1254. Sergeeva OA *et al.* (2010) [20511229]
 1255. Shah M *et al.* (2000) [10683185]
 1256. Shakkottai VG *et al.* (2001) [11527975]
 1257. Shao J *et al.* (2019) [31664615]
 1258. Sharma S *et al.* (2018) [29453757]
 1259. Sharma S *et al.* (2019) [30943030]
 1260. Sharma S *et al.* (2020) [31950584]
 1261. Shei RJ *et al.* (2018) [30340955]
 1262. Sheldon RS *et al.* (1986) [2431264]
 1263. Shemon AN *et al.* (2004) [15210579]
 1264. Shen D *et al.* (2012) [22415822]
 1265. Sheng S *et al.* (2007) [17522058]
 1266. Sheppard DN *et al.* (1992) [1281220]
 1267. Sherkheli MA *et al.* (2010) [20816009]
 1268. Shi S *et al.* (2021) [34329684]
 1269. Shimizu T *et al.* (2009) [18663466]
 1270. Shimkera RA *et al.* (1994) [7954808]
 1271. Shoshan-Barmatz V *et al.* (2010) [20346371]
 1272. Shteinberg M *et al.* (2021) [34090606]
 1273. Sidach SS *et al.* (2002) [11896142]
 1274. Sieghart W. (2006) [17175817]
 1275. Sieghart W *et al.* (2022) [35017178]
 1276. Sieghart W *et al.* (2018) [30275042]
 1277. Sigel E *et al.* (2011) [21189125]
 1278. Sigel E *et al.* (2012) [23038269]
 1279. Simó-Vicens R *et al.* (2017) [28925012]
 1280. Simonin C *et al.* (2015) [26457814]
 1281. Sine SM *et al.* (2006) [16554804]
 1282. Singleton DH *et al.* (2007) [17536794]
 1283. Sinninger-Brauns MJ *et al.* (2009) [19029287]
 1284. Skolnick P *et al.* (1997) [9353361]
 1285. Skov M *et al.* (2024) [38507469]
 1286. Small B *et al.* (1998) [9849663]
 1287. Smart D *et al.* (2001) [11301059]
 1288. Smith AJ *et al.* (2010) [20501796]
 1289. Smith ES *et al.* (2007) [17258862]
 1290. Smith JF *et al.* (2013) [23530196]
 1291. Smith MA *et al.* (2003) [12562896]
 1292. Smith PL *et al.* (2006) [16899456]
 1293. Smith RD *et al.* (1998) [9437003]
 1294. Smith SM *et al.* (2011) [22006335]
 1295. Smith SS. (2013) [24027497]
 1296. Smothers CT *et al.* (2007) [17502428]
 1297. Snyder DS *et al.* (2013) [23814642]
 1298. Sobolevsky AI *et al.* (2009) [19946266]
 1299. Søgaard R *et al.* (2001) [11245603]
 1300. Söhl G *et al.* (2005) [15738956]
 1301. Solt K *et al.* (2005) [16081679]
 1302. Song C *et al.* (2022) [35210842]
 1303. Soom M *et al.* (2001) [11172809]
 1304. Soto F *et al.* (1999) [10193905]
 1305. Spehr J *et al.* (2009) [19228965]
 1306. Spix B *et al.* (2022) [35031603]
 1307. Stallmeyer B *et al.* (2012) [21887725]
 1308. Starowicz K *et al.* (2007) [17349697]
 1309. Staruschenko A *et al.* (2007) [17443775]
 1310. Stewart A *et al.* (2003) [12623220]
 1311. Stewart CG *et al.* (2024) [39104314]
 1312. Stiber JA *et al.* (2012) [23054893]
 1313. Stieber J *et al.* (2005) [16043489]
 1314. Stieber J *et al.* (2006) [16387796]
 1315. Stocker JW *et al.* (2003) [12433690]
 1316. Stocker M *et al.* (2004) [15208027]
 1317. Stokłosa P *et al.* (2021) [34771564]
 1318. Stokes L *et al.* (2006) [17031385]
 1319. Stoltz DA *et al.* (2010) [20427821]
 1320. Stórustovu SI *et al.* (2006) [16272218]
 1321. Story GM *et al.* (2003) [12654248]
 1322. Straub I *et al.* (2013) [24006495]
 1323. Straub I *et al.* (2013) [23190005]
 1324. Strøbæk D *et al.* (2013) [22891655]
 1325. Strøbæk D *et al.* (2006) [16926279]
 1326. Strøbæk D *et al.* (2000) [10696100]
 1327. Strøbæk D *et al.* (2004) [15471565]
 1328. Strømgaard K *et al.* (2002) [11934578]
 1329. Strübing C *et al.* (2001) [11301024]
 1330. Strünker T *et al.* (2011) [21412338]
 1331. Stühmer W *et al.* (1989) [2555158]
 1332. Su Q *et al.* (2021) [34381056]
 1333. Su Q *et al.* (2018) [30093605]
 1334. Sun D *et al.* (2020) [32915133]
 1335. Sun H *et al.* (1999) [10381768]
 1336. Sun J *et al.* (2017) [28575668]
 1337. Sun J *et al.* (2020) [31883792]
 1338. Sun X *et al.* (2020) [31983608]
 1339. Sun X *et al.* (2019) [30918114]
 1340. Sun XY *et al.* (2018) [30108138]
 1341. Sun ZC *et al.* (2020) [32273889]
 1342. Surprenant A *et al.* (2009) [18851707]
 1343. Sutko JL *et al.* (1996) [8874493]
 1344. Sutton KA *et al.* (2006) [16883570]
 1345. Suzuki H *et al.* (2016) [27433342]
 1346. Suzuki M *et al.* (2004) [15010458]
 1347. Swanson DM *et al.* (2005) [15771431]
 1348. Swayne LA *et al.* (2009) [19575010]
 1349. Syeda R *et al.* (2015) [26001275]
 1350. Syme CA *et al.* (2000) [10712246]
 1351. Szallasi A *et al.* (2007) [17464295]
 1352. Török F *et al.* (2025) [39370994]
 1353. Ta CM *et al.* (2016) [26562072]
 1354. Takahashi N *et al.* (1994) [8083233]
 1355. Takaya J *et al.* (2015) [26630251]
 1356. Takezawa R *et al.* (2006) [16407466]
 1357. Takumi T *et al.* (1995) [7608203]
 1358. Talavera K *et al.* (2009) [19749751]
 1359. Talley EM *et al.* (2002) [11886861]
 1360. Taly A *et al.* (2009) [19721446]
 1361. Tan Q *et al.* (2011) [33179259]
 1362. Tanaka M *et al.* (2025) [40365404]
 1363. Tanemoto M *et al.* (2002) [11988170]
 1364. Tanemoto M *et al.* (2000) [10856114]
 1365. Tang C *et al.* (2024) [39656529]
 1366. Tang Q *et al.* (2018) [29700422]

1367. Tang QY *et al.* (2010) [19934650]
 1368. Tang W *et al.* (1994) [8034048]
 1369. Tao J *et al.* (2014) [24759175]
 1370. Tarabova B *et al.* (2007) [17651721]
 1371. Tatulian L *et al.* (2001) [11466425]
 1372. Teramoto N *et al.* (2003) [14623761]
 1373. Territo PR *et al.* (2017) [27765863]
 1374. Thiel G *et al.* (2013) [23511953]
 1375. Thiel G *et al.* (2017) [28720517]
 1376. Thompson AJ *et al.* (2011) [21505038]
 1377. Thompson AJ *et al.* (2011) [21059362]
 1378. Thompson AJ *et al.* (2010) [20849671]
 1379. Thompson AJ *et al.* (2008) [18311193]
 1380. Thompson AJ *et al.* (2006) [17073663]
 1381. Thompson AJ *et al.* (2007) [17373882]
 1382. Thompson AJ *et al.* (2011) [21708905]
 1383. Thompson CH *et al.* (2009) [19574231]
 1384. Thompson SA *et al.* (2002) [12367611]
 1385. Thompson SA *et al.* (1999) [10455284]
 1386. Thompson SA *et al.* (2004) [15100159]
 1387. Thorneloe KS *et al.* (2008) [18499743]
 1388. Tiapko O *et al.* (2019) [30997005]
 1389. Timmermann DB *et al.* (2007) [17625074]
 1390. Togashi K *et al.* (2006) [16601673]
 1391. Togashi K *et al.* (2008) [18204483]
 1392. Tomasinsig L *et al.* (2008) [18765670]
 1393. Tombola F *et al.* (2008) [18498736]
 1394. Tombola F *et al.* (2010) [20023640]
 1395. Tomita S. (2010) [20134027]
 1396. Toncheva D *et al.* (2014) [24949484]
 1397. Tóth B *et al.* (2012) [22847436]
 1398. Tóth B *et al.* (2015) [25918360]
 1399. Towart R *et al.* (2009) [19439185]
 1400. Tradtrantip L *et al.* (2017) [28258578]
 1401. Tradtrantip L *et al.* (2010) [19808995]
 1402. Trattinig SM *et al.* (2016) [26827532]
 1403. Traynelis SF *et al.* (1998) [9698310]
 1404. Traynelis SF *et al.* (2010) [20716669]
 1405. Trebak M *et al.* (2007) [17217081]
 1406. Tringham E *et al.* (2012) [22344687]
 1407. Truong EC *et al.* (2017) [28493701]
 1408. Tsumura T *et al.* (1998) [9769420]
 1409. Tu YH *et al.* (2018) [29371428]
 1410. Tzvetkov MV *et al.* (2007) [17010535]
 1411. Ugawa S *et al.* (2007) [17826743]
 1412. Ugawa S *et al.* (2001) [11588592]
 1413. Ullrich ND *et al.* (2005) [15670874]
 1414. Urban N *et al.* (2016) [26500253]
 1415. Usher S *et al.* (2025) [40085699]
 1416. Vaandrager AB *et al.* (1997) [9024297]
 1417. Vaelli P *et al.* (2024) [39322410]
 1418. Van den Eynde C *et al.* (2022) [35714693]
 1419. van Megen WH *et al.* (2022) [35022312]
 1420. Vandewauw I *et al.* (2018) [29539642]
 1421. Vandewiele F *et al.* (2022) [35822895]
 1422. Vega-Saenz de Miera E *et al.* (1992) [1381835]
 1423. Veldhuis NA *et al.* (2015) [25361914]
 1424. Veleiro AS *et al.* (2009) [19199916]
 1425. Velicelebi G *et al.* (2016) Patent number: WO2016138472A1.
 1426. Vennekens R *et al.* (2007) [17217063]
 1427. Vennekens R *et al.* (2008) [18220815]
 1428. Vergani P *et al.* (2005) [15729345]
 1429. Verkman AS *et al.* (2014) [24625825]
 1430. Verkman AS *et al.* (2009) [19153558]
 1431. Verkman AS *et al.* (2017) Patent number: US9790218B2.
 1432. Versteegen MMA *et al.* (2020) [33318612]
 1433. Vick JS *et al.* (2015) [25592215]
 1434. Vigil FA *et al.* (2020) [32636759]
 1435. Vincent F *et al.* (2009) [19737537]
 1436. Virginio C *et al.* (2022) [35058176]
 1437. Virginio C *et al.* (1998) [9614197]
 1438. Vithlani M *et al.* (2011) [21742794]
 1439. Voets T *et al.* (2004) [15306801]
 1440. Voets T *et al.* (2007) [17395625]
 1441. Voets T *et al.* (2007) [17217067]
 1442. Vogt A *et al.* (2005) [16143426]
 1443. Vogt-Eisele AK *et al.* (2007) [17420775]
 1444. Voilley N *et al.* (2001) [11588175]
 1445. Vriens J *et al.* (2009) [19297520]
 1446. Vriens J *et al.* (2014) [24390427]
 1447. Vriens J *et al.* (2011) [21555074]
 1448. Vriens J *et al.* (2018) [29305649]
 1449. Vukicevic M *et al.* (2006) [16282326]
 1450. Vulchanova L *et al.* (1997) [9364478]
 1451. Wafford KA *et al.* (2009) [18762200]
 1452. Wagner TF *et al.* (2008) [18978782]
 1453. Wahl P *et al.* (2001) [11125018]
 1454. Waldmann R *et al.* (1997) [9261094]
 1455. Waldmann R *et al.* (1997) [9062189]
 1456. Waldmann R *et al.* (1995) [7499195]
 1457. Waldmann R *et al.* (1996) [8631835]
 1458. Walker JR *et al.* (2012) [23077250]
 1459. Walker RL *et al.* (2002) [12388058]
 1460. Wallner M *et al.* (2006) [16814864]
 1461. Walsh KB *et al.* (1986) [2418195]
 1462. Walstab J *et al.* (2010) [20522555]
 1463. Walstab J *et al.* (2010) [20621123]
 1464. Wang B *et al.* (2019) [31636563]
 1465. Wang G *et al.* (1990) [2154667]
 1466. Wang G *et al.* (1992) [1331975]
 1467. Wang H *et al.* (2020) [32004513]
 1468. Wang H *et al.* (2009) [19516020]
 1469. Wang H *et al.* (2009) [19535918]
 1470. Wang HL *et al.* (2007) [17585751]
 1471. Wang HS *et al.* (2000) [10825393]
 1472. Wang HS *et al.* (1998) [9836639]
 1473. Wang J *et al.* (2023) [37836658]
 1474. Wang J *et al.* (2005) [16042391]
 1475. Wang JJ *et al.* (2020) [31980622]
 1476. Wang L *et al.* (2018) [30467180]
 1477. Wang L *et al.* (2019) [30655942]
 1478. Wang M *et al.* (2020) [32513195]
 1479. Wang RX *et al.* (2011) [21187320]
 1480. Wang S *et al.* (2016) [27797339]
 1481. Wang SY *et al.* (1998) [9482942]
 1482. Wang W *et al.* (2006) [16319075]
 1483. Wang X *et al.* (2020) [32094222]
 1484. Wang X *et al.* (2012) [23063126]
 1485. Wang XQ *et al.* (2006) [17046694]
 1486. Wang Y *et al.* (2017) [26648074]
 1487. Wang Y *et al.* (2018) [29610524]
 1488. Wang Y *et al.* (2023) [37541373]
 1489. Wang Y *et al.* (2002) [12237342]
 1490. Wang Y *et al.* (2010) [19842162]
 1491. Wang YZ *et al.* (2015) [26523449]
 1492. Wang Z *et al.* (2019) [31441214]
 1493. Ware AW *et al.* (2021) [33630195]
 1494. Warth R *et al.* (2004) [15141089]
 1495. Washburn DG *et al.* (2013) [23886683]
 1496. Watanabe H *et al.* (2003) [12879072]
 1497. Watanabe S *et al.* (2016) [26837049]
 1498. Watanabe TX *et al.* (1995) [7474554]
 1499. Waxman SG. (2023) [37530829]
 1500. Weatherall KL *et al.* (2010) [20359520]
 1501. Webb TI *et al.* (2007) [17692006]
 1502. Wehage E *et al.* (2002) [11960981]
 1503. Wei AD *et al.* (2005) [16382103]
 1504. Wei ZL *et al.* (2015) [26235950]
 1505. Weitz D *et al.* (2002) [12467591]
 1506. Wemmie JA *et al.* (2013) [23783197]
 1507. Werner S *et al.* (2019) [31746599]
 1508. Wescott SA *et al.* (2013) [22820171]
 1509. Wetzel C *et al.* (2017) [27941788]
 1510. Wickenden AD *et al.* (2000) [10953053]
 1511. Wiemuth D *et al.* (2014) [24365967]
 1512. Willegens K *et al.* (2022) [35768468]
 1513. Williams DK *et al.* (2011) [21575610]
 1514. Williams JA *et al.* (2008) [18518786]
 1515. Williams ME *et al.* (1994) [8071363]
 1516. Winter M *et al.* (2020) [33352987]
 1517. Wishka DG *et al.* (2006) [16821801]
 1518. Wissenbach U *et al.* (2007) [17217060]
 1519. Wittekindt OH *et al.* (2004) [14978258]
 1520. Witzgall R. (2007) [17217069]
 1521. Won J *et al.* (2025) [39478185]
 1522. Wong CO *et al.* (2010) [20233211]
 1523. Wood ML *et al.* (2012) [21843503]
 1524. Woudenberg-Vrenken TE *et al.* (2011) [20814221]
 1525. Wray S *et al.* (2021) [34867456]
 1526. Wrighton DC *et al.* (2015) [26045093]
 1527. Wu J *et al.* (2006) [16825297]
 1528. Wu J *et al.* (2011) [21787755]
 1529. Wu LJ *et al.* (2010) [20716668]
 1530. Wu SN *et al.* (2002) [11809993]
 1531. Wu SN *et al.* (1999) [10428416]
 1532. Wulff H *et al.* (2000) [10884437]
 1533. Wyllie DJ *et al.* (2013) [23376022]
 1534. Xia J *et al.* (2009) [19211808]
 1535. Xiao B *et al.* (2008) [18815250]
 1536. Xie J *et al.* (2011) [22180838]
 1537. Xiong W *et al.* (2011) [21460829]
 1538. Xiong ZG *et al.* (2007) [17127388]
 1539. Xu F *et al.* (2003) [12970074]
 1540. Xu H *et al.* (2005) [16192383]
 1541. Xu H *et al.* (2006) [16617338]
 1542. Xu H *et al.* (2007) [17989217]
 1543. Xu H *et al.* (2002) [12077604]
 1544. Xu H *et al.* (2015) [25668017]
 1545. Xu L *et al.* (2020) [33278032]
 1546. Xu SZ *et al.* (2005) [15806115]
 1547. Xu W *et al.* (2001) [11487617]
 1548. Xu X *et al.* (2013) *Biophys J* **104**: 454a
 1549. Xu Y *et al.* (2020) [33584647]
 1550. Yamada A *et al.* (2024) [39379156]
 1551. Yamada A *et al.* (2024) [38359825]
 1552. Yamada M *et al.* (1997) [9130167]
 1553. Yamaguchi N *et al.* (2003) [12707260]
 1554. Yamashita T *et al.* (1996) [8735700]
 1555. Yan J *et al.* (2021) [34087722]
 1556. Yang B. (2017) *In Advances in Experimental Medicine and Biology* Edited by Yang B: Springer: 1-276 [ISBN: 9789402410570]
 1557. Yang B *et al.* (2006) [16876206]
 1558. Yang B *et al.* (1997) [9195910]
 1559. Yang J *et al.* (1995) [7748552]
 1560. Yang KC *et al.* (2009) [19498417]

1561. Yang L *et al.* (2014) [25114023]
1562. Yang S *et al.* (2015) [26420335]
1563. Yang Z *et al.* (2008) [18755158]
1564. Yang Z *et al.* (2007) [17714449]
1565. Yang Z *et al.* (2025) [40172963]
1566. Yao J *et al.* (2005) [15695820]
1567. Yashiro K *et al.* (2025) [39981749]
1568. Ye L *et al.* (2021) [33352109]
1569. Yeh HI *et al.* (2017) [29079713]
1570. Yevenes GE *et al.* (2010) [20647311]
1571. Yevenes GE *et al.* (2006) [17040914]
1572. Yevenes GE *et al.* (2003) [12858180]
1573. Yevenes GE *et al.* (2011) [21557733]
1574. Yildirim E *et al.* (2003) [12601176]
1575. Yin Y *et al.* (2022) [36227998]
1576. Yoder N *et al.* (2020) [32496192]
1577. Yoder N *et al.* (2018) [29513651]
1578. You H *et al.* (2010) [20638393]
1579. Young GT *et al.* (2008) [18791069]
1580. Yu CR. (2015) [26157356]
1581. Yu D *et al.* (2025) [40485107]
1582. Yu FH *et al.* (2004) [15467096]
1583. Yu FH *et al.* (2005) [16382097]
1584. Yu J *et al.* (2021) [33567265]
1585. Yu L *et al.* (2010) [20142439]
1586. Yu L *et al.* (2020) [32128386]
1587. Yu M *et al.* (2019) [31749913]
1588. Yu P *et al.* (2019) [31216484]
1589. Yu X *et al.* (2021) [34788616]
1590. Yu Y *et al.* (2010) [20920791]
1591. Yuan JP *et al.* (2009) [19574740]
1592. Yue L *et al.* (2021) [34254641]
1593. Yuzaki M. (2003) [12725908]
1594. Zamponi GW *et al.* (2009) [19815411]
1595. Zdebik AA *et al.* (2008) [18063579]
1596. Zeevi DA *et al.* (2007) [17306511]
1597. Zeng D *et al.* (1996) [8967455]
1598. Zeng XH *et al.* (2011) [21427226]
1599. Zennaro MC *et al.* (2012) [21664233]
1600. Zezula J *et al.* (1996) [8773466]
1601. Zha XM *et al.* (2009) [19218436]
1602. Zhang D *et al.* (2001) [11239575]
1603. Zhang E *et al.* (2015) [25502473]
1604. Zhang G *et al.* (2022) [36351900]
1605. Zhang H *et al.* (2023) [36596769]
1606. Zhang H *et al.* (2018) [29723786]
1607. Zhang H *et al.* (2019) [30377214]
1608. Zhang H *et al.* (2021) [33784097]
1609. Zhang M *et al.* (2025) [40470184]
1610. Zhang S *et al.* (2024) [38167918]
1611. Zhang XF *et al.* (2010) [20566409]
1612. Zhang Y *et al.* (2022) [35563730]
1613. Zhao M *et al.* (1999) [10473538]
1614. Zhao Z *et al.* (2021) [34265294]
1615. Zheng J *et al.* (2002) [12467592]
1616. Zheng J *et al.* (2004) [15134638]
1617. Zheng QH. (2020) [32273217]
1618. Zheng X *et al.* (2022) [34969976]
1619. Zheng Y *et al.* (2022) [34767770]
1620. Zholos A. (2010) [20233227]
1621. Zhong G *et al.* (2021) [33786819]
1622. Zhong H *et al.* (2002) [12432397]
1623. Zhou H *et al.* (1996) [8997197]
1624. Zhou L *et al.* (2022) [35387979]
1625. Zhou LM *et al.* (1997) [8996224]
1626. Zhou T *et al.* (2020) [33155599]
1627. Zhou Y *et al.* (2017) [29217578]
1628. Zhou Z *et al.* (1998) [9449325]
1629. Zhou Z *et al.* (1999) [10376921]
1630. Zhu H *et al.* (2021) [34555840]
1631. Zhu MX. (2011) [22593967]
1632. Zhu Y *et al.* (2015) [25816897]
1633. Ziemann AE *et al.* (2009) [19945383]
1634. Zierler S *et al.* (2011) [21926172]
1635. Zifarelli G *et al.* (2009) [19131966]
1636. Zitt C *et al.* (1996) [8663995]
1637. Zong P *et al.* (2022) [35421327]
1638. Zong P *et al.* (2022) [35445217]
1639. Zufall F *et al.* (2005) [15971083]
Upioculto hosula nonosta mdinatum aus An