



Kent Academic Repository

Norton, Emily and Dudas, Olivier (2025) *Decomposition numbers for unipotent blocks with small sl_2 -weight in finite classical groups*. Journal of Combinatorial Algebra . ISSN 2415-6302. (In press)

Downloaded from

<https://kar.kent.ac.uk/113146/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.4171/jca>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

DECOMPOSITION NUMBERS FOR UNIPOTENT BLOCKS WITH SMALL $\mathfrak{sl}_2$ -WEIGHT IN FINITE CLASSICAL GROUPS

OLIVIER DUDAS AND EMILY NORTON

ABSTRACT. We show that parabolic Kazhdan-Lusztig polynomials of type A compute the decomposition numbers in certain Harish-Chandra series of unipotent characters of finite groups of Lie types B , C and D over a field of non-defining odd characteristic ℓ . Here, ℓ is a “unitary prime” – the case that remains open in general, in the sense that it cannot be reduced to a similar problem for q -Schur algebras. The bipartitions labeling the characters in these series are small with respect to d , the order of $q \bmod \ell$, although they occur in blocks of arbitrarily high defect. Our main technical tool is the categorical action of an affine Lie algebra on the category of unipotent representations, which identifies the branching graph for Harish-Chandra induction with the $\widehat{\mathfrak{sl}}_d$ -crystal on a sum of level 2 Fock spaces. Further key combinatorics has been adapted from Brundan and Stroppel’s work on Khovanov arc algebras to obtain the closed formula for the decomposition numbers in a d -small Harish-Chandra series.

INTRODUCTION

For a finite group G , decomposition numbers encode how ordinary irreducible representations behave after reduction modulo a prime number ℓ . In this paper, we study the case where G is one of the finite classical groups $\mathrm{SO}_{2n+1}(q)$, $\mathrm{Sp}_{2n}(q)$, $\mathrm{O}_{2n}^{\pm}(q)$ and ℓ is an odd prime that is coprime to an odd prime power q (non-defining characteristic).

Computing the decomposition matrix can be done blockwise, and using Lusztig’s Jordan decomposition one can often restrict to the unipotent blocks, which are the ones containing unipotent characters (see [1, Thm. 11.8]). However, even for those blocks, computing the decomposition numbers explicitly seems out of reach. Still, for finite general linear groups, they can be related to Kazhdan–Lusztig polynomials, at least when ℓ is large enough with respect to n , the rank of the underlying algebraic group. Our main result gives a similar interpretation for classical groups, see Corollary 4.2, but in a much more restricted case.

Let d be the multiplicative order of q in $\mathbb{F}_{\ell}^{\times}$. When d is odd (the *linear prime* case), the representation theory of unipotent blocks of classical groups is governed by that of finite general linear groups. When d is even (the *unitary prime* case), much less is known. For example, the decomposition numbers for unipotent blocks of $\mathrm{Sp}_{2n}(q)$ in the unitary case have been determined up to $n = 4$ only, and quite recently: in 1998 for $n = 2$ [23], in 2014 for $n = 3$ [19] and in 2022 for $n = 4$ [10]. The standard strategy for computing these numbers is the following:

- (1) Compute the projective cover of cuspidal representations;
- (2) Decompose the module obtained by Harish-Chandra induction of that projective cover.

We will work under two assumptions which will allow us to solve these two problems: first, we will restrict our ordinary irreducible characters to a given Harish-Chandra series. Second, we will impose a smallness condition with respect to d on the Lusztig symbols in the series. The exact condition is given in Section 3.1. In this setting, cuspidal representations, when they occur, are always the “smallest” within the series with respect to the partial order on symbols. Thus

problem (1) is no problem thanks to the unitriangular shape of the decomposition matrix [3]. Everything then hinges on problem (2) – and it turns out to be tractable for these series. Note that we end up working with blocks of any defect, even though in our situation the decomposition numbers turn out to be either 0 or 1.

Let us be more precise on the behaviour of (2). Using the $\widehat{\mathfrak{sl}}_d$ -action constructed in [13], one can break the Harish-Chandra induction functor into pieces, and obtain an i -induction functor F_i for each $i \in \mathbb{Z}/d\mathbb{Z}$. If P_S is the projective cover of a simple module S of highest weight (for the action of $\mathfrak{sl}_2$ corresponding to some $i \in \mathbb{Z}/d\mathbb{Z}$), then $F_i^n(P_S)$ is very close to being the projective cover of $F_i^n(S)$. The difference involves only characters which are in the image of F_i^{n+1} , see Proposition 1.3 for an exact statement. Under our assumption, all these extra characters lie in another Harish-Chandra series, hence do not contribute to the decomposition numbers we are interested in.

The explicit computation of the decomposition numbers now boils down to knowing exactly how to compute the i -induction on unipotent characters and simple modules in characteristic ℓ . Under our assumptions, one can attach to each irreducible character/module an up-down diagram and a cup diagram as in [5, 4] and obtain an elementary description of the action of F_i on these combinatorial data as in [6]. See Section 3, which takes its cue from the analogous situation for Hecke algebras at $d = \infty$ [6, 4]. This combinatorial set-up positions us to prove our main theorem, Theorem 4.1, in an inductive way using Proposition 1.3. The latter exploits the small highest weight of the simple $\mathfrak{sl}_2$ -modules in which such characters are found with respect to the categorical action on the category of representations. We observe that we get the exact same formula as in [4, 5], thus relating the decomposition numbers within the Harish-Chandra series to the parabolic Kazhdan–Lusztig polynomials for a maximal parabolic in type A .

1. CATEGORICAL $\mathfrak{sl}_2$ -ACTION

In this section we prove a formula relating the character of the projective cover of a simple module S and the character of the projective cover of its highest weight support. Only part of the character can be controlled, but this will be sufficient to show in Section 4 how to compute decomposition numbers from the case of cuspidal simple modules.

1.1. Recollection on categorical actions. We recall here some of the features of $\mathfrak{sl}_2$ -actions on categories as defined in [7, 24].

Let $\mathbb{A}$ be a ring with unit and $\mathcal{V}$ be a $\mathbb{A}$ -linear abelian category. A *categorical datum* on $\mathcal{V}$ is given by a pair of biadjoint exact endofunctors E and F of $\mathcal{V}$, together with two natural transformations $X \in \text{End}(E)$ and $T \in \text{End}(E^2)$ satisfying the relations given in [24, 3.3.3] in the case of $\mathfrak{sl}_2$. Equivalently, we require that X and T induce an action of the affine nil-Hecke algebra of $\mathfrak{S}_n$ on the functor E^n for all $n \geq 0$. Using that structure one can define the divided power functors $E^{(n)}$ and $F^{(n)}$ which are still exact and biadjoint. They satisfy

$$E^n \simeq (E^{(n)})^{\oplus n!} \quad \text{and} \quad F^n \simeq (F^{(n)})^{\oplus n!}.$$

Assume now that $\mathbb{A}$ is a field and that $\mathcal{V}$ has finite length. An $\mathfrak{sl}_2$ -categorical action on $\mathcal{V}$ is given by a categorical datum (E, F, X, T) and a decomposition

$$\mathcal{V} = \bigoplus_{\omega \in \mathbb{Z}} \mathcal{V}_\omega$$

of $\mathcal{V}$ into abelian categories (which we will call *weight categories*). Furthermore, the functors E and F should shift the weights by 2 and -2 respectively

$$\mathcal{V}_\omega \begin{array}{c} \xrightarrow{E} \\ \xleftarrow{F} \end{array} \mathcal{V}_{\omega+2}$$

such that in the Grothendieck group $K_0(\mathcal{V}_\omega)$, the commutator $[E][F] - [F][E]$ acts by multiplication by ω . In particular the classes $e = [E]$ and $f = [F]$ of the functors in the complexified Grothendieck group $V = \mathbb{C} \otimes_{\mathbb{Z}} K_0(\mathcal{V})$ induce an action of $\mathfrak{sl}_2$ for which the weight space of weight ω is exactly $V_\omega = \mathbb{C} \otimes_{\mathbb{Z}} K_0(\mathcal{V}_\omega)$. Note that we will always assume that this action is integrable, so that e and f are locally nilpotent.

For such a notion of $\mathfrak{sl}_2$ -action on a category, the divided power functors satisfy the following identity on weight spaces of weight $\omega \geq 0$, see [24, Lem. 4.8]

$$(1.1) \quad E^{(n)}F^{(n)}|_{\mathcal{V}_\omega} \simeq 1_{\mathcal{V}_\omega}^{\oplus \binom{\omega}{n}} \oplus (FE|_{\mathcal{V}_\omega})^{\oplus \binom{\omega-1}{n-1}} \oplus (F^{(2)}E^{(2)}|_{\mathcal{V}_\omega})^{\oplus \binom{\omega-2}{n-2}} \oplus \dots \oplus F^{(n)}E^{(n)}|_{\mathcal{V}_\omega}.$$

We now assume that $\text{End}(S) \simeq \mathbb{A}$ for all simple objects $S \in \text{Irr } \mathcal{V}$. In that case it is proven in [7, Prop. 5.20] that $E(S)$, if non-zero, has simple socle and head and that they are isomorphic. Successive applications of E give a highest weight semi-simple object. We will use the further following properties, which are also proved in [7, Prop. 5.20].

Lemma 1.2. *Let $S \in \text{Irr } \mathcal{V}_\omega$ and $n \geq 0$ be such that $E^{n+1}(S) = 0$ and $E^n(S) \neq 0$. Then:*

- (1) $E^{(n)}(S)$ is simple.
- (2) The socle and head of $F^{(n)}E^{(n)}(S)$ are isomorphic to S .
- (3) The simple module S occurs in $F^{(n)}E^{(n)}(S)$ with multiplicity $\binom{\omega+2n}{n}$ as a composition factor.

1.2. Decomposition numbers. Let $\mathbb{O}$ be a complete discrete valuation ring, with residue field $\mathbb{k}$ of positive characteristic and fraction field $\mathbb{K}$ of characteristic zero. Let $\{G_r\}_{r \in \mathbb{N}}$ be a family of finite groups. We consider the category

$$\bigoplus_{r \geq 0} \mathbb{O}G_r\text{-mod}$$

which is the sum of the categories of finitely generated representations of G_r over $\mathbb{O}$. We will work with a direct summand $\mathcal{V}$ of this category, which in our setting will consist of *unipotent representations*, see §2.6. If $\mathbb{k}$ is algebraically closed, such a category will satisfy the following conditions:

- For $\mathbb{A} = \mathbb{k}, \mathbb{K}$, the category $\mathbb{A}\mathcal{V}$ has finite length and $\text{End}(S) = \mathbb{A}$ for all $S \in \text{Irr}_{\mathbb{A}}\mathcal{V}$.
- Every $S \in \text{Irr}_{\mathbb{k}}\mathcal{V}$ has a projective cover P_S in $\mathbb{k}\mathcal{V}$, unique up to isomorphism.
- Every projective module P in $\mathbb{k}\mathcal{V}$ lifts uniquely to a projective module $\tilde{P}$ in $\mathbb{O}\mathcal{V}$.
- $\mathbb{K}\mathcal{V}$ is semisimple.

If $S \in \text{Irr}_{\mathbb{k}}\mathcal{V}$ and $\Delta \in \text{Irr}_{\mathbb{K}}\mathcal{V}$, the decomposition number of S in Δ is the multiplicity of Δ as a direct summand (equivalently, a composition factor) of $\mathbb{K} \otimes_{\mathbb{O}} \tilde{P}_S$. We denote it by

$$[P_S : \Delta]$$

in a way which will look familiar to the reader interested in highest weight categories.

We assume now that (E, F, X, T) is a categorical datum on $\mathcal{V}$ inducing an $\mathfrak{sl}_2$ -categorical action on $\mathbb{k}\mathcal{V}$. This is to ensure that the divided powers $E^{(n)}$ and $F^{(n)}$ defined in the previous section can be lifted to exact and biadjoint endofunctors of $\mathcal{V}$, even though $\mathcal{V}$ itself does not

have an action of $\mathfrak{sl}_2$ since its Grothendieck group might be too big. The image of these functors by extension of scalars will be still denoted by $E^{(n)}$ and $F^{(n)}$.

Proposition 1.3. *In the previous setting, let $S \in \text{Irr}_{\mathbb{K}}\mathcal{V}$ and $n \geq 0$ be such that $E^{n+1}(S) = 0$ and $E^n(S) \neq 0$. Then*

$$[P_S : \Delta] = [P_{E^{(n)}S} : E^{(n)}\Delta]$$

for all irreducible modules $\Delta \in \text{Irr}_{\mathbb{K}}\mathcal{V}$ such that $E^{n+1}\Delta = 0$.

Proof. Let ω be the weight of the simple module $T = E^{(n)}S$. Note that $\omega \geq 0$ since the class of T is a highest weight vector. Since S is the head of $F^{(n)}E^{(n)}S$ by Lemma 1.2(2) we have that $F^{(n)}P_T$ contains P_S as a direct summand. We write

$$(1.4) \quad F^{(n)}P_{E^{(n)}S} = P_S \oplus Q$$

for some projective module Q . We want to have some control on the character of Q , so for that we compute the image of (1.4) by $E^{(n)}$. Using (1.1) we have

$$E^{(n)}P_S \oplus E^{(n)}Q = E^{(n)}F^{(n)}P_T = P_T^{\oplus \binom{\omega}{n}} \oplus FEP_T^{\oplus \binom{\omega}{n-1}} \oplus \dots \oplus F^{(n)}E^{(n)}P_T.$$

Now we claim that $E^{(n)}P_S$ contains $\binom{\omega}{n}$ copies of P_T . Indeed, the weight of S equals $\omega - 2n$ and we have

$$\text{Hom}(E^{(n)}P_S, T) = \text{Hom}(E^{(n)}P_S, E^{(n)}S) \simeq \text{Hom}(P_S, F^{(n)}E^{(n)}S)$$

which has dimension $\binom{\omega}{n}$ by Lemma 1.2.(3). This proves that $E^{(n)}Q$ is a direct summand of a sum of modules of the form $F^{(k)}E^{(k)}(P_T)$ where $k \geq 1$. By the lifting property of projective modules, the same holds over $\mathbb{O}$.

Now, let Δ be an irreducible module of $\mathbb{K}\mathcal{V}$ such that $[Q : \Delta] \neq 0$. In other words, Δ is isomorphic to a submodule of $\mathbb{K} \otimes_{\mathbb{O}} \tilde{Q}$, and since $E^{(n)}$ is exact, $E^{(n)}\Delta$ is isomorphic to a submodule of $\mathbb{K} \otimes_{\mathbb{O}} E^{(n)}\tilde{Q}$. Furthermore, it must be non-zero since by definition $\tilde{Q}$ is a direct summand of $F^{(n)}\tilde{P}_T$, putting Δ in the image of $F^{(n)}$. Now by the previous paragraph, there is $k \geq 1$ such that $[F^{(k)}E^{(k)}(P_T) : E^{(n)}\Delta] \neq 0$, which forces $E^{(k)}E^{(n)}\Delta$, and therefore $E^{n+k}\Delta$ to be non-zero. We showed that $E^{n+1}\Delta = 0$ implies $[Q : \Delta] = 0$, which gives the result by (1.4). $\square$

Remark 1.5. The same result (with analogous proof) would hold for highest weight categories and standard objects Δ .

2. UNIPOTENT REPRESENTATIONS OF FINITE CLASSICAL GROUPS

By a *finite classical group* $G_n(q)$ we will always mean one of the following groups:

Group	$\text{SO}_{2n+1}(q)$	$\text{Sp}_{2n}(q)$	$\text{O}_{2n}^+(q)$	$\text{O}_{2n}^-(q)$
Type	B_n	C_n	D_n	${}^2D_n$

where $q > 1$ is a power of some odd prime. The convention for small values of n is that $\text{SO}_1(q) = \text{Sp}_0(q) = \text{O}_0^{\pm}(q) = 1$ is the trivial group, $\text{O}_2^+(q) = \text{GL}_1(q) \rtimes \mathbb{Z}/2$ and $\text{O}_2^-(q) = \text{GU}_1(q) \rtimes \mathbb{Z}/2$ are 1-dimensional tori (of respective orders $q-1$ and $q+1$) extended by $\mathbb{Z}/2$.

We will work with modular representations, therefore with different coefficient rings for our representations. As in Section 1.2 we fix a complete discrete valuation ring $\mathbb{O}$ with residue field $\mathbb{k}$ and fraction field $\mathbb{K}$. We assume that $\mathbb{K}$ has characteristic zero, $\mathbb{k}$ has characteristic $\ell > 0$ and that $\mathbb{k}$ is large enough for all the finite groups encountered. In particular, every irreducible unipotent representation over $\mathbb{K}$ or $\mathbb{k}$ will be absolutely irreducible. We will always work under

the assumption that ℓ is odd and does not divide q . The multiplicative order d of q in $\mathbb{k}^\times$ will be assumed to be even, so that we work in the *unitary prime* case.

Note that some of the results cited in this section for the groups $O_{2n}^\pm(q)$ have not been published yet. Still, we have decided to include these groups since the combinatorics is not much different from the groups of type B/C , and since there are work in progress by Li-Shan-Zhang [21] and Cia-Luvecce [8] which will contain all the results we need here.

2.1. Combinatorics. We recall here the combinatorics that is used to classify the unipotent characters and unipotent blocks of finite classical groups in the unitary prime case.

2.1.1. Partitions and symbols. Let m be a non-negative integer. A partition λ of m is a non-increasing sequence of non-negative integers $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq 0)$ which add up to m , the size $|\lambda|$ of λ . A bipartition $\boldsymbol{\lambda} = \lambda^1 \cdot \lambda^2$ of m is a pair of partitions (λ_1, λ_2) such that $|\lambda^1| + |\lambda^2| = m$. A charged bipartition $|\boldsymbol{\lambda}, \mathbf{s}|$ is the data of a bipartition and a pair of integers $\mathbf{s} = (s_1, s_2) \in \mathbb{Z}^2$, called the charge.

To a charged bipartition $|\boldsymbol{\lambda}, \mathbf{s}|$ one can attach a charged symbol $\Theta(\boldsymbol{\lambda}, \mathbf{s})$ corresponding to the pair of charged β -sets coming from the two partitions. More precisely, we have $\Theta(\boldsymbol{\lambda}, \mathbf{s}) = (X_1, X_2)$ where

$$X_k = \{s_k + \lambda_j - j + 1 \mid j \geq 1\}.$$

A symbol will be represented by the corresponding 2-abacus, with the first row X_1 on the bottom. The *defect* of a charged symbol $\Theta = \Theta(\boldsymbol{\lambda}, \mathbf{s})$ is $\text{def}(\Theta) = s_2 - s_1$.

Example 2.1. The charged symbol of charge $(-4, 3)$ and bipartition $1^3 \cdot 2^2 1$ will be represented as follows

row 2	● ● ● ● ● ● ● ● . ● . ● ● .
row 1	● . ● ● ●
	-7 -6 -5 -4 -3 -2 -1 0 1 2 3 4 5 6

2.1.2. Adding and removing boxes. The unipotent characters will be parametrized by such symbols. In order to explain how the induction and restriction of unipotent representations behave on symbols, we will need the notion of addable/removable boxes. Recall that d is a fixed even integer. Let $i \in \mathbb{Z}/d$ and $\Theta = (X_1, X_2)$ be a charged symbol. An addable i -box of Θ in the row $k \in \{1, 2\}$ is an integer x such that

- $x \equiv i + (k-1)\frac{d}{2} \pmod{d}$;
- $x \in X_k$ and $x+1 \notin X_k$.

Adding the i -box x in the symbol Θ consists in replacing x by $x+1$ in X_k . A removable i -box of Θ in the row $k \in \{1, 2\}$ is an integer x such that

- $x \equiv i + (k-1)\frac{d}{2} \pmod{d}$;
- $x \notin X_k$ and $x+1 \in X_k$.

Removing the i -box x in the symbol Θ consists in replacing $x+1$ by x in X_k .


There is a notion of *good* addable/removable i -box, see for example [20, Section 3], [15, Theorem 2.8]. Given i , there is at most one *good* addable/removable i -box in a charged symbol, and it can be used to describe the Kashiwara operators $\tilde{f}_i$ and $\tilde{e}_i$ on charged symbols (or charged bipartitions in other contexts).

Remark 2.2. The reader might be surprised by the occurrence of $d/2$ in the definition of addable/removable boxes. This comes from the fact that the charges of our symbols will come from parameters $(q^{s_1}, -q^{s_2})$ in the Hecke algebra. Using the fact that $-1 = q^{d/2}$, one should rather work with symbols of charge $(s_1, s_2 + d/2)$, but that will remove the symmetry from

Adding and removing boxes does not change the charge. Removing all the possible boxes yields a symbol containing in each row only consecutive integers. Such a symbol corresponds to the empty bipartition, and the charge equals $(\max(X_1), \max(X_2))$ where (X_1, X_2) is the β -set of the empty bipartition.

Figure 1 shows a 2D lattice with two rows and 13 columns. The columns are indexed from -7 to 6. Row 2 (top) contains black circles at columns -7 to 4, a grey circle at column 5, and a white circle at column 6. Row 1 (bottom) contains a grey circle at column -7, a white circle at column -6, and black circles at columns -5 to 0. Columns 1 to 6 are empty. Arrows point from the grey circle at (2,5) to the white circle at (1,6) and from the white circle at (1,-6) to the grey circle at (2,-7).

Example 2.4. Let us again consider the symbol drawn in Example 2.1. There are four 4-hooks, all being in the top row.



- $\mathrm{SO}_{2n+1}(q)$ and $\mathrm{Sp}_{2n}(q)$ have a cuspidal unipotent character if and only if $n = t^2 + t$ for some $t \in \mathbb{N}$. In that case it is unique, and we denote it by Δ_t .
- $\mathrm{O}_{2n}^+(q)$ has a cuspidal unipotent character if and only if $n = t^2$ for some $t \in 2\mathbb{N}$. If $t = 0$, there is a unique one, denoted by Δ_0 . If $t \neq 0$, there are exactly two, which we denote by Δ_t and Δ_{-t} .
- $\mathrm{O}_{2n}^-(q)$ has a cuspidal unipotent character if and only if $n = t^2$ for some $t \in 2\mathbb{N} + 1$. In that case there are exactly two, which we denote again by Δ_t and Δ_{-t} .

Let $n, m, t \geq 0$ be such that $G_n(q)$ has a cuspidal unipotent character Δ_t . Then the unipotent characters of $G_{n+m}(q)$ in the Harish-Chandra series above Δ_t are parametrized by bipartitions of m .

2.3. Classification of unipotent characters. In order to have a global treatment of all the Harish-Chandra series, we define, for each $t \in \mathbb{Z}$, the following charge

$$\sigma_t = \begin{cases} (t, -1-t) & \text{if } t \text{ is even and } G_n \text{ is of type } B/C, \\ (-1-t, t) & \text{if } t \text{ is odd and } G_n \text{ is of type } B/C, \\ (t, -t) & \text{if } t \text{ is even and } G_n \text{ is of type } D, \\ (-t, t) & \text{if } t \text{ is odd and } G_n \text{ is of type } {}^2D. \end{cases}$$

Then unipotent characters are classified by charged symbols of charge σ_t , see Section 2.1 for the definition or properties of symbols. In a series above Δ_t , the symbols will have defect $2t+1$, $-2t-1$, $2t$, $-2t$ depending on the type of groups and the parity of t .

Given a charged symbol Θ with charge σ_t ($t \in \mathbb{Z}$), we will denote by Δ_Θ the corresponding unipotent character. Note that there is no unique such parametrization, but the one we will choose comes from a categorical action and it will satisfy the properties given in the following sections.

2.4. Unipotent blocks. Recall that we work in the *unitary prime* case, where d , the multiplicative order of q in $\mathbb{k}^\times$, is even. A *unipotent ℓ -block* is an ℓ -block containing at least one unipotent character. The partition of unipotent characters into ℓ -blocks can be read off from the labelling of unipotent characters by symbols. By [16, 25], one can choose the parametrization such that two unipotent characters are in the same ℓ -block if and only if the corresponding symbols have the same $d/2$ -cocore.

2.5. Classification of unipotent simple modules over $\mathbb{k}$. Under the assumptions on q and ℓ , the decomposition matrix of the unipotent blocks of classical groups have a unitriangular shape. This is proven for finite classical groups coming from connected reductive algebraic groups in [3], and the case of $O_{2n}^\pm(q)$ follows for example from [14, Thm. 3.1]. In particular, the parametrization of unipotent characters by charged symbols yields a parametrization of the irreducible unipotent representations over $\mathbb{k}$ as well. Given a charged symbol Θ with charge σ_t ($t \in \mathbb{Z}$), we will denote by S_Θ the corresponding simple representation.

2.6. Categorical action. Let $G_n(q)$ be a finite classical group. We denote by $\mathcal{O}G_n(q)\text{-umod}$ the category of finitely generated unipotent representations of $G_n(q)$ over $\mathcal{O}$. It is the direct summand of $\mathcal{O}G_n(q)\text{-mod}$ corresponding to the sum of all unipotent blocks. The type of finite classical group being fixed, we denote by $\mathcal{V}$ the category

$$\mathcal{V} := \bigoplus_{n \geq 0} \mathcal{O}G_n(q)\text{-umod}.$$

In the case where d is even, there is a categorical datum on $\mathcal{V}$ for every $i \in \mathbb{Z}/d$ inducing an $\mathfrak{sl}_2$ -categorical action on $\mathbb{k}\mathcal{V}$. We will denote by E_i and F_i the corresponding functors (over any ring of coefficients amongst $\mathbb{K}$, $\mathcal{O}$ and $\mathbb{k}$). These functors are defined using Harish-Chandra induction and restriction functors, see [12, 13] for more details for their construction.

These various $\mathfrak{sl}_2$ -categorical actions come from an $\widehat{\mathfrak{sl}}_d$ -categorical action. We shall not use that fact since we will be working with each action separately. However, it is important to know that one can choose the parametrisation of unipotent characters in such a way that one can actually compute the action of each E_i , F_i on unipotent characters and unipotent Brauer characters. More precisely, if we define

$$(2.5) \quad \mathbf{s}_t = \sigma_t + (0, d/2)$$

then there exists an isomorphism of $\widehat{\mathfrak{sl}}_d$ -modules

$$(2.6) \quad \begin{aligned} \mathbb{C} \otimes_{\mathbb{Z}} K_0(\mathbb{k}\mathcal{V}) &\xrightarrow{\sim} \bigoplus_t F(\mathbf{s}_t) \\ [\Delta_{\Theta(\lambda, \sigma_t)}] &\longmapsto |\lambda, \mathbf{s}_t\rangle \end{aligned}$$

where $F(\mathbf{s}_t)$ is the level 2 Fock space of charge $\mathbf{s}_t$, and $[\Delta]$ denotes the class of any ℓ -reduction of the character Δ . Note that depending on the type of the classical group G_n , the integer t will run over the non-negative integers only (for type B/C), over all the even integers (for type D) or over all the odd integers (for type ${}^2D$).

Note that the action of $\widehat{\mathfrak{sl}}_d$ on the Fock space $F(\mathbf{s}_t)$ depends only on the charge up to shifts by $(0, d)$ and $(d, 0)$.

However, $\mathbf{s}_t \in \mathbb{Z}^2$ is chosen so that the map

$$S_{\Theta(\lambda, \sigma_t)} \longmapsto |\lambda, \mathbf{s}_t\rangle$$

induces an isomorphism between the crystals (which are sensitive to the charge, not only the residue class of the components of the charge). This is proven for type B/C in [11, Thm. 1.7], and the same proof should work for type D and ${}^2D$ once we have a good parametrization.

2.7. Explicit formulas. The categorification results contained in the previous section were proven to get an explicit description of the Harish-Chandra induction and restriction functors on the unipotent representations in both characteristic zero and ℓ . (For explicit formulas for the action on higher level Fock spaces, see for example the thesis of Yvonne, [28]). Given a charged symbol Θ with charge σ_t , we have

$$(2.7) \quad E_i(\Delta_{\Theta}) = \bigoplus_{\substack{j \equiv i \pmod{d} \\ \Theta \setminus \Psi = j}} \Delta_{\Psi},$$

where $\Theta \setminus \Psi = j$ means that Ψ is obtained from Θ by removing a j -box, see Section 2.1.2. Thus E_i removes all the possible removable i -boxes from the charged symbol Θ .

$$(2.8) \quad \text{soc } E_i(S_{\Theta}) = \text{hd } E_i(S_{\Theta}) = \begin{cases} 0 & \Theta \text{ has no good removable } i\text{-box,} \\ S_{\Psi} & \text{if } \Psi \text{ is obtained from } \Theta \text{ by removing the good } i\text{-box.} \end{cases}$$

Similar formulas are obtained for the action of F_i by adjunction.

3. THE DIAGRAMMATICS OF d -SMALL SYMBOLS

In this section, we adapt the combinatorics in [4, 5, 6] to an appropriate class of symbols labeling unipotent characters of types B , C , D and ${}^2D$. This will position us to prove the formula for decomposition numbers in Theorem 4.1 using Proposition 1.3.

3.1. d -small symbols. Given a symbol Θ , we work with its graphical representation as two rows of beads and spaces, and so we will often refer colloquially to the elements $\beta \in \Theta$ as *beads* and to the elements of $(\mathbb{Z}, \mathbb{Z}) \setminus \Theta$ as *spaces*.

Definition 3.1. Let $\Theta = (X_1, X_2)$ be a symbol. For each $i = 1, 2$, define the interval $J_i = [c_i, d_i] \subset \mathbb{Z}$ by

$$d_i = \max\{\beta \in \mathbb{Z} \mid \beta \in X_i\}$$

and

$$c_i = \max\{z \in \mathbb{Z} \mid \beta \in X_i \text{ for all } \beta < z\}.$$

Fix $d \in 2\mathbb{N}$. We say Θ is *d-small* if there exist intervals $I_1 := [a_1, b_1]$, $I_2 := [a_2, b_2] \subset \mathbb{Z}$ satisfying the following conditions:

Remark 3.5. Recall that if $\Theta = \Theta(\lambda, \sigma)$ is the symbol attached to the charged bipartition $|\lambda, \sigma\rangle$, then $|\lambda, (\sigma_1, \sigma_2 + \frac{d}{2})\rangle =: |\lambda, \mathbf{s}\rangle$ is the charged bipartition for the Fock space as in (2.6). Then Θ is d -small if and only if the set of residues of the charged contents of boxes in $|\lambda, \mathbf{s}\rangle$ is contained in a closed interval of length $\frac{d}{2} - 1$ modulo d .

3.2. Diagrams.

Definition 3.6. Fix $d \in 2\mathbb{N}$. Let Θ be a d -small symbol. Write the right region as $I_j = m + [1, \frac{d}{2}] \subset \mathbb{Z}$ for some $m \in \mathbb{Z}$. Let kd be the length of the middle region, $k \in \mathbb{N}$. To Θ we associate an *up-down diagram* $w_{\wedge\vee}(\Theta) = w_1 w_2 \cdots w_{\frac{d}{2}}$ consisting of a word of length $\frac{d}{2}$ in the alphabet $\{\wedge, \vee, \circ, \times\}$. Each $\beta_i = m + i \in I_j$ determines the i 'th letter w_i of $w_{\wedge\vee}(\Theta)$ according to the rule:


- if $\beta_i \in \Theta$ and $\beta_i - kd - \frac{d}{2}$ occurs in both rows of Θ , then $w_i = \times$,
- if $\beta_i \in \Theta$ and $\beta_i - kd - \frac{d}{2}$ occurs only once in Θ , then $w_i = \wedge$,
- if $\beta_i \notin \Theta$ and $\beta_i - kd - \frac{d}{2}$ occurs in both rows of Θ , then $w_i = \vee$,
- if $\beta_i \notin \Theta$ and $\beta_i - kd - \frac{d}{2}$ occurs only once in Θ , then $w_i = \circ$.

Thus, the up-down diagram of Θ records what happens in the left and right regions of Θ , in the opposite rows where a mix of beads and spaces is possible. Putting the right region on top of the left region, the rules correspond to the following possibilities:


$$(3.7) \quad \begin{array}{ll} \text{Right region} & \bullet \bullet \cdot \cdot \\ \text{Left region} & \bullet \cdot \bullet \cdot \\ & \times \wedge \vee \circ \end{array}$$

Definition 3.8. Given a d -small symbol Θ , we further associate to it a *cup diagram* $c_{\wedge\vee}(\Theta)$. It is uniquely determined by $w_{\wedge\vee}(\Theta)$ and consists of non-crossing counterclockwise arcs and rays attached to the $\vee$'s and $\wedge$'s of $w_{\wedge\vee}(\Theta)$ by the following recursive procedure. Start with a pair $\vee \cdots \wedge$ which has no $\wedge$ or $\vee$ symbols between them. Connect the $\vee$ to the $\wedge$ with a curved arc (a “cup”). Considering the $\vee$ and $\wedge$ symbols as directional, the cup is counterclockwise-oriented as its left endpoint is $\vee$ and its right endpoint is $\wedge$. Continue to connect with cups any $\vee \cdots \wedge$ pairs that have only $\times$ and $\circ$ symbols and previously constructed cups between them. When no more counterclockwise cups can be constructed by this rule, attach vertical rays below the remaining $\wedge$ and $\vee$ symbols.

Example 3.9. We continue with Example 3.2. We have $w_{\wedge\vee}(\Theta) = \wedge \vee \times \vee \wedge \wedge$ and the associated cup diagram $c_{\wedge\vee}(\Theta)$ is:



Example 3.10. Let $d = 28$ and consider the following d -small symbol:



Putting the right region on top of the left region we have



hence according to (3.7) we get the following up-down diagram:

$$w_{\wedge\vee}(\Theta) = \wedge \wedge \vee \vee \circ \wedge \vee \wedge \wedge \times \vee \vee \wedge \circ$$

and

$$c_{\wedge\vee}(\Theta) = \begin{array}{c} \uparrow \quad \uparrow \quad \vee \quad \vee \quad \circ \quad \uparrow \quad \vee \quad \uparrow \quad \uparrow \quad \times \quad \vee \quad \vee \quad \uparrow \quad \circ \\ \hline \uparrow \quad \uparrow \quad \vee \quad \vee \quad \circ \quad \uparrow \quad \vee \quad \uparrow \quad \uparrow \quad \times \quad \vee \quad \vee \quad \uparrow \quad \circ \end{array}$$

We remark that the rays will not play any role in computing decomposition numbers; what is important for our purposes are the cups in $c_{\wedge\vee}(\Theta)$. Likewise the $\times$ and $\circ$ symbols play a placeholder role. In the situation of Example 3.4 where there is more than one choice for the left and right regions, different choices are reflected in $w_{\wedge\vee}(\Theta)$ by replacing a prefix of $\times$'s with a suffix of the same number of $\circ$'s or vice versa.

3.3. The cocore of a d -small symbol.

Lemma 3.11. *Let Θ be a d -small symbol and let Θ° be its $\frac{d}{2}$ -cocore. Then Θ° is a d -small symbol, its middle region has length 0, and $w_{\wedge\vee}(\Theta^\circ)$ is obtained from $w_{\wedge\vee}(\Theta)$ by replacing all $\wedge$ by $\vee$.*

Proof. We will induct on k where kd is the length of the middle region of the d -small symbol Θ . Suppose $k = 0$, so the middle region has length 0. Given an occurrence of $\wedge$ in $w_{\wedge\vee}(\Theta)$, there is a bead in the corresponding position of the right region and a space $\frac{d}{2}$ to its left and in the opposite row. Thus each $\wedge$ yields a removable $\frac{d}{2}$ -cohook, and conversely, any removable $\frac{d}{2}$ -cohook is given in this way by the $\wedge$'s in $w_{\wedge\vee}(\Theta)$. Removing such a cohook moves the bead in question from the right region to the left region. Do this for each $\wedge$ in $w_{\wedge\vee}(\Theta)$, then flip the resulting symbol upside-down if the number of $\wedge$'s in $w_{\wedge\vee}(\Theta)$ is odd. No more cohooks can be removed from the resulting symbol, so we have found the $\frac{d}{2}$ -cocore Θ° of Θ . It is clear that Θ° has the desired description.

Next, suppose by induction that if Ψ is a d -small symbol whose middle region has length kd then Ψ° has the desired description. Let Θ be a d -small symbol whose middle region has length $(k+1)d$. Partition the middle region into equal segments of length $\frac{d}{2}$. For the leftmost such segment of length $\frac{d}{2}$ (adjacent to the left region), move the i 'th bead $\frac{d}{2}$ to the left and to the opposite row if there's a space in the i 'th position there, thus, if $\wedge$ or $\circ$ is the i 'th letter of $w_{\wedge\vee}(\Theta)$. For all other beads in the middle and right regions, they all belong to the same row. The opposite row has only empty spaces in the middle and right regions. We move these beads $\frac{d}{2}$ to the left and to the opposite row. Finally, flip the symbol upside down if the total number of beads moved was odd. Call this symbol Ψ . We have removed some number of cohooks from Θ to obtain Ψ , so Ψ has the same cocore as Θ . We observe that Ψ has the same pattern of beads and spaces in its left and right regions as Θ . Thus $w_{\wedge\vee}(\Psi) = w_{\wedge\vee}(\Theta)$. However, the middle region of Ψ has length kd . Applying induction, Θ° has the desired description. $\square$

A d -small symbol Θ is determined by the length of its middle region together with $w_{\wedge\vee}(\Theta)$, up to flipping the symbol upside-down. In type B , that is when the defect of the symbol is odd, by convention the flip of the symbol upside-down labels the same unipotent character. Thus when Θ is the symbol of a type B unipotent character, the preceding lemma completely characterizes the cocore of Θ . In the case of type D or ${}^2D$ unipotent characters, that is when the defect of the symbol is even, the lemma characterizes the cocore up to flipping it upside-down (these label two different unipotent characters in that case).

Example 3.12. Take $d = 12$ and the following d -small symbol Θ :

$$\begin{array}{ccccccc} \text{row 2} & \bullet & | & \bullet & \bullet & \bullet & \bullet & | & \cdot & \bullet & \bullet & \bullet & \bullet & | & \cdot \\ \text{row 1} & \bullet & | & \bullet & \bullet & \cdot & \cdot & \cdot & | & \cdot & \cdot & \cdot & \cdot & \cdot & | & \cdot \\ & -6 & & & & & & -1 & & & & & & & 4 \end{array}$$

3.4. Action of the crystal operators on d -small symbols. In this section, we explain how the action of the crystal operators $\tilde{e}_i$ on symbols translates to $\mathbf{w}_{\wedge\vee}(\Theta)$ for a d -small symbol Θ .

To describe the action of crystal operators using $\mathbf{w}_{\wedge\vee}(\Theta)$, we consider $i \in \mathbb{Z}/d\mathbb{Z}$ where i labels the residue mod d of a position in the right region if the right region lies in row 1 of Θ . Otherwise, the left region lies in row 1 of Θ , and then we let i label the residue mod d of a position in the left region. We note that if $i \in \mathbb{Z}/d\mathbb{Z}$ does not belong to the set of residues just described, then Θ has no removable i -box so necessarily $\tilde{e}_i$ acts by 0. Match up the letters $w_1, \dots, w_{\frac{d}{2}}$ in $\mathbf{w}_{\wedge\vee}(\Theta)$ from left to right with the residues of the appropriate region. In Example 3.12, we would have:

$$\begin{array}{cccccc} 1 & 2 & 3 & 4 & 5 & 6 \\ \circ & \times & \wedge & \circ & \vee & \wedge \end{array}$$

We then consider the two adjacent letters $w_j, w_{j+1} \in \{\wedge, \vee, \circ, \times\}$ in $\mathbf{w}_{\wedge\vee}(\Theta)$ corresponding to the residues $i, i+1$:

$$(3.15) \quad \begin{array}{cc} i & i+1 \\ w_j & w_{j+1} \end{array}$$

Continuing our example and taking $i = 5$, we would consider:

$$\begin{array}{cc} 5 & 6 \\ \vee & \wedge \end{array}$$

Note that when the residue i corresponds to $w_{\frac{d}{2}}$, then $\tilde{e}_i$ acts by 0 since there is no removable i -box of Θ .

Then the effect of $\tilde{e}_i$ on $\mathbf{w}_{\wedge\vee}(\lambda)$ is given by acting locally on these two letters as described in the next lemma.

Lemma 3.16. *Let Θ be a d -small symbol and let $\mathbf{w}_{\wedge\vee}(\Theta)$ be its up-down diagram. Label the letters of $\mathbf{w}_{\wedge\vee}(\Theta)$ with the residues of the appropriate region as described above. The action of the crystal operators $\tilde{e}_i$ and $\tilde{f}_i$ on Θ is computed locally in $\mathbf{w}_{\wedge\vee}(\Theta)$ on $w_j w_{j+1}$ as follows:*

$$(3.17) \quad \begin{array}{ccc} \star \times & \xrightleftharpoons[\tilde{f}_i]{\tilde{e}_i} & \times \star \\ \circ \star & \xrightleftharpoons[\tilde{f}_i]{\tilde{e}_i} & \star \circ \\ \circ \times & \xrightleftharpoons[\tilde{f}_i]{\tilde{e}_i} & \vee \wedge \xrightleftharpoons[\tilde{f}_i]{\tilde{e}_i} \times \circ \end{array}$$

for $\star \in \{\wedge, \vee\}$. The action of e_i on $\wedge\vee, \vee\vee, \wedge\wedge, \star\circ$, and $\times\star$ is zero.

Remark 3.18. The action of $\tilde{f}_i$ is obtained by adjunction whenever $\tilde{f}_i\Theta$ remains d -small.

Proof. It is clear that $\tilde{e}_i$ acts by 0 on $\vee\vee, \wedge\wedge, \star\circ$, and $\times\star$ since there is no removable i -box of Θ in these cases.

Suppose the Kashiwara i -word of Θ is non-empty for some i . Then it consists of at most two letters from the alphabet $\{+, -\}$, with at most one contributed by an addable or removable box in the left region and at most one contributed by an addable or removable box in the right region of Θ . If the Kashiwara i -word consists solely of $-$ then there is exactly one removable i -box and no addable i -box, we are in the situation $\circ\star$ and $\tilde{e}_i$ removes the i -box yielding $\star\circ$. For the remaining cases, we need to check that if the i -word has two letters, then the letter contributed by the right region is larger than the letter contributed by the left region.

Suppose that the right region occurs in row 2. Recall that $|\lambda, \mathbf{s}_t\rangle = |\lambda, \sigma_t + (0, \frac{d}{2})\rangle$. Then any addable or removable i -box in λ^2 has charged content at least d greater than any addable

or removable i -box of λ^1 . Thus when comparing addable and removable i -boxes of λ , the one in λ^2 is larger than the one in λ^1 . Next, suppose that the right region occurs in row 1. If the middle region has length at least d , then as in the previous case, an addable or removable i -box of λ^1 will have charged content at least d greater than an addable or removable i -box of λ^2 , so will be larger. Otherwise, the right region is $\frac{d}{2}$ to the right of the left region. Then in $|\lambda, \mathbf{s}_t\rangle$, the addable/removable i -boxes of λ^1 and λ^2 have the same charged content. In this situation, the box in λ^1 is considered larger. So again, the letter contributed to the i -word from the right region is larger. $\square$

The lemma implies that the class in the Grothendieck group of a simple module corresponding to a d -small symbol lies in a representation of $\mathfrak{sl}_2$ of dimension at most 3. The action on the standard modules is the same as for simple modules unless we are in the 3-dimensional representation of $\mathfrak{sl}_2$, where we have

$$(3.19) \quad e_i[\Delta_{\circ \times}] = [\Delta_{\vee \wedge}] + [\Delta_{\wedge \vee}] = f_i[\Delta_{\times \circ}]$$

in the Grothendieck group. Here the subscripts correspond to the changes in the $\wedge \vee$ -diagram of the corresponding symbol.

Example 3.20. Continuing with Example 3.12, the lemma tells us that

$$\tilde{e}_5(\circ \times \wedge \circ \vee \wedge) = \circ \times \wedge \circ \times \circ$$

and thus $\tilde{e}_5$ acts on Θ by moving the rightmost bead in the symbol one position to the left.

Remark 3.21. In the event that there are multiple choices for the left and right regions, these different choices involve small and ultimately unimportant differences in $w_{\wedge \vee}(\Theta)$ and the accompanying action of the $\tilde{e}_i$. The right region is determined by the left region. From the definition of the regions it is clear that if $w_1 = \wedge, \vee$ or $\circ$ then shifting the residues of the left and right regions by $+1$ is not possible. A shift of the residues in the left and right regions by $+1$ to obtain another valid choice of left and right regions corresponds to removing a prefix of $\times$ from $w_{\wedge \vee}(\Theta)$ and adding a suffix of $\circ$. Thus we cyclically rotate the letters of $w_{\wedge \vee}(\Theta)$ one position to the left, with the twist that a letter $\times$ that falls off the left end becomes a $\circ$ on the right end. Since $\tilde{e}_i$ acts by zero on $\star \circ$ and $\times \star$, the action is the same on both words.

3.5. Highest weights for the $\widehat{\mathfrak{sl}}_d$ -action. For every charge $\sigma = (\sigma_1, \sigma_2) \in \mathbb{Z}^2$, the symbol $|\emptyset, \emptyset, \sigma\rangle$ labels a cuspidal unipotent character of the appropriate group (depending on the charge of the symbol) in characteristic 0, see §2.3. When reduced modulo ℓ , it remains irreducible [9].

We now explain the classification of modular cuspidals labeled by non-empty d -small symbols. The bipartition of such a symbol is a rectangle concentrated in one component, with a condition on the charged content of the removable box of the rectangle. In the case that Θ is a d -small symbol, it follows from [13, 22] and the rule for computing the Heisenberg or $\widehat{\mathfrak{sl}}_\infty$ crystal [17, 18] that S_Θ is cuspidal if and only if $[S_\Theta]$ is a highest weight vector for $\widehat{\mathfrak{sl}}_d$, that is, if and only if $E_i S_\Theta = 0$ for all $i \in \mathbb{Z}/d$. However, we will not need the condition that S_Θ is cuspidal, but only the condition of its being annihilated by every E_i . The description of such Θ follows immediately from Lemma 3.16.

Corollary 3.22. *Let Θ be a d -small symbol. We have $\tilde{e}_i \Theta = 0$ for all $i \in \mathbb{Z}/d$ if and only if in $w_{\wedge \vee}(\lambda)$, all $\times$'s precede all $\wedge$'s which precede all $\vee$'s which precede all $\circ$'s.*

Remark 3.23. Let Θ be a d -small symbol on which $\tilde{e}_i$ acts by 0 for all $i \in \mathbb{Z}/d$. Set $w = \#\{\wedge\text{'s in } w_{\wedge \vee}(\lambda)\}$ and $h = \#\{\vee\text{'s in } w_{\wedge \vee}(\lambda)\}$. Set $\lambda = (w^h)$. Then $\Theta = \Theta(\lambda, \sigma)$ where $\lambda = \lambda \cdot \emptyset$ if the left region is in row 1 and $\lambda = \emptyset \cdot \lambda$ if the left region is in row 2. The charge $\sigma = (\sigma_1, \sigma_2)$ is given by:

In the configuration $\times \circ$ for Ψ' (and hence Θ'), suppose we are in the case $n = 1$. We must have the configuration $\vee \wedge$ for Θ , as $\Theta = \tilde{f}_i \tilde{e}_i \Theta = \tilde{f}_i \Theta'$. That is, in $\mathbf{w}_{\vee \wedge}(\Theta)$ we have $w_j w_{j+1} = \vee \wedge$, where w_j corresponds to residue i as in (3.15). Hence in $\mathbf{c}_{\vee \wedge}(\Theta)$ there is a cup connecting w_j and w_{j+1} . However, in $\mathbf{w}_{\vee \wedge}(\Theta')$ we have $w'_i w'_{i+1} = \times \circ$ so there is no cup

connecting w'_j and w'_{j+1} in $c_{\wedge\vee}(\Theta')$. Since all other letters in $w_{\wedge\vee}(\Theta)$ and $w_{\wedge\vee}(\Theta')$ are the same, the cup diagram $c_{\wedge\vee}(\Theta)$ is thus obtained from the cup diagram $c_{\wedge\vee}(\Theta')$ by inserting a cup at positions i and $i+1$. On the other hand, by (3.19) there are two possibilities for Ψ in positions i and $i+1$, namely $\vee\wedge$ and $\wedge\vee$. It follows that $w_{\wedge\vee}(\Psi)$ is obtained from $w_{\wedge\vee}(\Theta)$ by reversing the orientation on a subset of cups of $c_{\wedge\vee}(\Theta)$.

For all the other configurations, Ψ is uniquely determined from Ψ' . We conclude that in all the cases, we have that $w_{\wedge\vee}(\Psi)$ is obtained from $w_{\wedge\vee}(\Theta)$ by reversing the orientation on a subset of cups of $c_{\wedge\vee}(\Theta)$. $\square$

In the setting of Theorem 4.1, let h be the number of $\wedge$'s and w be the number of $\vee$'s in the up-down diagrams of d -small symbols in the same block. The rule for computing the decomposition numbers presented in Theorem 4.1 is the same as Brundan-Stroppel's rule computing the multiplicities of Verma modules in projective modules in the highest weight category of finite-dimensional representations of the Khovanov arc algebra K_h^w [5]. C. Stroppel showed that the latter module category is equivalent to the category of perverse sheaves on the Grassmannian $\mathrm{Gr}(h, h+w)$ of h -planes in $\mathbb{C}^{h+w}$, which in turn is equivalent to the principal block of the parabolic category $\mathcal{O}^{\mathfrak{p}}$ for the parabolic $\mathfrak{p}$ with Levi $\mathfrak{gl}_h \times \mathfrak{gl}_w$ in $\mathfrak{gl}_{h+w}$ [26]. This identifies the decomposition numbers in question with the value of parabolic Kazhdan-Lusztig polynomials evaluated at 1. In [2], another way of obtaining this rule uses an oriented version of the Temperley-Lieb algebra.

We deduce that the decomposition numbers can be computed from Kazhdan-Lusztig polynomials.

Corollary 4.2. *The entries of the square submatrix of the decomposition matrix given by the formula of Theorem 4.1 are given by parabolic Kazhdan-Lusztig polynomials of type $(W, P) = (S_{h+w}, S_h \times S_w)$ evaluated at 1.*

ACKNOWLEDGMENTS

We would like to thank Gunter Malle for his careful reading of a previous version of this paper. E. Norton acknowledges financial support for this project from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 677147).

REFERENCES

- [1] C. BONNAFÉ, R. ROUQUIER, Catégories dérivées et variétés de Deligne-Lusztig, *Publ. Math. Inst. Hautes Études Sci.* **97** (2003), 1–59.
- [2] C. BOWMAN, M. DE VISSCHER, N. FARRELL, A. HAZI, E. NORTON, Oriented Temperley-Lieb algebras and combinatorial Kazhdan-Lusztig theory, arXiv:2212.09402, 2022.
- [3] O. BRUNAT, O. DUDAS, J. TAYLOR, Unitriangular shape of decomposition matrices of unipotent blocks. *Ann. of Math. (2)* **192** (2020), 583–663.
- [4] J. BRUNDAN, An orthogonal form for level two Hecke algebras with applications, *Algebraic groups and quantum groups*, 29–53, *Contemp. Math.*, 565, Amer. Math. Soc., Providence, RI, 2012.
- [5] J. BRUNDAN, C. STROPPEL, Highest weight categories arising from Khovanov's diagram algebra I: cellularity, *Mosc. Math. J.* **11** (2011), no. 4, 685–722, 821–822.
- [6] J. BRUNDAN, C. STROPPEL, Highest weight categories arising from Khovanov's diagram algebra III: category $\mathcal{O}$. *Represent. Theory* **15** (2011), 170–243.
- [7] J. CHUANG, R. ROUQUIER, Derived equivalences for symmetric groups and $\mathfrak{sl}_2$ -categorification, *Ann. of Math. (2)* **167** (2008), 245–298.
- [8] S. CIA-LUVECCE, Categorical action on unipotent representations of finite orthogonal groups, Ph.D. Thesis, in preparation.
- [9] O. DUDAS, G. MALLE, Modular irreducibility of cuspidal unipotent characters, *Invent. Math.* **211** (2018) 579–589.

- [10] O. DUDAS, G. MALLE, Decomposition matrices for groups of Lie type in non-defining characteristic, to appear in *Memoirs of the AMS*, 2023.
- [11] O. DUDAS, E. NORTON, Decomposition numbers for the principal Φ_{2n} -block of $\mathrm{Sp}_{4n}(q)$ and $\mathrm{SO}_{4n+1}(q)$, to appear in *Annales de l'Institut Fourier*, 2023.
- [12] O. DUDAS, M. VARAGNOLO, E. VASSEROT, Categorical actions on unipotent representations of finite unitary groups, *Publ. Math. Inst. Hautes Études Sci.* **129** (2019), 129–197.
- [13] O. DUDAS, M. VARAGNOLO, E. VASSEROT, Categorical actions on unipotent representations of finite classical groups. In *Categorification and Higher Representation Theory*, 41–104, Contemp. Math. **683**, Amer. Math. Soc., Providence, RI, 2017.
- [14] Z. FENG, B. SPÄTH, Unitriangular basic sets, Brauer characters and coprime actions, *Represent. Theory* **27** (2023), 115–148.
- [15] O. FODA, B. LECLERC, M. OKADO, J.-Y. THIBON, AND T. WELSH, Branching functions of $A_{n-1}^{(1)}$ and Jantzen-Seitz problem for Ariki-Koike algebras, *Adv. Math.* **141** (1999), 322–365.
- [16] P. FONG, B. SRINIVASAN, The blocks of finite classical groups, *J. reine angew. Math.* **396** (1989), 122–191.
- [17] T. GERBER, Heisenberg algebra, wedges and crystals, *J. Alg. Combin.* **49** (2019), 99–124.
- [18] T. GERBER, E. NORTON The $\mathfrak{sl}_\infty$ -crystal combinatorics of higher level Fock spaces, *J. Combin. Algebra* **2** (2018), 103–145.
- [19] F. HIMSTEDT, F. NOESKE, Decomposition numbers of $\mathrm{SO}_7(q)$ and $\mathrm{Sp}_6(q)$. *J. Algebra* **413** (2014), 15–40.
- [20] M. JIMBO, K. C. MISRA, T. MIWA, AND M. OKADO, Combinatorics of representations of $U_q(\widehat{\mathfrak{sl}(n)})$ at $q = 0$, *Comm. Math. Phys.* **136(3)** (1991), 543–566.
- [21] P. LI, P. SHAN, J. ZHANG, Categorical action for finite classical groups and its applications: characteristic 0, arXiv:2311.15229, 2023.
- [22] E. NORTON, On modular Harish-Chandra series of finite unitary groups, *Represent. Theory* **24** (2020), 483–524.
- [23] T. OKUYAMA, K. WAKI, Decomposition numbers of $\mathrm{Sp}(4, q)$, *J. Algebra* **199** (1998), 544–555.
- [24] R. ROUQUIER, Quiver Hecke algebras and 2-Lie algebras, *Algebra Colloq.* **19** (2012), 359–410.
- [25] B. SRINIVASAN, Isolated blocks in finite classical groups, II. *J. Algebra* **319** (2008), 824–832.
- [26] C. STROPPEL, Parabolic category $\mathcal{O}$, perverse sheaves on Grassmannians, Springer fibres and Khovanov homology, *Compos. Math.* **145** (2009), 954–992.
- [27] J.-L. WALDSPURGER, Une conjecture de Lusztig pour les groupes classiques, *Mém. SMF* **96** (2004).
- [28] X. YVONNE, Base canoniques d'espaces de Fock de niveau supérieur. PhD thesis, University of Caen, 2005.

(O.D.) CNRS & AIX-MARSEILLE UNIVERSITÉ, INSTITUT DE MATHÉMATIQUES DE MARSEILLE, CAMPUS DE LUMINY, 13009 MARSEILLE, FRANCE

Email address: `olivier.dudas@univ-amu.fr`

(E.N.) SCHOOL OF ENGINEERING, MATHEMATICS AND PHYSICS, UNIVERSITY OF KENT, CANTERBURY, CT2 7NZ, UNITED KINGDOM

Email address: `E.Norton@kent.ac.uk`